



HAL
open science

Exposure assessment in ergonomic epidemiology: is there something specific to the assessment of biomechanical exposures?

Annette Leclerc

► To cite this version:

Annette Leclerc. Exposure assessment in ergonomic epidemiology: is there something specific to the assessment of biomechanical exposures?. *Occup Environ Med*, 2005, 62 (3), pp.143-4. 10.1136/oem.2004.017889 . inserm-00141369

HAL Id: inserm-00141369

<https://inserm.hal.science/inserm-00141369>

Submitted on 12 Apr 2007

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Exposure assessment in ergonomic epidemiology

Is there something specific to the assessment of biomechanical exposures?

Annette Leclerc

U88 INSERM-IFR69, Saint Maurice, France

Annette.leclerc@st-maurice.inserm.fr

Key words : ergonomics, work, exposure, cancer

In this issue of the Journal the authors of two articles in “ergonomic epidemiology” stress several necessary qualities of exposure data: they must be accurate and precise¹; the method of exposure measurement must be reliable². All epidemiologists in occupational epidemiology would agree with that: absence of systematic or random error is important, stability of the measure if repeated under identical conditions is important too. Among the expected qualities of exposure data one could add “relevant”; we expect that the exposure data are consistent with what is known (or suspected) about the mechanisms underlying their effect on disease. This is less obvious than it seems; for example, in many situations one can wonder whether the relevant exposure is that of today, or that of last week, or that of twenty years ago, or the cumulative exposure over the twenty last years. Another quality (or limit) has to do with feasibility. If exposure assessment, at an individual level, is very expensive (in terms of money or time) alternative solutions have to be found if the study sample is large. All that is common to all the fields of occupational epidemiology. Then, is there something specific to the assessment of biomechanical exposures? Are the problems met when recording the time spent with the upper arm elevated above 90° very different from those met when assessing the level of past exposure to (for example) formaldehyde or electromagnetic fields, considered as potential carcinogens?

At least the history of research in these domains is different: a decade ago, in the field of occupational risk factors for musculoskeletal disorders, some questions were not discussed: there was no debate on how to use the expertise of ergonomists or specialists in biomechanics for epidemiologic studies; there was almost no research activity on other methodological aspects such as comparisons between questionnaires and direct observation. These questions were almost absent from the first PREMUS (Prevention of Work-related Musculoskeletal disorders) conference in 1992 in Stockholm. One exception was an abstract by Winkel and colleagues from the MUSIC project, who had evaluated a questionnaire estimating physical

workload³. The situation was very different in occupational cancer research. The concept of JEM (Job Exposure Matrix) covering an array of chemical substances, job titles and industries, was described as early as 1980⁴. Ten years ago cancer epidemiologists applied specific methods for sampling and analysis of exposure data⁵. In the presentation of the results of a European concerted action on the retrospective evaluation of occupational exposures in epidemiology, in 1993 Goldberg and Hemon gave more than 50 references discussing the measurement of occupational exposures in epidemiological studies⁶. A reason for that early development of methods for exposure assessment could be that, for cancer, difficulties dealing with (past) assessment of exposure were obvious, which motivated occupational hygienists, epidemiologists, and also biostatisticians to find adapted solutions. For biomechanical exposures, the (false) idea that it was enough to use the tools provided by ergonomists and specialists in biomechanics was probably widespread. However, in the last ten years, there has been a remarkable development of research on methodological aspects of assessment of exposure to biomechanical risk factors. Several subjects have been widely studied and discussed: the limits of using job titles to assess exposure⁷ and various other aspects of variability of exposure measures¹, the validity of questionnaires versus observation or measures², the use of aggregated measures⁸, the retrospective assessment of exposure⁹, and other such as bias due to the presence of pain.

The two articles in this issue of the Journal bring interesting results in this field of research: Heinrich and al.² compared, in a study population of 87 computer workers, questionnaire data about exposure to postural load and duration of computer use with an observation of the workstation design and posture by a trained observer. They conclude to a low agreement between the two approaches. However, they raise the question of whether the “gold standard” for postural load is the observation technique, since observation is based on short periods of time, and being observed might modify the posture.

Svensson and her colleagues¹ explored an attractive approach to the assessment of upper arm elevation above 90° in machinists, car mechanics and painters. Workers filled diaries with approximately ten preprint tasks, and exposure was evaluated using a TEM (Task Exposure Matrix). The diary worked well, and could be used to take into account the variability of tasks between subjects in the same job, which is a positive result. However, there was considerable within-task variability; in addition the exposure contrast between tasks was relatively small in these jobs. In this situation, spending resources on obtaining task information does not seem to be the optimal strategy.

These two studies suggest important directions for future research:

The first one is the development of simple techniques for assessing biomechanical exposures. There is still no consensus about the validity of questionnaire data to assess postural load ². Improving the questionnaires by adding pictures to the questions is probably a good suggestion. It is also necessary to think about the qualities expected from a tool (such as a questionnaire) in the context of epidemiological studies. The most important qualities are not necessarily the same in epidemiology, ergonomics, or biomechanics.

The second direction deals with indirect measures such as JEMs (Job Exposure Matrices) or TEM (Task Exposure Matrix). The main limitations in the development of matrices for biomechanical exposures is the variability of exposures within a job and within a task. However, this is probably the case more for some risk factors than for others. In large scale studies, JEMs might be useful to evaluate levels of exposure among all the subjects in a first step, possibly in combination with more precise assessments in specific subsamples. In addition, adapted statistical methods should be used and developed. Multilevel models should be more widely used ¹⁰. They are adapted to situations where some data were collected at a group level, and also for repeated measures ¹¹. Methods have also been proposed for combining expert rating and exposure measurement ¹².

For these three aspects, the research needs are partly specific of the field of “ergonomic epidemiology”, but also common with other fields in occupational epidemiology.

Acknowledgements

I thank Maria Melchior for her editorial help.

Correspondence to: Annette Leclerc, INSERM U88, HNSM, 14 rue du val d’Osne, 94410 Saint-Maurice France

Annette.leclerc@st-maurice.inserm.fr

References

1- Svendsen SW, Mathiassen SE, Bonde JP. Task-based exposure assessment in ergonomic epidemiology: a study of upper arm elevation in the jobs of machinists, car mechanics, and house painters. *Occupational and Environmental Medicine*.

2 - Heinrich J, Blatter BM, Bongers PM. A comparison of methods for the assessment of postural load and duration of computer use. *Occupational and Environmental Medicine*.

3 - Winkel J, Dallner M, Ericson M, Fransson C, Karlqvist L, Nygard C-H, Selin k, Wigaeus Hjelm E, Wiktorin C and the Stockholm-MUSIC I Study Group. Evaluation of a questionnaire for the estimation of physical load in epidemiologic studies. *International*

Scientific Conference on Prevention of Work-related Musculoskeletal disorders PREMUS. *Arbete och halsa* 1992;17:343-345.

4 - Hoar SK, Morrisson AS, Cole P, Silverman DT. An occupation and exposure linkage system for the study of occupational carcinogenesis. *J Occup Med* 1980; 22:722-726.

5 - Kromhout H, Symanski E, Rappaport SM. A comprehensive evaluation of within-and between-worker components of occupational exposure to chemical agents. *Ann Occup Hyg* 1993;37(3):253-270.

6 - Goldberg M, Hémon D. Occupational epidemiology and assessment of exposure. *International Journal of Epidemiology* 1993;22,6(Suppl.2):S5-S9.

7 - Gardner LI, Landsittel DP, Nelson NA, Pan CS. Misclassification of physical work exposures as a design issue for musculoskeletal intervention studies. *Scandinavian Journal of Work, Environment and Health* 2000; 26(5):406-413.

8 - Punnett L, van der Beek AJ. A Comparison of approaches to modeling the relationship between ergonomic exposures and upper extremity disorders. *American Journal of Industrial Medicine* 2000; 37(6):645-655.

9 - Koster M, Alfredsson L, Michelsen H, Vingard E, Kilbom A. Retrospective versus original information on physical and psychosocial exposure at work. *Scandinavian Journal of Work, Environment and Health* 1999; 25(5):410-414.

10 – Diez-Roux AV. A glossary for multilevel analysis. *Journal of Epidemiology and community health* 2002;56(8):588-594.

11 – Jansen JP, Burdorf A, Steyerberg E. A novel approach for evaluating level, frequency and duration of lumbar posture simultaneously during work. *Scandinavian Journal of Work, Environment and Health* 2001; 27(6):373-380.

12 – Wild P, Sauleau EA, Bourgkard E, Moulin JJ. Combining expert ratings and exposure measurements: a random effect paradigm. *Annals of Occupational Hygiene* 2002; 46(5):479-487.