



HAL
open science

Flat nearly Kähler manifolds

Vicente Cortés, Lars Schaefer

► **To cite this version:**

| Vicente Cortés, Lars Schaefer. Flat nearly Kähler manifolds. 2007. hal-00143588

HAL Id: hal-00143588

<https://hal.science/hal-00143588>

Preprint submitted on 26 Apr 2007

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Flat nearly Kähler manifolds

Vicente Cortés and Lars Schäfer

Department Mathematik, Universität Hamburg, Bundesstraße 55, D-20146 Hamburg, Germany
cortes@math.uni-hamburg.de, schaefer@math.uni-hamburg.de

February 23, 2007

Abstract

We classify flat strict nearly Kähler manifolds with (necessarily) indefinite metric. Any such manifold is locally the product of a flat pseudo-Kähler factor of maximal dimension and a strict flat nearly Kähler manifold of split signature $(2m, 2m)$ with $m \geq 3$. Moreover, the geometry of the second factor is encoded in a complex three-form $\zeta \in \Lambda^3(\mathbb{C}^m)^*$. The first nontrivial example occurs in dimension $4m = 12$.

Keywords: Nearly Kähler manifolds, flat almost pseudo-Hermitian manifolds, pseudo-Riemannian manifolds, almost complex structures

MSC 2000: 53C50, 53C15

Contents

1	Introduction	1
2	Basic facts about flat nearly pseudo-Kähler manifolds	2
3	Classification results for flat nearly pseudo-Kähler manifolds	5

1 Introduction

Nearly Kähler geometry originated in the study of weak holonomy groups by Gray [G4]. In fact, nearly Kähler manifolds correspond to weak holonomy $U(n)$ and were intensively studied by Gray [G1, G2, G3]. These manifolds appear moreover in a natural way in the Gray-Hervella classification [GH] as one class of the sixteen classes of almost Hermitian manifolds.

Recent interest in nearly Kähler manifolds came from the fact, that in dimension 6 these manifolds are related to the existence of Killing spinors and that they admit a Hermitian connection with totally skew-symmetric torsion. Such connections were studied by Friedrich and Ivanov [FI] and are of interest in string theory. The classification of complete simply connected strict nearly Kähler manifolds was reduced to dimension 6 by Nagy [N1, N2]. Butruille [B] has shown that all strict homogenous nearly Kähler manifolds are 3-symmetric. These works are mainly concerned with Riemannian manifolds. In this paper we are especially interested in pseudo-Riemannian metrics.

The interest in flat nearly pseudo-Kähler manifolds is motivated by our study [S] of tt^* -structures (topological-antitopological fusion structures) on the tangent-bundle. In fact, flat nearly pseudo-Kähler manifolds provide an interesting class of tt^* -structures on the tangent-bundle. A second interesting class of solutions is given by *special Kähler manifolds* [CS]. In other words, one can interpret tt^* -structures on the tangent-bundle as a common generalisation of these two geometries. This duality can also be seen in this work. We construct flat nearly pseudo-Kähler manifolds of split signature from a certain constant three-form, while in [BC] flat special Kähler manifolds were constructed from a symmetric three-tensor.

Let us describe the structure and results of the paper. In the first section we summarize some basic facts about flat nearly pseudo-Kähler manifolds (M, g, J) . The essential points are the existence of a canonical connection ∇ with skew-symmetric torsion T and the properties of the tensor $\eta = \frac{1}{2}JDJ = D - \nabla = -\frac{1}{2}T$, where D is the Levi-Civita connection.

The classification results are then given in the second section. The first one is Theorem 1, which encodes a flat nearly pseudo-Kähler structure in a constant three-form η subject to two constraints. An explicit formula for J in terms of η is given. Next we analyze the constraints on η . It turns out that the first is equivalent to require that η has isotropic support (cf. Proposition 3) and the second is equivalent to a type condition on η (cf. Theorem 2). We explicitly solve the two constraints on the real three-form η (in $4m$

variables) in terms of a freely specifiable complex three-form $\zeta \in \Lambda^3(\mathbb{C}^m)^*$. In particular, any such form $\zeta \neq 0$ defines a complete simply connected strict flat nearly pseudo-Kähler manifold, see Corollary 4.

Further we show that any strict flat nearly pseudo-Kähler manifold is locally the product of a flat pseudo-Kähler factor of maximal dimension and a strict flat nearly pseudo-Kähler manifold of dimension $4m \geq 12$ and split signature (Theorem 3). This implies, in particular, the non-existence of strict flat nearly Kähler manifolds with positive definite metric. The work finishes with the classification of complete simply connected flat nearly Kähler manifolds up to isomorphism in terms of $GL_m(\mathbb{C})$ -orbits on $\Lambda^3(\mathbb{C}^m)^*$, see Corollary 5.

We thank Paul-Andi Nagy for useful discussions.

2 Basic facts about flat nearly pseudo-Kähler manifolds

In this section we discuss some basic properties of nearly pseudo-Kähler manifolds. Since in this paper we are concerned with *indefinite* nearly Kähler manifolds with *flat* Levi-Civita connection, our discussion includes indefinite metrics but specializes the general statements and proofs whenever possible using the flatness assumption.

Definition 1 *An almost complex manifold (M, J) is called almost pseudo-Hermitian if it is endowed with a pseudo-Riemannian metric g which is pseudo-Hermitian, i.e. which satisfies $J^*g(\cdot, \cdot) = g(J\cdot, J\cdot) = g(\cdot, \cdot)$. The nondegenerate two-form $\omega := g(J\cdot, \cdot)$ is called the fundamental two-form.*

An almost pseudo-Hermitian manifold (M, g, J) is called nearly pseudo-Kähler manifold, if its Levi-Civita connection D satisfies the nearly Kähler condition

$$(D_X J)Y = -(D_Y J)X, \quad \forall X, Y \in \Gamma(TM). \quad (2.1)$$

A nearly pseudo-Kähler manifold is called strict, if $DJ \neq 0$.

Proposition 1 *(cf. Friedrich and Ivanov [FI]) Let (M, g, J) be a nearly pseudo-Kähler manifold. Then there exists a unique connection ∇ with totally skew-symmetric torsion T^∇ satisfying $\nabla g = 0$ and $\nabla J = 0$.*

More precisely, it holds

$$T^\nabla = -2\eta \text{ with } \eta = \frac{1}{2}JDJ \quad (2.2)$$

and $\{\eta_X, J\} = 0$, for all vector fields X .

To prove the proposition we give two lemmas of independent interest.

Lemma 1 *Let (V, g, J) be a pseudo-Hermitian vector space and S a $(1, 2)$ tensor, such that*

(i) S_X is skew-symmetric (with respect to g) for all $X \in V$,

(ii) $T(X, Y, Z) := g(S_X Y, Z) - g(S_Y X, Z)$, with $X, Y, Z \in V$, is totally skew-symmetric.

Then S , or more precisely $(X, Y, Z) \mapsto g(S_X Y, Z)$, is totally skew-symmetric, too.

Proof: It holds

$$\begin{aligned} g(S_X Y, Z) - g(S_Y X, Z) &\stackrel{(ii)}{=} -(g(S_Z Y, X) - g(S_Y Z, X)) \\ &\stackrel{(i)}{=} -g(S_Z Y, X) - g(S_Y X, Z), \end{aligned}$$

which implies $g(S_X Y, Z) = -g(S_Z Y, X)$. Together with property (i), this shows that S is totally skew-symmetric. \square

Lemma 2 *Let (V, g, J) be a pseudo-Hermitian vector space and S a $(1, 2)$ tensor satisfying:*

(i) S is totally skew-symmetric and

(ii) $[S_X, J] = 0$ for all $X \in V$.

Then S vanishes.

Proof: With arbitrary $X, Y, Z \in V$ we show

$$\begin{aligned} g(S_X Y, Z) &= g(JS_X Y, JZ) \stackrel{(ii)}{=} g(S_X JY, JZ) \\ &\stackrel{(i)}{=} -g(S_{JZ} JY, X) \stackrel{(ii)}{=} -g(JS_{JZ} Y, X) \\ &= g(S_{JZ} Y, JX) \stackrel{(i)}{=} -g(S_Y JZ, JX) \\ &\stackrel{(ii)}{=} -g(S_Y Z, X) \stackrel{(i)}{=} -g(S_X Y, Z). \end{aligned}$$

This shows $S = 0$. \square

Proof: (of Proposition 1) First we show the uniqueness:

Let ∇ and ∇' be two such connections and $S := \nabla - \nabla'$ their difference tensor.

Then from $\nabla J = \nabla' J = 0$ we obtain $[S_X, J] = 0$ and from $\nabla g = \nabla' g = 0$ we get the skew-symmetry of S_X (with respect to g) for all vector fields X .

In addition

$$(T^\nabla - T^{\nabla'})(X, Y, Z) = g(S_X Y - S_Y X, Z)$$

is the difference of two totally skew-symmetric tensors and hence totally skew-symmetric. Lemma 1 implies that the tensor S is totally skew-symmetric and Lemma 2 shows the uniqueness, i.e. the vanishing of S .

To prove the existence we define

$$\nabla := D - \eta \text{ with } \eta = \frac{1}{2} J D J.$$

The skew-symmetry of J yields that $D_X J$ is skew-symmetric (with respect to g). Further we have $\{J, D_X J\} = 0$, as follows from deriving $J^2 = -\mathbb{1}$, which shows that $\{\eta_X, J\} = 0$ and that $\eta_X = JD_X J$ is skew-symmetric for all vector fields X . From the skew-symmetry of η_X and $Dg = 0$ we obtain $\nabla g = 0$. Further we compute

$$\begin{aligned}\nabla_X J &= D_X J - \frac{1}{2}[JD_X J, J] = D_X J - \frac{1}{2}(JD_X J J - J^2 D_X J) \\ &= D_X J + J^2 D_X J = 0 \quad \forall X.\end{aligned}$$

This means $\nabla J = 0$. Finally we calculate the torsion

$$\begin{aligned}T^\nabla(X, Y) &= D_X Y - D_Y X - \eta_X Y + \eta_Y X - [X, Y] \\ &= T^D(X, Y) - (\eta_X Y - \eta_Y X) = -\eta_X Y + \eta_Y X \\ &= -2\eta_X Y,\end{aligned}$$

since DJ and consequently η is skew-symmetric by the nearly Kähler condition. Since η_X is skew-symmetric for all X and $\eta_X Y = -\eta_Y X$ for all X, Y , η is totally skew-symmetric and $T^\nabla = -2\eta$ is totally skew-symmetric, too. \square

Proposition 2 *Let (M, g, J) be a flat nearly pseudo-Kähler manifold. Then*

- 1) $\eta_X \circ \eta_Y = 0$ for all X, Y ,
- 2) $D\eta = \nabla\eta = 0$.

Lemma 3 *(cf. Gray [G2] Prop. 2.1) Let (M, g, J) be a nearly pseudo-Kähler manifold. Then for all $X, Y, Z, W \in TM$ it is*

$$g(R^D(X, Y)JZ, JW) - g(R^D(X, Y)Z, W) = g((D_X J)Y, (D_Z J)W). \quad (2.3)$$

Proof: (of Proposition 2)

1) From the last lemma we have

$$\begin{aligned}0 = g((D_X J)Y, (D_Z J)W) &= -g((D_Z J)(D_X J)Y, W) \\ &= -g(J(D_Z J)J(D_X J)Y, W) \\ &= -4g(\eta_Z \eta_X Y, W).\end{aligned}$$

This shows $\eta_X \eta_Y = 0$ for all X, Y and finishes the proof of part 1).

2) With two vector fields X, Y we calculate

$$\begin{aligned}(D_X \eta)_Y &= D_X(\eta_Y) - \eta_{D_X Y} \\ &\stackrel{D=\nabla+\eta}{=} \nabla_X(\eta_Y) + [\eta_X, \eta_Y] - \eta_{D_X Y} \\ &= (\nabla_X \eta)_Y + \eta_{[\nabla_X Y - D_X Y]} + [\eta_X, \eta_Y] \\ &= (\nabla_X \eta)_Y - \eta_{\eta_X Y} + [\eta_X, \eta_Y] \\ &\stackrel{n.K.}{=} (\nabla_X \eta)_Y + \eta \eta_X Y + [\eta_X, \eta_Y] \stackrel{1)}{=} (\nabla_X \eta)_Y.\end{aligned}$$

Using $\eta = \frac{1}{2}JDJ$ and $\nabla J = 0$ we obtain

$$\nabla\eta = \frac{1}{2}J\nabla(DJ).$$

Therefore it is sufficient to show $\nabla(DJ) = 0$. We calculate for D -parallel vector fields:

$$\begin{aligned} g(\nabla_X(DJ)_Y Z, W) &\stackrel{\nabla=D-\eta}{=} g(D_X(DJ)_Y Z, W) - g([\eta_X, D_Y J]Z, W) \\ &\stackrel{(*)}{=} g(D_X(DJ)_Y Z, W) \\ &\stackrel{DW=0}{=} Xg((DJ)_Y Z, W) \\ &= X[(D_Y\omega)(Z, W)] \\ &= D_X[(D_Y\omega)(Z, W)] \\ &\stackrel{DY=DZ=DW=0}{=} (D_{X,Y}^2\omega)(Z, W), \end{aligned}$$

where $\omega = g(J\cdot, \cdot)$. The second term in $(*)$ vanishes by part 1), since by $\{\eta_X, J\} = 0$ we get

$$J[\eta_X, D_Y J] = -\{\eta_X, JD_Y J\} = -2\{\eta_X, \eta_Y\} = 0.$$

The next lemma finishes the proof. □

Lemma 4 (compare Gray [G3] Prop. 2.3)

Let (M, g, J) be a flat nearly pseudo-Kähler manifold, then $D^2\omega = 0$.

3 Classification results for flat nearly pseudo-Kähler manifolds

We denote by $\mathbb{C}^{k,l}$ the complex vector space (\mathbb{C}^n, J_{can}) , $n = k + l$, endowed with the standard J_{can} -invariant pseudo-Euclidian scalar product g_{can} of signature $(2k, 2l)$.

Let (M, g, J) be a flat nearly pseudo-Kähler manifold. Then there exists for each point $p \in M$ an open set $U_p \subset M$ containing the point p , a connected open set U_0 of $\mathbb{C}^{k,l}$ containing the origin $0 \in \mathbb{C}^{k,l}$ and an isometry

$$\Phi : (U_p, g) \xrightarrow{\sim} (U_0, g_{can}),$$

such that at the point p we have:

$$\Phi_* J_p = J_{can} \Phi_*.$$

In other words, we can suppose, that locally M is a connected open subset of $\mathbb{C}^{k,l}$ containing the origin 0 and that $g = g_{can}$ and $J_0 = J_{can}$.

From Proposition 1 and 2 we obtain:

Corollary 1 Let $M \subset \mathbb{C}^{k,l}$ be an open neighborhood of the origin endowed with a nearly pseudo-Kähler structure (g, J) such that $g = g_{can}$ and $J_0 = J_{can}$. Then the $(1, 2)$ -tensor

$$\eta := \frac{1}{2}JDJ$$

defines a constant three-form on $M \subset \mathbb{C}^{k,l} = \mathbb{R}^{2k,2l}$ defined by

$$\eta(X, Y, Z) := g(\eta_X Y, Z)$$

satisfying

- (i) $\eta_X \eta_Y = 0, \quad \forall X, Y,$
- (ii) $\{\eta_X, J_{can}\} = 0, \quad \forall X.$

Conversely, we have the

Lemma 5 *Let η be a constant three-form on an open connected neighborhood $M \subset \mathbb{C}^{k,l}$ of 0 satisfying (i) and (ii) of Corollary 1. Then there exists a unique almost complex structure J on M such that*

- a) $J_0 = J_{can},$
- b) $\{\eta_X, J\} = 0, \quad \forall X,$
- c) $DJ = -2J\eta,$

where D stands for the Levi-Civita connection of the pseudo-Euclidian vector space $\mathbb{C}^{k,l}$. With $\nabla := D - \eta$ and assuming b), the last equation is equivalent to

$$c') \quad \nabla J = 0.$$

Proof: The equivalence of c) and c') follows from a straightforward calculation. First we show the uniqueness of J :

Given two almost complex structures J and J' satisfying a)-c) we find

- $J_0 = J'_0$ and
- $\nabla J = \nabla J' = 0,$

which shows $J = J'$.

To show the existence we define

$$J = \exp \left(2 \sum_{i=1}^{2n} x^i \eta_{\partial_i} \right) J_{can} \tag{3.1}$$

$$\stackrel{(i)}{=} \left(Id + 2 \sum_{i=1}^{2n} x^i \eta_{\partial_i} \right) J_{can}, \tag{3.2}$$

where x^i are linear coordinates of $\mathbb{C}^{k,l} = \mathbb{R}^{2k,2l} = \mathbb{R}^{2n}$ and $\partial_i = \frac{\partial}{\partial x^i}$.

Claim: J is an almost complex structure satisfying a)-c).

a) From $x^i(0) = 0$ we obtain $J_0 = J_{can}$.

- b) follows from equation (3.2) using (i) and (ii).
c) One computes

$$\begin{aligned} D_{\partial_j} J &= 2 \exp \left(2 \sum_{i=1}^{2n} x^i \eta_{\partial_i} \right) \eta_{\partial_j} J_{can} \\ &\stackrel{(ii)}{=} \underbrace{-2 \exp \left(2 \sum_{i=1}^{2n} x^i \eta_{\partial_i} \right) J_{can}}_J \eta_{\partial_j} = -2J \eta_{\partial_j}. \end{aligned}$$

It remains to prove $J^2 = -Id$.

$$\begin{aligned} J^2 &= \left(Id + 2 \sum_{i=1}^{2n} x^i \eta_{\partial_i} \right) J_{can} \left(Id + 2 \sum_{i=1}^{2n} x^i \eta_{\partial_i} \right) J_{can} \\ &\stackrel{(ii)}{=} - \left(Id + 2 \sum_{i=1}^{2n} x^i \eta_{\partial_i} \right) \left(Id - 2 \sum_{i=1}^{2n} x^i \eta_{\partial_i} \right) \\ &= - \left[Id - 4 \left(\sum_{i=1}^{2n} x^i \eta_{\partial_i} \right)^2 \right] \stackrel{(i)}{=} -Id. \end{aligned}$$

This finishes the proof. \square

Theorem 1 *Let η be a constant three-form on a connected open set $U \subset \mathbb{C}^{k,l}$ containing 0 which satisfies (i) and (ii) of Corollary 1. Then there exists a unique almost complex structure*

$$J = \exp \left(2 \sum_{i=1}^{2n} x^i \eta_{\partial_i} \right) J_{can} \quad (3.3)$$

on U such that

- a) $J_0 = J_{can}$,
b) $M(U, \eta) := (U, g = g_{can}, J)$ is a flat nearly pseudo-Kähler manifold.

Any flat nearly pseudo-Kähler manifold is locally isomorphic to a flat nearly pseudo-Kähler manifold of the form $M(U, \eta)$.

Proof: (M, g) is a flat pseudo-Riemannian manifold. Due to Lemma 5, J is an almost complex structure on M and $J_0 = J_{can}$.

In addition it holds

$$J = J_{can} + \left(2 \sum_{i=1}^{2n} x^i \eta_{\partial_i} \right) J_{can},$$

where $\{\eta_{\partial_i}, J\} = 0$ and η_{∂_i} is g -skew-symmetric. This implies that J is g -skew-symmetric. Finally from Lemma 5 c) and the skew-symmetry of η it follows the skew-symmetry of

DJ. Therefore (M, g, J) is nearly pseudo-Kähler.

The remaining statement follows from Corollary 1 and Lemma 5. \square

Now we discuss the general form of solutions of (i) and (ii) of Corollary 1. In the following we shall freely identify the real vector space $V := \mathbb{C}^{k,l} = \mathbb{R}^{2k,2l} = \mathbb{R}^{2n}$ with its dual V^* by means of the pseudo-Euclidian scalar product $g = g_{can}$.

Proposition 3 *A three-form $\eta \in \Lambda^3 V^* \cong \Lambda^3 V$ satisfies (i) of Corollary 1 if and only if there exists an isotropic subspace $L \subset V$ such that $\eta \in \Lambda^3 L \subset \Lambda^3 V$. If η satisfies (i) and (ii) of Corollary 1 then there exists a J_{can} -invariant isotropic subspace $L \subset V$ with $\eta \in \Lambda^3 L$.*

Remark: From the Proposition 3 we conclude that there are no strict flat nearly pseudo-Kähler manifolds of dimension less than 8. We shall see later that the dimension cannot be smaller than 12, see Corollary 2.

We define the **support** of $\eta \in \Lambda^3 V$ by

$$\Sigma_\eta := \text{span}\{\eta_X Y \mid X, Y \in V\} \subset V. \quad (3.4)$$

Proof: (of Proposition 3) The proposition follows from the next two lemmas by taking $L = \Sigma_\eta$. \square

Lemma 6 *Σ_η is isotropic if and only if η satisfies (i) of Corollary 1. If η satisfies (ii) of Corollary 1, then Σ_η is J_{can} -invariant.*

Proof: First the isotropy of Σ_η is equivalent to $g(\eta_X Y, \eta_Z W) = 0$ for all $X, Y, Z, W \in V$. From

$$g(\eta_X Y, Z) = \eta(X, Y, Z) = -\eta(X, Z, Y) = -g(\eta_X Z, Y) = -g(Y, \eta_X Z), \quad \forall X, Y, Z$$

we get

$$g(\eta_X Y, \eta_Z W) = -g(\eta_Z \eta_X Y, W). \quad (3.5)$$

Equation (3.5) shows $\eta_X \eta_Y = 0$ for all $X, Y \in V$ if and only if Σ_η is isotropic. The last assertion follows from

$$J\Sigma_\eta = \text{span}\{J\eta_X Y \mid X, Y \in V\} \stackrel{(ii)}{=} \text{span}\{-\eta_X JY \mid X, Y \in V\} = \Sigma_\eta.$$

\square

Lemma 7 *Let $\eta \in \Lambda^3 V$. Then $\eta \in \Lambda^3 \Sigma_\eta$.*

Proof: We take a complement $W \subset V$ of $L = \Sigma_\eta$. The decomposition

$$\Lambda^3 V = \bigoplus_{p+q=3} \Lambda^p L \wedge \Lambda^q W$$

induces a decomposition

$$\eta = \sum_{p+q=3} \eta^{p,q}.$$

Taking $X, Y \in L^\perp$ yields $L \ni \eta_X Y = \eta_X^{0,3} Y + \eta_X^{1,2} Y$. Now since $\eta_X^{0,3} Y \in W$ and $\eta_X^{1,2} Y \in L$, we get $\eta^{0,3} = 0$. Further the choice $X \in L^\perp$ and $Y \in W^\perp$ yields $\eta^{1,2} = 0$ and then the choice $X, Y \in W^\perp$ yields $\eta^{2,1} = 0$. This shows $\eta = \eta^{3,0}$. \square

Any three-form η on (V, J_{can}) decomposes with respect to the grading induced by the decomposition

$$V_{\mathbb{C}} = V^{1,0} \oplus V^{0,1}$$

into

$$\eta = \eta^+ + \eta^- \tag{3.6}$$

with

$$\eta^+ \in \Lambda^+ V := (\Lambda^{2,1} V + \Lambda^{1,2} V)^\rho$$

and

$$\eta^- \in \Lambda^- V := (\Lambda^{3,0} V + \Lambda^{0,3} V)^\rho,$$

where ρ is the canonical real structure on $V_{\mathbb{C}}$ with real-points V which extends to the exterior algebra.

Theorem 2 *A three-form $\eta \in \Lambda^3 V^* \cong \Lambda^3 V$ satisfies (i) and (ii) of Corollary 1 if and only if there exists an isotropic J_{can} -invariant subspace $L \subset V$ such that $\eta \in \Lambda^- L = (\Lambda^{3,0} L + \Lambda^{0,3} L)^\rho \subset \Lambda^3 L \subset \Lambda^3 V$. (The smallest such subspace L is Σ_η .)*

Proof: By Proposition 3, the conditions (i) and (ii) of Corollary 1 imply the existence of an isotropic J_{can} -invariant subspace $L \subset V$ such that $\eta \in \Lambda^3 L$. The next lemma shows that the condition (ii) is equivalent to $\eta \in \Lambda^- V$. Therefore $\eta \in \Lambda^3 L \cap \Lambda^- V = \Lambda^- L$. The converse statement follows from the same argument. \square

Lemma 8 *It is*

$$(\Lambda^{3,0} V + \Lambda^{0,3} V)^\rho = \{\eta \in \Lambda^3 V \mid \eta(\cdot, J\cdot, J\cdot) = -\eta(\cdot, \cdot, \cdot)\} = \{\eta \in \Lambda^3 V \mid \{\eta_X, J\} = 0, \forall X \in V\}.$$

Proof: We have the decomposition

$$\Lambda^2 V = (\Lambda^{1,1} V)^\rho \oplus (\Lambda^{2,0} V + \Lambda^{0,2} V)^\rho,$$

where

$$(\Lambda^{1,1} V)^\rho = \{\alpha \in \Lambda^2 V \mid \alpha(J\cdot, J\cdot) = \alpha\} \cong \{A \in \mathfrak{so}(V) \mid [A, J] = 0\}$$

and

$$(\Lambda^{2,0}V + \Lambda^{0,2}V)^\rho = \{\alpha \in \Lambda^2V | \alpha(J\cdot, J\cdot) = -\alpha\} \cong \{A \in \mathfrak{so}(V) | \{A, J\} = 0\}.$$

This induces the following direct decomposition:

$$V \otimes \Lambda^2V = V \otimes (\Lambda^{1,1}V)^\rho + V \otimes (\Lambda^{2,0}V + \Lambda^{0,2}V)^\rho.$$

We claim that

$$(V \otimes (\Lambda^{2,0}V + \Lambda^{0,2}V)^\rho) \cap \Lambda^3V = (\Lambda^{3,0}V + \Lambda^{0,3}V)^\rho.$$

The claim implies the lemma. To see the claim, let us first observe the following obvious inclusion:

$$(V \otimes (\Lambda^{2,0}V + \Lambda^{0,2}V)^\rho) \cap \Lambda^3V \supset (\Lambda^{3,0}V + \Lambda^{0,3}V)^\rho.$$

To show the equality we observe that an element of $V \otimes (\Lambda^{2,0}V + \Lambda^{0,2}V)^\rho$ is totally skew if and only if its four components in

$$V^{1,0} \otimes \Lambda^{2,0}V, \quad V^{1,0} \otimes \Lambda^{0,2}V, \quad V^{0,1} \otimes \Lambda^{2,0}V \quad \text{and} \quad V^{0,1} \otimes \Lambda^{0,2}V$$

are totally skew. To finish we notice that

$$(V^{1,0} \otimes \Lambda^{2,0}V + V^{0,1} \otimes \Lambda^{0,2}V) \cap \Lambda^3V = (\Lambda^{3,0}V + \Lambda^{0,3}V)^\rho$$

and

$$(V^{1,0} \otimes \Lambda^{0,2}V + V^{0,1} \otimes \Lambda^{2,0}V) \cap \Lambda^3V = 0.$$

□

Corollary 2 *There are no strict flat nearly pseudo-Kähler manifolds of dimension less than 12.*

Proof: By Theorem 1 and 2 any flat nearly pseudo-Kähler manifold M is locally of the form $M(U, \eta)$, where $\eta \in \Lambda^-L$ for an isotropic J_{can} -invariant subspace $L \subset V$ and $U \subset V$ is an open subset. $M(U, \eta)$ is strict if and only if $\eta \neq 0$, which is possible only for $\dim_{\mathbb{C}} L \geq 3$, i.e. for $\dim M \geq 12$. □

Theorem 3 *Any strict flat nearly pseudo-Kähler manifold is locally a pseudo-Riemannian product $M = M_0 \times M(U, \eta)$ of a flat pseudo-Kähler factor M_0 of maximal dimension and a strict flat nearly pseudo-Kähler manifold $M(U, \eta)$ of (real) signature $(2m, 2m)$, $4m = \dim M(U, \eta) \geq 12$. The J_{can} -invariant isotropic support Σ_η has complex dimension m .*

Corollary 3 *Let (M, g, J) be a flat nearly Kähler manifold with a (positive or negative) definite metric g then $\eta = 0$, $\nabla = D$ and $DJ = 0$, i.e. (M, g, J) is a Kähler manifold.*

Proof: (of Theorem 3) By Theorem 1 and 2, M is locally isomorphic to an open subset of a manifold of the form $M(V, \eta)$, where $\eta \in \Lambda^3 V$ has a J_{can} -invariant and isotropic support $L = \Sigma_\eta$. We choose a J_{can} -invariant isotropic subspace $L' \subset V$ such that $V' := L + L'$ is nondegenerate and $L \cap L' = 0$ and put $V_0 = (L + L')^\perp$. Then $\eta \in \Lambda^3 V' \subset \Lambda^3 V$ and $M(V, \eta) = M(V_0, 0) \times M(V', \eta)$. Notice that $M(V_0, 0)$ is simply the flat pseudo-Kähler manifold V_0 and that $M(V', \eta)$ is strict and of split signature $(2m, 2m)$, where $m = \dim_{\mathbb{C}} L \geq 3$. \square

For the rest of this paper we consider the case $V \cong \mathbb{C}^{m,m}$ and denote a maximal J_{can} -invariant isotropic subspace by L . We will say that a complex three-form $\zeta \in \Lambda^3(\mathbb{C}^m)^*$ has *maximal support* if $\text{span}\{\zeta(Z, W, \cdot) | Z, W \in \mathbb{C}^m\} = (\mathbb{C}^m)^*$.

Corollary 4 *Any non-zero complex three-form $\zeta \in \Lambda^{3,0}L \cong \Lambda^3(\mathbb{C}^m)^*$ defines a complete flat simply connected strict nearly pseudo-Kähler manifold $M(\eta) := M(V, \eta)$, $\eta = \zeta + \bar{\zeta} \in \Lambda^3 L \subset \Lambda^3 V$, of split signature. $M(\eta)$ has no pseudo-Kähler de Rham factor if and only if ζ has maximal support.*

Conversely, any complete flat simply connected nearly pseudo-Kähler manifold without pseudo-Kähler de Rham factor is of this form.

Proof: This follows from the previous results observing that the support of η is maximally isotropic if and only if ζ has maximal support. \square

Corollary 5 *The map $\zeta \mapsto M(\zeta + \bar{\zeta})$ induces a bijective correspondence between $GL_m(\mathbb{C})$ -orbits on the open subset $\Lambda_{reg}^3(\mathbb{C}^m)^* \subset \Lambda^3(\mathbb{C}^m)^*$ of three-forms ζ with maximal support and isomorphism classes of complete flat simply connected nearly pseudo-Kähler manifolds $M(\zeta + \bar{\zeta})$ of real dimension $4m \geq 12$ and without pseudo-Kähler de Rham factor.*

Examples:

1) The case $m \leq 5$.

For $m = 3, 4, 5$ the group $GL_m(\mathbb{C})$ acts transitively on $\Lambda_{reg}^3(\mathbb{C}^m)^* = \Lambda^3(\mathbb{C}^m)^* \setminus \{0\}$. Therefore there exists precisely one complete flat simply connected strict nearly pseudo-Kähler manifold of dimension 12, 16 and 20 respectively.

2) The case $m = 6$.

$GL_6(\mathbb{C})$ has precisely one open orbit in $\Lambda_{reg}^3(\mathbb{C}^6)^*$. This orbit consists of the stable three-forms $\Lambda_{stab}^3(\mathbb{C}^6)^*$ in the sense of Hitchin [H]. A three-form ζ on \mathbb{C}^6 is stable if and only if $\zeta = e_1^* \wedge e_2^* \wedge e_3^* + e_4^* \wedge e_5^* \wedge e_6^*$ for some basis (e_1, e_2, \dots, e_6) of \mathbb{C}^6 . $\Lambda^3(\mathbb{C}^6)^* \setminus \Lambda_{stab}^3(\mathbb{C}^6)^*$ is precisely the zero-set of the unique homogeneous quartic $SL_6(\mathbb{C})$ -invariant and we have the following strict inclusions:

$$\Lambda_{stab}^3(\mathbb{C}^6)^* \subset \Lambda_{reg}^3(\mathbb{C}^6)^* \subset \Lambda^3(\mathbb{C}^6)^* \setminus \{0\}.$$

An example of an instable regular form is

$$e_1^* \wedge e_2^* \wedge e_3^* + e_1^* \wedge e_4^* \wedge e_5^* + e_2^* \wedge e_4^* \wedge e_6^*.$$

References

- [B] J.-B. Butruille, *Classification des variétés approximativement kähleriennes homogènes*, Ann. Global Anal. Geom. **27** (2005) 201–225.
- [BC] O. Baues and V. Cortés, *Abelian simply transitive groups of symplectic type*, Annales de l’Institut Fourier **52**, no. 6 (2002), 1729–1751.
- [CS] V. Cortés and L. Schäfer, *Topological-antitopological fusion equations*, pluriharmonic maps and special Kähler manifolds, in ‘Complex, Contact and Symmetric Manifolds’, eds. O. Kowalski, E. Musso and D. Perrone, Progress in Mathematics 234, Birkhäuser 20, 59–74.
- [FI] Th. Friedrich and S. Ivanov, *Parallel spinors and connections with skew-symmetric torsion in string theory*, Asian J. Math. **6** (2002), no. 2, 303–335.
- [G1] A. Gray, *Almost complex submanifolds of the six sphere*, Proc. Amer. Math. Soc. **20** (1969) 277–279.
- [G2] A. Gray, *Nearly Kähler manifolds*, J. Diff. Geom. **4** (1970), 283–309.
- [G3] A. Gray, *Riemannian manifolds with geodesic symmetries of order 3*, J. Diff. Geom. **7** (1972), 343–369.
- [G4] A. Gray, *Weak holonomy groups*, Math. Z. **125** (1971), 290–300.
- [GH] A. Gray and L. M. Hervella *The sixteen classes of almost Hermitian manifolds and their linear invariants*, Ann. Mat. Pura Appl. **123** (1980), 35–58.
- [H] N. Hitchin, *The geometry of three-forms in six dimensions*, J. Differ. Geom. **55** (2000), no. 3, 547–576.
- [N1] P.A. Nagy, *On nearly-Kähler geometry*, Ann. Global Anal. Geom. **22** (2002), no. 2, 167–178.
- [N2] P.A. Nagy, *Nearly Kähler geometry and Riemannian foliations*, Asian J. Math. **6** (2002), no. 3, 481–504.
- [S] L. Schäfer, *tt^* -geometry on the tangent bundle of an almost complex manifold*, to appear in J. Geom. Phys.