

Historical reconstruction of the Atlantic Meridional Overturning Circulation from the ECMWF operational ocean reanalysis

Magdalena Balmaseda, Gregory Smith, Keith Haines, David P. Anderson, Tim Palmer, Arthur Vidard

▶ To cite this version:

Magdalena Balmaseda, Gregory Smith, Keith Haines, David P. Anderson, Tim Palmer, et al.. Historical reconstruction of the Atlantic Meridional Overturning Circulation from the ECMWF operational ocean reanalysis. Geophysical Research Letters, 2007, 34 (23), pp.L23615. 10.1029/2007GL031645 . inria-00181739

HAL Id: inria-00181739 https://inria.hal.science/inria-00181739

Submitted on 24 Oct 2007

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Historical reconstruction of the Atlantic Meridional Overturning Circulation from the ECMWF operational ocean reanalysis

Magdalena A. Balmaseda¹, Gregory C. Smith², Keith Haines², David Anderson¹, T.N.
Palmer¹ and Arthur Vidard³

¹European Centre for Medium Range Weather Forecasts, Shinfield Park, Reading RG2
9AX, UK.

² ESSC, Harry Pitt Bld, Earley Gate, Reading University, Reading RG6 6AL, UK.

8 ³ INRIA, LJK - B.P. 53 - 38041 - Grenoble cedex 9, France.

9 A reconstruction of the Atlantic Meridional Overturning Circulation (MOC) for the 10 period 1959-2006 has been derived from the ECMWF operational ocean reanalysis. The 11 reconstruction shows a wide range of time-variability, including a downward trend. At 12 26N, both the MOC intensity and changes in its vertical structure are in good agreement 13 with previous estimates based on trans-Atlantic surveys. At 50N, the MOC and strength 14 of the subpolar gyre are correlated at interannual time scales, but show opposite secular 15 trends. Heat transport variability is highly correlated with the MOC but shows a smaller 16 trend due to the warming of the upper ocean, which partially compensates for the 17 weakening of the circulation. Results from sensitivity experiments show that although 18 the time-varying upper boundary forcing provides useful MOC information, the 19 sequential assimilation of ocean data further improves the MOC estimation by 20 increasing both the mean and the time variability.

21 **1. Introduction**

The Atlantic meridional overturning circulation (MOC) is composed of a warm near-surface branch flowing northward as part of the Gulf Stream and a return flow of cold waters at depth. It plays a major role in the heat transport of the ocean, in turn 25 affecting the climate of Europe and North America (e.g. Cubash et al 2001), and its 26 variability plays an important role in future climate change scenarios. However, reliable 27 estimates and understanding of the variability remain elusive. Bryden et. al. (2005), 28 (hereafter BLC05) using density measurements from five transatlantic research cruises 29 at approximately 26°N between 1957-2004, found a 30% decrease in MOC intensity, with a notable reduction in the southward flow of the lower North Atlantic Deep Water 30 (NADW) coming from high latitudes, although these conclusions were based on very 31 32 limited temporal sampling. In contrast, estimates relying on ocean model simulations 33 have produced an intensification of the MOC (e.g. Böning et. al. 2006), which could be 34 attributed to the prevailing positive phase of the North Atlantic Oscillation (NAO) since 35 1980's (Eden and Willebrand (2001).

36 The contradictory results between observational and model estimates illustrate the 37 underlying uncertainties in the different methodologies: the observational BLC05 data 38 clearly have insufficient temporal sampling to estimate trends, and the model results can 39 be affected by errors in the forcing fluxes and model formulation. A hybrid approach is 40 the synthesis of ocean model and observations using data assimilation techniques, to 41 produce an ocean analysis (for a summary of ongoing activities see 42 http://www.clivar.org/organization/gsop/synthesis/synthesis.php). In theory, the error in 43 the MOC from an ocean analysis should be smaller than the errors in ocean model or observational estimates alone. In practice, some new uncertainties may be introduced 44 45 from different assimilation techniques or observations of varying density/accuracy.

46 Ocean analyses such as ECCO have previously been used to derive 47 reconstructions of the MOC (Wunsch and Heimbach (2006) and Köhl and Stammer 48 (2007)). These ECCO analyses are based on long-window adjoint methods, and 49 typically rely on the correction of the ocean initial conditions and surface forcing to get 50 close to the observed ocean data. Here we present a 48 year historical reconstruction of 51 the MOC (for the period 1959-2006) from the ECMWF operational ocean reanalysis 52 System 3 (ORAS3 in what follows), which uses a sequential assimilation method to 53 directly correct the density field, which is critical to circulation indices such as the 54 MOC.

The paper is organized as follows: we describe the ocean analysis system, and the sensitivity experiments in secn 2, the reconstruction of the MOC, including the meridional and vertical structure in secn 3 and the implications for the meridional heat transports in secn 4. Results from sensitivity experiments are presented in secn 5 and conclusions in secn 6.

60 **2** The data assimilation system

61 The analysis of the ocean state is obtained by integrating a global ocean model with 62 atmospheric surface fluxes acting as time-dependent upper boundary conditions. The ocean model is HOPE (Wolff et al. 1997, Balmaseda 2004), 1° x 1° resolution, with a 63 64 tropical enhancement to $1/3^{\circ}$, and 29 vertical levels, with partial step topography and 65 explicit free surface. From 1959 to August 2002, the forcing fluxes are from the 66 ERA40 atmospheric reanalysis with corrected freshwater fluxes, and from the 67 operational atmospheric analysis thereafter (ERA40/OPS in what follows). The ocean 68 observations are assimilated sequentially via an optimal interpolation (OI) method, 69 which imposes dynamical and physical constraints. The analysis cycle is repeated every 10 days. A detailed description of the system is given in Balmaseda et al 2007a. 70 71 The subsurface observations consist of vertical profiles of temperature and salinity from 72 Bathythermographs (MBT, XBT), Conductivity Temperature Depth (CTD) sensor measurements 73 from scientific cruises, TAO/TRITON and PIRATA moorings, and more recently Argo floats.

74 Historical salinity data are scarce, and it is only with the advent of Argo floats that a near-global

coverage of salinity observations is available (from 2000 onwards). For the period 1959-2004 the

subsurface data are from the comprehensive quality-controlled data set ENACT-ENSEMBLES
(Ingleby and Huddleston, 2006), which contains 5.1 million temperature and 1.4 million salinity
profiles. From 2005 onwards, the subsurface data are from the ECMWF operational archive, and
are subject to a different automatic quality control procedure. For the later period, a typical 10-day
assimilation window contains 2500 profiles of temperature and 1100 profiles of salinity. Maps of

81 sea surface temperature (Reynolds et al 2002) are also assimilated and, from 1993 onwards,

82 satellite-derived sea level anomaly maps (Le Traon et al 1998) are used. Supplementary figure 1

83 shows a timeseries of the number of temperature profile observations used in a 10-day

84 assimilation cycle in the North Atlantic (20N-50N) as a function of depth. The observation

85 coverage maps for the individual assimilation cycles can be seen at

86 http://www.ecmwf.int/products/forecasts/d/charts/ocean/reanalysis/obsmap/.

87 The ORAS3 is part of the operational monthly and seasonal forecasting system, where a 88 reliable reconstruction of the time variability of the ocean is required to improve the 89 skill of the system. Special attention has been paid to the tuning of the error 90 covariances, where the correlation scales and the diagonal elements have been chosen 91 so as to improve both the mean state and the interannual variability. In addition, to 92 reduce spurious time-variability resulting from the changing nature of the observing 93 system, ORAS3 uses low frequency bias-corrections to both the pressure gradient and 94 the temperature and salinity fields (Balmaseda et al 2007b). Only a weak relaxation to 95 the full temperature and salinity climatology is used (10-year time scale), which does 96 not significantly damp the interannual variability.

97 To assess the impact of assimilating data, a control experiment (ORA-nobs) is 98 conducted by integrating the ocean model with the ERA40/OPS fluxes but without 99 assimilating profiles or altimeter data. Everything else (spin up, relaxation to SST and 100 3D climatology) is the same as in ORAS3. To assess the impact of the forcing fluxes 101 and spin up an additional experiment is conducted, identical to ORA-nobs but using a climatology of the daily fluxes as forcing. The effect of initial conditions on the MOC
in ORAS3 at the beginning of the record is explored by a set of 10-year assimilation
experiments starting from perturbed initial conditions in 1956.

3. The historical reconstruction of the MOC

106 Balmaseda et al 2007a show that the ORAS3 reanalysis is consistent with the 107 observed profile data, and quantitatively reproduces the expected mean circulations and 108 time variations in temperature, salinity and surface currents. The Atlantic meridional 109 heat transports in ORAS3 are in good agreement with WOCE estimates (Ganachaud 110 and Wunsch 2003, supplementary table1). Figure 1 shows the Atlantic MOC at 26°N 111 for ORAS3, calculated by integrating the zonal-mean velocity from the surface to 112 1200m (chosen as the depth of maximum overturning in the model). The agreement 113 between ORAS3 and the BLC05 values is remarkably good for 1981, 1992 and 1998, but differs in 2004, where the BLC05 value is substantially lower. However, more 114 115 recent estimates from the RAPID array (Cunningham et al 2007) yield an average MOC 116 value of 18.7Sv for 2004, which is in good agreement with ORAS3. Although the 117 agreement is very encouraging, one should remember that there are only four points and 118 there are likely substantial uncertainties in both the section/array estimates and model 119 values.

Figure 1 also indicates the large seasonal (1.8 Sv) and interannual (1.9Sv) variability of the MOC. The seasonal variability of the MOC at 26°N can be attributed mainly to the seasonality of the Ekman transport, which has a standard deviation of 1.9Sv. Ekman transport makes up about 25% of the time-mean and interannual transports (4.9 Sv and 0.56 Sv respectively). The MOC at 26°N in ORAS3 shows a small decrease over the 48-year period which amounts to -0.07 ± 0.01 Sv/yr, equivalent to a reduction of 4% per decade, although from figure 1 it is clear that this trend is not constant (e.g. the trend 127 after the mid-1970's is only 2% per decade). The weakening MOC is associated with 128 changes in vertical structure of the circulation (figure 2a). Consistent with BLC05, there 129 is a reduction in the southward transport of the lower North Atlantic Deep Water 130 (NADW) in ORAS3, associated with a shallower and weaker recirculation cell. This is 131 an important difference from the 11-year ECCO-GODAE reanalysis (Wunsch and 132 Heimbach, 2006), which also shows a slow-down of the MOC, but with an intensification of the southward NADW flow. (The differences between ORA-S3 and 133 134 ECCO-GODAE are likely to stem from the different assimilation methods). IThe 135 coherent changes in the vertical structure of the circulation occur at low frequency, and 136 do not seem to be affected by the seasonal variability of the Ekman transport. This 137 implies that vertical structure comparisons with BLC05 are more robust, since they are 138 not contaminated by high frequency variability.

139 Figure 2a also shows a reduction of the northward transport within the 140 thermocline which, according to Cunningham and Alderson 2007, results from an 141 intensified southward geostrophic transport caused by the increased east-west 142 thermocline slope, and is consistent with the changes in the vertical density structure in 143 ORAS3. There is a general warming and salinification in the upper subtropical ocean, 144 indicative of thermocline deepening, which is more pronounced in the western part of 145 the basin. ORAS3 also reproduces an increase in temperature and salinity (0.42 K and 146 0.07 psu respectively at 450m) in the Eastern Atlantic between 1992 and 2002, noted by 147 Vargas-Yáñez et al (2004) from a cruise survey at 24°N.

148

The time variability of the MOC in ORAS3 changes considerably as a function of latitude (figure 2b). Within the subtropical gyre (south of 30N) the interannual variability is dominant, while in subpolar latitudes decadal variability is stronger. A 152 reduction in the MOC (2-4% per decade, supplementary table 2 and supplementary 153 figure 4) is apparent in most of the North Atlantic domain, and is particularly 154 pronounced after 1995, with a visible reduction in the meridional extension of the 155 MOC. Häkkinen and Rhines (2004) attribute this reduction of the MOC after 1995 to 156 the weakening of the subpolar gyre (SPG), characterized by a decrease in sea level 157 gradients from satellite altimetry. The intensity of the SPG in ORAS3 (measured by the 158 sea level differences between 40N and 60N) is correlated with the MOC at 50N at 159 interannual time scales (r=0.8), in agreement with Böning et al (2006), with the MOC in 160 ORAS3 lagging the subpolar gyre by 18 months (figure 3). But contrary to other model 161 studies, the secular trends of the MOC and the SPG found here are of opposite sign. 162 There are several possible reasons for this: i) the atmospheric forcing fluxes (ORAS3 163 uses ERA40/OPS instead of NCEP) ii) the surface heat flux closure (in ORAS3 there is 164 strong relaxation to time-varying SST, which may compensate for errors in the heat 165 fluxes, thus contributing to a better simulation of the upper ocean warming); and iii) the 166 representation of the overflows. For instance, Böning et al (2006) impose climatological 167 boundary conditions at 70N, while ORAS3 overflow properties may vary in time and be 168 affected by the assimilation of ocean observations.

169 **4 Heat transports**

170 It has been suggested that any slowdown of the MOC could have significant 171 implications for the climate of Europe (Vellinga and Wood 2002) due to a resulting 172 reduction in heat transport in the northward flowing upper limb. In ORAS3, the 173 interannual variability in the heat transport at 26°N follows closely the MOC variability 174 (correlated at r = 0.9), and also shows a small downward trend of -0.0029 ± 0.0007 175 PW/yr, equivalent to a reduction of 2.7% per decade. This fractional trend in heat 176 transport is weaker than for the MOC over the whole North Atlantic domain 177 (supplementary table 2 and supplementary figure 4). This is a consequence of the

increased vertical temperature gradient resulting from a general upper ocean warming (Fig. 4). At 26°N there is a modest warming trend in the upper 300 m of 0.05 ± 0.01 K/decade, while at 40°N this increases to 0.26 ± 0.04 K/decade. The increased upper ocean temperatures in the poleward moving branch of the MOC intensify the poleward heat transport, partially cancelling the effect of the weakening MOC, in agreement with the simulations of Drijfhout and Hazeleger (2006).

184

5. Sensitivity experiments

185 The time variability of the MOC reconstruction could be affected by variations in 186 the observing system and spin-up effects. Here we use sensitivity experiments to assess 187 the robustness of the ORAS3 results. The agreement with the observed temperature and 188 salinity profiles is better for ORAS3 than for ORA-nobs (about 30% in the North 189 Atlantic, supplementary figure 2). The improved representation of the density field 190 affects both the mean overturning strength and the amplitude of the variability, 191 improving dramatically the agreement with the BLC05 values relative to the ORA-nobs 192 (fig 1), as well as the heat transports, which are underestimated in ORA-nobs 193 (supplementary table 1). The coherence between ORAS3 and ORA-nobs is also 194 apparent at 50N, where the MOC and the SPG intensity in ORA-nobs show positively 195 correlated interannual variability and opposite secular trends (not shown).

The large degree of coherence between the time evolution of the MOC in ORAS3 and ORA-nobs is indicative of the atmospherically-driven component. ORA-nobs simulates the same large MOC values during the 60's, increased variability during the 80's, and the quasi-biennial signals after 2000. ORA-nobs also shows a decline in MOC intensity, although of a smaller magnitude than ORAS3 (2% per decade), suggesting that some trend is directly linked to changes in the atmospheric forcing. In contrast, the experiment with climatological forcing (supplementary figure 3), shows no significant trend after an initial adjustment, supporting the attribution of part of the MOC decline tothe time-varying upper boundary forcing.

205 Direct comparison with the BLC05 value for 1957, outside the ORAS3 record, is 206 not possible. Additional experiments, similar to ORAS3 but starting from 1956 were 207 conducted. Prior to 1958 there is no ERA40 forcing, and so climatological forcing was 208 used. Different ocean initial conditions were used: a) ORAS3 spin up, b) ORAS3 (1 Jan 209 1962) and c) ORAS3 (1 Jan 1965). None of these experiments reproduced the BCL05 210 MOC value for 1957, probably because of the scarcity of information (both 211 observational and forcing). Results also show that the MOC converges to the ORAS3 212 value by 1962, suggesting that the spin up is not a determining factor in ORAS3 after 213 1962.

Additional experiments show that the estimated MOC trend and the specific agreement with the BLC05 values remain unchanged even if all the specific section data used by BLC05 are withdrawn from the ORAS3 reanalysis. This illustrates the ability of data assimilation systems to propagate observational information either directly, via the prescribed error correlation functions, or via physical processes represented by the ocean model.

6. Summary

These results show that assimilating data in ORAS3 improves the representation of the Atlantic MOC against section-based estimates, and permits a 48-year reconstruction, for the period 1959-2006, which exhibits a wide range of time variability (seasonal, interannual and secular trends). ORAS3 results suggest a slowdown of the MOC (2-4% per decade) for most of the North Atlantic basin, although the trends are not constant, being much smaller in the second half of the record. The MOC variability in the subtropical gyre is highly correlated with the heat transport variability, but the trends in heat transport are weaker, due to slow changes in the vertical thermal structure, with the pronounced upper ocean warming partially compensating for the reduction in the MOC.

231 Sensitivity experiments suggest that either ERA40 atmospheric forcing and/or 232 the strong constraint on the SST can explain some of the reduction of the MOC, but that 233 the trend is enhanced by the assimilation of in situ ocean data. The results presented 234 here support the paradigm of the North Atlantic Oscillation (NAO) as providing the 235 primary forcing for the MOC at 50N on interannual timescales (Eden and Willebrand, 236 2001), with positive NAO conditions leading to the intensification of the MOC. 237 However, the reduction of the MOC at 50N in ORAS3 under prevailing positive NAO 238 conditions during recent decades, accompanied by the decline in the southward 239 transport of the lower NADW, suggest that other factors are more important for the 240 MOC on longer timescales.

241 These results illustrate the potential of ocean reanalysis for the study of ocean 242 climate. In the latest IPCC Assessment Report, it was stated that due to the conflict 243 between model and observational studies, "no coherent evidence" of a trend in the 244 MOC over the last 50 years existed, and hence no baseline comparison was possible for 245 climate model simulations. It is shown here that data assimilation can reconcile model 246 and observations, giving a self consistent MOC timeseries which agrees with traditional 247 section-based estimates where available. Sensitivity experiments can test robustness and 248 further reanalyses based on other models and methods are underway (within CLIVAR-249 GSOP panel) that will further reduce the uncertainty in these estimations of the MOC. 250 Ocean reanalysis should be able to provide a past baseline for MOC estimates, and more 251 generally, a valuable gauge on the quality of climate models used for future climate

- 252 projections. The uncertainties in the ocean reanalysis will be reduced, as the quality of
- the assimilation methods, ocean model and atmospheric reanalyses improves.

254

- 255 References.
- 256 Balmaseda, M.A., 2004: Ocean data assimilation for seasonal forecasts. ECMWF
- 257 Seminar Proceedings. Seminar on Recent developments in data assimilation for
- atmosphere and ocean, 8-12 September 2003, 301-326.
- 259 Balmaseda, M., Vidard, A. & Anderson, D. 2007a. The ECMWF System 3 ocean
- analysis system. ECMWF Technical Memorandom, 508.
- 261 (http://www.ecmwf.int/publications/library/ecpublications/_pdf/tm/501-600/tm508.pdf).
- 262 Balmaseda, M.A., D. Dee, A. Vidard and D.L.T. Anderson, 2007b. A Multivariate Treat
- 263 ment of Bias for Sequential Data Assimilation: Application to the Tropical Oceans. Q.
- 264 J. R. Meteorol. Soc., 133, 167-179.
- 265 Bindoff, N.L.& Willebrand, J., 2007: Observations: Climate Change and Sea Level,
- 266 IPCC WG1 AR4 Final Report.
- 267 http://ipccwg1.ucar.edu/wg1/Report/AR4WG1_Ch05.pdf.
- 268 Boning, C. W., M. Scheinert, J. Dengg, A. Biastoch, and A. Funk, 2006: Decadal
- 269 variability of subpolar gyre transport and its reverberation in the North Atlantic
- 270 overturning. Geophys. Res. Let., Vol. 33, L21S01.
- 271 Bryden, H. L., Longworth, H. L. & Cunningham, S. A., 2005. Slowing of the Atlantic
- 272 Meridional Overturning Circulation at 25°N. Nature 438, 655-657..
- 273 Cubasch, U. et al. 2001. Climate Change 2001: The Scientific Basis (ed. Houghton, J.
- 274 *T.) Ch* 9, 525-585 (Cambridge Univ. Press, Cambridge, UK, 2001).

- 275 Cunninham S.A. and S. Alderson. 2007. Transatlantic temperature and salinity changes
- 276 at 24.5N from 1957 to 2004. *Geosphys. Res. Lett.* 34, L14505.
- 277 Cunningham, S., Kanzow T., Rayner D., Baringer M.O., Johns W.E., Marotzke J.,
- 278 Longworth H.R., Grant E.M., Hirschi J.J-M., Beal L.M, Meinen C.S. and Bryden H.L.,
- 279 2007. Temporal Variability of the Atlantic Meridional Overturning Circulation at
- 280 26.5°N. Science, **317**, 938-941.
- 281 Drijfhout, S. S. and W. Hazeleger, 2006: Changes in MOC and gyre-induced Atlantic
- 282 Ocean heat transport. *Geophys. Res. Let.*, **33**, L07707.
- Eden C., and J. Willebrand, 2001: Mechanism of interannual to decadal variability of
- the North Atlantic circulation. J. Climate, 14, 2266-2280.
- 285 Ganachaud, A. & Wunsch, C., 2003. Large scale ocean heat and freshwater transports
- during the World Ocean Circulation Experiment. J. Clim. 16, 696-705.
- 287 Häkkinen, S. & Rhines, P. B., 2004. Decline of subpolar North Atlantic circulation
- during the 1990s. *Science*, **304**, 555-559.
- 289 Ingleby, B. & Huddelston, M., 2006. Quality control of ocean temperature and salinity
- 290 profiles historical and real-time data. J. Mar. Sys., 65, 158-175.
- 291 Köhl A. and D. Stammer, 2007. Variability of the Meridional Overturning in the North
- Atlantic from the 50 years GECCO State Estimation. The ECCO report series, 43.
- 293 Available at www.ecco-group.org.
- Le Traon, P.-Y., F. Nadal, and N. Ducet., 1998. An improved mapping method of
- 295 multisatellite altimeter data. J. Atmos. Oceanic Technol., 15, 522-534.
- 296 Reynolds R., N Rayner, T Smith, D Stokes, W Wang., 2002. An improved in situ and
- satellite SST analysis for climate. J. Clim, 15, 1609-1625.

- 298 Vargas-Yáñez, M., Parilla, G., Lavin, A., Vélez-Belchi, P. & González-Pola, C., 2004.
- 299 Temperature and salinity increase in the eastern North Atlantic along the 24.5°N in the
- 300 last ten years. *Geophys. Res. Lett.*, **31**, L06210, doi:10.1029/2003GL019308.
- 301 Vellinga, M. & Wood, 2002. R. A. Global climatic impacts of a collapse of the Atlantic
- 302 thermohaline circulation. *Clim. Change* **54**, 251-267.
- 303 Wolff, J., E. Maier-Reimer and S. Legutke, 1997. The Hamburg Ocean Primitive
- 304 Equation Model. *Deutsches Klimarechenzentrum*, Hamburg, Technical Report No. 13.
- 305 Wunsch, C. & Heimbach, P., 2006. Estimated Decadal Changes in the North Atlantic
- 306 Meridional Overturning Circulation and Heat Flux 1993-2004. J. Phys. Ocean., 36,

307 2012-2024.

308

309

310 Figures



Figure 1 | Meridional overturning circulation (MOC) variability at 26°N. The time evolution of the MOC for both ORAS3 (black) and ORA-nobs (blue) is shown using monthly values (thin lines) and annual means (thick lines). Over-plotted are the annual-mean MOC values from BLC05 (red circles) and Cunningham et al 2007 (green circle).



Figure2 | Vertical (a) and meridional (b) structure Atlantic MOC as a function of time. In (a) the vertical structure of the MOC is represented by the zonally integrated meridional velocity at 26° N, and units are 10^{3} m²/s. Both the poleward transport within the upper 1000 m and the equatorward transports below 2000 m are decreasing with time. In (b), the MOC is calculated as the integrated meridional velocity above a reference depth of 1200m in units of Sv.



Figure 3 | Normalized timeseries of the subpolar gyre index (black) and MOC at 50N (red) from ORAS3. Overplotted are the linear trend estimates. The subpolar gyre index is computed as the sea level differences at 40N and 60N. The decrease in subpolar gyre intensity during the 90's is consistent with Hakkinen and Rhines 2004. The subpolar gyre variability leads the MOC variability at interannual time scales, but the trends are opposite.



Figure 4 | Trends in the vertical temperature structure from the 48-year ORAS3 analysis, at 26°N (red) and 40°N (black). Units are K/yr. Shaded are the trend values within the 95% C.I.