



HAL
open science

Unsuspected consequences of the adolescent overweight epidemic.

Sylvie Mesrine, Agnès Fournier, Marie-Christine Boutron-Ruault, Françoise Clavel-Chapelon

► **To cite this version:**

Sylvie Mesrine, Agnès Fournier, Marie-Christine Boutron-Ruault, Françoise Clavel-Chapelon. Unsuspected consequences of the adolescent overweight epidemic.. *Pediatrics*, 2007, 120 (4), 924-5; author reply 925-6. 10.1542/peds.2007-1698 . inserm-00193057

HAL Id: inserm-00193057

<https://inserm.hal.science/inserm-00193057>

Submitted on 30 Sep 2008

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Unsuspected Consequences of the Adolescent Overweight Epidemic

Sylvie Mesrine, MD, Agnès Fournier, MSc, Marie-Christine Boutron-Ruault, MD, PhD, Françoise Clavel-Chapelon, PhD

Institut National de la Santé et de la Recherche Médicale ERI 20 Gustave Roussy Institute F-94805 Villejuif, France

We read with great interest the article entitled “Weight Status in Young Girls and the Onset of Puberty,” which was published in the March issue of *Pediatrics*.¹ The authors showed, from longitudinal follow-up of 354 girls, that a higher BMI z score at 36 months of age and a higher rate of change of BMI between 36 months of age and first grade were associated with earlier puberty. They concluded that increasing rates of obesity in the United States may result in an earlier average age of onset of puberty and in a number of adverse outcomes, in particular reproductive cancers.

That early menarche is predictive of elevated breast carcinoma risk is well known.² More unexpected and, indeed, paradoxical are the results of a number of studies, including our prospective cohort study (the E3N cohort) of 98 995 French women (mainly teachers, aged 40–65 years at recruitment in 1990–1991) who were followed, on average, for 12 years.³ We showed that, indeed, women who had breast cancer had significantly earlier menarche than controls; however, when adjusting for age at menarche, significant trends of decreasing risk with increasing body silhouettes at age 8 and at menarche were observed (*P* trend: <.001 to <.0005, respectively, with relative risks of 0.86 [95% confidence interval: 0.75–0.99] and 0.89 [95% confidence interval: 0.80–0.99]) for women who were overweight (ie, who reported a silhouette equal or greater than the fifth silhouette according to the drawings of Sørensen et al⁴) at age 8 and at menarche, respectively. We observed no clear effect modification according to menopausal status, age at menarche, delay between age at menarche and regular cycling, regularity of cycles in adult life, or BMI at baseline. Our results suggest that overweight during childhood or adolescence reduces breast cancer risk and that this reduction is not fully explained by menstrual characteristics after menarche or by adult BMI but, rather, by hormonal mechanisms in the peripubertal period, when mammary tissue develops.

Several contradictory explanations may be put forward. According to Stoll,⁵ excess adiposity around menarche can alter the production of ovarian hormones and increase the frequency of anovulatory cycles, notably through insulin pathways, which lead to a decrease in the promotion of mammary carcinogenesis. However, others have shown that the leanest body silhouette at age 10 was associated with both higher peak growth velocities during adolescence and elevated breast carcinoma risk.⁶ Baer et al⁷ concluded that rapid adolescent growth may increase breast cancer risk by increasing levels of growth hormones and epithelial proliferation in the breast or by decreasing the amount of time for repair of DNA damage. Finally, alternative explanations have been proposed. Obesity in preadolescent and adolescent girls is associated with higher basal insulin levels, hyperandrogenia, decreased plasma levels of sex hormone-binding globulin, and, consequently, increased levels of unbound testosterone and estradiol.⁷ Higher prepubertal estrogenic exposures might paradoxically reduce breast cancer risk through differentiation of terminal end buds, mammary gland structures that are known to be the sites for malignant transformation.⁸ Additional research is required, therefore, to fully assess the consequences of the adolescent overweight epidemic.

References

1. Lee JL, Appugliese D, Kaciroti N, Corwyn RF, Bradley RH, Lumeng JC. Weight status in young girls and the onset of puberty [published correction appears in *Pediatrics*. 2007;120: 251]. *Pediatrics*. 2007;119(3).
2. Kelsey JL. A review of the epidemiology of human breast cancer. *Epidemiol Rev*. 1979;1:74–109
3. Tehard B, Kaaks R, Clavel-Chapelon F. Body silhouette, menstrual function at adolescence and breast cancer risk in the E3N cohort study. *Br J Cancer*. 2005;92:2042–2048
4. Sørensen TI, Stunkard AJ, Teasdale TW, Higgins MW. The accuracy of reports of weight: children’s recall of their parents’ weights 15 years earlier. *Int J Obes*. 1983;7:115–122
5. Stoll BA. Teenage obesity in relation to breast cancer risk. *Int J Obes Relat Metab Disord*. 1998;22:1035–1040
6. Berkey CS, Frazier AL, Gardner JD, Colditz GA. Adolescence and breast carcinoma risk. *Cancer*. 1999;85:2400–2409
7. Baer HJ, Colditz GA, Rosner B, et al. Body fatness during childhood and adolescence and incidence of breast cancer in premenopausal women: a prospective cohort study. *Breast Cancer Res*. 2005;7:R314–R325

8. Hilakivi-Clarke L, Wang C, Kalil M, Riggins R, Pestell RG. Nutritional modulation of the cell cycle and breast cancer. *Endocr Relat Cancer*. 2004;11:603–622