



HAL
open science

The fate of new production from N₂ fixation

M. R. Mulholland

► **To cite this version:**

M. R. Mulholland. The fate of new production from N₂ fixation. Biogeosciences Discussions, 2006, 3 (4), pp.1049-1080. hal-00297834

HAL Id: hal-00297834

<https://hal.science/hal-00297834>

Submitted on 18 Jun 2008

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Biogeosciences Discussions is the access reviewed discussion forum of *Biogeosciences*

The fate of new production from N₂ fixation

M. R. Mulholland

Department of Ocean, Earth and Atmospheric Sciences, Old Dominion University, 4600
Elkhorn Avenue, Norfolk, Virginia 23 529-0276, USA

Received: 8 May 2006 – Accepted: 19 May 2006 – Published: 19 July 2006

Correspondence to: M. R. Mulholland (mmulholl@odu.edu)

BGD

3, 1049–1080, 2006

Fate of N₂ fixation

M. R. Mulholland

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

Abstract

While we now know that marine N_2 fixation is a significant source of new nitrogen (N) in the marine environment, little is known about the fate of this production, despite the importance of diazotrophs to global carbon and nutrient cycles. Specifically, does new production from N_2 fixation fuel autotrophic or heterotrophic growth, facilitate carbon (C) export from the euphotic zone, or contribute primarily to microbial productivity and respiration in the euphotic zone? For *Trichodesmium*, the diazotroph we know the most about, the transfer of recently fixed N_2 (and C) appears to be primarily through dissolved pools. The release of N appears to vary among and within populations and, probably as a result of the changing physiological state of cells and populations. The net result of trophic transfers appears to depend on the complexity of the colonizing community and co-occurring organisms. In order to understand the impact of diazotrophy on carbon flow and export in marine systems, we need a better assessment of the trophic flow of elements in *Trichodesmium* communities dominated by different species, various free and colonial morphologies, and in various defined physiological states. Nitrogen and carbon fixation rates themselves vary by orders of magnitude within and among studies highlighting the difficulty in extrapolating global rates of N_2 fixation from direct measurements. Because the stoichiometry of N_2 and C fixation does not appear to be in balance with the stoichiometry of particles, and the relationship between C and N_2 fixation rates is also variable, it is equally difficult to derive global rates of one from the other. A better understanding of the physiology and physiological ecology of *Trichodesmium* and other marine diazotrophs is necessary to understand and predict the effects of increased or decreased diazotrophy in the context of the carbon cycle and global change.

BGD

3, 1049–1080, 2006

Fate of N_2 fixation

M. R. Mulholland

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

1 Introduction

Although a variety of marine cyanobacteria and bacteria are now known to fix dinitrogen (N_2) in marine environments, *Trichodesmium* spp. remain the most studied and most quantitatively significant pelagic nitrogen fixer based on available information. *Trichodesmium* spp. occur throughout the subtropical and tropical ocean where it can represent up to half of the primary production (Carpenter et al., 2004). Based on direct rate measurements, *Trichodesmium* accounts for a quarter to half of geochemically derived estimates of marine N_2 fixation (Mahaffey et al., 2005). In addition to *Trichodesmium*, pelagic nitrogen fixers include other filamentous cyanobacteria, unicellular cyanobacteria, bacterioplankton, and cyanobacterial endosymbionts (Carpenter and Capone, 2006)¹ Global estimates of nitrogen fixation and possible controls on marine nitrogen fixation have been recently summarized and reviewed (LaRoche and Breitbarth, 2005; Mahaffey et al., 2005; Carpenter and Capone, 2006¹) and so will not be re-reviewed here. While we now know that marine N_2 fixation is significant source of new nitrogen (N) in the marine environment, little is known about the fate of this production, despite the importance of diazotrophs to global carbon and nutrient cycles (Karl et al., 2002; LaRoche and Breitbarth, 2005). Growth rates of these organisms vary by orders of magnitude as do rates of N_2 and carbon fixation (see Mulholland et al., 2006) and reasons for this variability are not well understood. Inputs of N and carbon (C) via N_2 fixation and associated carbon fixation have been measured directly or extrapolated in a variety of systems, however, the quantification of loss terms is poorly constrained. *Trichodesmium* are rarely found in sediment traps and are positively buoyant (Walsby, 1992) and so sinking appears to be a minor loss term compared with cell lysis (Ohki, 1999; Hewson et al., 2004), extracellular release (Capone et al., 1994; Glibert and Bronk, 1994; Mulholland et al., 2004a), and grazing (O'Neil et al., 1996; O'Neil, 1999)

¹Carpenter, E. J. and Capone, D. G.: Nitrogen fixation in the marine environment, in: Nitrogen in the Marine Environment, edited by: Capone, D. G., Bronk, D. A., Mulholland, M. R., and Carpenter, E. J., in review, 2006.

BGD

3, 1049–1080, 2006

Fate of N_2 fixation

M. R. Mulholland

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

each of which are discussed below. In this paper, the fate of production from N₂ fixation will be examined based on observations of N₂ fixation and associated carbon fixation rates, N and C release, and trophodynamics associated with nutrient cycling.

2 N₂ fixation

5 Pelagic N₂ fixation is an important source of new N to marine systems. The most widely studied pelagic marine diazotrophs, *Trichodesmium* spp., play a pivotal role in marine elemental cycles in otherwise oligotrophic tropical and subtropical seas (Capone et al., 1997; Karl et al., 2002; LaRoche and Breitbarth, 2005). Globally, based on direct measurements, oceanic N₂ fixation by *Trichodesmium* has been estimated to be 80 Tg N
10 year⁻¹ (Capone and Carpenter, 1999), and represents upwards of 50% of the new production in some oligotrophic tropical and subtropical oceans (Karl et al., 1997; Capone et al., 2005; Mahaffey et al., 2005). Based on observed and derived N₂ fixation rates by *Trichodesmium*, N₂ fixation by these species alone is comparable in magnitude to estimated nitrate flux across the base of the euphotic zone in tropical and subtropical
15 systems (Karl et al., 1997; Capone, 2001; Capone et al., 2005).

However, *Trichodesmium* still represent only 40 to 59% of the geochemically inferred N₂ fixation for the North Atlantic and Pacific (Mahaffey et al., 2005). The recent discoveries of diazotrophic unicellular cyanobacteria and bacterioplankton in marine systems (e.g., Zehr et al., 2001; Falcón et al., 2004; Montoya et al., 2004) suggest that there
20 are additional sources of N₂ fixation that may yet reconcile measured with geochemically predicted rates of N₂ fixation in the ocean. Although, the full range of diazotrophic marine organisms is as yet unknown, it is thought that unicellular diazotrophs may contribute up to 10% of global new production (Montoya et al., 2004). *Richelia intracel-*
25 *lularis*, an endosymbiotic cyanobacterium that can inhabit a diverse group of diatoms, fix significant amounts of nitrogen where these associations occur (Carpenter et al., 1999). Based on the available rate measurements and regional and global abundance estimates in the euphotic zone, endosymbiotic and free-living unicellular cyanobacteria

BGD

3, 1049–1080, 2006

Fate of N₂ fixation

M. R. Mulholland

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

and bacteria are now believed to fix at least as much nitrogen as *Trichodesmium* in the ocean (Table 1, Carpenter et al., 1999; Zehr et al., 2001; Montoya et al., 2004). As a result, recent estimates for total pelagic marine N₂ fixation are now estimated to be between 100 and 200 Tg N year⁻¹ (Karl et al., 2002; Galloway et al., 2004).

In addition to extrapolations from direct measurements of N₂ fixation rates, global estimates of marine N₂ fixation have been inferred based on geochemical arguments that rely on elemental stoichiometry of particles and dissolved nutrients in the ocean (see Mahaffey et al., 2005, for a more complete discussion). There are limitations to both of these approaches, however, because of methodological constraints and the physiological peculiarities of the dominant marine N₂ fixer, *Trichodesmium*, discussed below. The physiology of more recently identified N₂ fixers is still being elucidated and so it is premature to speculate on how these groups may influence estimates of global new production and carbon export.

Extrapolation of N₂ fixation rates made in laboratory or field populations of *Trichodesmium* to the world's ocean can yield a wide range of global marine N₂ fixation rates. For example, rates of N₂ fixation by *Trichodesmium* from field populations vary by six orders of magnitude (LaRoche and Breitbart, 2005; Mulholland et al., 2006). Laboratory estimates vary only by about 4 orders of magnitude, but still, which rates do we choose for our global estimate? Based on laboratory studies, rates of N₂ fixation vary according to physiological state and yet the physiological state of natural populations is impossible to assess at the time of sampling. It is thought that rates of N₂ fixation and growth by *Trichodesmium* are limited by phosphorus (P), iron (Fe), or light (Sañudo-Wilhelmy et al., 2001; Mills et al., 2004; Fu et al., 2005; Mulholland and Bernhardt, 2005). However, the range of responses to these variables and their interactions is unknown (Mulholland and Bernhardt, 2005).

In addition to real physiological variability, rates vary depending on the method used to estimate N₂ fixation. The two most commonly used methods are the acetylene reduction method and ¹⁵N₂ uptake; the former measures gross N₂ fixation and the latter measures net N₂ uptake into biomass. The acetylene reduction method relies

BGD

3, 1049–1080, 2006

Fate of N₂ fixation

M. R. Mulholland

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

on a conversion factor to convert moles of acetylene to moles N₂ reduced and the value of this conversion factor has been a matter of debate (see Mulholland et al., 2006; LaRoche and Breitbarth, 2005). Paired comparisons between the two methods, used to calibrate one against the other, demonstrate that the ratio between acetylene reduction and N₂ uptake varies widely both within and among systems and studies (Table 2). Consequently, we are left with an unsatisfying set of data with which to make direct estimates of global N₂ fixation.

3 Carbon fixation

Global estimates of carbon fixation by marine diazotrophs based on direct measurements have not been attempted to my knowledge. There are fewer published estimates of carbon fixation by *Trichodesmium* and global carbon fixation by this genus is generally estimated by multiplying the nitrogen fixation rate by some average C:N for *Trichodesmium* biomass. Modeling results assume N₂ fixation equals denitrification, and this corresponds to 480 to 960 Tg C year⁻¹ (Mahaffey et al. 2005). Fortunately, the C:N ratio of *Trichodesmium* biomass, unlike the N:P ratios, fall within a narrow range (e.g., 4.7 to 7.3; LaRoche and Breitbarth, 2005) with an average value of 6.3, very near the Redfield ratio. Unfortunately, as for N₂ fixation, direct rate measurements of carbon fixation and carbon specific turnover times by *Trichodesmium* vary by orders of magnitude (Mulholland et al., 2006). Further, there is no consistent stoichiometric relationship between the ratio of C to N₂ fixation. Available paired estimates of N₂ and C fixation suggest that in general, C:N fixation ratios far exceed the C:N ratio of cells (Table 3; Mulholland et al., 2006). Consequently, geochemical estimates that rely on elemental stoichiometry to extrapolate of N₂ fixation from observations of carbon drawdown or the carbon cycle in general may be grossly in error (Mahaffey et al., 2005). For example, at the Bermuda Atlantic Time-Series Study (BATS) site, the observed rates of C drawdown were much higher than that which can be accounted for based on the observed rates of N₂ fixation and Redfield stoichiometry (e.g., C:N of 6.6:1).

BGD

3, 1049–1080, 2006

Fate of N₂ fixation

M. R. Mulholland

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

However, when the average observed ratio of carbon to N₂ fixation rates measured at BATS (C:N₂ fixation rates of 128) were used, the observed low rates of N₂ fixation could indeed account for the observed carbon drawdown at BATS (Orcutt et al., 2001). Interestingly, the extrapolation of N₂ fixation rates necessary to close C budgets may be seriously biased (overestimated) if the actual rate relationships between N₂ and carbon fixation are not considered. The relationship between N and P may be even more complex.

There are a variety of reasons why there may be higher-than-stoichiometrically-expected carbon to N₂ fixation ratios in nature; these include factors resulting in underestimates of N₂ fixation rates and rationalizations as to why *Trichodesmium* may have unusually high carbon fixation rates. Gross N₂ fixation rates can be underestimated in ¹⁵N₂ incubations if there is substantial N release (Glibert and Bronk, 1994; Mulholland et al., 2004a, 2006; see Sect. 4 below) or gross N utilization may be underestimated if alternative N sources are taken up (Mulholland and Capone, 1999; Mulholland et al., 1999a, 1999b). On the other hand, carbon fixation rates may be stoichiometrically higher than expected, based on the elemental ratio of cells, if carbon is used as ballast for vertical migration (Villareal and Carpenter, 1990; Romans et al., 1994; Gallon et al., 1996), if substantial carbon is excreted as mucilage or extracellular polymeric substances (Stal, 1995; Sellner, 1997), to support the high observed respiration rates by *Trichodesmium* (Carpenter and Roenneberg, 1995; Kana, 1993), or if cells “over-fix” carbon to support Mehler reactions to reduce cellular oxygen concentrations or support the production of ATP (Kana, 1992, 1993). Kana (1993) estimated that 48% of gross photosynthetic electron flow went to oxygen reduction. *Trichodesmium* also make poly-beta-hydroxybutyric acid as a storage product (Siddiqui et al., 1992) and this may be important in carbohydrate ballasting (Romans et al., 1994) but would require additional cellular carbon reserves. In addition to these physiological reasons why carbon might be “over-fixed” relative to nitrogen, active release of carbon compounds and photosynthate has been observed and will be discussed below in Sect. 5. Alternatively, N and C uptake may not be tightly coupled in diazotrophic cyanobacteria (Gallon et al., 2002).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Fate of N₂ fixation

M. R. Mulholland

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Another interesting genomic finding is that *Trichodesmium erythraeum* is unusual among cyanobacteria in that it lacks any genes encoding known high-affinity carbon concentrating mechanisms (Badger and Price, 2003). While it is not clear how this affects photosynthetic C acquisition by *Trichodesmium*, it suggests this species is vulnerable to C limitation. Continuous carbon fixation or storage of fixed carbon compounds, even in excess of their growth requirements, may protect them from carbon limitation in nature. Although carbon limitation has not been demonstrated for these species in nature, the interior of *Trichodesmium* colonies have been shown to exhibit oxygen dynamics that may be an important for aerobic N₂ fixation (Paerl and Bebout, 1988; Carpenter et al., 1990; Gallon, 1992).

Although the ultimate biogeochemical fate of *Trichodesmium*-fixed elements is not fully understood (Mahaffey et al., 2005; Mulholland et al., 2006), any fraction of new production from diazotrophy that is exported to underlying waters will contribute to sequestering atmospheric carbon and so it is important that we gain a better understanding of the coupled N and C cycles for these organisms.

4 N release

Trichodesmium sp. can fix N₂ at high rates, thereby introducing new nitrogen N into nutrient impoverished areas of the tropical and subtropical ocean. Because *Trichodesmium* also release fixed N as dissolved organic N (DON) (Glibert and Bronk, 1994), amino acids (Capone et al., 1994), and ammonium (Mulholland and Capone, 2001; Mulholland et al., 2004a), they may contribute to regenerated production where they occur. Elevated NH₄⁺ and/or DON concentrations have also been observed in and around *Trichodesmium* blooms in the Pacific (Devassy et al., 1978, 1979; Karl et al., 1992, 1997), the Gulf of Mexico (Lenes et al., 2001), along the coast of Australia (Glibert and O'Neil, 1999) and in aging *Trichodesmium* cultures (Mulholland and Capone, 2001). However, nutrient concentrations within and around blooms may not always be high if the released N is rapidly taken up by organisms growing on and

around colonies (e.g., see Sellner, 1992, 1997) or co-occurring in the water column (Mulholland et al., 2006).

In addition to direct release of labile N, viral cell lysis (Ohki, 1999; Hewson et al., 2004), grazing (O'Neil et al., 1996), and cell death (Berman-Frank et al., 2004) may contribute dissolved or particulate N to the available N pool. *Trichodesmium* biomass may also be degraded via bacterial activity and extracellular enzymes thereby rendering large organic compounds into smaller utilizable compounds (Paerl et al., 1989; Nausch, 1996)

N release from *Trichodesmium* was first suggested by Devassy et al. (1978) who observed substantial enrichment of phosphate, nitrate and ammonium within, during, and after *Trichodesmium* blooms and relative to non-bloom sites or at times prior to blooms. Based on changes in nutrient concentrations within incubations, Devassy et al. (1978) estimated 1.5 $\mu\text{mol N}$ (as inorganic N) and 6.8 $\mu\text{mol inorganic P}$ was released per g of *Trichodesmium*. These authors also suggested that release rates would have been much higher if DOP, urea and amino acids had been measured.

N release has been difficult to estimate using isotopic tracer or other methods for a number of reasons:

(1) release products may be diverse and so it is often difficult to isolate and measure all relevant dissolved pools, (2) release products are rapidly taken up by organisms in oligotrophic environments, and (3) intracellular pools of intermediate metabolites can accumulate before their release for variable amounts of time.

To get around these problems, it has been suggested that the difference between net and gross N_2 fixation measured using $^{15}\text{N}_2$ uptake and acetylene reduction techniques, respectively, might be a good metric of N release (Gallon et al., 2002; Mulholland et al., 2004a, 2006).

High N release rates would seem to argue for high cellular N turnover, however, if N is released prior to its assimilation into biomass, this would contribute to gross N_2 fixation (e.g., reduction of N_2 to NH_4^+) but not net uptake into biomass. In numerous paired comparisons between acetylene reduction and $^{15}\text{N}_2$ uptake, ratios of acetylene

BGD

3, 1049–1080, 2006

Fate of N_2 fixation

M. R. Mulholland

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

(C₂H₂) reduced to N₂ taken up have varied by at least an order of magnitude (Table 2). Because C₂H₂ reduction measures just the reduction step, it is a measure of gross N₂ fixation while movement of ¹⁵N₂ from the dissolved to the particulate pool measures net N assimilation (see Gallon et al., 2002; Mulholland et al., 2004a, 2006; Mulholland and Bernhardt, 2005). Release of recently reduced N₂ and the difficulty in chemically recovering all possible dissolved pools into which products of N₂ fixation might be released, may make intercalibration between the two methods impossible. However, the difference between N₂ reduction (gross N₂ fixation) and net N₂ assimilation may prove to be an excellent index of the release of recently fixed N₂ (Mulholland et al., 2004a, 2006; Mulholland and Bernhardt, 2005). If this is the case, and the theoretical ratio, three, is assumed to be correct (see Mulholland et al., 2006, for a discussion of this assumption) in estimating N₂ fixation from C₂H₂ reduction, paired comparisons in which C₂H₂:N₂ reduction ratios of approximately 3:1 are observed, would indicate no N release while C₂H₂:N₂ reduction ratios of six would translate into a release rate of about 50%, and so on.

Examining paired comparisons of C₂H₂ reduction and ¹⁵N₂ uptake from recent studies and the literature (Table 1), rates of release of recently fixed N can be compared across systems and with respect to temperature. Results demonstrate that release rates are highly variable on a variety of temporal and spatial scales. Release rates appeared to be high in populations collected from a South Pacific lagoon, varied seasonally in the North Atlantic along a latitudinal gradient, and on a daily and interannual basis in the Gulf of Mexico (Table 2). In cultures, release rates varied with the growth rate (Mulholland and Bernhardt, 2005) and with the time of day (Table 2). Based on various studies where ¹⁵N₂ and acetylene reduction were compared directly, it appears that N release from *Trichodesmium* is common but varies with physiological state and among environments. One paired comparison is available for a diatom/*Richelia* association (Table 2; Carpenter et al., 1999) and suggests that there was N release from N₂ fixation for this association as well. No published paired comparisons of C₂H₂:N₂ reduction for unicells are available and so release by these organisms cannot be as-

BGD

3, 1049–1080, 2006

Fate of N₂ fixation

M. R. Mulholland

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

sessed at this time.

While there have been observations that *Trichodesmium* release recently fixed N₂ as DON in natural populations (Capone et al., 1994; Glibert and Bronk, 1994) and as NH₄⁺ in cultures of *Trichodesmium* IMS101 (Mulholland et al., 2004a; Mulholland and Bernhardt, 2005) it is unclear why cells do this. Previous speculation suggested that this is a mechanism for the extracellular transfer of fixed N between cells that fix N₂ and those that do not have that capability (Mulholland and Capone, 1999, 2000). Another possible fate for regenerated N is co-occurring organisms (O'Neil et al., 1996; Mulholland et al., 2004b, 2006). Release rates averaged about 52% of the recently fixed N₂ or 0.29 nmol col⁻¹ h⁻¹ in a recent study in the Gulf of Mexico (Mulholland et al., 2006), much of this may have fueled production of co-occurring phytoplankton (Mulholland et al., 2004b).

N release from cellular material can also be mediated through cell lysis. Viruses and a lytic cycle have been observed in natural populations and cultures of *Trichodesmium* (Ohki, 1999). Hewson et al. (2004) estimated lysis rates of 0.3 to 6.5% trichomes d⁻¹, representing a release of 3 to 65% of the production for *Trichodesmium* growing at 0.1 d⁻¹. While these authors report this as 3 to 65% of recently fixed N d⁻¹, this also applies to C (see below). At an average rate of 43 pmol N fixed trichome⁻¹ d⁻¹, this represents a release rate of 1.3 to 28 pmol trichome⁻¹ d⁻¹ or (using an average colony size of 100 trichomes col⁻¹) 13 to 280 pmol col⁻¹ d⁻¹. For C, this could be substantially greater than a factor of 6 if stoichiometrically more C is fixed per day. These estimates agree well with calculated mortality rates for *Trichodesmium* derived from models at BATS and in the equatorial Atlantic estimate 2.1 to 2.5% d⁻¹ (Hood et al., 2002, 2004).

5 C release

Dissolved organic N also contains C and so it is therefore logical to assume that *Trichodesmium* also release substantial amounts of photosynthetic products as dissolved organic carbon (DOC). In fact, Shimura et al. (1978) first quantified the extracellular re-

BGD

3, 1049–1080, 2006

Fate of N₂ fixation

M. R. Mulholland

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

Fate of N₂ fixation

M. R. Mulholland

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

lease of photosynthate from ¹⁴C incubations and calculated that about 8% of the total photosynthetic products were released during incubation experiments (range 0–18%). Similarly, Gallon et al. (1996) estimated that DOC excretion by *Trichodesmium* in the western North Atlantic and eastern Caribbean Sea represented only 7% of the primary productivity. As for N, the amount of C released changed depending on light conditions and the physiological status of cells. More recently, Renaud et al. (2005) estimated a much lower value (1%) for DOC release by *Trichodesmium*. However, they suggested that tight coupling between organisms in the *Trichodesmium* consortium might cause underestimates of actual release rates. Thus, the same methodological limitations that make it difficult to estimate N release from tracer studies make it difficult to make estimates of C release; just as ¹⁵N₂ uptake can underestimate gross N₂ fixation, ¹⁴C (or ¹³C) incorporation can underestimate the gross rate of photosynthetic carbon fixation (Gallon et al., 2002).

Cyanobacteria release C compounds such as glycolate (Renström-Kellner et al., 1989), as well as, amino acids which contain N and C (Capone et al., 1994). Amino acid release as glutamine and glutamate (molar C:N ratios of 5:2 and 5:1, respectively) represented only 3% of the C fixed by *Trichodesmium* (Capone et al., 1994). However, *Trichodesmium* have a carbohydrate mucoid matrix, which is colonized by other organisms (Stal, 1995; Sellner, 1997; Sheridan et al., 2002) and so there is a constant production of glucose- and mannose-rich mucilage that could account for more DOC release (e.g., Sellner, 1997). Cyanobacteria in general can exude as much as 80% of the CO₂ they fix as extracellular polymeric substances (mainly polysaccharides).

Production of colored dissolved organic matter (CDOM) by *Trichodesmium* was recently been observed (Steinberg et al., 2004). Production of DOC ranged from 0.04 to 0.32 μg C col⁻¹ h⁻¹. Assuming an average of 11.3 μg C col⁻¹ (McCarthy and Carpenter, 1979), this represents between 0.4 and 2.8% h⁻¹ or up to 67% d⁻¹, although it is unclear whether this production is confined to the dark or light periods. The CDOM had absorption spectra similar to microsporin-like amino acids, compounds that *Trichodesmium* are known to have and serve in photoprotection (Subramaniam et

al., 1999).

Although globally, we are interested in the fate of new production from N₂ fixation as a means to export C, little has been done to quantify or characterize DOC release from *Trichodesmium* or other marine N₂ fixers. If we are to extrapolate export from production of N₂ fixers, it will be important to determine the primary pathways of C flow through these organisms.

6 Trophic interactions

It is impossible to discuss the fate of new production by diazotrophs without discussing trophic interactions. Colonies of *Trichodesmium* provide stable “homes” for a numerous and diverse association of organisms (Siddiqui et al., 1992; Sellner, 1997; O’Neil, 1999; Sheridan et al., 2002). This creates a complex microenvironment with multifarious pathways for internal nutrient cycling. Sheridan et al. (2002) estimated that 85% of *Trichodesmium* colonies were inhabited by other organisms. Colonizing organisms include bacteria, other cyanobacteria, fungi, pennate and centric diatoms, heterotrophic and autotrophic dinoflagellates, chrysophytes, ciliates, amoebae, hydroids, different life stages of harpacticoid copepods and juvenile decapods. Bacteria and dinoflagellates were the most common associates. Despite the fact that colonies are rich microenvironments, there is a variety of evidence suggesting that *Trichodesmium* themselves go largely ungrazed and so viral cell lysis and decomposition are the likely fates for many of these populations (nutrient accumulation), and that the importance of higher trophic levels in processing *Trichodesmium* biomass is minimal as compared to recycled primary production and bacterial productivity.

Consistent with this idea is the observation that a variety of phytoplankton, bacteria, and higher trophic levels co-occur or occur in the water column subsequent to blooms of *Trichodesmium* sp. (Devassy, 1978, 1979; Revelante et al., 1982; Furnas and Mitchell, 1996; Walsh and Steidinger, 2001; Mulholland et al., 2006). It is thought that these communities are relieved from N limitation as a result of N release from *Trichodesmium*.

BGD

3, 1049–1080, 2006

Fate of N₂ fixation

M. R. Mulholland

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

Fate of N₂ fixation

M. R. Mulholland

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Trichodesmium occur as variously sized and shaped aggregates or colonies but also as free filaments or trichomes. Large colonies may contain hundreds of trichomes. However, the average colony size and colony abundance can vary from day to day (Devassy et al., 1978). Colonies take the form of bundles with trichomes arranged in parallel (tufts) or radially (puffs). Little is known about the causes of bundle formation but, the distributions of free filaments and bundles vary regionally and apparently with the degree of turbulence (Bryceson and Fay, 1981; Mahaffey et al., 2005). The purpose of bundle formation is unclear but, there has been speculation that it may be a behavioral strategy for minimizing the exposure of nitrogenase (an oxygen sensitive protein) to oxygen (e.g., Paerl et al., 1989; Gallon, 1992). Regardless of the reasons colonies form, the trophodynamics of *Trichodesmium* varies depending on the form it takes and the amount of stable surface area and interfilamental space available for colonization.

There are few direct measurements of the trophic transfer of recently fixed N or C. Bryceson and Fay (1981) first demonstrated that the trophic transfer of recently fixed N₂ might be important in communities dominated by *Trichodesmium* and they subsequently demonstrated enrichment in non-*Trichodesmium* size fractions after incubation of *Trichodesmium* and natural marine communities with ¹⁵N₂. They did not have control incubations to account for N₂ fixation by smaller diazotrophic cyanobacteria and bacterioplankton but, nevertheless, they report enrichment in the 2 to 30 μm and 0.2 to 2.0 μm size fractions (Bryceson and Fay, 1981). Subsequently, the only other direct estimates of the trophic transfer of recently fixed N₂ demonstrated that up to 11% of recently fixed N₂ was transferred to non-N₂ fixing cells in whole water samples even in short (2 h) incubations (Mulholland et al., 2004b). This suggests that *Trichodesmium* may support community productivity in the upper water column and the growth of co-occurring organisms, including heterotrophs, rather than substantial direct sinking flux.

Despite the idea that dissolved nutrients may be the primary route of trophic transfer of recently fixed N₂, isotopically “light” zooplankton were collected from the tropical Atlantic (Montoya et al., 2002) and isotopically light sediment trap material was collected under a station experiencing a *Trichodesmium* bloom in the Indian Ocean (Capone et

al., 1998) indicating that recently fixed N_2 , which has an isotopic signature similar to atmospheric N, is being transferred to higher trophic levels.

6.1 Bacteria

Bacterial associates with *Trichodesmium* colonies have been widely observed (Paerl et al., 1989; Nausch, 1996; Sheridan et al., 2002; Renaud et al., 2005; Mulholland et al., unpublished data). *Trichodesmium* colonies are inhabited by both rod-shaped and filamentous bacteria, as are many other filamentous cyanobacteria (Paerl et al., 1989). Bacteria associates, including heterotrophic N_2 fixers, were located around and within aggregates where they took up carbohydrates and amino acids.

Varying degrees of enrichment of bacteria have been found on and around colonies. Nausch (1996) reported that bacteria were 2 to 5 times higher on colonies of *Trichodesmium* than in the surrounding water, however, during her study, *Trichodesmium* were not abundant, the water column was turbulent, and colonies were small. At BATS, Sheridan et al. (2002) report that bacteria were enriched on average 401 and 1709 times on *Trichodesmium* puffs and tufts, respectively. Carpenter and Price (1977) found that up to 8.3% of *Trichodesmium* were populated by bacteria in the Sargasso Sea. So, it appears that there is high variability in the degree of colonization of bacterial colonization of *Trichodesmium* aggregates.

In terms of their productivity, Nausch (1996) found thymidine incorporation to be enhanced in colonies of *Trichodesmium* relative to that of the water column and comparable to the enrichment found in marine snow. However, productivity per unit bacteria appeared to be lower in colonies than in the surrounding water. In the Gulf of Mexico, leucine uptake increased by up to 72% in association with *Trichodesmium* colonies relative to the surrounding water column (Mulholland et al., unpublished data). Similarly, Tseng et al. (2005) found that bacterial productivity and abundance was higher but productivity per unit bacterial biomass was lower, in association with *Trichodesmium* populations. In addition, they found that populations became more autotrophic during times of the year when *Trichodesmium* was abundant (lower bacterial productiv-

BGD

3, 1049–1080, 2006

Fate of N_2 fixation

M. R. Mulholland

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

ity: primary productivity ratio). The authors attribute this to N release and alleviation of competition between bacteria and phytoplankton for scarce NH_4^+ . However, they also note that *Trichodesmium* occurred as free filaments in the Kuroshio and therefore lacked harpacticoid grazer populations and associated organisms observed in other communities (Tseng et al., 2005).

High rates of amino acid oxidase activity (Mulholland et al., 1998; Glibert and O'Neil, 1999), peptide hydrolysis (Mulholland et al., unpublished data), and hydrolytic enzyme activity have also been found in association with *Trichodesmium* colonies, suggesting bacteria and organisms associated with colonies actively cycle nutrients. Nausch (1996) calculated C and N release rates between 30.5 and 1086 $\text{ng C col}^{-1} \text{ h}^{-1}$ and 4.6 to 209 $\text{ng N col}^{-1} \text{ h}^{-1}$, respectively, based on hydrolytic enzyme activities associated with *Trichodesmium* colonies.

6.2 Phytoplankton

In some coastal systems, blooms of dinoflagellates and diatoms have been observed during and subsequent to *Trichodesmium* blooms (Devassy et al., 1978; Revelante et al., 1982; Furnas and Mitchell, 1996). For example, Devassy et al. (1979) found that as blooms of *Trichodesmium* decayed, *Chaetoceras* populations increased, followed by a succession of cladocerans, dinoflagellates, green algae (*Noctiluca*), copepods, and finally, carnivores. On the West Florida shelf, dense *Karenia brevis* blooms occur during and subsequent to *Trichodesmium* blooms and it has been hypothesized that they provide a source of new N to fuel destructive red tides (Walsh and Steidinger, 2001, Mulholland et al., 2006). Based on direct estimates of N_2 fixation, N release, and in situ water column N uptake rates, *Trichodesmium* produced ample dissolved N to fuel *K. brevis* population growth in the Gulf of Mexico (Mulholland et al., 2006).

Experiments suggest that *Tetraselmis* grew well on decaying *Trichodesmium* (Devassy et al., 1978). Similarly, *Karenia brevis* cultures grew well on culture medium enriched in *Trichodesmium* exudates as the sole source of nitrogen (Mulholland et al., unpublished data). While direct evidence of trophic transfer from *Trichodesmium*

BGD

3, 1049–1080, 2006

Fate of N_2 fixation

M. R. Mulholland

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

to phytoplankton in nature are lacking, Bryceson and Fay (1981) and Mulholland et al. (2004b), demonstrated that ^{15}N derived from $^{15}\text{N}_2$ additions moved into the co-occurring plankton, which presumably included a variety of phytoplankton.

6.3 Zooplankton and higher trophic levels

5 The fate of recently fixed N_2 and transfer of *Trichodesmium* biomass to higher trophic levels is poorly understood. Although a variety of herbivores are thought to graze on *Trichodesmium* (e.g., Sellner, 1997), *Trichodesmium* sp. are not grazed by many of the dominant zooplankton in marine systems and are toxic to many copepods (Hawser and Codd, 1992; O'Neil, 1999). Some specialized harpacticoid copepods do graze
10 on and inhabit *Trichodesmium* colonies but these do not produce fecal pellets that would rapidly remove grazed material from the euphotic zone (O'Neil and Roman, 1994; O'Neil et al., 1996).

O'Neil et al. (1996) estimated that the harpacticoid copepod, *Macrosetella*, could consume 33–45% of total colony N, or 100% of the new N_2 fixed each day. The copepod then excretes 48% of its body N per day, mainly as NH_4^+ , thereby recycling much of the N in the water column (O'Neil et al., 1996). Further, Roman (1978) found that *Macrosetella* could ingest from 90 to 126% of its body carbon per day when feeding on *Trichodesmium*. Based on stoichiometric arguments, O'Neil (1999) calculated 30% of the recently fixed C from *Trichodesmium* flowed into grazers and because *Macrosetella*
15 appear to have a higher C:N ratio than the *Trichodesmium* themselves, they are likely to excrete excess N. Therefore, the major flux of recently fixed N and C through zooplankton may also be through extracellular release and dissolved nutrient pools. In addition to excretory release, zooplankton grazers can mediate the transfer of N through additional release from sloppy feeding (O'Neil and Roman, 1996).

25 Not much is known about higher trophic levels, although isotopic evidence suggests that there are other grazers of *Trichodesmium*. In general, it has been observed that there is a low quality of fish associated with blooms although *Trichodesmium* do not appear to be directly toxic to fish (Devassy et al., 1978). Fish and some other higher

BGD

3, 1049–1080, 2006

Fate of N_2 fixation

M. R. Mulholland

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

trophic levels have been observed to graze on *Trichodesmium* (see Carpenter and Capone, 2006¹).

7 Implications

While we appear to be making strides in our ability to derive global estimates for marine N₂ fixation, we have a long way to go before we understand the role of N₂ fixation in the context of C dynamics in the ocean. Because many direct estimates of global N₂ fixation are based on highly spatially, temporally, and physiologically variable data, and because many geochemical estimates rely on stoichiometric relationships of nutrient standing stocks without considering the imbalances between rate estimates of C and N₂ fixation we should proceed cautiously when inferring one from the other (see examples in Sect. 3 above). Based on the observed C drawdown from the atmosphere, we may be trying to find “too much” N₂ fixation if we use Redfield stoichiometry versus the observed relative rates of C and N₂ fixation.

It is also difficult to determine the effect of N₂ fixation on system trophic status. In some systems *Trichodesmium* appears to fuel primary productivity and make the system more autotrophic (e.g., Tseng et al., 2005). In other systems dominated by *Trichodesmium*, heterotrophic processes appear to dominate (Fig. 1; also see Sect. 6.3 above). For the Kenyan coast, primary productivity, even during *Trichodesmium* blooms, could barely sustain the observed bacterial productivity (Kromkamp et al., 1997).

The preponderance of filaments versus colonial morphology can also seriously bias our understanding of trophodynamics associated with *Trichodesmium* and the net outcome of elemental cycling (e.g., recycling and respiratory losses versus export). Not only would it influence colony specific estimates of N and C fixation, but the degree of aggregation and size of colonies affect the degree to which *Trichodesmium* is colonized and thereby recycled to bacteria and copepods, and possibly other taxa. Free filaments often dominate populations in the Pacific (Saino and Hattori, 1978, 1980; Letelier and

BGD

3, 1049–1080, 2006

Fate of N₂ fixation

M. R. Mulholland

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

Karl, 1996; Tseng et al., 2005) and the Atlantic, at least seasonally (Orcutt et al., 2001) although in many systems, colonies appear to be more common (Capone et al., 1997; Carpenter et al., 2004).

Finally, diazotrophs may have different fates. Diatom/*Richelia* assemblages may be prone to gravitational settling while unicellular cyanobacteria may be more readily grazed. Not enough is known about community dynamics associated with these populations to speculate at this time.

Acknowledgements. Funding for this project was provided by the National Science Foundation, grants OCE-0095923 and OCE-0136367 to MRM. The author also wishes to thank C. Dupuyot and A. Leboutellier for their invitation and support to participate in field work in New Caledonia.

References

- Badger, M. R. and Price, G. D.: CO₂ concentrating mechanisms in cyanobacteria: molecular components, their diversity and evolution, *J. Exp. Bot.*, 54, 609–622, 2003.
- Berman-Frank, I., Bidle, K. D., Haramaty, L., and Falkowski, P. G.: The demise of the marine cyanobacterium, *Trichodesmium* spp., via an autocatalyzed cell death pathway, *Limnol. Oceanogr.*, 49, 997–1005, 2004.
- Bryceson, I. and Fay, P.: Nitrogen fixation in *Oscillatoria (Trichodesmium) erythroa* in relation to bundle formation and trichome differentiation, *Mar. Biol.*, 61, 159–166, 1981.
- Capone, D. G.: Marine nitrogen fixation: what's the fuss?, *Current Opinion in Microbiology*, 4, 341–348, 2001.
- Capone, D. G., Burns, J. A., Montoya, J. P., Subramaniam, A., Mahaffey, C., Gunderson, T., Michaels, A. F., and Carpenter, E. J.: Nitrogen fixation by *Trichodesmium* spp.: an important source of new nitrogen to the tropical and subtropical North Atlantic Ocean, *Global Biogeochem. Cycles*, 19, GB2024, doi:10.1029/2004GB002331, 2005.
- Capone, D. G. and Carpenter, E. J.: Nitrogen fixation by marine cyanobacteria: Historical and global perspectives, *Bull. Inst. Oceanogr. Monaco*, 19, 235–256, 1999.
- Capone, D. G., Ferrier, M. D., and Carpenter, E. J.: Amino acid cycling in colonies of the planktonic marine cyanobacterium, *Trichodesmium thiebautii*, *Appl. Environm. Microbiol.*, 60, 3989–3995, 1994.

BGD

3, 1049–1080, 2006

Fate of N₂ fixation

M. R. Mulholland

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

Fate of N₂ fixation

M. R. Mulholland

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Capone, D. G., Subramaniam, A., Montoya, J. P., Voss, M., Humborg, C., Johansen, A. M., Siefert, R. L., and Carpenter, E. J.: An extensive bloom of the N₂-fixing cyanobacterium *Trichodesmium erythraeum* in the central Arabian Sea, *Mar. Ecol. Prog. Ser.*, 172, 281–292, 1998.
- 5 Capone, D. G., Zehr, J. P., Paerl, H. W., Bergman, B., and Carpenter, E. J.: *Trichodesmium*, a globally significant marine cyanobacterium, *Science*, 276, 1221–1229, 1997.
- Carpenter, E. J., Chang, J., Cottrell, M., Schubauer, J., Paerl, H. W., Bebout, B. M., and Capone, D. G.: Re-evaluation of nitrogenase oxygen-protective mechanisms in the planktonic marine cyanobacterium *Trichodesmium*, *Mar. Ecol. Prog. Ser.*, 65, 151–158, 1990.
- 10 Carpenter, E. J. and McCarthy, J. J.: Nitrogen fixation and uptake of combined nitrogenous nutrients by *Oscillatoria (Trichodesmium) thiebautii* in the western Sargasso Sea, *Limnol. Oceanogr.*, 20, 389–401, 1975.
- Carpenter, E. J., Montoya, J. P., Burns, J., Mulholland, M. R., Subramaniam, A., and Capone, D. G.: Extensive bloom of a N₂-fixing symbiotic association in the tropical Atlantic Ocean, 15 *Mar. Ecol. Prog. Ser.*, 185, 273–283, 1999.
- Carpenter, E. J. and Price IV, C. C.: Nitrogen fixation, distribution, and production of *Oscillatoria (Trichodesmium)* spp. in the western Sargasso and Caribbean Seas, *Limnol. Oceanogr.*, 22, 60–72, 1977.
- Carpenter, E. J. and Roenneberg, T.: The marine planktonic cyanobacteria *Trichodesmium* spp.: photosynthetic rate measurements in the SW Atlantic Ocean, *Mar. Ecol. Prog. Ser.*, 20 118, 267–273, 1995.
- Carpenter, E. J., Subramaniam, A., and Capone, D. G.: Biomass and primary productivity of the cyanobacterium *Trichodesmium* spp. in the tropical N Atlantic Ocean, *Deep-Sea Res. I*, 51, 173–203, 2004.
- 25 Devassy, V. P., Bhattathiri, P. M. A., and Qasim, S. Z.: *Trichodesmium* phenomenon, *Ind. J. Mar. Sci.*, 7, 168–186, 1978.
- Devassy, V. P., Bhattathiri, P. M., and Qasim, S. Z.: Succession of organisms following *Trichodesmium* phenomenon, *Ind. J. Mar. Sci.*, 8, 89–93, 1979.
- Dore, J. E., Brum, J. R., Tupas, L. M., and Karl, D. M.: Seasonal and interannual variability in sources of nitrogen supporting export in the oligotrophic subtropical North Pacific Ocean, 30 *Limnol. Oceanogr.*, 47, 1595–1607, 2002.
- Falcón, L. I., Carpenter, E. J., Cipriano, F., Bergman, B., and Capone, D. G.: N₂ fixation by unicellular bacterioplankton from the Atlantic and Pacific oceans: phylogeny and in situ rates,

- Appl. Environ. Microbiol., 70, 765–770, 2004.
- Fu, F.-X, Zhang, Y., Leblanc, K., Sañudo-Wilhelmy, S. A., and Hutchins, D. A.: The biological and biogeochemical consequences of phosphate scavenging onto phytoplankton cell surfaces, *Limnol. Oceanogr.*, 50, 1459–1472, 2005.
- 5 Furnas, M. J. and Mitchell, A. W.: Pelagic primary production in the Coral and southern Solomon Seas, *Mar. Freshw. Res.*, 47, 395–705, 1996.
- Gallon, J. R.: Reconciling the incompatible: N_2 fixation and O_2 , *New Phytol.*, 122, 571–601, 1992.
- Gallon, J. R., Evans, A. M., Jones, D. A., Albertano, P., Congestri, R., Bergman, B., Gundersen, K., Orcutt, K. M., von Bröckel, K., Fritsche, P., Meyerhöfer, M., Nachtigall, K., Ohlendieck, U.,
10 te Lintel Hekkert, S., Sivonen, K., Repka, S., Stal, L. J., and Staal, M.: Maximum rates of N_2 fixation and primary production are out of phase in a developing cyanobacterial bloom in the Baltic Sea, *Limnol. Oceanogr.*, 47, 1514–1521, 2002.
- Gallon, J. R., Jones, D. A., and Page, T. S.: *Trichodesmium*, the paradoxical diazotroph, *Algalogical Studies*, 83, 215–243, 1996.
- 15 Galloway, J., Dentener, F. J., and Capone, D. G.: Nitrogen Cycles: Past, Present and Future, *Biogeochemistry*, 70, 153–226, 2004.
- Glibert, P. M. and Bronk, D. A.: Release of dissolved organic nitrogen by marine diazotrophic cyanobacteria, *Trichodesmium* spp., *Appl. Environm. Microbiol.*, 60, 3996–4000, 1994.
- 20 Glibert P. M. and O’Neil, J. M.: Dissolved organic nitrogen release and amino acid oxidase activity by *Trichodesmium* spp., *Bull. Inst. Océanogr. Monaco*, 19, 265–271, 1999.
- Gruber, N. and Sarmiento, J.: Global patterns of marine nitrogen fixation and denitrification, *Global Biogeochem. Cycles*, 11, 235–266, 1997.
- Hawser, S. P. and Codd, G. A.: The toxicity of *Trichodesmium* blooms from Caribbean waters, in: *Marine Pelagic Cyanobacteria: Trichodesmium and other Diazotrophs*, edited by: Carpenter, E. J., Capone, D. G., and Rueter, J. G., Kluwer Academic Publishers, 319–329, 1992.
- 25 Hewson, I., Govil, S. R., Capone, D. G., Carpenter, E. J., and Fuhrman, J. A.: Evidence of *Trichodesmium* viral lysis and potential significance for biogeochemical cycling in the oligotrophic ocean, *Aquatic Microbial Ecol.*, 36, 1–8, 2004.
- 30 Hood, R. R., Michaels, A. F., and Capone, D. G.: Answers sought to the enigma of marine nitrogen fixation, *EOS, Transactions, Am. Geophys. Union*, 81(133), 138–139, 2000.
- Hood, R. R., Bates, N. R., Capone, D. G., and Olson, D. B.: Modeling seasonal and interannual

BGD

3, 1049–1080, 2006

Fate of N_2 fixation

M. R. Mulholland

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

biogeochemical and N₂ fixation cycles at BATS, Deep-Sea Res. Part II, 48, 1609–1648, 2002.

Hood, R. R., Coles, V. J., and Capone, D. G.: Modeling the distribution of *Trichodesmium* and nitrogen fixation in the Atlantic Ocean, J. Geophys. Res., 109, C06006, doi:10.1029/2002JC001753, 2004.

Kana, T. M.: Oxygen cycling in cyanobacteria with specific reference to oxygen protection in *Trichodesmium* spp., in: Marine Pelagic Cyanobacteria: *Trichodesmium* and other Diazotrophs, edited by: Carpenter, E. J., Capone, D. G., and Rueter, J. G., Kluwer Academic Publishers, 29–41, 1992.

Kana, T. M.: Rapid oxygen cycling in *Trichodesmium thiebautii*, Limnol. Oceanogr., 38, 18–24, 1993.

Karl, D. M., Letelier, R., Hebel, D. V., Bird, D. F., and Winn, C. D.: *Trichodesmium* blooms and new nitrogen in the North Pacific Gyre, in: Marine Pelagic Cyanobacteria: *Trichodesmium* and other Diazotrophs, edited by: Carpenter, E. J., Capone, D. G., and Rueter, J. G., Kluwer Academic Publishers, 219–237, 1992.

Karl, D., Michaels, A., Bergman, B., Capone, D., Carpenter, E., Letelier, R., Lipschultz, F., Paerl, H., Sigman, D., and Stal, L.: Nitrogen fixation in the world's oceans, Biogeochemistry, 57/58, 47–98, 2002.

Karl, D. M., Letelier, R., Hebel, D., Tupas, L., Dore, J., Christian, J., and Winn, C.: Ecosystem changes in the North Pacific subtropical gyre attributed to the 1991–1992 El Nino, Nature, 373, 230–234, 1995.

Karl, D. M., Letelier, R., Tupas, L., Dore, J., Christian, J., and Hebel, D.: The role of nitrogen fixation in biogeochemical cycling in the subtropical North Pacific, Nature, 388, 533–538, 1997.

Kromkamp, J., de Bie, M., Goosen, N., Peen, J., Van Rijswijk, P., Sinke, J., and Duineveld, G. C. A.: Primary production by phytoplankton along the Kenyan coast during the SE monsoon and November intermonsoon 1992, and the occurrence of *Trichodesmium*, Deep-Sea Res. II, 44, 1195–1212, 1997.

LaRoche, J. and Breitbarth, E.: Importance of the diazotrophs as a source of new nitrogen in the ocean, Netherlands J. Sea Res., 53, 67–91, 2005.

Lenes, J. M., Darrow, B. P., Catrall, C., Heil, C. A., Callahan, M., Vargo, G. A., Byrne, R. H., Prospero, J. M., Bates, D. E., and Walsh, J. J.: Iron fertilization and the *Trichodesmium* response on the West Florida shelf, Limnol. Oceanogr., 46, 1261–1277, 2001.

BGD

3, 1049–1080, 2006

Fate of N₂ fixation

M. R. Mulholland

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

Fate of N₂ fixation

M. R. Mulholland

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Letelier, R. M. and Karl, D. M.: Role of *Trichodesmium* spp. in the productivity of the subtropical North Pacific Ocean, *Mar. Ecol., Prog. Ser.*, 133, 263–273, 1996.
- Li, W. K. W., Glover, H. E., and Morris, I.: Physiology of carbon photoassimilation by *Oscillatoria theibautii* in the Caribbean Sea, *Limnol. Oceanogr.*, 25, 447–456, 1980.
- 5 Lugomela, C., Lyimo, T. J., Bryceson, I., Semesi, A. K., and Bergman, B.: *Trichodesmium* in coastal waters of Tanzania: diversity, seasonality, nitrogen and carbon fixation, *Hydrobiologia*, 477, 1–13, 2002.
- Mague, T. H., Weare, N. M., and Holm-Hansen, O.: Nitrogen fixation in the North Pacific Ocean, *Mar. Biol.*, 24, 109–119, 1974.
- 10 Mague, T. H., Mague, F. C., and Holm-Hansen, O.: Physiology and chemical composition of nitrogen-fixing phytoplankton in the central North Pacific Ocean, *Mar. Biol.*, 41, 213–227, 1977.
- Mahaffey, C., Michaels, A., and Capone, D. G.: The conundrum of marine nitrogen fixation, *Am. J. Sci.*, 305, 546–595, 2005.
- 15 McCarthy, J. J. and Carpenter, E. J.: *Oscillatoria (Trichodesmium) theibautii* (Cynaophyta) in the central North Atlantic Ocean, *J. Phycol.*, 15, 75–82, 1979.
- Mills, M. M., Ridame, C., Davey, M., LaRoche, J., and Geider, R. J.: Iron and phosphorus co-limit nitrogen fixation in the eastern tropical North Atlantic, *Nature*, 429, 292–294, 2004.
- Montoya, J. P., Voss, M., Kähler, P., and Capone, D. G.: A simple, high precision tracer assay for dinitrogen fixation, *Appl. Environm. Microbiol.*, 62, 986–993, 1996.
- 20 Montoya, J. P., Carpenter, E. J., and Capone, D. G.: Nitrogen fixation and isotope abundance in zooplankton of the oligotrophic North Atlantic, *Limnol. Oceanogr.*, 47, 1617–1628, 2002.
- Montoya, J. P., Holl, C. M., Zehr, J. P., Hansen, A., Villareal, T. A., and Capone, D. G.: High rates of N₂ fixation by unicellular diazotrophs in the oligotrophic Pacific Ocean, *Nature*, 430, 1027–1032, 2004.
- 25 Mulholland, M. R. and Bernhardt, P. W.: The effect of growth rate, phosphorus concentration and temperature on N₂ fixation and N regeneration rates in continuous cultures of *Trichodesmium* IMS101, *Limnol. Oceanogr.*, 50, 839–849, 2005.
- Mulholland, M. R., Bernhardt, P. W., Heil, C. A., Bronk, D. A., and O’Neil, J. M.: Nitrogen fixation and regeneration in the Gulf of Mexico, *Limnol. Oceanogr.*, 51, 1762–1776, 2006.
- 30 Mulholland, M. R., Bronk, D. A., and Capone, D. G.: N₂ fixation and regeneration of NH₄⁺ and dissolved organic N by *Trichodesmium* IMS101, *Aquatic Microb. Ecol.*, 37, 85–94, 2004a.
- Mulholland, M. R. and Capone, D. G.: Nitrogen utilization and metabolism relative to patterns

- of N₂ fixation in populations of *Trichodesmium* from the North Atlantic Ocean and Caribbean Sea, *Mar. Ecol. Prog. Ser.*, 188, 33–49, 1999.
- Mulholland, M. R. and Capone, D. G.: The physiology of the marine N₂ fixing cyanobacteria *Trichodesmium*, *Trends in Plant Science*, 5, 148–153, 2000.
- 5 Mulholland, M. R. and Capone, D. G.: The stoichiometry of N and C utilization in cultured populations of *Trichodesmium* IMS101, *Limnol. Oceanogr.*, 46, 436–443, 2001.
- Mulholland, M. R., Glibert, P. M., Berg, G. M., Van Heukelem, L., Pantoja, S., and Lee, C.: Extracellular amino acid oxidation by phytoplankton and cyanobacteria: a cross-ecosystem comparison, *Aquat. Microb. Ecol.*, 15, 141–152, 1998.
- 10 Mulholland, M. R., Heil, C. A., Bronk, D. A., O’Neil, J. M., and Bernhardt, P. W.: Does nitrogen regeneration from the N₂ fixing cyanobacteria, *Trichodesmium* spp. fuel *Karenia brevis* blooms in the Gulf of Mexico?, in: *Harmful Algae 2002*, edited by: Steidinger, K. A., Lansberg, J. H., Tomas, C. R., and Vargo, G. A., Florida Fish and Wildlife Conservation Commission, Florida Institute of Oceanography, and Intergovernmental Oceanographic Commission of UNESCO, St. Petersburg, Florida, USA, 47–49, 2004b.
- 15 Mulholland, M. R., Ohki, K., and Capone, D. G.: Nitrogen utilization and metabolism relative to patterns of N₂ fixation cultures of *Trichodesmium* NIBB1067, *J. Phycol.*, 35, 977–988, 1999a.
- Mulholland, M. R., Shoemaker, C., Ohki, K., and Capone, D. G.: Utilization of combined forms of N in cultures and field populations of *Trichodesmium*, *Bull. Inst. Océanogr. Monaco*, 19, 279–286, 1999b.
- 20 Nausch, M.: Microbial activities on *Trichodesmium* colonies, *Mar. Ecol. Prog. Ser.*, 141, 173–181, 1996.
- Ohki, K.: A possible role of temperate phage in the regulation of *Trichodesmium*, *Bull. Inst. Oceanogr. Monaco*, 19, 235–256, 1999.
- 25 O’Neil, J. M.: The colonial cyanobacterium *Trichodesmium* as a physical and nutritional substrate for the harpacticoid copepod *Macrosetella gracilis*, *J. Plank. Res.*, 20, 43–59, 1998.
- O’Neil, J. M.: Grazer interactions with nitrogen-fixing marine Cyanobacteria: adaptation for N-acquisition?, *Bull. Inst. Oceanogr. Monaco*, 19, 293–317, 1999.
- 30 O’Neil, J. M., Metzler, P., and Glibert, P. M.: Ingestion of ¹⁵N₂-labelled *Trichodesmium*, and ammonium regeneration by the pelagic harpacticoid copepod *Macrosetella gracilis*, *Mar. Biol.*, 125, 89–96, 1996.
- O-Neil, J. M. and Roman, M. R.: Ingestion of the cyanobacterium *Trichodesmium* spp. by the

BGD

3, 1049–1080, 2006

Fate of N₂ fixationM. R. Mulholland

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

pelagic harpacticoid copepods *Macrosetella*, *Miracia* and *Oculosetella*, *Hydrobiologia*, 292–293, 235–240, 1994.

Orcutt, K. M., Lipschultz, F., Gundersen, K., Arimoto, R., Michaels, A. F., Knap, A. H., and Gallon, J. R.: A seasonal study of the significance of N₂ fixation by *Trichodesmium* spp. at the Bermuda Atlantic Time-series Study (BATS) site, *Deep-Sea Res. II*, 48, 1583–1608, 2001.

Paerl, H. W. and Bebout, B. M.: Direct measurement of O₂-depleted microzones in marine *Oscillatoria*: relation to N₂ fixation, *Science*, 241, 442–445, 1988.

Paerl, H. W., Bebout, B. M., and Prufert, L. E.: Bacterial associations with marine *Oscillatoria* sp. (*Trichodesmium* sp.) populations: ecophysiological implications, *J. Phycol.*, 25, 773–784, 1989.

Renaud, F., Pringault, O., and Rochelle-Newall, E.: Effects of the colonial cyanobacterium *Trichodesmium* spp. on bacterial activity, *Aquatic Microbial Ecol.*, 41, 261–270, 2005.

Renström-Kellner, E., Rai, A. N., and Bergman, B.: Glycolate metabolism in cyanobacteria II. Evidence for a mediated transport of glycolate in *Anabaena*, *Physiol. Plant.*, 75, 144–150, 1989.

Revelante, N., Williams, W. T., and Bunt, J. S.: Temporal and spatial distribution of diatoms, dinoflagellates and *Trichodesmium* in waters of the Great Barrier Reef, *J. Exp. Mar. Biol. Ecol.*, 63, 27–45, 1982.

Roman, M. R.: Ingestion of the blue-green alga *Trichodesmium* by the harpacticoid copepod *Macrosetella gracilis*, *Limnol. Oceanogr.*, 23, 1245–1248, 1978.

Romans, K. M., Carpenter, E. J., and Bergman, B.: Buoyancy regulation in the colonial diazotrophic cyanobacterium *Trichodesmium tenue*: ultrastructure and storage of carbohydrate, polyphosphate and nitrogen, *J. Phycol.*, 30, 935–942, 1994.

Saino, T.: Biological nitrogen fixation in the ocean with emphasis on the nitrogen fixing blue-green alga *Trichodesmium* and its significance in the nitrogen cycling in the low latitude sea areas, PhD thesis, Ocean Research Institute, University of Tokyo, 1977.

Saino, T. and Hattori, A.: Diel variation in nitrogen fixation by a marine blue-green alga, *Trichodesmium thiebautii*, *Deep-Sea Res.*, 25, 1259–1263, 1978.

Saino, T. and Hattori, A.: Nitrogen fixation by *Trichodesmium* and its significance in nitrogen cycling in the Kuroshio area and adjacent waters, in: *The Kuroshio IV*, edited by: Takenouti, Y., Saikon Publishing A, 697–709, 1980.

Sañudo-Wilhelmy, S. A., Kustka, A. B., Gobler, C. J., Hutchins, D. A., Yang, M., Lwiza, K., Burns,

BGD

3, 1049–1080, 2006

Fate of N₂ fixation

M. R. Mulholland

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

Fate of N₂ fixation

M. R. Mulholland

- J., Capone, D. G., Raven, J. A., and Carpenter, E. J.: Phosphorus limitation of nitrogen fixation by *Trichodesmium* in the central North Atlantic Ocean, *Nature*, 411, 66–69, 2001.
- Scranton, M. I.: The role of the cyanobacterium *Oscillatoria (Trichodesmium) thiebautii* in the marine hydrogen cycle, *Mar. Ecol. Prog. Ser.*, 11, 79–87, 1983.
- 5 Scranton, M. I.: Hydrogen cycling in the waters near Bermuda: the role of the nitrogen fixer, *Oscillatoria thiebautii*, *Deep-Sea Res.*, 31, 133–143, 1984.
- Scranton, M. I., Novelli, P. C., Michaels, A., Horrigan, S. G., and Carpenter, E. J.: Hydrogen production and nitrogen fixation by *Oscillatoria thiebautii* during in situ incubations, *Limnol. Oceanogr.*, 32, 998–1006, 1987.
- 10 Sellner, K. G.: Trophodynamics of marine cyanobacteria blooms, 75–94, in: *Marine Pelagic Cyanobacteria: Trichodesmium and other Diazotrophs*, edited by: Carpenter, E. J., Capone, D. G., and Rueter, J. G., Kluwer Academic Publisher, 1992.
- Sellner, K. G.: Physiology, ecology, and toxic properties of marine cyanobacteria blooms, *Limnol. Oceanogr.*, 42, 1089–1104, 1997.
- 15 Sheridan, C. C., Steinberg, D. K., and Kling, G. W.: The microbial and metazoan community associated with colonies of *Trichodesmium* spp.: a quantitative survey, *J. Plank. Res.*, 24, 913–922, 2002.
- Shimura, S., Yamaguchi, T., Aruga, Y., Fujita, Y., and Ichimura, S.: Extracellular release of photosynthetic products by a pelagic blue green alga *Trichodesmium thiebautii*, *J. Oceanogr. Soc. Japan*, 34, 181–188, 1978.
- 20 Siddiqui, P. J. A., Bergman, B., and Carpenter, E. J.: Filamentous cyanobacterial associates of the marine planktonic cyanobacterium *Trichodesmium*, *Phycologia*, 31, 326–337, 1992.
- Stal, L. J.: Tansley Review No. 84. Physiological Ecology of Cyanobacteria in Microbial Mats and Other Communities, *New Phytologist*, 131, 1–32, 1995.
- 25 Steinberg, D. K., Nelson, N. B., Carlson, C. A., and Prusak, A. C.: Production of chromophoric dissolved organic matter (CDOM) in the open ocean by zooplankton and the colonial cyanobacterium *Trichodesmium* spp., *Mar. Ecol. Prog. Ser.*, 267, 45–56, 2004.
- Subramaniam, A., Carpenter, E. J., Karentz, D., and Falkowski, P. G.: Bio-Optical Properties of the Marine Diazotrophic Cyanobacteria *Trichodesmium* spp. I. Absorption and Photosynthetic Action Spectra, *Limnol. Oceanogr.*, 44, 608–617, 1999.
- 30 Tseng, Y.-F., Lin, F.-J., Chiang, K.-P., Kao, S.-J., and Shiah, F.-K.: Potential impacts of the N₂-fixing *Trichodesmium* on heterotrophic bacterioplankton turnover rates and organic carbon transfer efficiency in the subtropical oligotrophic ocean system, *Terrestrial, Atmos. Ocean*

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Sci., 16, 361–376, 2005.

Villareal, T. A. and Carpenter, E. J.: Diel buoyancy regulation in the marine diazotrophic cyanobacterium *Trichodesmium thiebautii* Ehr, *Limnol. Oceanogr.*, 35, 1832–1837, 1990.

Voss, M., Croot, P., Lochte, K., Mills, M., and Peeken, I.: Patterns of nitrogen fixation along 10oN in the tropical Atlantic, *Geophys. Res. Lett.*, 31, L23S09, doi:10.1029/2004GL020127, 2004.

Walsby, A. E.: The gas vesicles and buoyancy of *Trichodesmium*, in: *Marine Pelagic Cyanobacteria: Trichodesmium and other Diazotrophs*, edited by: Carpenter, E. J., Capone, D. G., and Rueter, J. G., Kluwer Academic Publishers, 141–161, 1992.

Walsh, J. J. and Steidinger, K. A.: Saharan dust and red tides: the cyanophyte connection, *J. Geophys. Res.*, 106, 11 597–11 612, 2001.

Zehr, J. P., Waterbury, J. B., Turner, P. J., Montoya, J. P., Omoregie, E., Steward, G. F., Hansen, A., and Karl, D. M.: Unicellular cyanobacteria fix N₂ in the subtropical North Pacific Ocean, *Nature*, 412, 635–638, 2001.

BGD

3, 1049–1080, 2006

Fate of N₂ fixation

M. R. Mulholland

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

Table 1. Ranges of water column N₂ fixation rates. Rates are presented as hourly rates because it is unclear whether N₂ fixation by unicellular diazotrophs exhibit diel periodicity. For comparison, rates of N₂ fixation by *Trichodesmium* range from <0.1 to 21.4 nmol N col⁻¹ d⁻¹ (N₂ fixation is confined to the light period) and colony abundance can range from <1 to >1000 colonies L⁻¹ (Mulholland et al., 2006).

Date	Location	Depth	N ₂ fixation	Method	Reference
			(nmol L ⁻¹ h ⁻¹)		
2001–2003	Gulf of Mexico	Surface	0.011–0.23	¹⁵ N ₂	Mulholland et al. (2006)
2003	Gulf of Mexico	Pigment maximum	0.044–0.063	¹⁵ N ₂	Mulholland et al. (2006)
2002	New Caledonia	Surface	0.23–0.85	¹⁵ N ₂	Mulholland et al. (unpublished)
2002	N Atlantic	25 m (nighttime)	~0.147	AR	Falcon et al. (2004)
2001	N. Atlantic	Upper 100 m	0.025–0.045	¹⁵ N ₂	Falcon et al. (2004)
2002	N Pacific	Upper 100 m (nighttime)	~0.003	AR	Falcon et al. (2004)
2000–2001	Station ALOHA & Kaneohe Bay	25 m & Surface	0.01–0.15	¹⁵ N ₂	Montoya et al. (2004)
2002	Eastern N. Pacific	Mixed layer & pigment maximum	0.047–1.85(0.72)	¹⁵ N ₂	Montoya et al. (2004)
1999	Arafura Sea	Pigment maximum	20–62	¹⁵ N ₂	Montoya et al. (2004)
2000	Station ALOHA	25 m	0.010–0.016	¹⁵ N ₂	Zehr et al. (2001)
2002	Tropical Atlantic	Upper 100 m	up to 3.1	¹⁵ N ₂	Voss et al. (2004)
2000–2001	Station ALOHA	Upper 100 m	0–0.09	¹⁵ N ₂	Dore et al. (2002) ¹

¹Converted from daily rate assuming N₂ fixation persisted for 24 h per day.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Fate of N₂ fixation

M. R. Mulholland

Table 2. Results from paired comparisons of C₂H₂:N₂ measurements. Numbers are reported as molar ratios and N release is estimated as the observed molar ratio minus the theoretical ratio (3) divided by the observed molar ratio. Modified and expanded from Mulholland et al. (2006).

Location	C ₂ H ₂ :N ₂ (molar ratio)		Average N release %	Reference
	Range	Average		
<i>Trichodesmium</i> :				
Gulf of Mexico	3.3–15.8	7.3	52	Mulholland et al. (2006)
North Atlantic (latitudinal gradient) – Aug	3.1–7.5	4.6	29	Mulholland et al. (unpublished)
North Atlantic (latitudinal gradient) – Mar	6.3–52.7	28.3	79	Mulholland et al. (unpublished)
North Atlantic	0.9–7.3	3.6	17	Capone et al. (2005)
Sargasso Sea		6.0	50	Carpenter and McCarthy (1975)
Sargasso Sea		6.3	52	Carpenter and Price (1977)
Sargasso Sea		2.9		Scranton (1984)
Caribbean and Sargasso Seas		4.1	27	Scranton et al. (1987)
Caribbean Sea		3.4	12	Glibert and Bronk (1994)
BATS (net tows)		4.9	39	Orcutt et al. (2001)
BATS (SCUBA)		1.4		Orcutt et al. (2001)
BATS		3.0		Orcutt et al. (2001)
North Pacific		1.9		Mague et al. (1977)
Bay of Bengal & South China Sea	3–10	2.9		Saino (1977)
New Caledonia lagoon – S Pacific	4.8–19.5		21–97	Mulholland et al. (in prep.)
<i>Trichodesmium</i> IMS101 (batch)	1.7–9.8	5.6	46	Mulholland et al. (2004)
<i>Trichodesmium</i> IMS101 (continuous)	3.0–22.2	1.4	74	Mulholland and Bernhardt (2005)
<i>Trichodesmium</i> IMS101 (semi-continuous) – morning	4.9–14.9	8.9	59.7	Mulholland et al. (unpublished)
<i>Trichodesmium</i> IMS101 (semi-continuous) – afternoon	14.3–45.6	23.5	81.6	Mulholland et al. (unpublished)
<i>Trichodesmium</i> GBRTRL1101 – afternoon	8.8–24.9	16.7	76.5	Mulholland et al. (unpublished)
Other marine cyanobacteria:				
<i>Rhizosolenia/Richelia</i> association		9.3		Mague et al. (1974)
Mixed cyanobacteria – <i>Nodularia spumigena/Anabaena/Aphanizomenon</i>		3.8–20		Gallon et al. (2002)

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 3. Paired comparisons of C and N₂ fixation. Modified and expanded from Mulholland et al. (2006).

Location	C:N ₂ fixation Range	Average	Reference
<i>Trichodesmium</i> :			
Gulf of Mexico	5.4–42.7	13.1	Mulholland et al. (2006)
New Caledonia (lagoon) – morning	3.7–51.3		Mulholland et al. (unpublished)
North Atlantic (latitudinal gradient) – Aug	11.0–30.6	20.1	Mulholland et al. (unpublished)
North Atlantic (latitudinal gradient) – Mar	5.2–22.4	12.8	Mulholland et al. (unpublished)
North Pacific	1.2–2.1		Mague et al. (1977)
Sargasso Sea		16	Carpenter and Price (1977)
Sargasso Sea	1.5–87		McCarthy and Carpenter (1979)
BATS (puffs)	13–437	128	Orcutt et al. (2001)
BATS (tufts)	15–703	198	Orcutt et al. (2001)
N. Atlantic (May–June 1994)		47.1	Carpenter et al. (2004); Capone et al. (2005)
N. Atlantic (April 1996)		37.7	Carpenter et al. (2004); Capone et al. (2005)
N. Atlantic (October 1996)		24.6	Carpenter et al. (2004); Capone et al. (2005)
Indian Ocean (Tanzania)		20	Lugomela et al. (2002) ¹
<i>Trichodesmium</i> IMS101 (batch)	4.6–132.5	28	Mulholland and Capone (2001) ²
<i>Trichodesmium</i> IMS101 (batch)	6.5–15.2	9.6	Mulholland and Capone (2001) ³
<i>Trichodesmium</i> IMS101 (continuous)	13.4–20.0		Mulholland and Bernhardt (2005) ⁴
<i>Trichodesmium</i> IMS101 (semi-continuous) – morning	3.2–10.0	5.4	Hutchins et al. (unpublished)
<i>Trichodesmium</i> IMS101 (semi-continuous) – afternoon			Hutchins et al. (unpublished)
<i>Trichodesmium</i> GBRTLI101	2.0–12.4	6.5	Hutchins et al. (unpublished)
Other marine cyanobacteria:			
<i>Hemiaulus/Richelia</i> association		12.5	Carpenter et al. (1999) ⁵
Mixed cyanobacteria – <i>Nodularia</i>		17.6 (0–7 m)	Gallon et al. (2002)
<i>spumigena/Anabaena/Aphanizomenon</i>		5.1 (7–14 m)	
		1.5 (14–21 m)	

¹Using study averages and recalculating with a conversion factor of 3:1. ²Mid-day estimate during exponential growth. ³Cumulative estimate over a growth or diel cycle. Ratio increased during stationary phase growth. ⁴Lower at faster growth rates. ⁵Calculated using average N₂ fixation rate of 0.2 mg N m⁻³ h⁻¹ and average C fixation rate at bloom stations of 2.14 mg C m⁻³ h⁻¹.

Fate of N₂ fixation

M. R. Mulholland

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 1. Nitrogen and carbon cycling in the oligotrophic ocean with and without N₂ fixation. Panel (A) represents an ocean without N₂ fixation where new nitrate upwelled from below the euphotic zone balances export of material out of the euphotic zone. Panel (B) represents a low nutrient/high chlorophyll (HNLC) area fertilized with iron to stimulate drawdown and sequestration of atmospheric in the deep ocean. Panel (C) represents carbon drawdown from the atmosphere associated with N₂ fixation with alternative pathways whereby new production is respired in the euphotic zone resulting in no net drawdown or export of carbon from the euphotic zone (red) or drawdown and export of C from the atmosphere to the deep ocean. Redrawn and expanded from Hood et al. (2000).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 2. Pathways of trophic transfer promoting (a) sinking flux and (b) carbon flow through a microbial loop with no net carbon export.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)