



Transport and chemical transformations influenced by shallow cumulus over land

J. Vilà-Guerau de Arellano, S.-W. Kim, M. C. Barth, E. G. Patton

► To cite this version:

J. Vilà-Guerau de Arellano, S.-W. Kim, M. C. Barth, E. G. Patton. Transport and chemical transformations influenced by shallow cumulus over land. *Atmospheric Chemistry and Physics Discussions*, 2005, 5 (5), pp.8811-8849. hal-00303939

HAL Id: hal-00303939

<https://hal.science/hal-00303939>

Submitted on 18 Jun 2008

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Transport and chemical transformations influenced by shallow cumulus over land

J. Vilà-Guerau de Arellano¹, S.-W. Kim², M. C. Barth², and E. G. Patton²

¹Meteorology and Air Quality Section, Wageningen University, The Netherlands

²Mesoscale and Microscale Meteorology Division, National Center for Atmospheric Research, USA

Received: 5 August 2005 – Accepted: 24 August 2005 – Published: 16 September 2005

Correspondence to: J. Vilà-Guerau de Arellano (jordi.vila@wur.nl)

© 2005 Author(s). This work is licensed under a Creative Commons License.

J. Vilà-Guerau de Arellano et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Print Version](#)

[Interactive Discussion](#)

EGU

Abstract

ACPD

5, 8811–8849, 2005

The distribution and evolution of reactive species in a boundary layer characterized by the presence of shallow cumulus over land is studied by means of two large-eddy simulation models: the NCAR and WUR codes. The study focuses on two physical processes that can influence the chemistry: the enhancement of the vertical transport by the buoyant convection associated with cloud formation and the perturbation of the photolysis rates below, in and above the clouds. It is shown that the dilution of the reactant mixing ratio caused by the deepening of the atmospheric boundary layer is an important process and that it can decrease reactant mixing ratios by 10 to 50 percent compared to very similar conditions but with no cloud formation. Additionally, clouds transport chemical species to higher elevations in the boundary layer compared to the case with no clouds which influences the reactant mixing ratios of the nocturnal residual layers following the collapse of the daytime boundary layer. Estimates of the rate of reactant transport based on the calculation of the integrated flux divergence range from to -0.2 ppb hr^{-1} to -1 ppb hr^{-1} , indicating a net loss of sub-cloud layer air transported into the cloud layer. A comparison of this flux to a parameterized mass flux shows good agreement in mid-cloud, but at cloud base the parameterization underestimates the mass flux. Scattering of radiation by cloud drops perturbs photolysis rates. It is found that these perturbed photolysis rates substantially (10–40%) affect mixing ratios locally (spatially and temporally), but have little effect on mixing ratios averaged over space and time. We find that the ultraviolet radiance perturbation becomes more important for chemical transformations that react with a similar order time scale as the turbulent transport in clouds. Finally, the detailed intercomparison of the LES results shows very good agreement between the two codes when considering the evolution of the reactant mean, flux and (co-)variance vertical profiles.

Transport and transformation in shallow cumulus

J. Vilà-Guerau de Arellano et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Print Version](#)

[Interactive Discussion](#)

EGU

1. Introduction

Shallow cumulus (Cu) clouds over land are a manifestation of convection in the atmospheric boundary layer (ABL). Their formation and evolution depend strongly on the partitioning between the sensible- and latent-heat fluxes, the wind shear and the vertical structure of the thermodynamic state variables. Because the structure of the boundary layer with shallow cumulus differs from the clear boundary layer, the dynamics driven by the presence of Cu leads to variations of the boundary layer characteristics, which also determines the evolution and distribution of atmospheric reactive compounds. Shallow cumulus clouds generally form in the same meteorological situations that favor the accumulation of pollutants in the ABL. As a consequence, shallow Cu can affect the variability of chemical species in four ways: (a) the enhancement of vertical transport of the chemical species, (b) the turbulent mixing of atmospheric compounds, (c) the perturbation of ultraviolet radiation, and therefore of photodissociation rates, below, in and above the clouds and (d) the chemical reactions occurring within the cloud droplets. In this study, we focus on the first three of these effects.

Although shallow Cu clouds are a common feature in atmospheric boundary layers over land, little is known about their role on the transport, mixing, and photodissociation of chemical compounds. Shallow cumulus have the potential to enhance vertical transport of atmospheric compounds such that the compounds reach higher altitudes. The vertical transport is dependent on the strength of the clouds and on the temperature capping inversion. Mid-troposphere observations ([Ching and Alkezweeny, 1986](#); [Ching et al., 1988](#); [Baumann et al., 2000](#); [Angevine, 2005](#)) indicate that the presence of shallow cumuli leads to a reduction of the pollutants in the sub-cloud layer due to dilution.

In these cloudy boundary layers, highly turbulent regions are localized in the cloud updraft whereas the compensating subsidence regions are characterized by less turbulent downward motions. As a result, reactive species may become segregated such that turbulent mixing controls the reactivity of second-order chemical reactions ([Schu-](#)

Transport and transformation in shallow cumulus

J. Vilà-Guerau de Arellano et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Print Version](#)

[Interactive Discussion](#)

EGU

mann, 1989; Molemaker and Vilà-Guerau de Arellano, 1998). This limitation of reactivity is particularly important if the chemical time scales are of similar order to the turbulent transport time scale.

Finally, clouds disrupt the ultraviolet radiation field by perturbing the contributions of direct and diffuse radiation. In a seminal paper, Madronich (1987) investigated how cloud layers modify the vertical profiles of actinic flux, and therefore the photodissociation rates. Madronich (1987) showed that compared to the clear-sky photodissociation values, the perturbed photodissociation rates decrease in the sub-cloud layer, increase linearly in the cloud, and are enhanced above the cloud. In a previous numerical study on the reactions in a homogeneous stratocumulus cloud deck, Vilà-Guerau de Arellano and Cuijpers (2000) showed that the influence of the physical processes (turbulence and radiation) leads to an increase/decrease of the reactant's concentration gradient and consequently to a departure from photochemical equilibrium.

Our objective is to study the roles of transport and radiation in fair weather cumulus on reactive species. By means of a large-eddy simulation technique, we investigate the daytime evolution of reactive species in shallow Cu. We define a series of numerical experiments to (1) investigate the modification on the reactant distribution because of the enhancement of the boundary layer growth (defined by the cloud top height) by shallow Cu, (2) test the validity of current parameterizations of the vertical transport, (3) quantify the control of turbulent mixing on the species reactivity and (4) study how perturbed photodissociation rates by clouds affect the transformation of species. The study is based on the simulations carried out by the following two large-eddy simulation (LES) models: National Center for Atmospheric Research (NCAR) and Wageningen University Research (WUR). The intercomparison of the two codes allows us to assess the use of the LES technique to calculate diagnostic fields for turbulent reacting flows. In the absence of reliable and systematic observational field data, the LES method could provide a unique data base to study the influence of cloudy boundary layer dynamics on chemical transformations. To our knowledge, this is the first time that such an intercomparison study of reacting flows in the ABL has been carried out.

Transport and transformation in shallow cumulus

J. Vilà-Guerau de Arellano et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Print Version](#)

[Interactive Discussion](#)

EGU

2. Description of the case

Our study is based on the meteorological situation discussed by Brown et al. (2002). The same case has also been used to test the ability of single column cloud models to reproduce shallow cumulus over land (Lenderink et al., 2004). Below, we provide a brief description of the observations, the set up of the numerical experiment, and the validation of the diagnostic fields associated with the thermodynamic and reactant variables.

2.1. Observations

A cloudy boundary layer characterized by the presence of shallow cumulus was observed on 21 June 1997 at the Southern Great Plains (SGP) site of the Atmospheric Radiation Measurement (ARM) program in Oklahoma (Fig. 1). Based on the surface and on the upper air observations, Brown et al. (2002) proposed an idealized numerical experiment to study the dynamics of shallow cumulus convection over land by means of large-eddy simulations. The case was characterized by a strong diurnal cycle of both sensible and latent heat fluxes (Fig. 2), which were the driving forcing of the formation of clouds. The friction velocity, which was almost constant with time, had an average value of 0.5 m s^{-1} . The observed radiosonde vertical profiles of potential temperature showed first a stable boundary layer at 11:30 UTC (05:30 LT) which rapidly became a well mixed boundary layer by 14:30 UTC (08:30 LT). Measurements of the cloud fraction showed the presence of scattered clouds after 14:30 UTC with cloud cover ranging from 0.2 to 0.45. The cloud base height was around 1000 m after this time.

2.2. Numerical experiment

Two large-eddy simulation (LES) codes are used to study the transport and the transformation of reactants in a cloudy boundary layer. Both codes participated previously at the intercomparison exercise described in Brown et al. (2002). In the WUR-LES

Transport and transformation in shallow cumulus

J. Vilà-Guerau de Arellano et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Print Version](#)

[Interactive Discussion](#)

EGU

code, the filtered governing equations are solved using a finite-volume technique. All the resolved terms in the governing equations are discretized using straightforward second-order central differences, except for the advective terms that are discretized using the method of Piacsek and Williams (1970). The leap-frog scheme is used for the time integration of the momentum equations. The conservation equation of mass is obtained by solving a diagnostic equation for the pressure. For the advection and diffusion of temperature, moisture and reactants we use the limited $\kappa=1/3$ scheme (Koren, 1993) for the spatial discretization and a two-stage Runge-Kutta method for the time integration. For the time advancement of chemistry we apply the routine Twostep (Verwer, 1994). The entire numerical discretization for all the scalars satisfies the following three properties: it is conservative, monotone and positive. The subgrid-scale terms that appear in the filtered equations are parameterized as a function of the resolved local gradients and an exchange diffusivity coefficient which depends on the subgrid-scale kinetic energy and the characteristic grid size. The condensation scheme is based on the all-or-nothing assumption: cloud water exists when relative humidity is greater than 100%. The WUR code has been previously used to study the boundary layer dynamics of shallow cumulus (Cuijpers and Duynkerke, 1993; Siebesma and Cuijpers, 1995) and atmospheric turbulent reacting flows (Petersen et al., 1999; Vilà-Guerau de Arellano and Cuijpers, 2000).

The NCAR LES code has been described in Moeng (1984), Moeng (1986) and Sullivan et al. (1996). The code has recently been modified to run on massively parallel machines using the Message Passing Interface (MPI) and is described in Patton et al. (2005). The NCAR code uses a pseudospectral and a second-order finite difference method for horizontal and vertical advection, respectively. Vertical scalar advection is performed using a limited $\kappa=1/3$ scheme (Koren, 1993). Time integration proceeds using a third-order Runge-Kutta method (Spalart et al., 1991). An Euler backward iteration method (Barth et al., 2003) is used to solve the chemistry. Similar to the WUR LES, subgrid-scale terms are parameterized with subgrid-scale turbulent kinetic energy and stability dependent length scale. The condensation scheme is also based on the

Transport and transformation in shallow cumulus

J. Vilà-Guerau de Arellano et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Print Version](#)

[Interactive Discussion](#)

EGU

all-or-nothing assumption.

The initial conditions and surface forcing are the same as those prescribed by Brown et al. (2002). Figure 3 shows the initial profiles (11:30 UTC or 05:30 LT) of the potential temperature, specific humidity, and reactants A, B and C. Following the suggestion of Brown et al. (2002), the initial potential temperature profile has been slightly modified (see Fig. 10 in their paper) to simulate a cloud top height that rises faster than their standard case. The surface fluxes of moisture and potential temperature vary with time following the measurements presented in Fig. 2. The only other external forcing is a prescribed westerly geostrophic wind equal to 10 m s^{-1} . Small external tendencies representing the advection of heat and moisture are not included. Initial profiles of specific humidity for two sensitivity simulations are shown in Fig. 3d. The model domain size is $6400 \text{ m} \times 6400 \text{ m} \times 4400 \text{ m}$ with a horizontal grid length equal to 66.7 m and a vertical grid length equal to 40 m . The simulations are integrated from 11:30 UTC (05:30 LT) to 24:30 UTC (18:30 LT).

The chemical system is based on the atmospheric cycle of nitrogen dioxide, nitric oxide, and ozone, which are represented as species A, B and C. By using a simple chemical system, the influence of shallow cumulus on the transport of the species and on the photodissociation rates below, in and above the clouds can be clearly determined. The LES codes include the following two reactions:



where j and k are first- and second-order chemical reaction rates, respectively. The initial mixing ratio profiles of species A, B and C are shown in Fig. 3c. We also simulate an inert tracer whose initial profile is zero. The surface fluxes are constant with time using the following values: inert tracer (0.1 ppb m s^{-1}), reactant A (0.1 ppb m s^{-1}), reactant B ($0.05 \text{ ppb m s}^{-1}$) and reactant C (0.0 ppb m s^{-1}). By varying the values of j and k , one can study how the control of turbulence and the departure of chemical equilibrium influences the reactivity of the system. In the control simulation we use

Transport and transformation in shallow cumulus

J. Vilà-Guerau de Arellano et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Print Version](#)

[Interactive Discussion](#)

$$j = 8.3 \times 10^{-3} \text{ (s}^{-1}\text{)} \text{ and } k = 4.75 \times 10^{-4} \text{ (ppb s}^{-1}\text{).}$$

An important aspect of this study is to investigate the modification of the photolysis rates by the presence of clouds. Because clouds alter the different proportions of direct and diffuse ultraviolet radiation, the actinic flux (and therefore the photolysis rates) has different values below, in, and above the clouds (Madronich, 1987). We have implemented this effect by calculating at every time step a factor below and above the clouds and by applying this factor to the clear sky value of the photolysis rate j_{clear} following Chang et al. (1987):

$$j_{\text{clouds}} = F j_{\text{clear}} \quad (3)$$

Above the cloud, the factor (F) is defined as:

$$F = 1 + \alpha (1 - t_r) \cos(\chi_o). \quad (4)$$

While, below the cloud, F is defined as:

$$F = 1.6 t_r \cos(\chi_o). \quad (5)$$

Here, t_r is the energy transmission coefficient for normally incident light, χ_o is the solar zenith angle, and α is a reaction dependent coefficient (for nitrogen dioxide $\alpha=1.2$). To simplify the calculation, a linear interpolation is assumed inside the cloud. Based on measurements of j_{NO_2} (Fröhlich et al., 2000), the linear interpolation assumption likely overestimates the photolysis rate in the middle to lower regions of the cloud while underestimates the photolysis rate near cloud top.

The energy transmission coefficient t_r depends on the cloud optical depth and a scattering phase function asymmetry factor (f) equal to 0.86 for the typical cloud particle size ranges under study. The expression reads:

$$t_r = \frac{5 - e^{-\tau}}{[4 + 3\tau(1 - f)]}. \quad (6)$$

Transport and transformation in shallow cumulus

J. Vilà-Guerau de Arellano et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Print Version](#)

[Interactive Discussion](#)

The cloud optical depth (τ) is calculated according to the expression given by Stephens (1984)

$$\tau = \frac{3}{2} \frac{W}{\rho_l} r_e^{-1}, \quad (7)$$

where W is the vertically integrated liquid water (kg m^{-2}), ρ_l is the water density (kg m^{-3}) and r_e is the effective radius. Here, we have used a constant value of $r_e=10 \mu\text{m}$. For clouds characterized by values of $\tau < 5$ and for regions between clouds, we have assumed the photolysis rate of clear sky conditions. Regions between clouds may actually have enhanced photolysis rates due to cloud scattering. Here, we are simply investigating the importance of cloud scattering in the cloud column. Our control simulation includes the modification of photolysis rates due to cloud scattering, while a sensitivity simulation excludes this effect on the chemistry.

2.3. Model validation: dynamics

The simulated and observed profiles of the potential temperature and the total water specific humidity are shown in Fig. 4. Differences between observations and LES results are due to two reasons: (a) the LES simulations are based on an idealization of the initial conditions and of the surface boundary conditions, and (b) one-hour averaged LES results are compared with instantaneous soundings. The comparison between both LES results (Fig. 4) reveals small differences in the range of the inter-comparison study of Brown et al. (2002). Note that the aim of the validation with measurements is not to compare the exact values (as shown in Fig. 4 the vertical profiles of the radiosoundings at the site C1 and B also have large differences), but to evaluate the ability of the LES in reproducing the dynamics of a shallow cumulus convection situation and to capture the observed vertical profile evolution.

The evolution of the vertical profiles of the state variables is as follows. At 17:30 UTC (11:30 LT), at the onset of clouds, observations and LES results show three-distinct

Transport and transformation in shallow cumulus

J. Vilà-Guerau de Arellano et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Print Version](#)

[Interactive Discussion](#)

EGU

layers. In the sub-cloud layer (up to 1000 m), there is a well-mixed layer in both observations and LES results. Above this layer, the cloud layer is still not conditionally unstable. Above cloud top (at around 1300 m) there is an absolutely stable boundary layer. This three layer structure is maintained at 20:30 UTC (14:30 LT). At this stage 5 the clouds have matured. However, the cloud layer has deepened to ~3000 m and is conditionally unstable. The two LES codes predict similar boundary layer evolution.

The other cloud parameters (cloud cover, cloud base and cloud top heights, and liquid water path) evolve similarly to those reported by Brown et al. (2002) (Fig. 5). We find very good agreement of these cloud properties calculated by both LES codes. In 10 concluding, in view of the evolution of the vertical profiles of the state variables and their fluxes (not shown), we are confident of the ability of both LES codes to reproduce a daily evolution of shallow cumulus over land.

2.4. Model validation: chemistry

The evolution and the distribution of the reactants in this cloudy boundary layer are assessed by analyzing the mean vertical profiles, the fluxes, and the variances. Figure 15 6 shows the time evolution of the mean profiles of the inert scalar and the three reactants. The agreement between the results of both LES is very good. Both models show the growth of the boundary layer during the day and the enhancement of the transport in the cloud. Slight differences are found in the vertical profiles near the cloud top once 20 the clouds are mature (see the profiles at 20:30 and 23:30). The NCAR-LES simulates larger vertical transport in the sub-cloud layer and inside the cloud layer compared to WUR-LES due to a larger buoyancy flux in the sub-cloud layer.

The evolution of the vertical flux profile of the species is demonstrated in Fig. 7. At 25 14:30 UTC (08:30 LT), both model results show the importance of the entrainment flux of reactants in the initial period of the clear convective boundary layer growth. At 20:30 and 23:30 UTC (14:30 and 17:30 LT), larger fluxes occur inside the cloud layer due to the buoyant updraft motions driven by the release of latent heat. The flux enhancement of the species is also related to the vertical variation of the photodissociation rates per-

Transport and transformation in shallow cumulus

J. Vilà-Guerau de Arellano et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Print Version](#)

[Interactive Discussion](#)

turbed by the clouds (Eq. 3). Both model results show the enhancement of the flux of species *B* at 20:30 UTC, with a maximum value around $z=1600$ m. As shown in Fig. 3, species *B* has the lowest mixing ratio compared to *A* and *C*. Thus, the modification of the photolysis rate in the cloud will have a strong influence on the vertical distribution of species *B*. Since the clouds are mature at 20:30 UTC, the difference in the photolysis rate both in and below the cloud are larger. Larger gradients are created due to the different values of the first-order reaction rate, yielding to the increase of the flux.

Satisfactory agreement between both LES models is also found for the evolution of the vertical profiles of the variances (Fig. 8). Before cloud formation (14:30 UTC, 10 08:30 LT), the variances between the models differ significantly in the entrainment zone as a result of the different subgrid-scale models used in each LES code. The NCAR and WUR-LES results show that the largest concentration fluctuations occur at the interface between the sub-cloud and cloud layer. Because of the cloud-induced updrafts, the scalar variance becomes more vertically distributed within the cloud layer as the 15 clouds reach maximum development (around 20:30 UTC, 14:30 LT). In Sect. 4, we will discuss how the turbulence driven by the release of latent heat affects the mixing of reactants and their chemical reactivity.

3. Enhancement of the vertical transport

Large uncertainties in atmospheric chemistry models are associated with the crude 20 representation of boundary layer clouds in regional and large scale models (Dabberdt et al., 2004). In this section, we study the potential effects of cloud-induced changes on boundary layer structure, and consequently on the reactant concentrations, and evaluate a preliminary representation of the vertical transport in the clouds. Fair weather cumuli strongly modify the boundary layer structure by deepening its height, which in 25 turn transports the sub-cloud compounds to the cloud layer.

In order to investigate the impact of clouds, we conduct an LES experiment where shallow cumulus clouds do not form (i.e. clear sky conditions). In this experiment, we

Transport and transformation in shallow cumulus

J. Vilà-Guerau de Arellano et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Print Version](#)

[Interactive Discussion](#)

EGU

prescribe the same initial and boundary conditions as in the control simulation (see Sect. 2.2) except that we reduce the initial profile of the total water mixing ratio by 3 g kg^{-1} (see Fig. 3d). In order to have an independent criterion to estimate the maximum vertical transport of the reactants, we define a height (h) as a function of the mixing ratio of an inert tracer. h is defined as the height where the horizontally-averaged mixing ratio becomes 0.5% of the horizontally-averaged mixing ratio in the sub-cloud layer. This height indicates maximum elevation where the scalar and reactants are vertically transported.

The evolution of h in the cloudy control simulation (CL) and the clear sky boundary layer (CS) is shown in Fig. 9a. In the cloudy boundary layer, h closely follows the cloud top height, which is defined as the highest elevation where liquid water is found in the entire domain, although at early times (e.g., 19:00 UTC), h falls below the cloud-top height because only a limited number of clouds has formed. At later times (e.g., 23:00 UTC) h persists at the highest elevation attained by the clouds during the day even though the clouds have decayed. The shallow cumulus clouds act to transport scalars to elevated regions in the cloud layer that remain there following the clouds' dissipation. Consequently, the presence of clouds has an impact on the mixing ratio in the early stages of the formation of the nocturnal residual layer. In general, and depending on the strength of the thermal inversion above the cloud top, the reactants remain in the boundary layer and they are not transported to regions above the cloud layer in agreement with Cotton et al. (1995).

The difference in h between the CS and CL simulations reaches its maximum at around 20:00 UTC (14:00 LT) and the difference is approximately 600 m (Fig. 9a). This different evolution of the boundary layer growth has a direct impact on the scalar distribution and evolution. To illustrate this, we calculate the boundary-layer averaged inert scalar mixing ratio for the CS and CL cases (Fig. 9b), which we call A_{CS} and A_{CL} , respectively. Due to the larger PBL-volume in the cloudy layer, A_{CL} begins to decrease with the onset of clouds. When Cu are fully developed (after 20:00 UTC), the difference between A_{CS} and A_{CL} can reach 50%.

Transport and transformation in shallow cumulus

J. Vilà-Guerau de Arellano et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Print Version](#)

[Interactive Discussion](#)

EGU

Compared to the clear sky situation, clouds redistribute scalar species vertically (Fig. 10). Similar vertical profiles of carbon monoxide were shown by (Angevine, 2005) which indicate the enhanced transport of species due to the shallow cumulus presence. In Fig. 10, before the onset of the clouds, the *CL* and *CS* scalar profiles are similar. Following cloud development, the compounds are mixed through a larger vertical depth such that sub-cloud mixing ratios are reduced in the cloudy boundary layer. For instance, for reactant *A* in the *CL* case, the mixing ratio is reduced at 500 m by a factor 0.5%, 7% and 12% at, respectively, 17:30, 20:30, 23:30 UTC compared to those in *CS*.

Clouds tend to enhance the volume of the cloud layer and consequently dilute the concentration of boundary layer compounds. It is therefore very important to accurately describe the development of shallow cumulus and their influence on the boundary layer dynamics in large-scale atmospheric chemistry models. In addition, small variations in specific humidity profiles (as in our study) dramatically influence cloud formation and intensity. Therefore, it is also essential to accurately determine the state variables and other related boundary conditions (soil moisture, temperature lapse rate) (Ek and Holstlag, 2004) to determine if clouds will form in the ABL.

To emphasize the impact of vertical scalar transport by shallow cumulus, we now investigate conditionally-averaged fields. Here, we define within-cloud conditionally-averaged fields by spatially averaging the compounds over the grid points containing liquid water (liquid water content greater than zero). Figure 11 presents one-hour averaged within-cloud field along with the horizontally averaged field over the whole domain for species *A* at 20:30 UTC (14:30 LT). Two simulations are shown: (a) using photolysis rate perturbed by clouds and (b) using the photolysis rate under clear sky conditions. The comparison of these two simulations will be discussed later in Sect. 5. Note that the results of the two LES models agree very well. As expected, the results indicate that the updrafts associated with the buoyant cumulus clouds are the main transport mechanism.

To test current parameterizations of cumulus convection used in large models, we

Transport and transformation in shallow cumulus

J. Vilà-Guerau de Arellano et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Print Version](#)

[Interactive Discussion](#)

EGU

compare horizontally-averaged vertical scalar fluxes from the LES results to those estimated by the mass-flux parameterization (Betts, 1973; Arakawa and Schubert, 1974; Cotton et al., 1995). Briefly, in mass-flux schemes, the ABL is represented by strong updrafts inside the clouds (c) surrounded by the environment downdrafts (e). In this approach, the average values of a generic scalar ψ is calculated for both updrafts and downdrafts. The average value of the entire layer is decomposed as follows:

$$\bar{\psi} = a\psi_c + (1 - a)\psi_e, \quad (8)$$

where a denotes the cloud cover. The flux (F_ψ) of the scalar ψ can be written according to:

$$F_\psi = \rho w' \bar{\psi}' \approx M(\psi_c - \bar{\psi}), \quad (9)$$

where M is by definition the mass flux which reads:

$$M \equiv \rho a_c w_c. \quad (10)$$

Here, a_c and w_c denote the cloud cover and vertical velocity calculated by the previously described conditional averaging method, respectively. Equation (9) is an approximation of the total flux and takes into account only the organized turbulent transport in the cloud and the environment (see Siebesma and Cuijpers, 1995).

The results of the mass-flux approach for reactant A are plotted in Fig. 12. We found very consistent results in the comparison of the total flux and the mass-flux approximation calculated by the NCAR-LES and the WUR-LES. There is an underestimation by a factor 1.5 at cloud base whereas in the middle of the cloud the mass-flux approach represents the scalar flux quite well. Near cloud top, the mass-flux approach overestimates by a factor 1.5 compared to the LES-calculated total flux. These results are consistent with the previous study carried out by Siebesma and Cuijpers (1995) where they evaluate the mass-flux scheme for heat and total water. A key aspect that still requires further research is the exchange flux between the sub-cloud layer and the cloud layer. As shown in Fig. 12 this flux is underestimated by the mass-flux approximation. For chemically reactive species, the exchange flux is also modified by the

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

cloud-imposed photolysis rate dependence (see the flux profile at Fig. 7c). Greenhut (1986) quantified the transport of ozone by cumulus clouds from aircraft observations. They found that cloud turbulence accounts for ~30% of the total cloud flux, whereas our LES results show that the organized turbulence in the clouds contributes more than

5 50% of the total cloud flux.

We also examine the species transport rate per unit time by integrating the flux divergence term included in the reactants' conservation equation over height. The integral reads:

$$\frac{1}{z_{ct} - z_{cb}} \int_{z_{cb}}^{z_{ct}} \frac{\partial \overline{w' c'_i}}{\partial z} \delta z, \quad (11)$$

10 where z_{cb} and z_{ct} denote the cloud base and cloud top height, respectively, and $\overline{w' c'_i}$ is the total turbulent flux calculated by the model. This rate gives us an indication of the buoyancy driven transport through cloud. For species A in the control case, we obtain values ranging from 1 ppb hr⁻¹ at cloud onset to 0.2 ppb hr⁻¹ once the cloud has fully developed. Calculation of this rate for species A from the simulation with
15 greater cloud cover (≤ 0.5 ; the simulation depicted by the dotted line in the initial profile of the total specific humidity, Fig. 3d) results in similar rates. This fact indicates that shallow cumulus are more vigorous and just as effective at transporting species when the sky is not completely cloud covered compared to when clouds cover most of the sky. Vukovich and Ching (1990) estimated similar rates of species transport, but their
20 calculations revealed greater cloud cover dependency.

Currently, for state variables like the liquid water potential temperature and the total water mixing ratio, the exchange flux is parameterized by combining a turbulent closure scheme in the sub-cloud layer (i.e., a function of an exchange coefficient and the gradient of the variable) with the mass-flux closure in the cloud layer. As discussed thoroughly by Lenderink et al. (2005), there are still large uncertainties and discrepancies in the representation of these fluxes in large scale models. Based on our LES results, we conclude that schemes which describe shallow cumulus should first accu-

Transport and transformation in shallow cumulus

J. Vilà-Guerau de Arellano et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Print Version](#)

[Interactive Discussion](#)

rately estimate the evolution of the boundary layer depth and second to describe the enhanced vertical transport of reactants. The dilution and the mixing ratio increase in the cloud layer can influence nocturnal chemistry in the residual layer once the convective boundary layer has collapsed.

5 4. Control of the reactivity by turbulent mixing

Organized turbulence can segregate reactants by transporting them in opposite directions (see for instance the initial profiles of reactants B and C in Fig. 3) and can therefore limit their chemical reactivity: reactants must be mixed before the reaction takes place. In particular, turbulent control of reactivity becomes important if the turbulent mixing time scale (τ_t) is similar to the time scale of chemical reaction (τ_c), i.e. when the Damköhler number ($Da = \tau_t / \tau_c$) is $\mathcal{O}(1)$. The turbulent time scale is calculated by the expression w_*/z_i , where w_* is the characteristic convective velocity scale and z_i is the boundary layer height. In our calculations, w_* is determined following Frisch et al. (1995) and z_i is found by the local gradient method (Sullivan et al., 1998). The time scale of chemistry is calculated as $(j)^{-1}$ for a first-order reaction and as $(kC)^{-1}$ for a second-order reaction, where C is the mixing ratio of the reactant.

In dry convective boundary layers (Schumann, 1989; Molemaker and Vilà-Guerau de Arellano, 1998), this control is quantified by defining the intensity of segregation $/s$ which is the ratio of the covariance between reactants normalized by the ensemble horizontal mixing ratio average. This dimensionless variable reads:

$$20 \quad /s = \frac{\overline{c'_i c'_j}}{\overline{c_i} \overline{c_j}}. \quad (12)$$

For second-order chemical reactions, the covariance between reactants appears explicitly in their governing equation. Therefore, $/s$ can be a measure the concentration fluctuation correlation's importance in determining their ensemble means.

Transport and transformation in shallow cumulus

J. Vilà-Guerau de Arellano et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Print Version](#)

[Interactive Discussion](#)

EGU

In Fig. 13, we show vertical profiles of $/s$ between species B and C and the liquid water content. The results of both LESs show the close relation between the vertical cloud development and the $/s$ -evolution. Two distinct turbulent layers are found under the presence of Cu (see for instance Fig. 6c in Brown et al., 2002). In the sub-cloud layer, even though the variances of reactive species are large (Fig. 8), the covariance (or $/s$) between species B and C is almost negligible. This implies that reactants B and C are well mixed before reaction occurs. However, the cloud layer resembles a two-stream layer where species transported in opposite directions could remain segregated.

The maximum values of $/s$ range from 5% at the early stages of the cloud formation (around 17:30 UTC, or 11:30 LT) to 3% when clouds are mature (around 20:30 UTC, or 14:30 LT). The absolute values of these $/s$ -profiles are rather low compared to ones in the previous studies concerning dry convective boundary layers. Because w_* and z_i change during the simulations, the turbulent timescale and therefore the Damköhler number have a range of values, $\tau_t=10\text{--}40$ min and $Da=4\text{--}20$. Thus, our chemical system is characterized by a $Da>1$, which indicates that the chemistry is much faster than the turbulent mixing. One may expect higher absolute values of $/s$ in more complex chemical systems where Damköhler numbers of some reactions are $\mathcal{O}(1)$ and in situations where the chemistry departs from the photochemical equilibrium state.

20 5. Perturbation of the photolysis rate by the clouds

To study the effect of clouds on the photolysis rate, we carry out two simulations using the two LES codes. The first one, the control simulation discussed in the previous sections, takes into account the perturbation of the j -values by clouds using Eq. (3) (j_{cloudy}). The second simulation assumes a constant photolysis rate with height and time and therefore the photodissociation rates are not perturbed by the Cu (j_{clear}). Our aim is to: 1) analyze the modification of the reactant distribution resulting from the cloud radiation perturbation, and 2) investigate the dependence of this modification

Transport and transformation in shallow cumulus

J. Vilà-Guerau de Arellano et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Print Version](#)

[Interactive Discussion](#)

EGU

to spatial and temporal averaging. In Fig. 14, we show the following variables: 1) an instantaneous cross section of the liquid water content, and 2) the mixing ratio difference of reactant A calculated using j_{cloudy} minus that calculated using j_{clear} . The vertical cross section corresponds to the time when the shallow cumulus were fully developed (around 20:30 UTC, 14:30 LT). In the figure, we have also included the sensitivity of the results to the Damköhler number by repeating the same simulation using the same initial and boundary conditions, but changing the chemical reaction rate. The following first- and second-reaction rates are prescribed:

| Figure | Da | $j(\text{s}^{-1})$ | $k(\text{ppb s}^{-1})$ |
|---------------|------------|----------------------|------------------------|
| 14a | 0.4 – 2.0 | 8.3×10^{-4} | 4.75×10^{-5} |
| 14b (Control) | 4.0 – 20. | 8.3×10^{-3} | 4.75×10^{-4} |
| 14c | 40. – 200. | 8.3×10^{-2} | 4.75×10^{-3} |

In Fig. 14, the three cases reveal similar spatial patterns with regards to the cloud effect on the photodissociation rates. In the lower portion of the cloud and in the sub-cloud layer, we find larger values of the mixing ratio calculated by using j_{cloudy} . In this region where $j_{\text{cloudy}} < j_{\text{clear}}$, clouds decrease the reactivity resulting in $A_{\text{cl}} > A_{\text{cs}}$. In the upper part of the cloud, the opposite effect is found; due to the enhancement of the actinic flux ($j_{\text{cloudy}} > j_{\text{clear}}$) $A_{\text{cl}} < A_{\text{cs}}$. As shown, the order of magnitude of this effect is a function of the Damköhler number. The case where $Da=0.4\text{--}2.0$ (Fig. 14a), shows differences of about $\pm 10\%$ at cloud top and cloud base compared to the case using a constant j -value. When $Da=40\text{--}200$, this difference increases considerably (Figs. 14b and c): 40% in the sub-cloud layer and -20% in the upper cloud region. This sensitivity reveals a coupling between chemistry driven by ultraviolet radiation and the turbulent transport under the presence of the clouds.

In reactive turbulent flows characterized by $Da < 1$, the turbulent transport time scale is faster than the chemical reactivity scale, implying that turbulence can efficiently transport and mix the species before chemical reactions occur. As a result, the mixing ratio gradients tend to decrease by turbulent mixing. However, in flows with $Da > 1$, because

Transport and transformation in shallow cumulus

J. Vilà-Guerau de Arellano et al.

[Title Page](#)

| | |
|-----------------------------|------------------------------|
| Abstract | Introduction |
| Conclusions | References |
| Tables | Figures |

| | |
|-----------------------------------|-----------------------|
| ◀ | ▶ |
| ◀ | ▶ |
| Back | Close |
| Full Screen / Esc | |

| |
|--|
| Print Version |
| Interactive Discussion |

EGU

the chemical reactivity is faster than the mixing, the reactivity maintains scalar gradients which increase the effect of the perturbed photolysis rates. It is important to note that the Damköhler number can vary spatially in cloudy boundary layers. Because the buoyant convection enhances vertical velocities in the cloud cores, the convective velocity scale becomes spatially heterogeneous inducing a large variability in Da . As a consequence, effects of the modified photolysis rates can become larger in certain regions depending on the local value of Da .

By doing spatial and temporal averaging, we can further study the impact of different photolysis rates (j_{cloudy} and j_{clear}) on the reactant distribution. In Fig. 15, we plot the differences of the mixing ratio for species A calculated using j_{cloudy} (A_{cl}) and j_{clear} (A_{cs}) from the surface to cloud top (30-min average). Except for the two peaks at around 19:00 and 20:00 UTC calculated by NCAR-LES results, the two LES codes give variations of around 2%. This value contrasts with the larger values of ~40% reported in Fig. 14. Thus, the photolysis rate disturbed by clouds has a large impact on the instantaneous mixing ratios locally, but its impact is smoothed by averaging in time and space. Since our chemical system is close to an equilibrium, the impact of local variations becomes minimal by averaging.

To complete this study, we calculate the conditionally averaged mixing ratios computed using j_{cloudy} and j_{clear} (presented in Fig. 9). The one-hour average vertical profiles show different values for the profiles A_{cl} and A_{cs} . The results from the two LESs show very similar structure: large depletion of species A near cloud base by the simulation using j_{clear} and larger depletion of species A in the upper cloud region by the simulation using j_{cloudy} . The differences are higher than the ones shown in Fig. 15. Near cloud base, they range from 4% to 12% and near cloud top (around 2500 m) are –15%.

In view of the different results shown in Figs. 9, 14 and 15, it is very difficult to provide a conclusive statement as to the effect of modified photodissociation in clouds on the reactant distribution. Locally and at short time scales, the perturbation can modify the chemistry by enhancing/diminishing the concentration fields and by perturbing the establishment of possible chemical equilibrium. However, this perturbation by clouds

Transport and transformation in shallow cumulus

J. Vilà-Guerau de Arellano et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Print Version](#)

[Interactive Discussion](#)

EGU

6. Conclusions

The effects of shallow cumulus dynamics and radiation on three chemical species are studied by analyzing and comparing results from two large-eddy simulation codes:

5 NCAR and WUR. The chemical system is composed by a first- and second-order reaction near a chemical equilibrium. Both LES codes satisfactorily reproduce the diurnal evolution of shallow cumulus over land based on a situation observed on 21 June 1997. With respect to the diagnostic fields of the three reactants, we find very good agreement in the evolution of the mean, fluxes, and (co-)variances. The presence of clouds
10 increases the boundary layer height. Consequently, reactants are more diluted than in cloudless situations because the species are distributed through a greater vertical depth. Therefore, although cloud venting to the free troposphere played a minor role in the studied situation, the dilution of the mixing ratio due to the deepening of the boundary layer is a process that has to be taken into account in large-scale models if shallow
15 cumulus form. We found that the presence of clouds could lead to a decrease of 50% of the reactant mixing ratio compared to the clear sky situation. This decrease depends on the cloud cover evolution and values. We also note that reactants transported to elevated regions remain at those levels following the clouds' dissipation, which should have an effect on simulating nocturnal chemistry of the residual layer.

20 Analysis of the enhancement of the vertical transport reveals that the buoyant upward motions are responsible for transporting the reactants in the cloud. Current parameterizations based on the mass-flux approach adequately describe the transport in mid-cloud, but they tend to underestimate the flux at cloud base by a factor 2, showing the need to improve the parameterization of reactant exchange between the sub-cloud
25 and cloud layers. The variability of the species quantified by the mixing ratio variance profile is closely associated with the cloud development and is characterized by two maxima, one in the sub-cloud layer and the other within the cloud layer. Our analysis

Transport and transformation in shallow cumulus

J. Vilà-Guerau de Arellano et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Print Version](#)

[Interactive Discussion](#)

EGU

also shows that species are uniformly mixed in the sub-cloud layer, but that in the cloud layer reactants transported in opposite direction are somewhat segregated and therefore the turbulence might control their reactivity. The control is expected to be larger when the time scale of turbulence is similar to the time scale of chemistry, namely when the Damköhler number is $\mathcal{O}(1)$. We found that the reaction rate is reduced between 2% and 5% at different stages of the cloud development. We note that these absolute values can increase if the reactants depart from photochemical equilibrium.

The perturbation of photodissociation rates due to the presence of clouds is discussed comparing a simulation that uses photodissociation rate perturbed by the cloud with one that uses clear sky values. By analyzing the instantaneous fields of the mixing ratio, we find differences up to a 40% below the cloud base height and up to 20% close to cloud top. These differences are dependent on the capacity of the vertical transport to smooth out the gradients created by the perturbed photodissociation rates, i.e. depending on the Damköhler number. For turbulent reacting flows characterized by values $Da \sim \mathcal{O}(1)$ and larger, we find that turbulence decreases mixing ratio gradients created by the different photodissociation rates below, in and above the cloud. These differences are smaller when averaged over space and time. Further studies will attempt to investigate and to complete these findings in more complex chemical mechanisms and to explore the effect of cloud penetration into the free troposphere on the distribution of reactants.

Acknowledgements. J. Vilà was partially supported for his visit at NCAR by the Netherlands Organization for Scientific Research NWO (project 76/241). The WUR-LES numerical simulations were financially supported by NCF (SG-132). We appreciate the fruitful discussions with C.-H. Moeng. The National Center for Atmospheric Research is operated by the University Corporation for Atmospheric Research under the sponsorship of the National Science Foundation.

Transport and transformation in shallow cumulus

J. Vilà-Guerau de Arellano et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Print Version](#)

[Interactive Discussion](#)

EGU

- Angevine, W. M.: An integrated turbulence scheme for boundary layers with shallow cumulus applied to pollutant transport, *J. Appl. Meteorol.*, in press, 2005. [8813](#), [8823](#)
- Arakawa, A. and Schubert, W. H.: Interaction of a cumulus cloud ensemble with the large scale environment, *J. Atmos. Sci.*, 31, 674–701, 1974. [8824](#)
- Barth, M. C., Sillman, S., Hudman, R., Jacobson, M. Z., Kim, C.-H., Monod, A., and Liang, J.: Summary of the cloud chemistry modeling intercomparison: Photochemical box model simulation, *J. Geophys. Res.*, 108, 4214, doi:10.1029/2002JD002673, 2003. [8816](#)
- Baumann, K., Willimas, E. J., Angevine, W. M., Roberts, J. M., Norton, R. B., Frost, G. J., Fehsenfeld, F. C., Springston, S. R., Bertman, S. B., and Hartsell, B.: Ozone production and transport in Nashville, Tennessee: results from the 1994 study at New Hendersonville, *J. Geophys. Res.*, 105, 9137–9153, 2000. [8813](#)
- Betts, A. K.: Non precipitating cumulus convection and its parameterization, *Quart. J. Roy. Meteorol. Soc.*, 99, 178–196, 1973. [8824](#)
- Brown, A. R., Cederwall, R. T., Chlond, A., Duynkerke, P. G., Golaz, J., Khairoutdinov, J. M., Lewellen, D., Lock, A., Macveen, M. K., Moeng, C., Neggers, R. A. J., Siebesma, A. P., and Stevens, B.: Large-eddy simulation of the diurnal cycle of shallow cumulus convection over land, *Quart. J. Roy. Meteorol. Soc.*, 128(B), 1075–1094, 2002. [8815](#), [8817](#), [8819](#), [8820](#), [8827](#)
- Chang, J., Brost, R., Isaksen, I., Madronich, S., Middleton, P., Stockwell, W., and Walcek, C. J.: A three-dimensional Eulerian acid deposition model: physical concepts and formulation, *J. Geophys. Res.*, 92, 14 681–14 700, 1987. [8818](#)
- Ching, J. K. S. and Alkezweeny, A. J.: Trace study of vertical exchange by cumulus clouds, *J. Clim. Appl. Meteorol.*, 25, 1702–1711, 1986. [8813](#)
- Ching, J. K. S., Shipley, S. T., and Browell, E. V.: Evidence for cloud venting of mixed layer ozone and aerosols, *Atmos. Environ.*, 22, 225–242, 1988. [8813](#)
- Cotton, W. R., Alexander, G., Hertenstein, R., Walko, R., McAnelly, R., and Nicholls, M.: Cloud venting- a review and some new global annual estimates, *Earth Sci. Rev.*, 39, 169–206, 1995. [8822](#), [8824](#)
- Cuijpers, J. W. M. and Duynkerke, P. G.: Large eddy simulations of trade wind with cumulus clouds, *J. Atmos. Sci.*, 50, 3894–3908, 1993. [8816](#)
- Dabberdt, W. F., Carroll, M. A., Baumgardner, D., Carmichael, G., Cohen, R., Dye, T., Ellis,

5, 8811–8849, 2005

Transport and transformation in shallow cumulus

J. Vilà-Guerau de Arellano et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

EGU

- J., Grell, G., Grimmond, S., Hanna, S., Irwin, J., Lamb, B., Madronich, S., McQueen, J., Meagher, J., Odman, T., Pleim, J., Schmidt, H. P., and Westphal, D. L.: Meteorological research needs for improved air quality forecasting, Bull. Amer. Meteorol. Soc., 85, 563–586, 2004. [8821](#)
- 5 Ek, M. and Holstlag, A. A. M.: Influence of soil moisture on boundary layer development, J. Hydromet., 5, 86–99, 2004. [8823](#)
- Frisch, A. S., Lenschow, D. H., Fairall, C. W., Schubert, W. H., and Gibson, J. S.: Doppler radar measurements of turbulence in marine stratiform cloud during ASTEX, J. Atmos. Sci., 52, 2800–2808, 1995. [8826](#)
- 10 Fröhlich, B., Trautmann, T., Wendisch, M., and Keil, A.: Comparison of observed and simulated NO₂ photodissociation frequencies in a cloudless atmosphere and in continental boundary layer clouds, J. Geophys. Res., 105, 9843–9857, 2000. [8818](#)
- Greenhut, G.: Transport of ozone between boundary layer and cloud layer by cumulus clouds, J. Geophys. Res., 91, 8613–8622, 1986. [8825](#)
- 15 Koren, B.: A robust upwind discretization method for advection, diffusion and source terms, in: Notes on Numerical Fluid Mechanics, edited by: Vrengdenhil, C. B. and Koren, B., vol. 45, chap. 5, pp. 117–138, Vieweg-Braunschweig, 1993. [8816](#)
- Lenderink, G., Siebesma, A. P., Cheinet, S., Irons, S., Jones, C. G., Marquet, P., Müller, F., Olmeda, D., Calvo, J., Sanchez, E., and Soares, P. M. M.: The diurnal cycle of shallow cumulus over land: a single column model intercomparison study, Quart. J. Roy. Meteorol. Soc., 130, 3323–3338, 2004. [8815](#)
- 20 Lenderink, G., Siebesma, A. P., Cheinet, S., Irons, S., Jones, C. G., Marquet, P., Müller, F., Olmeda, D., Calvo, J., Sanchez, E., and Soares, P. M. M.: The diurnal cycle of shallow cumulus clouds over land: a single column model intercomparison study, Quart. J. Roy. Meteorol. Soc., 123, 223–242, 2005. [8825](#)
- 25 Madronich, S.: Photodissociation in the atmosphere: 1. Actinic fluxes and the effects of ground reflections and clouds, J. Geophys. Res., 92, 9740–9752, 1987. [8814](#), [8818](#)
- Moeng, C.-H.: A large-eddy simulation model for the study of planetary boundary-layer turbulence, J. Atmos. Sci., 41, 2052–2062, 1984. [8816](#)
- 30 Moeng, C. H.: Large-eddy simulation of a stratus-topped boundary layer: Part I. Structure and budgets, J. Atmos. Sci., 43, 2886–2900, 1986. [8816](#)
- Molemaker, M. J. and Vilà-Guerau de Arellano, J.: Turbulent control of chemical reactions in the convective boundary layer, J. Atmos. Sci., 55, 568–579, 1998. [8814](#), [8826](#)

Transport and transformation in shallow cumulus

J. Vilà-Guerau de Arellano et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Print Version](#)

[Interactive Discussion](#)

EGU

Transport and transformation in shallow cumulus

J. Vilà-Guerau de Arellano et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Print Version](#)

[Interactive Discussion](#)

EGU




Fig. 1. GOES-8 satellite visible image at 17:45 UTC (11:45 LT) 21 June 1997. The image shows shallow cumulus fields above the radiosonde sites C1 and B5. The locations of the central facility of the Southern Great Plains (SGP) site C1 (36.605° N, 97.485° W, elevation 315 m) of the Atmospheric Radiation Measurement (ARM) program and the radiosounding B5 (35.969° N, 95.856° W, elevation 345 m) are shown.

Transport and transformation in shallow cumulus

J. Vilà-Guerau de Arellano et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Print Version](#)

[Interactive Discussion](#)

Transport and transformation in shallow cumulus

J. Vilà-Guerau de Arellano et al.




Fig. 2. Time evolution of the sensible heat flux (solid line) and the latent heat flux (dashed line).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)




Fig. 3. Initial profiles of: **(a)** potential temperature, **(b)** total water mixing ratio and **(c)** reactant mixing ratio (ppb) and **(d)** the initial profile of the total mixing ratio to simulate a clear boundary layer (dashed line) and a more cloudy boundary layer than the control simulation (dotted line). Notice the different vertical axis in panel (c).

[Print Version](#)[Interactive Discussion](#)[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)

Transport and transformation in shallow cumulus

J. Vilà-Guerau de Arellano et al.




Fig. 4. Time evolution (at 17:30 UTC (11:30 LT) and 20:30 UTC (14:30 LT)) of vertical profiles of **(a)** potential temperature and **(b)** total-water mixing ratio. The thick solid lines show the one-hour averaged profiles obtained by the WUR-LES, the thick dashed lines show the one-hour averaged profiles obtained by the NCAR-LES. The thin solid lines are the observations taken by the radiosonde profile at the Southern Great Plains ARM C1 site and the thin dotted lines represent the values of the radiosounding launched at the site B5.

| | |
|--|------------------------------|
| Title Page | |
| Abstract | Introduction |
| Conclusions | References |
| Tables | Figures |
| ◀ ▶ | |
| ◀ | ▶ |
| Back | Close |
| Full Screen / Esc | |
| Print Version | |
| Interactive Discussion | |

Transport and transformation in shallow cumulus

J. Vilà-Guerau de Arellano et al.




Fig. 5. Time evolution of: (a) cloud top height and cloud base height; of (b) cloud cover and of (c) liquid water path. Black: WUR-LES, Red: NCAR-LES.

- [Title Page](#)
- [Abstract](#) [Introduction](#)
- [Conclusions](#) [References](#)
- [Tables](#) [Figures](#)
- [◀](#) [▶](#)
- [◀](#) [▶](#)
- [Back](#) [Close](#)
- [Full Screen / Esc](#)
- [Print Version](#)
- [Interactive Discussion](#)

Transport and transformation in shallow cumulus

J. Vilà-Guerau de Arellano et al.




Fig. 6. Time evolution of the vertical profiles one-hour slab averaged **(a)** inert scalar, **(b)** reactant A, **(c)** reactant B and **(d)** reactant C mixing ratios. Solid line: WUR-LES, Dashed line: NCAR-LES.

[Print Version](#)[Interactive Discussion](#)[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)

Transport and transformation in shallow cumulus

J. Vilà-Guerau de Arellano et al.




Fig. 7. Time evolution of the one-hour slab averaged vertical profiles of the resolved flux of: **(a)** inert scalar, **(b)** reactant A, **(c)** reactant B and **(d)** reactant C. Solid line: WUR-LES, Dashed line: NCAR-LES.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Print Version](#)

[Interactive Discussion](#)

Transport and transformation in shallow cumulus

J. Vilà-Guerau de Arellano et al.




Fig. 8. Time evolution of vertical profiles of the one-hour slab averaged resolved variances of (a) inert scalar, (b) reactant A, (c) reactant B and (d) reactant C. Solid line: WUR-LES, Dashed line: NCAR-LES.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Print Version](#)

[Interactive Discussion](#)

Transport and transformation in shallow cumulus

J. Vilà-Guerau de Arellano et al.




Fig. 9. Time evolution of the: (a) inert tracer height with clouds (black), inert tracer height without clouds (red), cloud top height (solid blue line) and cloud base height (blue dashed line), (b) the PBL-averaged inert scalar mixing ratio for the cloudy case (CL , solid black) and the clear sky case (CS , red).

Transport and transformation in shallow cumulus

J. Vilà-Guerau de Arellano et al.




Fig. 10. Time evolution of the vertical profiles one-hour slab averaged **(a)** inert scalar, **(b)** reactant A, **(c)** reactant B and **(d)** reactant C mixing ratios. These profiles come solely from the WUR-LES. Solid line: *CL*. Dashed line: *CS*.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Print Version](#)

[Interactive Discussion](#)

Transport and transformation in shallow cumulus

J. Vilà-Guerau de Arellano et al.




Fig. 11. One-hour averaged vertical profiles at 20:30 UTC of the mixing ratio of species A (horizontally-averaged over the whole simulation domain) and of the mixing ratio of species A (conditionally average by calculating the average fields over grid points with a nonzero liquid water content). Solid lines represent the mixing ratio calculated with the photolysis rate j_{cloud} and the dotted lines represent the mixing ratio of species A calculated with the photolysis rate j_{clear} . Black: WUR-LES. Red: NCAR-LES.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

Transport and transformation in shallow cumulus

J. Vilà-Guerau de Arellano et al.




Fig. 12. One-hour average vertical profiles of the total flux of species A (solid line) (horizontally-averaged over the whole simulation domain) and of the parameterized mass flux of species A (dotted line). Black: WUR-LES. Red: NCAR-LES.

| | |
|--|------------------------------|
| Title Page | |
| Abstract | Introduction |
| Conclusions | References |
| Tables | |
| Figures | |
| ◀ | ▶ |
| ◀ | ▶ |
| Back | Close |
| Full Screen / Esc | |
| Print Version | |
| Interactive Discussion | |

Transport and transformation in shallow cumulus

J. Vilà-Guerau de Arellano et al.




Fig. 13. Time evolution of the vertical profile (one-hour slab averaged) of the (a) intensity of segregation of species B and C and of the (b) liquid water content. WUR-LES: solid line and NCAR-LES dashed line.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)




Fig. 14. Instantaneous vertical cross section at 20:30 UTC (14:30 LT) of the liquid water mixing ratio (blue shading) and the % difference of the mixing ratio of species A calculated with the photolysis rate j_{cloud} (A_{cl}) minus the mixing ratio of species A calculated using the photolysis rate j_{clear} (A_{cs}). The red lines show positive differences (mixing ratio calculated with j_{cloud} is higher than the one calculated with j_{clear}). The black lines show negative differences. Panels: (a) Damköhler number range of 0.4–2.0, (b) Damköhler number range of 4–20, and (c) Damköhler number range of 40–200.

Transport and transformation in shallow cumulus

J. Vilà-Guerau de Arellano et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Print Version](#)

[Interactive Discussion](#)




Fig. 15. Time evolution of the percent difference of the volume averaged mixing ratio of species A from surface to cloud top (cross) calculated with the photolysis rate j_{cloud} (A_c) minus calculated using the photolysis rate j_{clear} (A_{cs}). Black: WUR-LES. Red: NCAR-LES.

Transport and transformation in shallow cumulus

J. Vilà-Guerau de Arellano et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Print Version](#)

[Interactive Discussion](#)