



HAL
open science

Drawing disconnected graphs on the Klein bottle

Laurent Beaudou, Antoine Gerbaud, Roland Grappe, Frédéric Palesi

► **To cite this version:**

Laurent Beaudou, Antoine Gerbaud, Roland Grappe, Frédéric Palesi. Drawing disconnected graphs on the Klein bottle. 2008. hal-00363494

HAL Id: hal-00363494

<https://hal.science/hal-00363494>

Preprint submitted on 23 Feb 2009

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Institut Fourier

Institut Fourier

Unité Mixte de Recherche 5582
CNRS – Université Joseph Fourier

Drawing disconnected graphs on the Klein bottle

Laurent Beaudou¹, Antoine Gerbaud¹, Roland
Grappe² and Frédéric Palesi¹

14 mars 2008

Drawing disconnected graphs on the Klein bottle

Laurent Beaudou¹, Antoine Gerbaud¹, Roland Grappe²
and Frédéric Palesi¹

¹Institut Fourier
100, rue des Maths
38 402 St-Martin d'Hères – FRANCE

²Laboratoire G-Scop
46 avenue Félix Viallet
38 000 Grenoble – FRANCE.

laurent.beaudou@ujf-grenoble.fr
antoine.gerbaud@ujf-grenoble.fr
roland.grappe@g-scop.inpg.fr
frederic.palesi@ujf-grenoble.fr

Abstract/Résumé

We prove that two disjoint graphs must always be drawn separately on the Klein bottle in order to minimize the crossing number of the whole drawing.

Keywords: Klein bottle, topological graph theory, crossing number.

Dans ce rapport, nous prouvons que deux graphes disjoints doivent toujours être dessinés séparément sur la bouteille de Klein lorsque le nombre de croisements du dessin est minimal.

Mots-clés: bouteille de Klein, graphes topologiques, croisements.

Introduction

All graphs in this paper are finite, undirected and without loops. A *path* of G is a sequence of vertices v_0, \dots, v_k of G such that for each integer i between 1 and $k - 1$, $v_i v_{i+1}$ is an edge of G and all edges are distinct. A *circuit* of G is a path v_0, \dots, v_k such that $v_0 = v_k$. A graph with a circuit that visits each of its edges exactly once is called *eulerian*. A graph is *connected* if for every pair of vertices u and v there is a path v_0, \dots, v_k such that $v_0 = u$ and $v_k = v$. We refer to [1] for an introduction to graph theory.

A *surface* is a two-dimensional manifold, with or without boundary. According to [2], there are two infinite classes of compact connected surfaces without boundary: the orientable surfaces homeomorphic to a sphere with handles attached, and the non-orientable surfaces homeomorphic to a connected sum of projective planes. For an orientable surface, the number of handles is called the *orientable genus*. For a non-orientable surface, the number of projective planes is called the *non-orientable genus*. The non-orientable surfaces of genus 1 and 2 are the projective plane and the Klein bottle, respectively. Formal definitions of these surfaces can be found in [13].

Every curve considered throughout this paper is undirected and we do not distinguish between a curve and its image. A *drawing* of a graph G on a surface Σ is a representation Ψ of G on Σ where vertices are distinct points of Σ , and edges are curves of Σ joining the points corresponding to their endvertices. A drawing is *proper* if edges are simple curves without vertices of the graph in their interiors. A *crossing* is a transversal intersection of two curves on Σ . In this paper, we restrict our attention to proper drawings where two incident edges do not cross each other, two non-incident edges cross at most once and no more than two edges cross at a single point. The *crossing number* of a drawing Ψ , denoted by $\text{cr}(\Psi)$, is the number of crossings between each pair of curves in Ψ . The crossing number of a graph G on a surface Σ is the minimum crossing number among all drawing of G on Σ . A drawing that achieves the crossing number of a graph is said *optimal*. By definition, a drawing with no crossing is an embedding. For background material about topological graph theory, the reader can refer to [9].

The crossing number of a graph on a surface leads to many unsolved problems, see [4, 10]. DeVos, Mohar and Samal conjectured the following in [3].

Conjecture 1. *Let G be the disjoint union of two connected graphs H and K and let Σ be a surface. For every optimal drawing of G on Σ , the restrictions to H and K do not intersect.*

This conjecture is obviously true for the sphere or equivalently for the Euclidean plane. It was announced proved for the projective plane in [3]. The problem remains open in the general case. In this paper, we prove that Conjecture 1 holds if Σ is the Klein bottle.

Theorem 2. *Let G be the disjoint union of two connected graphs H and K . For every optimal drawing of G on the Klein bottle, the restrictions to H and K do not cross.*

We introduce the following notations. A closed curve is *one-sided* if its neighborhood is a Möbius strip, *two-sided* otherwise. There exist two non freely homotopic one-sided simple curves a and b on the Klein bottle, a two-sided

simple curve m that cuts open the Klein bottle into a cylinder, and a two-sided simple curve e that separates the Klein bottle into two Möbius strips. A closed curve not contractible is called *essential*. According to Kawrencenko and Negami in [6], each essential simple closed curve on the Klein bottle is freely homotopic to either a , b , m or e .

For each curve c on Σ , $[c]$ denotes the set of curves freely homotopic to c . For each couple of curves (c, d) , $\text{cr}([c], [d])$ denotes the minimum number of crossings, counting multiplicities, taken over all couples of $[c] \times [d]$. Let c be a curve on Σ and I a collection of curves. The number of crossings between c and I is denoted by $\text{cr}(c, I)$. The minimum of $\text{cr}(c', I)$ taken over all curves c' in $[c]$ is denoted by $\text{cr}([c], I)$. If I is a drawing of a graph G , the minimum $\text{cr}([c], I)$ is taken on the curves in $[c]$ that do not contain any vertex of G .

We define two relations on freely homotopy classes of closed curves on the Klein bottle. Two classes $[c]$ and $[d]$ are said to be orthogonal if $\text{cr}([c], [d]) \geq 1$, otherwise disjoint. These definitions slightly differ from those of Luo in [8]. Let Ψ be a drawing on the Klein bottle. The circuits c of Ψ orthogonal to $[a]$ and disjoint from $[b]$ are called a -circuits. The circuits orthogonal to $[b]$ and disjoint from $[a]$ are called b -circuits. The circuits orthogonal to $[a]$ and $[b]$ are called m -circuits. Finally, the circuits orthogonal to $[m]$ and disjoint from $[a]$ and $[b]$ are called e -circuits.

We will apply the following result.

Theorem 3. (*De Graaf, Schrijver [5]*) *Let Ψ be an embedding of an eulerian graph on a metrizable surface Σ . Then Ψ can be decomposed into a collection of circuits I such that for each closed curve c on Σ ,*

$$\text{cr}([c], \Psi) = \sum_{d \in I} \text{cr}([c], [d]).$$

Decomposing a drawing Ψ of a graph G into a collection of circuits I means that each edge of G in Ψ is visited by exactly once by one single element of I . A related result has been obtained by Lins [7] for the projective plane and generalized by Schrijver [11] for the Klein bottle: the maximum number of pairwise edge-disjoint one-sided circuits equals the minimum number of edges intersecting all one-sided circuits. We provide another expression of this number in 7. Similarly, we express the maximum number of edge-disjoint a -circuits in 8.

This paper is organized as follows. We express the maximum number of circuits of given homotopy class in section 2. Therefore we get a lower bound on the crossing number of a drawing. A case study in Section 3 guaranties that suitable transformations, defined in Section 1, yield a representation with strictly fewer crossings.

1 Drawing graphs on surfaces of smaller genus

In this section, starting with a drawing of a graph on a surface, we define new drawings of the same graph on surfaces of smaller genus. We compute the crossing numbers of these drawings.

1.1 Removing a crosscap

Let Σ be the non-orientable surface of genus g and Σ' be the non-orientable surface of genus $g - 1$.

Proposition 4. *Let Ψ denote a drawing of a graph G on Σ . Let c be a simple closed one-sided curve on Σ which does not contain any vertex of G . There is a drawing Ψ' of G on Σ' such that*

$$\text{cr}(\Psi') = \text{cr}(\Psi) + \frac{\text{cr}(c, \Psi)(\text{cr}(c, \Psi) - 1)}{2}.$$

Proof. We cut open Σ along c and we obtain the non-orientable surface Σ' with one hole. We can glue a disk D along the boundary component to obtain Σ' . Let Ψ' be the drawing of G defined by restricting Ψ to $\Sigma \setminus c$ and redrawing the edges of Ψ that crossed c on D , crossing exactly once pairwise. The crossings of these edges add to the crossings of Ψ to give the correct number of crossings of Ψ' stated in Proposition 4. \square

The non-orientable surface of genus g can be seen as a sphere with g crosscaps attached. Attaching a crosscap to a surface Σ means removing an open disk D of Σ and identifying opposite points on the boundary of D .

Corollary 5. *Let G be the disjoint union of two eulerian connected graphs H and K . If G has a drawing on the projective plane such that the restrictions to H and K are embeddings that cross each other, then we can find another drawing of G on the projective plane with strictly fewer crossings such that the restrictions to H and K do not cross.*

Proof. Let Ψ be a drawing of G on the projective plane such that the restriction Ψ_H to H and the restriction Ψ_K to K are embeddings.

All one-sided simple essential closed curves on the projective plane are freely homotopic. Let c be such a curve. By a theorem of Lins [7], the maximum number of edge-disjoint one-sided circuits of Ψ_H and Ψ_K are $\text{cr}([c], \Psi_H)$ and $\text{cr}([c], \Psi_K)$, respectively. We may assume that $\text{cr}([c], \Psi_H)$ is smaller than $\text{cr}([c], \Psi_K)$.

Two one-sided circuits cross at least once. Hence, each one-sided circuit of Ψ_H crosses each one-sided circuit of Ψ_K , and

$$\text{cr}(\Psi) \geq \text{cr}([c], \Psi_H) \times \text{cr}([c], \Psi_K).$$

Let c' be an one-sided closed curve on the projective plane that achieves $\text{cr}([c], \Psi_H)$. By Proposition 4, there exists a drawing Ψ'_H of G on the Euclidean plane such that

$$\begin{aligned} \text{cr}(\Psi'_H) &= \frac{1}{2} \text{cr}(c', \Psi_H) \times (\text{cr}(c', \Psi_H) - 1) \\ &< \text{cr}([c], \Psi_H) \times \text{cr}([c], \Psi_K) \\ &\leq \text{cr}(\Psi). \end{aligned}$$

Let Ψ' denote the drawing of G on the projective plane obtained by disjoint union of the drawings Ψ'_H and Ψ_K . The drawings Ψ'_H and Ψ_K do not cross each other and since Ψ_K is an embedding, all crossings of Ψ' are crossings of Ψ'_H . It follows that the drawing Ψ' of G on the projective plane has strictly fewer crossings than Ψ and the restrictions to H and K do not cross. \square

1.2 Removing the two crosscaps of the Klein bottle

Proposition 6. *Let Ψ denote a drawing of a graph G on the Klein bottle. Let a' be a simple curve freely homotopic to a and m' a simple curve freely homotopic to m such that neither a' nor m' contains any vertex of G , and such that a' and m' cross only once. Then there is a drawing Ψ' of G on the Euclidean plane such that*

$$\text{cr}(\Psi') = \text{cr}(\Psi) + \text{cr}(a', \Psi) \times \text{cr}(m', \Psi) + \frac{1}{2} \text{cr}(m', \Psi) \times (\text{cr}(m', \Psi) - 1).$$

Proof. We cut the Klein bottle open along m' , disconnecting $\text{cr}(m', \Psi)$ edges of Ψ . By definition of m , the resulting surface is a cylinder. We reconnect the cut edges such that their new part remains in a small neighborhood of a' , creating exactly $\text{cr}(a', \Psi)$ crossings for each cut edge. Moreover, we can draw the $\text{cr}(m', \Psi)$ edges so that they cross each other only once. We obtain a drawing of G on the cylinder with the desired crossing number, therefore a drawing Ψ' on the Euclidean plane with the same crossing number. \square

2 Maximum number of pairwise edge-disjoint circuits

In this section we estimate the number of essential circuits of the representation.

2.1 One-sided circuits

Proposition 7. *Let Ψ be an embedding of an eulerian graph on the Klein bottle. Then the maximum number of pairwise edge-disjoint one-sided circuits equals*

$$\min \{ \text{cr}([a], \Psi) + \text{cr}([b], \Psi), \text{cr}([m], \Psi) \}.$$

Moreover, we can decompose Ψ into a collection of circuits I that achieves the maximum number of one-sided circuits and such that the number of m -circuits in I is

$$\frac{1}{2} (\text{cr}([a], \Psi) + \text{cr}([b], \Psi) - \min \{ \text{cr}([a], \Psi) + \text{cr}([b], \Psi), \text{cr}([m], \Psi) \}).$$

Proof. Let Ψ be an embedding of an eulerian graph G on the Klein bottle. Consider a collection I of edge-disjoint one-sided circuits of Ψ . Every one-sided circuit intersects either a or b . Consequently, for each circuit c in I , $\text{cr}([a], c) \geq 1$ or $\text{cr}([b], c) \geq 1$. Hence,

$$\text{cr}([a], \Psi) + \text{cr}([b], \Psi) \geq \sum_{c \in I} (\text{cr}([a], [c]) + \text{cr}([b], [c])) \geq |I|.$$

Similarly, every one-sided circuit intersects m . Hence,

$$\text{cr}([m], \Psi) \geq \sum_{c \in I} \text{cr}([m], [c]) \geq |I|.$$

Therefore the maximum number of pairwise edge-disjoint one-sided circuits is smaller than $\min \{ \text{cr}([a], \Psi) + \text{cr}([b], \Psi), \text{cr}([m], \Psi) \}$.

To complete the proof of Proposition 7, it remains to decompose Ψ into a collection of circuits that contains

$$\min \{ \text{cr}([a], \Psi) + \text{cr}([b], \Psi), \text{cr}([m], \Psi) \}$$

one-sided circuits and

$$\frac{1}{2} \text{cr}([a], \Psi) + \text{cr}([b], \Psi) - \min \{ \text{cr}([a], \Psi) + \text{cr}([b], \Psi), \text{cr}([m], \Psi) \}$$

m -circuits.

Let I be a collection of circuits given by Theorem 3, with n_a a -circuits, n_b b -circuits, n_m m -circuits and n_e e -circuits. By definition of I , the following equalities hold:

$$\begin{aligned} \text{cr}([a], \Psi) &= n_a + n_m, \\ \text{cr}([b], \Psi) &= n_b + n_m, \\ \text{cr}([m], \Psi) &= n_a + n_b + 2n_e. \end{aligned} \tag{1}$$

If n_m or n_e equals zero, then the result follows. Now assume that n_m and n_e are positive.

Let $r = \min \{n_m, n_e\}$. Consider r distinct m -circuits m_1, \dots, m_r and r distinct e -circuits e_1, \dots, e_r in I . For every integer i between 1 and r , the circuits m_i and e_i intersect and can be decomposed into an a -circuit a_i and an b -circuit b_i . Thus, we get $n_a + r$ a -circuits, $n_b + r$ b -circuits, $n_m - r$ m -circuits and $n_e - r$ e -circuits. The resulting collection of circuits I' still decomposes Ψ . By (1),

$$(n_a + r) + (n_b + r) = \min \{ \text{cr}([a], \Psi) + \text{cr}([b], \Psi), \text{cr}([m], \Psi) \},$$

and

$$2(n_m - r) = \text{cr}([a], \Psi) + \text{cr}([b], \Psi) - \min \{ \text{cr}([a], \Psi) + \text{cr}([b], \Psi), \text{cr}([m], \Psi) \}.$$

Thus I' is the desired collection of circuits. \square

2.2 a -circuits

Proposition 8. *Let Ψ be an embedding of an eulerian graph on the Klein bottle. Then the maximum number of edge-disjoint a -circuits equals*

$$\min \{ \text{cr}([a], \Psi), \text{cr}([m], \Psi) \}.$$

Proof. Let Ψ be an embedding of an eulerian graph on the Klein bottle. Consider a collection I of edge-disjoint a -circuits of Ψ . Every a -circuit intersects a . Hence,

$$\text{cr}([a], \Psi) \geq \sum_{c \in I} \text{cr}([a], c) \geq |I|.$$

Similarly, every a -circuit intersects m . Hence,

$$\text{cr}([m], \Psi) \geq \sum_{c \in I} \text{cr}([m], c) \geq |I|.$$

Therefore the maximum number of pairwise edge-disjoint a -circuits is smaller than $\min \{ \text{cr}([a], \Psi) + \text{cr}([b], \Psi), \text{cr}([m], \Psi) \}$.

To complete the proof of Proposition 8, it remains to exhibit $\min \{ \text{cr}([a], \Psi), \text{cr}([m], \Psi) \}$ a -circuits.

Let I be a collection of circuits of Ψ as stated in Theorem 3, with n_a a -circuits, n_b b -circuits, n_m m -circuits and n_e e -circuits. If n_a or n_m equals zero, then the result follows. Now assume that n_a and n_m are positive.

Let $r = \min \{ n_m, n_e \}$. Consider r distinct m -circuits m_1, \dots, m_r and r distinct e -circuits e_1, \dots, e_r in I . For every integer i between 1 and r , the circuits m_i and e_i intersect and can be decomposed into an a -circuit a_i and an b -circuit b_i . Consequently, we get $n_a + r$ a -circuits, $n_b + r$ b -circuits, $n_m - r$ m -circuits and $n_e - r$ e -circuits. The resulting collection of circuits I' still decomposes Ψ .

Let $s = \min \{ n_m - r, n_b + r \}$. If $n_m \leq n_e$, then s equals zero and we have found $\text{cr}([a], \Psi)$ a -circuits. Otherwise, consider s distinct b -circuits b_1, \dots, b_s and s distinct m -circuits m'_1, \dots, m'_s of I' . For every integer i between 1 and r , the circuits b_i and m'_i intersect and can be decomposed into an a -circuit a'_i . And so, we get $n_a + r + s$ a -circuits.

By (1),

$$\begin{aligned} n_a + r + m &= n_a + r + \min \{ n_m - r, n_b + r \} \\ &= n_a + \min \{ n_m, n_b + 2n_e \} \\ &= \min \{ \text{cr}([a], \Psi), \text{cr}([m], \Psi) \}. \end{aligned}$$

Proposition 8 is proved. \square

Note that Proposition 8 still holds when replacing a by b .

3 Main result

This section is devoted to the proof of our main result. First, we need to prove the following special case of the theorem.

Lemma 9. *Let G be the disjoint union of two eulerian connected graphs H and K . If G has a drawing on the Klein bottle such that the restrictions to H and K are embeddings that cross each other, then we can find another drawing of G on the Klein bottle with strictly fewer crossings such that the restrictions to H and K do not cross.*

Proof. Let Ψ be a drawing of G on the Klein bottle such that the restrictions Ψ_H to H and Ψ_K to K are embeddings.

To prove Lemma 9, it is enough to find two drawings Ψ'_H and Ψ'_K of H and K on two disjoint subsurfaces of the Klein bottle such that the sum of the crossings of Ψ'_H and the crossings of Ψ'_K is strictly less than the crossings of Ψ . Indeed, let Ψ'_H and Ψ'_K be such drawings and let Ψ' denote the drawing of G on the Klein bottle plane obtained by disjoint union of the drawings Ψ'_H and Ψ'_K . Then the number of crossings of Ψ' is the sum of the crossings of Ψ'_H and the crossings of Ψ'_K . It follows that the drawing Ψ' of G on the Klein bottle has strictly fewer crossings than Ψ and the restrictions to H and K do not cross.

For convenience, we denote $\text{cr}([a], \Psi_H)$, $\text{cr}([b], \Psi_H)$ and $\text{cr}([m], \Psi_H)$ by h_a , h_b and h_m , respectively. Similarly, we denote $\text{cr}([a], \Psi_K)$, $\text{cr}([b], \Psi_K)$ and $\text{cr}([m], \Psi_K)$ by k_a , k_b and k_m , respectively.

Assume without loss of generality that $h_m \leq k_m$.

By Proposition 7, there exist a decomposition of Ψ_H into pairwise edge-disjoint circuits with $\min\{h_m, h_a + h_b\}$ one-sided circuits and $(h_a + h_b - \min\{h_m, h_a + h_b\})/2$ m -circuits. Each one-sided circuit crosses Ψ_K at least $\min\{k_a, k_b\}$ times, and each m -circuit crosses Ψ_K at least k_m times. Counting the crossings between Ψ_H and Ψ_K gives the following inequality.

$$\text{cr}(\Psi) \geq \min\{h_m, h_a + h_b\} \times \min\{k_a, k_b\} + \frac{1}{2}(h_a + h_b - \min\{h_m, h_a + h_b\}) \times k_m. \quad (2)$$

With a similar decomposition of Ψ_K we obtain

$$\text{cr}(\Psi) \geq \min\{k_m, k_a + k_b\} \times \min\{h_a, h_b\} + \frac{1}{2}(k_a + k_b - \min\{k_m, k_a + k_b\}) \times h_m. \quad (3)$$

Beside, by Proposition 8, there exist $\min\{k_a, k_m\}$ pairwise edge-disjoint a -circuits of Ψ_K . Each of them crosses Ψ_H at least h_a times, therefore

$$\text{cr}(\Psi) \geq \min\{k_a, k_m\} \times h_a. \quad (4)$$

Similarly, considering b -circuits gives

$$\text{cr}(\Psi) \geq \min\{k_b, k_m\} \times h_b. \quad (5)$$

Let $m_1 \leq m_2 \leq m_3 \leq m_4$ be an ordering of the numbers h_a, h_b, k_a, k_b .

(Case 1) If $k_m \geq m_2$, then applying twice Proposition 4 provides a drawing Ψ'_H of H and a drawing Ψ'_K of K on disjoint subsurfaces of the Klein bottle such that

$$\text{cr}(\Psi'_H) + \text{cr}(\Psi'_K) = \frac{1}{2}m_1 \times (m_1 - 1) + \frac{1}{2}m_2 \times (m_2 - 1).$$

By definition of m_2 and since $k_m \geq m_2$,

$$m_2 \times m_2 \leq \max(\min\{k_a, k_m\} \times h_a, \min\{k_b, k_m\} \times h_b).$$

Hence, by (4) and (5),

$$\text{cr}(\Psi'_H) + \text{cr}(\Psi'_K) < m_2 \times m_2 \leq \text{cr}(\Psi).$$

(Case 2) If $k_m < m_2$, then

$$h_m \leq k_m \leq m_1 + m_2 \leq h_a + h_b.$$

Thus, (2) becomes

$$\text{cr}(\Psi) \geq h_m \times \min\{k_a, k_b\} + \frac{1}{2}(h_a + h_b - h_m) \times k_m. \quad (6)$$

Since $k_m \leq k_a + k_b$, the (3) becomes

$$\text{cr}(\Psi) \geq k_m \times h_a + \frac{1}{2}(k_a + k_b - k_m) \times h_m. \quad (7)$$

(Case 2.1) If $h_m \leq k_a + k_b - k_m$, then by Proposition 6 there exists a drawing Ψ'_H of H on the Euclidean plane such that

$$\text{cr}(\Psi'_H) = h_m \times h_a + \frac{1}{2}h_m \times (h_m - 1).$$

We get by (7)

$$\begin{aligned} \text{cr}(\Psi'_H) + \text{cr}(\Psi_K) &= h_m \times h_a + \frac{1}{2}h_m \times (h_m - 1) \\ &\leq k_m \times h_a + \frac{1}{2}(k_a + k_b - k_m) \times (h_m - 1) \\ &< \text{cr}(\Psi). \end{aligned}$$

(Case 2.2) If $h_m > k_a + k_b - k_m$, then $k_m < m_2$ implies

$$h_m + \max\{k_a, k_b\} \geq h_m + k_m > k_a + k_b.$$

Hence $h_m > \min\{k_a, k_b\} = m_1$.

(Case 2.2.1) If $m_1 < k_m/2$ then we apply Proposition 6. There exists a drawing Ψ'_K of K on the Euclidean plane such that

$$\text{cr}(\Psi'_K) = k_m \times m_1 + \frac{1}{2}k_m \times (k_m - 1).$$

By (6),

$$\begin{aligned} \text{cr}(\Psi_H) + \text{cr}(\Psi'_K) &\leq k_m \times m_1 + \frac{1}{2}k_m \times (k_m - 1) \\ &\leq h_m \times m_1 + (k_m - h_m) \times m_1 + \frac{1}{2}k_m \times (k_m - 1) \\ &< h_m \times m_1 + (2k_m - h_m) \times \frac{1}{2}k_m \\ &< h_m \times \min\{k_a, k_b\} + \frac{1}{2}(h_a + h_b - h_m) \times k_m \\ &< \text{cr}(\Psi). \end{aligned}$$

(Case 2.2.2) If $m_1 \geq k_m/2$ and $m_2 < 2k_m$, then applying twice Proposition 4 provides a drawing Ψ'_H of H and a drawing Ψ'_K of K on disjoint subsurfaces of the Klein bottle such that

$$\text{cr}(\Psi'_H) + \text{cr}(\Psi'_K) = \frac{1}{2}m_1 \times (m_1 - 1) + \frac{1}{2}m_2 \times (m_2 - 1).$$

Since

$$m_1 < h_m, \quad m_2 > k_m, \quad m_2 \leq h_a \quad \text{and} \quad m_1 + m_2 \leq k_a + k_b,$$

we get, by (7),

$$\begin{aligned} \text{cr}(\Psi'_H) + \text{cr}(\Psi'_K) &= \frac{1}{2}m_2 \times (m_2 - 1) + \frac{1}{2}m_1 \times (m_1 - 1) \\ &< \frac{1}{2}m_2 \times 2k_m + \frac{1}{2}m_1 \times (h_m - 1) \\ &< h_a \times k_m + \frac{1}{2}(m_1 + m_2 - k_m) \times h_m \\ &< h_a \times k_m + \frac{1}{2}(k_a + k_b - k_m) \times h_m \\ &< \text{cr}(\Psi). \end{aligned}$$

(**Case 2.2.3**) If $m_1 \geq k_m/2$ and $m_2 \geq 2k_m$ then we apply Proposition 6. There exists a drawing Ψ'_K of K on the Euclidean plane such that

$$\text{cr}(\Psi'_K) = k_m \times m_1 + \frac{1}{2}k_m \times (k_m - 1).$$

Hence, by (4),

$$\begin{aligned} \text{cr}(\Psi_H) + \text{cr}(\Psi'_K) &= k_m \times m_1 + \frac{1}{2}k_m \times (k_m - 1) \\ &< k_m \times \left(m_1 + \frac{1}{2}k_m\right) \\ &< k_m \times (2m_1) \\ &< m_1 \times m_2 \\ &< \text{cr}(\Psi). \end{aligned}$$

□

Now, we may prove the Theorem.

Theorem 10. *Let G be the disjoint union of two connected graphs H and K . For every optimal drawing of G on the Klein bottle, the restrictions to H and K do not cross.*

Proof. Let G be the disjoint union of two connected graphs H and K . Let Ψ be an optimal drawing of G on the Klein bottle.

First, assume that the restrictions to H and K are embeddings. Duplicate each edge of G and denote by G' , H' , K' the resulting eulerian graphs and by Ψ' the resulting drawing. The drawing Ψ' has $4\text{cr}(\Psi)$ crossings. Since G' is eulerian, by Lemma 9, we can find a drawing Ψ'' of G' , where the restrictions to H' and K' do not cross each other, with strictly less than $4\text{cr}(\Psi)$ crossings. Moreover, we can assume that every pair of parallel edges are drawn close enough to have the same crossings. Therefore, two pairs of parallel edges have either four crossings or none. Thus, by deleting one copy of each edge, we get a drawing of G with strictly less than $\text{cr}(\Psi)$ crossings. Moreover, the drawings of H and K do not intersect.

Secondly, suppose that the restrictions to H and K are not embeddings. Consider the graphs H'' and K'' obtained from H and K by adding a vertex for each internal crossing. The corresponding drawings are embeddings and we apply what was shown just above. Theorem 2 is proved when we replace the new vertices by the former crossings. □

References

- [1] B. Bollobás, *Modern Graph Theory*, Springer, New-York, 1998
- [2] H. R. Brahana, Systems of circuits of two-dimensional manifolds, *Ann. of Math.*, **30**(1923), 234–243
- [3] M. DeVos, B. Mohar and R. Samal, *Open Problems Garden*, http://garden.irmacs.sfu.ca/?q=op/drawing_disconnected_graphs_on_surfaces, 2007

- [4] P. Erdős and R. K. Guy, Crossing number problems, *The American Mathematical Monthly*, **80**(1973), 52–57
- [5] M. de Graaf and A. Schrijver, Decomposition on surfaces, *J. Combin. Theory Ser. B*, **70**(1997), 157–165
- [6] S. Kawrencenko and S. Negami, Irreducible triangulations of the Klein Bottle, *J. Combin. Theory Ser. B*, **70** (1997), 265–291, doi:10.1006/jctb.1997.9999
- [7] S. Lins, A minimax theorem on circuits in projective graphs, *J. Combin. Theory Ser. B*, **30**(1981), 253–262, doi:10.1016/0095-8956(81)90042-3
- [8] F. Luo, On non-separating simple closed curves in a compact surface, *Topology*, **36**(2)(1997), 381–410
- [9] B. Mohar and C. Thomassen, Graphs on surfaces, *Johns Hopkins University Press, Baltimore*, 2001
- [10] J. Pach and G. Tóth, Thirteen problems on crossing numbers, *Geombinatorics*, **9**(2000), 195–207
- [11] A. Schrijver, The Klein bottle and multicommodity flows, *Combinatorica*, **9**(1989), 375–384, doi:10.1007/BF02125349
- [12] A. Schrijver, Decomposition of graphs on surfaces and a homotopic circulation theorem, *J. Combin. Theory Ser. B*, **51**(1991), 161–210
- [13] J. Stillwell, Classical topology and combinatorial group theory, *Springer-Verlag, New York*, 1980, 62–67