



HAL
open science

Polo-like kinase 1 (PLK1) regulates IFN induction by MAVS.

Damien Vitour, Stéphanie Dabo, Malek Ahmadi Pour, Myriam Vilasco, Pierre-Olivier Vidalain, Yves Jacob, Mariana Mezel-Lemoine, Suzanne Paz, Meztli Arguello, Rongtuan Lin, et al.

► **To cite this version:**

Damien Vitour, Stéphanie Dabo, Malek Ahmadi Pour, Myriam Vilasco, Pierre-Olivier Vidalain, et al.. Polo-like kinase 1 (PLK1) regulates IFN induction by MAVS.. Journal of Biological Chemistry, 2009, epub ahead of print. 10.1074/jbc.M109.018275 . pasteur-00398965



HAL Id: pasteur-00398965

<https://pasteur.hal.science/pasteur-00398965>








Submitted on 25 Jun 2009

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.




Time (hr) : 0 2 4 6 8 10 12 14




% cells in	G1	S	G2
0 hr	78.2	19.3	0.0
2 hr	77.7	17.7	0.0
4 hr	92.5	0.91	0.0
6 hr	88.0	13.9	0.0
8 hr	4.2	9.4	85.3
10 hr	36.3	9.1	58.5
12 hr	74.3	12.7	10.6
14 hr	70.1	22.4	4.71

FL2-A: DNA-Area

1 MPFAEDKTYKYICRNFSNFCNVDVVEILPYLPCLTARDQDRPRATCTLSG 50
51 NRDTLWHLFNTLQRRPGWVEYFIAALRGCELVDLADEVASVYQSYQPRTS 100
101 DRPPDPLEPPSLPAERPGPPTPAAAHSIPYNSCREKEPSYPMPVQETQAP 150
151 ESPGENSEQALQTLSPRAIPRNPDGGPLESSSDLAALSPLTSSSGHQEQDT 200
201 ELGSTHTAGATSSLTPSRGPVSPSVSFQPLAR**STP**RASRLPGPTGSVVST 250
251 GTSFSSSSPGLASAGAAEGKQGAESDQAEPIICSSGAEAPANSLPSKVPT 300
301 TLMPVNTVALKVPANPASVSTVPSKLP TSSKPPGAVPSNALTNPAPSKLP 350
351 INSTRVGMVPSKVPTSMVLTKVPASTVPTDGSSRNEETPAAPT PAGATGG 400
401 SSAWLDSSSEN RGLGSELSKPGVLASQVDSPFSGCFEDLAISASTSLGMG 450
451 PCHGPEENEYKSEGTFGIH VAENPSIQLLEGNP GPPADPDGGPRPQADRK 500
501 FQEREVPCHRPSPGALWLQVAVTGVLVVTLLVVLYRRRLH 540





SUPPLEMENTARY INFORMATION

Supplementary Fig 1. Respective expression of PLK1 and MAVS as function of the cell cycle.

Hela cells were synchronized by double thymidine block as described (58). The medium was removed and cells were washed three times in growth medium. At different times after thymidine release, cells were trypsinized and fixed in ethanol for FACS analysis to study DNA content, and protein extracts were prepared in RIPA buffer (10mM Tris-HCl pH 7.4, 150mM NaCl, 5mM EDTA, 0.1% SDS, 0.5% sodium deoxycholate, 1% NP-40) for immunoblot analysis of PLK1, MAVS and Hsp60 (as control of mitochondria expression). The percentage of cells in the different phases of cell cycle are listed in a table.

Supplementary Fig 2. MAVS amino acid sequence.

The domains 1-157 and 470-540 are represented in grey and correspond to regions that do not bind PLK1. In the 157-470 domain, the 180-280 sequence that specifically recruits the phosphopeptide binding site of PLK1 is framed, as well as the STP motif containing the T234 residue.

Supplementary Fig 3. Requirement of MAVS T234 for specific binding to PLK1-PBD.

HEK 293T cells were transfected with pFLAG-PLK1-PBD WT in the presence of pcDNA3.1(+) (Empty vector) or pDEST-GST-MAVS1-360, in which the sequence was either unmodified (WT) or carrying the mutation T234A. 24 hrs after transfection, the cells were lysed in buffer B minus glycerol and submitted to GST pull-down of the GST-MAVS constructs. The presence of PLK1-PBD and MAVS retained in the GST pull-down and expression of PLK1 in the total cell extracts was revealed by immunoblot using anti-FLAG and anti-MAVS antibodies.

Supplementary Fig 4. PLK1 does not affect MAVS dimerization

HEK 293T cells were transfected with pMyc-MAVS, pFLAG-MAVS and increasing amounts of pEGFP-PLK1-PBD. After 24 hr, Myc-MAVS was immunoprecipitated as in Fig5A. The presence of FLAG-MAVS (upper panel), Myc-MAVS and PLK1-PBD (middle panels) in the immunocomplexes was revealed by immunoblot using anti-FLAG, anti-c-Myc or anti-PLK1 antibodies. Expression of FLAG-MAVS was controlled in the total cell extracts using anti-FLAG antibody (lower panel).