



HAL
open science

A dynamical approach to von Neumann dimension

Antoine Gournay

► **To cite this version:**

Antoine Gournay. A dynamical approach to von Neumann dimension. *Discrete and Continuous Dynamical Systems - Series S*, 2010, 26 (3), pp.967-987. 10.3934/dcds.2010.26.967 . hal-00473487

HAL Id: hal-00473487

<https://hal.science/hal-00473487>

Submitted on 15 Apr 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A dynamical approach to Von Neumann dimension

Antoine Gournay

e-mail: gournay@math.kyoto-u.ac.jp
Department of Mathematics, Faculty of Science
Kyoto University,
Kyoto 606-8502, Japan

Abstract

Let Γ be an amenable group and V be a finite dimensional vector space. Gromov pointed out that the von Neumann dimension (with respect to Γ) can be obtained by looking at the semi-axes of certain ellipsoids. This metric point of view does not require a Hilbertian structure. It is used in this article to associate to a Γ -invariant linear subspace Y of $\ell^p(\Gamma; V)$ a real positive number $\dim_{\ell^p} Y$ (which is the von Neumann dimension when $p = 2$). By analogy with von Neumann dimension, we explore the properties of this number to conclude that there can be no injective Γ -equivariant linear map of finite-type from $\ell^p(\Gamma; V) \rightarrow \ell^p(\Gamma; V')$ if $\dim V > \dim V'$. A generalization of the Ornstein-Weiss lemma is developed along the way.

1 Introduction

Let Γ be a discrete group, then it is possible to associate to certain unitary representations a positive real number called von Neumann dimension (see [11, §1] or [14, §1]). More precisely, let $f : \Gamma \rightarrow X$ be a map. The natural (right) action of Γ on spaces of maps means is, in the present text, the action given by $\gamma f(\cdot) = f(\gamma^{-1}\cdot)$. Now, let H be a Hilbert space and consider the space $\ell^2(\Gamma) \otimes H$ where Γ acts naturally on the first factor and trivially on the second. Then, the von Neumann dimension is defined for Γ -invariant subspaces of $\ell^2(\Gamma) \otimes H$.

Let V be a *finite* dimensional vector space and $\|\cdot\|$ a norm (the choice of which will not matter as the dimension is finite). The subject matter in this article are Γ -invariant linear subspaces of

$$\ell^p(\Gamma; V) = \ell^p(\Gamma) \otimes V = \{f : \Gamma \rightarrow V \mid \sum_{\gamma \in \Gamma} \|f(\gamma)\|^p \text{ is finite}\}$$

for the natural action of Γ . From now on Γ will be assumed amenable. There are reasons to exclude non-amenable groups Γ . Indeed, D. Gaboriau pointed out that if a

⁰AMS Subject classification: 37A35, 37C45, 41A46, 43A07, 43A15, 43A65, 46E30, 54C25, 70G60

notion of dimension existed in the ℓ^p setting (that is, a quantity satisfying properties P1-P10 listed below), then there would be a formula for the Euler characteristic of Γ as the alternate sum of the dimensions of ℓ^p cohomology spaces. On one hand, torsion-free cocompact lattices in $SO(4, 1)$ have positive Euler characteristic. On the other, for p big enough, their ℓ^p cohomology vanishes in all degrees but the first (see [15, Theorem 2.1]). This would lead to a contradiction.

We are looking for a notion of dimension for such subspaces, which would increase under injective equivariant linear maps. Inspired by an argument of [6, §1.12] (and partially answering a question found therein), we shall introduce a quantity \dim_{ℓ^p} which, when $p = 2$, coincides with definition of von Neumann dimension. This quantity is obtained by a process similar to that of metric entropy or mean dimension, that is an asymptotic growth factor. The definition relies *a priori* on an exhaustion of Γ , but a generalization of the Ornstein-Weiss lemma in section 5 implies the result is independent of this choice.

Though we prove many properties of \dim_{ℓ^p} , important properties are still lacking. Nevertheless, the results obtained in this paper suffice to establish a non existence result for maps of finite type. We recall their construction.

Let $D \subset \Gamma$ be a finite set and let $g : V^D \rightarrow V'$ be a continuous map. This data enables the definition of a Γ -equivariant continuous map g_D from $Z \subset \ell^p(\Gamma; V)$ to $\ell^p(\Gamma; V')$ as follows

$$g_D(z)(\gamma) = g(z(\gamma\delta))_{\delta \in D}.$$

Remark that what we denote here as $\ell^p(\Gamma; V)$ is more frequently written $\ell^p(\Gamma) \otimes V$.

Theorem 1.1. *Let Γ be an amenable discrete group. Let V and V' be finite dimensional vector spaces. If $f : \ell^p(\Gamma; V) \rightarrow \ell^p(\Gamma; V')$ is an injective Γ -equivariant linear map of finite type then $\dim V \leq \dim V'$.*

2 Definition and properties of \dim_{ℓ^p}

Given a positive number ε , a notion of dimension up to scale ε for (X, τ, δ) a topological space equipped with a pseudo-distance will be needed. Data compression problems turn out to be a good source of inspiration. When one is interested in compression algorithms, it is not only important that the compression map has “small” fibers (so that not too much data is lost) but also has an image which is “small” in some sense (so that the compression is effective).

A slight variant of the one used in [5], [6], or [18] shall be employed, namely one that is defined for pseudo-distances only. As such, it will be useful to use a topology τ that does not come from the pseudo-distance. Please note that the term diameter (denoted Diam) will continue to be used even if it is defined using a pseudo-distance (thus a set of diameter 0 may contain more than one point).

Definition 2.1. *Let (X, τ, δ) be a metric space. Call $\text{wdim}_{\varepsilon}(X, \tau, \delta)$ the smallest integer k such that there exists a continuous (for τ) map $f : X \rightarrow K$ where K is a k -dimensional polyhedron such that $\forall k \in K, \text{Diam } f^{-1}(k) \leq \varepsilon$.*

$$\text{wdim}_{\varepsilon}(X, \tau, \delta) = \inf_{f: X \rightarrow K} \{ \dim K \mid f \text{ is continuous for } \tau \text{ and } \forall k \in K, \text{Diam } f^{-1}(k) < \varepsilon \}.$$

We will sometimes omit to mention τ when it is the topology induced by δ .

Definition 2.2. Let (X, τ, δ) be a space endowed with a topology τ and a pseudo-distance δ . Let Γ be a countable group which acts on X and let $\{\Omega_i\}$ be an increasing sequence of finite subsets of Γ . The $\ell^p(\Gamma)$ width growth coefficient of X for the sequence $\{\Omega_i\}$ is

$$\text{Wgc}_{\ell^p}(X, \tau, \{\Omega_i\}) = \lim_{\varepsilon \rightarrow 0} \limsup_{i \rightarrow \infty} \frac{\text{wdim}_{\varepsilon}(X, \tau, \delta_{\ell^p(\Omega_i)})}{|\Omega_i|} \in [0, +\infty].$$

where $\delta_{\ell^p(\Omega)}(x, x') = \left(\sum_{\gamma \in \Omega} \delta(\gamma x, \gamma x')^p \right)^{1/p}$ when $p < \infty$ and $\delta_{\ell^\infty(\Omega)}(x, x') = \sup_{\gamma \in \Omega} \delta(\gamma x, \gamma x')$.

When δ is a distance, $\delta_{\ell^\infty(\Omega)}$ is but the dynamical distance. If furthermore τ is the topology this distance induces, then this is the (metric) mean dimension (see [6, §1.5] or [9, §4]). In the present text, this is an intermediate definition and will only be used in a particular context, namely when X is a subset of $\ell^\infty(\Gamma; V)$. The pseudo-metric will be given by evaluation at the neutral element e_Γ of Γ : $ev(x, x') = \|x(e_\Gamma) - x'(e_\Gamma)\|_V$. Lastly, τ^* will denote the product topology induced from $X \subset V^\Gamma$ (which coincides with the weak-* topology, when defined).

Definition 2.3. Let V be a finite-dimensional normed vector space. Let $Y \subset \ell^\infty(\Gamma; V)$ be a subset invariant by the natural action of Γ , an amenable countable group. Let Ω_i be a Følner sequence for Γ . Then, the ℓ^p von Neumann dimension of Y is defined by

$$\dim_{\ell^p}(Y, \{\Omega_i\}) = \sup_{r \in \mathbb{R}_{\geq 0}} \text{Wgc}_{\ell^p}(B_r^{Y,p}, ev, \{\Omega_i\})$$

where $B_r^{Y,p} = Y \cap B_r^{\ell^p(\Gamma; V)}$.

Note that $B_r^{Y,p}$ is defined by an intersection rather than a projection, as the former are not always easy to define in ℓ^p . Also, the choice of τ^* as a topology comes from the fact that it is the weakest topology that is stronger than the topologies induced by $\delta_{\ell^p(\Omega)}$ for any p or Ω .

In the rest of this article, Y will often be a linear subspace. In these cases, one does not need to take the sup on r . Indeed, $\text{Wgc}_{\ell^p}(B_r^{Y,p}, ev, \{\Omega_i\})$ does not depend on r (as can be seen using dilation and a change of variable $\varepsilon \mapsto r\varepsilon$).

When Y is a Γ -invariant linear subspace of $\ell^\infty(\Gamma; V)$,

- P1 (Independence) $\dim_{\ell^p}(Y, \{\Omega_i\})$ is actually independent of the choice of Følner sequence $\{\Omega_i\}$ (cf. corollary 5.2);
- P2 (Normalization) $\dim_{\ell^p} \ell^p(\Gamma; V) = \dim V$ (cf. example 3.5);
- P3 (Invariance) If $f : Y_1 \rightarrow Y_2$ is an injective Γ -equivariant linear map of finite type, then $\dim_{\ell^p} Y_1 \leq \dim_{\ell^p} Y_2$ (cf. proposition 3.8 and example 3.9);
- P4 (Completion) If \overline{Y} is the completion of Y in $\ell^p(\Gamma; V)$ for the ℓ^p norm, then $\dim_{\ell^p} Y = \dim_{\ell^p} \overline{Y}$ (cf. proposition 3.10);

P5 (Reduction) If $\Gamma_1 \subset \Gamma_2$ is of finite index, and if $Y \subset \ell^p(\Gamma_2; V)$ is seen by restriction as a subspace of $\ell^p(\Gamma_1; V^{[\Gamma_2:\Gamma_1]})$ then $[\Gamma_2 : \Gamma_1] \dim_{\ell^p}(Y, \Gamma_2) = \dim_{\ell^p}(Y, \Gamma_1)$ (cf. proposition 3.11).

P6 If $Y \subset \ell^2(\Gamma; V)$, $\dim_{\ell^2} Y$ coincides with the von Neumann dimension (cf. corollary A.2);

In light of P6, when $p = 2$ the following further properties of \dim_{ℓ^2} are listed by Cheeger and Gromov in [1, §1].

P7 (Non-triviality) $Y \subset \ell^2$ is trivial if and only if $\dim_{\ell^2} Y = 0$.

P8 (Additivity) $\dim_{\ell^2} Y_1 \oplus Y_2 = \dim_{\ell^2} Y_1 + \dim_{\ell^2} Y_2$;

P9 (Continuity) If $\{Y_i\}$ is a decreasing sequence of closed linear subspaces then $\dim_{\ell^2}(\cap Y_i) = \lim_{i \rightarrow \infty} \dim_{\ell^2} Y_i$

P10 (Reciprocity) If $\Gamma_1 \subset \Gamma_2$ and if $Y_2 \subset \ell^2(\Gamma_2; V)$ is the subspace induced by $Y_1 \subset \ell^2(\Gamma_1; V)$ then $\dim_{\ell^2}(Y_2, \Gamma_2) = \dim_{\ell^2}(Y_1, \Gamma_1)$;

Proposition 4.1 also establishes P7 for \dim_{ℓ^1} . On the other hand, the continuity property (P9) of the Von Neumann dimension does not hold if $p = 1$ (see example 4.2).

For linear subspaces $Y \subset \ell^\infty$ non-triviality (P7) is false, though it might be true for $Y \subset c_0(\Gamma, V)$, the latter being the space of all $x \in \ell^\infty(\Gamma; V)$ tending to 0 at infinity (i.e. $\|x\|_{\ell^\infty(\Gamma \setminus F_i)} \rightarrow 0$ for all exhaustive increasing sequence of (finite) subsets $\{F_i\}$).

Finally, the existence of an element of finite support in Y implies P7. By using a similar but less convenient definition of \dim_{ℓ^2} , the author is also aware of a proof of P8 and P10 (when the index is finite) for $p = 2$ without using P6 and the previously known properties of von Neumann dimension.

Though these properties are stated for Γ -invariant linear subspaces, some remain true for more general subsets Y : P1 and P5 hold for any Γ -invariant subset, and P4 is also true when Y is not Γ -invariant.

These properties offer a partial answer to the question discussed at the beginning of the present article.

Proof of theorem 1.1. It is but a simple consequence of P2 and P3. □

Being crucial to the proof above and less technical, we shall begin by proving properties P2-P5. Section 4 then discusses P7 and P9 for $p = 1$ or ∞ . The proof of P1 requires some technical lemmas on amenable groups and is thus relegated to section 5. As for P6, it relies mostly on a result of Gromov and is discussed in appendix A.

3 Proof of properties P2-P5

Before the properties of \dim_{ℓ^p} can be established, the basic properties of wdim_ε must be mentioned.

3.1 Properties of wdim_ε

Most of the content of this subsection may be found in [2, §4.5], [3, §3], [5, Proposition 2.1] and [6, §1.1].

Proposition 3.1. *Let X be space endowed with a topology τ and a pseudo-distance δ .*

- a. *The function $\varepsilon \mapsto \text{wdim}_\varepsilon(X, \tau, \delta)$ is non-increasing.*
- b. *Suppose δ is a distance and τ the topology it induces. Let $\dim X$ be the covering dimension of X , then $\text{wdim}_\varepsilon(X, \tau, \delta) \leq \dim X$.*
- c. $\text{wdim}_\varepsilon(X, \tau, \delta) = 0 \Leftrightarrow \varepsilon \geq \text{Diam} X$

Except for b, the proof of these properties are simple. Before moving on, let us recall two (fundamental) examples.

Example 3.2: Let X be a normed vector space with the distance $\delta(x, x') = \|x - x'\|$ and τ the norm topology. Let $A = B_1^X$ be its unit ball. Then $\text{wdim}_\varepsilon(A, \tau, \delta) = \dim X$ if $\varepsilon < 1$ (see [6, §1.1B] or, for more details, [5, Lemma 2.5]) and $\text{wdim}_\varepsilon(A, \tau, \delta) = 0$ if $\varepsilon \geq 2$ (consider the map which sends all of A to one point).

The second example comes from a question which arises naturally in the context of compressed sensing, namely we look at a ball for some norm but we endow with a metric coming from another norm.

Example 3.3: Let $\ell^p(n)$ denote \mathbb{R}^n with its ℓ^p norm. Then one can look at $B_1^{\ell^q(n)} \subset \ell^p(n)$, and try to compute its wdim. (In compression theory, it is frequent to consider a ball for some metric endowed with a different metric; see [4].) When $q \geq p$ then we the behaviour is essentially as in the previous example. However, if $q < p$ then one finds that, for $1 \leq k < n$,

$$\text{wdim}_\varepsilon(B_1^{\ell^p(n)}, \ell^q) \begin{cases} = 0 & \text{if } 2 \leq \varepsilon, \\ \leq k & \text{if } 2(k+1)^{1/q-1/p} \leq \varepsilon, \\ \geq k & \text{if } \varepsilon < k^{1/q-1/p}, \\ = n & \text{if } \varepsilon < n^{1/q-1/p}. \end{cases}$$

We briefly mention how to obtain these. The first line is a consequence of 3.1.c. The second is found by using an explicit map described in [5, proposition 1.3] and [18]. This maps takes a vector, keeps only the k biggest coordinates (in absolute value), then from these k coordinates take the smallest and subtract (or add, so as to reduce in absolute value) it to the others. Finally, the third line comes from the presence of an ℓ^q ball of dimension k and radius $k^{1/q-1/p}$ in $\ell^p(n)$. The fourth line is also obtained using this argument (for n) together with proposition 3.1.b.

This second set of properties are crucial to what follows.

Proposition 3.4. *For $i = 1, 2$, let X_i be spaces endowed with topologies τ_i and pseudo-metric δ_i .*

- a. *Let $f : X_1 \rightarrow X_2$ be a continuous map such that $\delta_1(x, x') \leq C\delta_2(f(x), f(x'))$ where $C \in]0, \infty[$. Then $\text{wdim}_\varepsilon(X_1, \tau_1, \delta_1) \leq \text{wdim}_{\varepsilon/C}(X_2, \tau_2, \delta_2)$.*

- b. A dilation has the expected effect, i.e. let $f : X_1 \rightarrow X_2$ be a homeomorphism such that $\delta_1(x, x') = C\delta_2(f(x), f(x'))$. Then $\text{wdim}_\varepsilon X_1 = \text{wdim}_{\varepsilon/C} X_2$.
- c. For $q \in [1, \infty]$, let $X := X_1 \times_q X_2$ be the space $X_1 \times X_2$ endowed with the product topology and the pseudo-metric $\delta := \delta_1 \times_q \delta_2$ given by $\delta(x, x')^q = \delta_1(x, x')^q + \delta_2(x, x')^q$. Also for $q = \infty$, let $\delta(x, x') = \max(\delta_1(x, x'), \delta_2(x, x'))$. Then $\text{wdim}_{2^{1/q}\varepsilon} X \leq \text{wdim}_\varepsilon X_1 + \text{wdim}_\varepsilon X_2$.

The proofs can be found in [2, §4.5], [3, Lemma 3.2] or [5, Proposition 2.1]. For example, the third is obtained by looking at the size of the fibers of the map $f = f_1 \oplus f_2$, where $f_i : X_i \rightarrow K_i$ satisfy the conditions of definition 2.1 and $\dim K_i = \text{wdim}_\varepsilon(X_i, \tau_i, \delta_i)$.

A useful way of stating 3.4.a is that a continuous map that does not reduce distance will not make wdim_ε smaller.

3.2 Properties of \dim_{ℓ^p}

Let us begin by two basic examples.

Example 3.5: If $1 \leq q < p \leq \infty$, and $Y = B_1^{\ell^q(\Gamma; \mathbb{R})}$ then $\dim_{\ell^p}(Y, \{\Omega_i\}) = 0$ (independently of the choice of sequence $\{\Omega_i\}$). Indeed, $B_r^{Y,p} \subset B_r^{\ell^q(\Gamma; \mathbb{R})}$, and $\text{wdim}_\varepsilon(B_r^{Y,p}, \text{ev}_{\ell^p(\Omega_i)}) = \text{wdim}_\varepsilon(B_r^{\ell^q(n_i)}, \ell^p)$ where $n_i = |\Omega_i|$. However using example 3.3 (and dilations to get back to a unit ball, see proposition 3.1.b), $\text{wdim}_\varepsilon(B_r^{\ell^q(n_i)}, \ell^p)$ is, for fixed ε , bounded above and below by two functions that do not depend on n_i . Thus,

$$\limsup_{i \rightarrow \infty} \frac{\text{wdim}_\varepsilon(B_r^{Y,p}, \tau^*, \text{ev}_{\Omega_i})}{|\Omega_i|} \leq \limsup_{i \rightarrow \infty} \frac{\text{wdim}_\varepsilon(B^{\ell^q(n_i)}, \ell^p)}{|\Omega_i|} = 0.$$

Example 3.6: By direct computation, we now show that

$$\dim_{\ell^p} \ell^q(\Gamma; V) = \dim V.$$

For $q \in [1, \infty]$, let $Y' = \ell^q(\Gamma; V)$. Then $(B_1^{Y',p}, \text{ev}_{\ell^p(\Omega)})$ is “isometric” to $(B_1^{\ell^p(\Omega; V)}, \text{ev}_{\ell^p(\Omega)})$. Indeed, the restriction map to Ω has a kernel of “diameter” 0, so property 3.4.a applies with $C = 1$. On the other hand, inclusion of $\ell^p(\Omega, V)$ in $\ell^p(\Gamma, V)$ (by extending the functions by 0) is also a linear map and property 3.4.a holds again with $C = 1$. Consequently, $(B_1^{Y',p}, \text{ev}_{\ell^p(\Omega)})$ will have the same wdim_ε as $(B_1^{\ell^p(\Omega; V)}, \text{ev}_{\ell^p(\Omega)})$, $\forall \varepsilon$. This later being a ball with its proper metric, if $\varepsilon < 1$ its wdim_ε will be the dimension of the space, $|\Omega| \dim V$.

In what follows the total vector space will be $Y \subset \ell^p(\Gamma; V)$. Thus we will abbreviate by $(B_r^{Y,p}, \text{ev}_{\ell^p(\Omega)})$ to mean the set $B_r^{Y,p} \subset \ell^p(\Gamma; V)$ with the pseudo-norm $\text{ev}_{\ell^p(\Omega)}$ and the topology induced from the product topology. We stress that $B_r^{Y,p}$ is not the ball for the pseudo-norm $\text{ev}_{\ell^p(\Omega)}$; it is the intersection of Y with the ball of radius r in $\ell^p(\Gamma)$ (endowed with its actual norm). The next property is a corollary of a generalization of the Ornstein-Weiss lemma described in section 5. Even if proposition 5.2 is a very important property, weaker version can be sufficient for some of our needs. Indeed, the following simple lemma is actually all that we need to show that \dim_{ℓ^p} is preserved under Γ -equivariant maps of finite type.

Lemma 3.7. *Let Y be as above, and let $\{\Omega_i\}$ and $\{\Omega'_i\}$ be such that*

$$\lim_{i \rightarrow \infty} \frac{|\Omega_i \cup \Omega'_i \setminus \Omega_i \cap \Omega'_i|}{|\Omega_i \cup \Omega'_i|} = 0,$$

then $\text{Wgc}_{\ell^p}(B_1^{Y,p}, \text{ev}, \{\Omega_i\}) = \text{Wgc}_{\ell^p}(B_1^{Y,p}, \text{ev}, \{\Omega'_i\})$

Proof. It suffices to note that, when $\Omega \subset \Omega'$,

$$\begin{aligned} \frac{\text{wdim}_{\varepsilon}(B_1^Y, \text{ev}_{\ell^p(\Omega)})}{|\Omega|} &\leq \frac{\text{wdim}_{\varepsilon}(B_1^Y, \text{ev}_{\ell^p(\Omega')})}{|\Omega'|} \\ &\leq \frac{\text{wdim}_{\varepsilon}(B_1^Y, \text{ev}_{\ell^p(\Omega)})}{|\Omega|} \frac{|\Omega|}{|\Omega'|} + \dim V \frac{|\Omega' \setminus \Omega|}{|\Omega'|}. \end{aligned}$$

Furthermore, $\frac{|\Omega|}{|\Omega'|} = 1 - \frac{|\Omega' \setminus \Omega|}{|\Omega'|}$. Thus, computing Wgc with respect to the sequences $\{\Omega_i \cap \Omega'_i\}$, $\{\Omega_i\}$ or $\{\Omega'_i\}$ will yield the same result as a computation made using $\{\Omega_i \cup \Omega'_i\}$. \square

Proposition 3.8. *Let $Y \subset \ell^\infty(\Gamma; V)$ and $Y' \subset \ell^\infty(\Gamma; V')$ be Γ -invariant linear subspaces. Let $f : Y \rightarrow Y'$ be a Γ -equivariant map continuous for τ^* and such that there exists a real $c_f \in \mathbb{R}_{>0}$ and a finite subset $D_f \subset \Gamma$ satisfying $\text{ev}(x, y) \leq c_f \text{ev}_{\ell^p(D_f)}(f(x), f(y))$ then*

$$\dim_{\ell^p}(Y, \{\Omega_i\}) \leq \dim_{\ell^p}(Y', \{\Omega_i\})$$

Proof. The case $p = \infty$ is simpler, we shall only describe the case $p < \infty$. Here $B_r^{Y',p} = Y' \cap \tilde{B}_r^{\ell^p(\Gamma; V')}$. On one hand, since f is continuous for τ^* (the product topology or the weak-* topology), $\exists r_f \in \mathbb{R}_{>0}$ such that $f(B_1^Y) \subset B_{r_f}^{Y',p}$. Indeed, since the image is weakly-* compact (in particular, weakly-* bounded) it is bounded (cf. [16, theorem 3.18]). On the other hand, the assumption satisfied by f on distances propagates by equivariance to different evaluations:

$$\begin{aligned} \text{ev}(\gamma x, \gamma y) &\leq c_f \text{ev}_{\ell^p(D_f)}(f(\gamma x), f(\gamma y)) = c_f \text{ev}_{\ell^p(D_f)}(\gamma f(x), \gamma f(y)) \\ &= c_f \text{ev}_{\ell^p(D_f \gamma)}(f(x), f(y)). \end{aligned}$$

This implies that $\text{ev}_{\ell^p(\Omega)}(x, y) \leq c_f |D_f| \text{ev}_{\ell^p(\Omega D_f)}(f(x), f(y))$ and, incidentally, that f is injective. Lastly, since the image of the ball (of radius 1) is contained in a ball (of radius r_f)

$$\begin{aligned} \text{wdim}_{\varepsilon}(B_1^{Y,p}, \text{ev}_{\ell^p(\Omega_i)}) &\leq \text{wdim}_{\varepsilon/c_f |D_f|}(\tilde{B}_{r_f}^{Y',p}, \text{ev}_{\ell^p(D_f \Omega_i)}) \\ &\leq \text{wdim}_{\varepsilon/c_f |D_f| r_f}(B_1^{Y',p}, \text{ev}_{\ell^p(\Omega_i D_f)}). \end{aligned}$$

The first inequality comes from 3.4.a. Dividing by $|D_f \Omega_i| = \frac{|D_f \Omega_i|}{|\Omega_i|} |\Omega_i|$ and passing to the limit yields that

$$\text{Wgc}_{\ell^p}(B_1^{Y,p}, \text{ev}, \{\Omega_i\}) \lim_{i \rightarrow \infty} \frac{|D_f \Omega_i|}{|\Omega_i|} \leq \text{Wgc}_{\ell^p}(B_1^{Y',p}, \text{ev}, \{\Omega_i D_f\}).$$

Since $\{\Omega_i\}$ is a Følner sequence, the limit on the left-hand side is 1. Furthermore, the hypothesis of lemma 3.7 are satisfied; the right-hand term is nothing else than $\dim_{\ell^p}(Y', \{\Omega_i\})$. \square

From now on, we will drop the explicit reference to the Følner sequence.

Since the assumptions of the previous proposition are quite abstract, it is good to check that they hold in certain categories of maps. The main constraint is the existence of c_f and D_f . Let f be a map to which proposition 3.8 applies. Let $f^{-1} : Y' = \text{Im } f \rightarrow Y$ the inverse of f on its image, then the condition

$$ev(x, y) \leq c_f ev_{\ell^p(D_f)}(f(x), f(y))$$

can be read as a condition on the modulus of continuity of f^{-1} . More precisely, $f^{-1} : (Y', ev_{\ell^p(D_f)}) \rightarrow (Y, ev)$ must be continuous with a linear modulus of continuity (i.e. that f^{-1} must be Lipschitz). If the function f^{-1} is continuous for the product topology, weakening the topology on its image is evidently not restrictive. Things are not so direct on the domain.

For $\Omega \subset \Gamma$, denote by $R_\Omega : V^\Gamma \rightarrow V^\Omega$ the restriction of functions to the domain Ω . Let $U \subset (Y, ev)$ be an open set; then if Y is seen as a subset of V^Γ , $R_{\{e\}}U$ is an open set on the factor $R_{\{e\}}Y$, and all of Y on the other factors: $R_{\Gamma \setminus \{e\}}U = R_{\Gamma \setminus \{e\}}Y$. It is then possible that on a finite number of factors of $Y' \subset V^\Gamma$ (the required set D_f) $f(U)$ will not be all the image of f : $R_D f(U) \neq R_D Y'$. For example, for $F \subset \Gamma$ a finite subset and $f = f_F$ of finite type, the condition is that $f : (Y, ev) \rightarrow (Y, ev_{\ell^p(F)})$ be open (on its image) and of Lipschitz inverse. Remember that the condition on the distances in proposition 3.8 and Γ -equivariance imply injectivity of the map. Here is the major application of proposition 3.8.

Corollary 3.9. *Let $Y \subset \ell^\infty(\Gamma; V)$ and $Y' \subset \ell^\infty(\Gamma; V')$ be Γ -invariant linear subspaces. Let $f : Y \rightarrow Y'$ be a Γ -equivariant injective linear map of finite type. Then*

$$\dim_{\ell^p} Y \leq \dim_{\ell^p} Y'.$$

Proof. Let $F \subset \Gamma$ be a finite subset which can be used to define f as a map of finite type, i.e. $f = f_F$. If f is a linear map, injectivity of f implies that it is open on its image (Banach-Schauder theorem or open mapping theorem) for the norm topologies. This remains true for the topology of ev on the domain and $ev_\ell^p(F)$ on the image as the first is weaker (its open sets are described above) and the image of open sets is of the form $R_F^{-1}U'$ for $U' \subset V^F$. So $f : (Y, ev) \rightarrow (Y, ev_{\ell^p(F)})$ is open (on its image).

Next, write the Γ -equivariant linear map of finite type f as

$$x \mapsto f(x) \text{ such that } f(x)(\gamma) = \sum_{\gamma' \in F} a_{\gamma'}(x(\gamma'\gamma)),$$

where $a_{\gamma'} \in \text{Hom}(V, V')$. Since it is injective, it possesses a Γ -equivariant linear inverse (on its image) $f^{-1} = g$:

$$x \mapsto g(x) \text{ such that } g(x)(\gamma) = \sum_{\gamma' \in G} b_{\gamma'}(x(\gamma'\gamma)),$$

where $b_\gamma \in \text{Hom}(V', V)$ and $G \subset \Gamma$ might not be finite.

Then proposition 3.8 can be invoked with $D_f = F$, and c_f the Lipschitz constant of $g : (Y', \text{ev}_{F^{-1}}) \rightarrow (Y, \text{ev})$. Thus $c_f \leq \|\oplus_{\gamma \in F^{-1} \cap G} b_\gamma\|$. \square

Proposition 3.10. *Let $Y \subset \ell^\infty(\Gamma; V)$ be an open linear subspace and let \bar{Y} be its completion in $\ell^p(\Gamma; V)$, then $\dim_{\ell^p} Y = \dim_{\ell^p} \bar{Y}$.*

Proof. The argument is identical to that of example 3.5: when restricted to a finite $\Omega \subset \Gamma$, these two spaces cannot be distinguished (being of finite dimension they are closed). In other words, there exists a linear map, given by the restriction R_Ω , and whose kernel is in the “ball” of radius 0:

$$R_\Omega : (B_1^{\bar{Y}}, \text{ev}_{\ell^p(\Omega)}) \rightarrow (R_\Omega B_1^Y, \text{ev}_{\ell^p(\Omega)}).$$

Thus, $\forall \varepsilon \in [0, 1]$, $\text{wdim}_\varepsilon(B_1^{\bar{Y}, p}, \text{ev}_{\ell^p(\Omega)}) \leq \text{wdim}_\varepsilon(R_\Omega B_1^{Y, p}, \text{ev}_{\ell^p(\Omega)})$. On the other hand, let $s : R_\Omega B_1^{Y, p} \rightarrow B_1^{Y, p}$ such that $R_\Omega \circ s = \text{Id}$ be determined by an inverse of $R_\Omega Y \rightarrow Y$, then s is a linear map which increases distances. Consequently, $\text{wdim}_\varepsilon(R_\Omega B_1^{Y, p}, \text{ev}_{\ell^p(\Omega)}) \leq \text{wdim}_\varepsilon(B_1^{Y, p}, \text{ev}_{\ell^p(\Omega)})$. Finally, by inclusion $Y \subset \bar{Y}$, we have $\text{wdim}_\varepsilon(B_1^{Y, p}, \text{ev}_{\ell^p(\Omega)}) \leq \text{wdim}_\varepsilon(B_1^{\bar{Y}, p}, \text{ev}_{\ell^p(\Omega)})$. \square

If $[\Gamma_2 : \Gamma_1] = |G| < \infty$, a set $Y \subset \ell^p(\Gamma_2; V)$ is also a set of $\ell^p(\Gamma_1; V^G)$. Indeed, to $y \in \ell^p(\Gamma_2; V)$ one can associate $i(y)$ where $i(y)(\gamma) = (y(\gamma g))_{g \in G} \in V^G$. This operation behaves nicely with \dim_{ℓ^p} .

Proposition 3.11. *Let $\Gamma_1 \subset \Gamma_2$ be amenable groups and $G = \Gamma_2/\Gamma_1$ where $|G| < \infty$, if $Y \subset \ell^p(\Gamma_2; V)$ is seen by restriction as a linear subspace of $\ell^p(\Gamma_1; V^G)$ then $|G| \dim_{\ell^p}(Y, \Gamma_2) = \dim_{\ell^p}(Y, \Gamma_1)$.*

Proof. Let $\{\Omega_i^{(1)}\}$ be a Følner sequence for Γ_1 and let $\{\Omega_i^{(2)}\} = \{\Omega_i^{(1)} G\}$ be the corresponding Følner sequence in Γ_2 . It is then sufficient to see that $(B_1^{\Omega_i^{(2)}}, \text{ev}_{\Omega_i^{(2)}})$ is by construction isometric to $(B_1^{\Omega_i^{(1)}}, \text{ev}_{\Omega_i^{(1)}})$. \square

Let us mention a typical problem when one deals with ℓ^p spaces, for $p \neq 2$, that is the existence of linear subspaces which are not the image of projection (cf. [12] and [17]). A characterization of subspaces of ℓ^p possessing a projection of norm 1 can be found in [10, I.§2]. We shall briefly discuss the case where $Y \subset \ell^p(\Gamma; \mathbb{R})$ is a Γ -invariant linear subspace on which there exists a Γ -equivariant projection, P_Y . Then let $y = P_Y \delta_{e_\Gamma}$ where δ_{e_Γ} is the Dirac mass at $e_\Gamma \in \Gamma$, and let $q \leq p$ be such that $y \in \ell^q(\Gamma; \mathbb{R})$. For a $x \in \ell^p(\Gamma; \mathbb{R})$, write $x = \sum k_\gamma \delta_\gamma$. By linearity and Γ -equivariance of P_Y ,

$$P_Y x = P_Y \sum_{\gamma \in \Gamma} k_\gamma \delta_\gamma = \sum_{\gamma \in \Gamma} k_\gamma P_Y(\gamma \delta_{e_\Gamma}) = \sum_{\gamma \in \Gamma} k_\gamma \gamma y$$

Thus $(P_Y x)(e_\Gamma) = \sum_{\gamma \in \Gamma} k_\gamma y(\gamma^{-1})$. Taking $k_\gamma = |y(\gamma^{-1})|^{\frac{q}{p}-1} y(\gamma^{-1})$, it appears that $(P_Y x)(e_\Gamma) = \sum |y(\gamma^{-1})|^{\frac{q}{p}+1}$. This forces $\frac{q}{p} + 1 \leq q$, in other words $q \leq p'$ (where p' is the conjugate exponent to p). When $p > \frac{p}{2}$, the existence of such a projection means that there exists in Y an element of $\ell^{p'}(\Gamma; \mathbb{R})$, which is quite restrictive.

4 Further properties in special cases

We now discuss property P7, that is if Y is non-trivial then $\dim_{\ell^p} Y$ is positive. This question is difficult as an intuitive proof only works for $p = 1$. Before we move to this proof, let us argue why the three following assumptions seem necessary for it to hold: Y must be a linear subspace, Y must be Γ -invariant, and Y must be contained in $\ell^p(\Gamma; V)$ for finite p or in $c_0(\Gamma; V)$ if $p = \infty$. Here are some cases of non-trivial Y for which one of the assumptions does not hold and where \dim_{ℓ^p} is 0.

First, suppose Y is not a linear subspace. In example 3.5 the ℓ^q balls where $q < p$ are shown to have their \dim_{ℓ^p} equal to 0. Alternatively, one could also take Y to be the subset of $\ell^\infty(\Gamma; V)$ given by function with support of cardinality less than k (for a fixed $k \in \mathbb{Z}_{>0}$).

Second, if Y is a linear subspace of $\ell^\infty(\Gamma; V)$ but is not Γ -invariant, it could be of finite dimension, and consequently \dim_{ℓ^p} will be trivial.

Last, when p is finite, the existence of a $y \in Y$ whose ℓ^p norm is finite is only guaranteed if $Y \subset \ell^p$. Without this assumption, it could happen that $Y \cap B_r^{\ell^p(\Gamma; V)} = \{0\}, \forall r$. On the other hand, if $p = \infty$, take $Y \subset \ell^\infty(\Gamma; V)$ the (Γ -invariant) line generated by a constant function y (i.e. such that $\exists v \in V, \forall \gamma \in \Gamma, y(\gamma) = v$). Y is 1-dimensional, and consequently $\dim_{\ell^\infty} Y = 0$. But Y is not trivial. However, the question for a Γ -invariant linear subspace $Y \subset c_0(\Gamma; V)$ remains interesting.

Fortunately, in the ℓ^1 case things can be proved without difficulties. As noted before this method does not extend to $p > 1$.

Proposition 4.1. *Let $Y \subset \ell^1(\Gamma; V)$ be a Γ -invariant linear subspace, then $\dim_{\ell^1} Y = 0$ if and only if Y is trivial.*

Proof. This proof requires some results on amenable groups; these can be found in section 5. If one wants, it is possible to think of Γ as \mathbb{Z}^n and take finite sets to be rectangles.

If Y is trivial then $\dim_{\ell^1} Y$ is obviously 0. Otherwise, let $0 \neq y \in Y$ and renormalize it so that $\|y\|_{\ell^1(\Gamma)} = 1$. For all $\varepsilon \in]0, 1/2[$, $\exists F \subset \Gamma$ finite (which depends on y and ε) such that $\|y\|_{\ell^1(F)} > 1 - \varepsilon$ (and consequently $\|y\|_{\ell^1(\Gamma \setminus F)} \leq \varepsilon$). Then let \tilde{y} be identical to y on F and 0 elsewhere.

For i sufficiently big, Ω_i contains a non-empty ρ -quasi-tiling by F (see definition 5.4), since $F \subset \Omega_i$ and $\alpha(\Omega_i; F)$ tends to 0. Applying lemma 5.5 to find translates of F which are ρ -disjoint, where $\rho = 1/2|F|$, we obtain a quasi-tiling whose elements are actually disjoint since $\rho < |F|^{-1}$, and the number of such translates is at least $(1 - \alpha(\Omega_i; F))|\Omega_i|/2|F|$.

Let γ_j for $j \in J_i \subset \mathbb{Z}_{>0}$ be the elements by which the sets F are translated for a ρ -quasi-tiling of Ω_i (since the Ω_i form an increasing sequence and that lemma 5.5 applies to all maximal ρ -quasi-tiling, it can be assumed that the J_i are increasing). Let $V_i = \langle \gamma_j y | j \in J_i \rangle$ be the linear subspace generated by the corresponding translates of y . Trivially $B_1^{V_i} \subset B_1^Y$, and we will construct a map from a ball to $B_1^{V_i}$. Let

$$\begin{aligned} \pi : \ell^1(J_i; \mathbb{R}) &\rightarrow V_i & \tilde{\pi} : \ell^1(J_i; \mathbb{R}) &\rightarrow V_i \\ (a_j)_{j \in J_i} &\mapsto \sum_{j \in J_i} a_j \gamma_j y & \text{and} & & (a_j)_{j \in J_i} &\mapsto \sum_{j \in J_i} a_j \gamma_j \tilde{y} \end{aligned}$$

With these notations,

$$\begin{aligned}\|\tilde{\pi}(a)\|_{\ell^1(\Gamma)} &= \sum_{k \in J_i} \left\| \sum_{j \in J_i} a_j \gamma_j \tilde{y} \right\|_{\ell^1(\gamma_k F)} = \sum_{k \in J_i} \left\| a_k \gamma_k \tilde{y} \right\|_{\ell^1(\gamma_k F)} \\ &= \sum_{k \in J_i} |a_k| \|\tilde{y}\|_{\ell^1(F)} = \|y\|_{\ell^1(F)} \sum_{k \in J_i} |a_k|.\end{aligned}$$

On the other hand,

$$\begin{aligned}\|\tilde{\pi}(a) - \pi(a)\|_{\ell^1(\Gamma)} &= \left\| \sum_{j \in J_i} a_j \gamma_j (\tilde{y} - y) \right\|_{\ell^1(\Gamma)} = \sum_{\gamma \in \Gamma} \left| \sum_{j \in J_i} a_j \gamma_j (\tilde{y} - y) \right| \\ &\leq \sum_{\gamma \in \Gamma} \sum_{j \in J_i} |a_j \gamma_j (\tilde{y}(\gamma) - y(\gamma))| = \sum_{\gamma \in \Gamma} \sum_{j \in J_i} |a_j| |\tilde{y}(\gamma) - y(\gamma)| \\ &= \sum_{j \in J_i} |a_j| \left(\sum_{\gamma \in \Gamma} |\gamma_j (\tilde{y}(\gamma) - y(\gamma))| \right) = \|y\|_{\ell^1(\Gamma \setminus F)} \sum_{j \in J_i} |a_j|.\end{aligned}$$

The last two computations mean that $\|\tilde{\pi}(a)\|_{\ell^1(\Gamma)} = \|y\|_{\ell^1(F)} \|a\|_{\ell^1(J_i)}$ and $\|\tilde{\pi}(a) - \pi(a)\|_{\ell^1(\Gamma)} \leq \|y\|_{\ell^1(\Gamma \setminus F)} \|a\|_{\ell^1(J_i)}$. Thus

$$\left(\|y\|_{\ell^1(F)} - \|y\|_{\ell^1(\Gamma \setminus F)} \right) \|a\|_{\ell^1(J_i)} \leq \|\pi(a)\|_{\ell^1(\Gamma; V)} \leq \left(\|y\|_{\ell^1(F)} + \|y\|_{\ell^1(\Gamma \setminus F)} \right) \|a\|_{\ell^1(J_i)}$$

This means that $(B_1^Y, \mathcal{E}_{\ell^1(\Omega_i)})$ contains, with a controlled distortion, a ℓ^1 ball (with its ℓ^1 metric) of radius 1 and of dimension $\frac{1}{2|F|}(1 - \alpha(\Omega_i; F))|\Omega_i|$, whence

$$\begin{aligned}\dim_{\ell^1} Y &= \lim_{\varepsilon' \rightarrow 0} \limsup_{i \rightarrow \infty} \frac{\text{wdim}_{\varepsilon'}(B_1^Y, \mathcal{E}_{\ell^1(\Omega_i)})}{|\Omega_i|} \\ &\geq \lim_{\varepsilon' \rightarrow 0} \limsup_{i \rightarrow \infty} \frac{1}{2|F|}(1 - \alpha(\Omega_i; F)) = \frac{1}{2|F|}.\end{aligned}$$

As required $\dim_{\ell^1} Y > 0$. \square

This result can be extended to $p > 1$ in the special case that $Y \subset \ell^p(\Gamma; V)$ contains an element in ℓ^1 (in particular an element of finite support). By P6, positivity (P7) is also true for $p = 2$. Positivity means that if one looks at a p -summable two-sided sequence $y \in \ell^p(\mathbb{Z}; \mathbb{R})$, the dimension of the space generated by y and sequences obtained by shifting y up to n times left or right grows linearly with n . The above result is a simple consequence that this is true for $p = 1$, and one is then lead to ask if this can be true for other values of $p \neq \infty$ or in c_0 .

Even if we cannot show continuity, the following example is worthy of interest. The sequence of vector subspaces discussed there will not satisfy the continuity property (P9). This is quite unfortunate, as ℓ^1 is among the few cases where positivity can be shown.

Example 4.2: We exhibit a decreasing sequence of closed linear subspaces of $\ell^1(\mathbb{Z}, \mathbb{R})$, $\{Y_i\}$, such that

$$1 = \lim_{i \rightarrow \infty} \dim_{\ell^1} Y_i \neq \dim_{\ell^1} \bigcap_{i \rightarrow \infty} Y_i = 0.$$

Define $\forall k \in \mathbb{Z}_{>0}, \pi_k : \ell^1(\mathbb{Z}; \mathbb{R}) \rightarrow \ell^\infty(\mathbb{Z}/k\mathbb{Z}; \mathbb{R})$ in the following way: for $n \in \mathbb{Z}/k\mathbb{Z}$

$$\pi_k(x)(n) = \sum_{i \equiv n \pmod k} x(i).$$

Continuous linear maps between Banach spaces have a closed kernel (for τ , the norm topology in ℓ^1), thus $Y_j = \bigcap_{k=1}^j \ker \pi_k$ is a decreasing sequence of closed sets (for τ). To compute \dim_{ℓ^1} , choose the Følner sequence $\Omega_i = [-i, i] \cap \mathbb{Z}$. For a $N \in \mathbb{N}$, let $y_N \in Y_1$ be such that $y_N(0) = 1/2$, $y_N(N) = -1/2$ and which is zero elsewhere. Let $N_j = \text{lcm}(1, 2, \dots, j)$. For all j , $y_{N_j} \in B_1^{Y_j}$. These elements give a map $(B_{1/2}^{\ell^1(\mathbb{Z}; \mathbb{R})}, \nu_{\ell^1(\Omega)})$ to $(B_1^{Y_j}, \nu_{\ell^1(\Omega)})$ which possesses fibers of “diameter” 0. They are defined as follows, $y \in B_{1/2}^{\ell^1(\mathbb{Z}; \mathbb{R})}$ is restricted to Ω then extended by 0 outside Ω . Then, let $k \in \mathbb{Z}_{>0}$ be such that kN_j is bigger than the diameter of $\Omega \subset \mathbb{Z}$, then $\tilde{y}(m) = \sum_{n \in \Omega} 2y_{kN_j}(m-n)y(n)$ is an element of $B_1^{Y_j}$. Thence $\dim_{\ell^1} Y_j \geq 1$, and as the other inequality is automatic, $\dim_{\ell^1} Y_j = 1$.

We claim that $Y_\infty = \bigcap Y_j = \{0\}$. If this were false, then a non-trivial element $y \in Y_\infty$ would have the property that

$$\forall i \in \mathbb{Z}, \forall n \in \mathbb{Z}, -y(i) = \sum_{0 \neq k \in \mathbb{Z}} y(i+kn).$$

To get a contradiction, take the limit when $n \rightarrow \infty$ and show that it is equal to 0. First we normalize y so that it is of norm 1 and suppose that $|y(i)| > \delta$ for some i . As an absolutely convergent sequence, y should be concentrated on some set: there exists n_δ such that $\|y\|_{\ell^1(\Omega_{n_\delta})} \geq 1 - \delta/2$. However when $n > 2n_\delta + 1$

$$|y(i)| = \left| \sum_{0 \neq k \in \mathbb{Z}} y(i+kn) \right| \leq \delta/2,$$

which is a contradiction. Thus $\dim_{\ell^1} Y_\infty = 0$ whereas $\lim_{n \rightarrow \infty} \dim_{\ell^1} Y_n = 1$.

Such an possibility is fortunately confined to ℓ^1 ; more generally the above construction can be described as follows. Let $\Gamma' \subset \Gamma$ be a subgroup of finite index. Let $\pi: Y \rightarrow W$ be a Γ' -invariant linear map from $Y \subset \ell^p(\Gamma; V)$ to a finite dimensional vector space W . Then the existence of such a map implies the existence of $\dim W$ elements of $\ell^{p'}(\Gamma; V^*)$ which are invariant by Γ' . This is impossible if $p' \neq \infty$, as such elements would not be decreasing at infinity. For such spaces to exist, p' must be ∞ .

5 P1 and Ornstein-Weiss' Lemma

The aim of this section is to show independence (P1) on the choice Følner sequence. This will be achieved by extending Ornstein-Weiss' lemma to meet our needs.

Theorem 5.1. *Let Γ be a discrete amenable group, and let $a: \mathbb{R}_{\geq 0} \times \mathcal{P}_{\text{finite}}(\Gamma) \rightarrow \mathbb{R}_{\geq 0}$ be a function such that, $\forall \Omega, \Omega' \subset \Gamma$ are finite and $\forall \varepsilon \in \mathbb{R}_{>0}$*

- (a) *a is Γ -invariant, i.e.* $\forall \gamma \in \Gamma, \quad a(\varepsilon, \gamma\Omega) = a(\varepsilon, \Omega)$
- (b) *a is decreasing in ε , i.e.* $\forall \varepsilon' \leq \varepsilon, \quad a(\varepsilon', \Omega) \geq a(\varepsilon, \Omega)$
- (c) *a is K -sublinear in Ω , i.e.* $\exists K \in \mathbb{R}_{>0}, \quad a(\varepsilon, \Omega) \leq K|\Omega|$
- (d) *a is c -subadditive in Ω , i.e.* $\exists c \in]0, 1], \quad a(\varepsilon, \Omega \cup \Omega') \leq a(c\varepsilon, \Omega) + a(c\varepsilon, \Omega')$

then, for any Følner sequence $\{\Omega_i\}$,

$$\lim_{\varepsilon \rightarrow 0} \limsup_{i \rightarrow \infty} \frac{a(\varepsilon, \Omega_i)}{|\Omega_i|} = \lim_{\varepsilon \rightarrow 0} \liminf_{i \rightarrow \infty} \frac{a(\varepsilon, \Omega_i)}{|\Omega_i|}.$$

Furthermore, these limits are independent of the chosen sequence $\{\Omega_i\}$.

Remark that the K -sublinear hypothesis (c) is equivalent to another statement. Indeed, using c -subadditivity (d), Γ -invariance (a) and monotonicity in ε (b), for all Ω , $a(\varepsilon, \Omega) \leq a(c^{|\Omega|}\varepsilon, e_\Gamma)|\Omega|$ where $e_\Gamma \in \Gamma$ is the neutral element. Thus if (a), (b) and (d) hold, then (c) $\Leftrightarrow \lim_{\varepsilon \rightarrow 0} a(\varepsilon, e_\Gamma) < \infty$.

This understood, the previous theorem is a generalization of the Ornstein-Weiss lemma. Indeed, the assumptions of the latter, are that $a(\varepsilon, \Omega) = a(\Omega)$ is sub-additive: then monotonicity (b) always hold, being K -sublinear (c) is automatic (see above), and c -subadditivity (d) is equivalent to usual subadditivity ($c = 1$).

The proof of theorem 5.1 being quite technical, let us first show why P1 is a consequence of this theorem.

Corollary 5.2. \dim_{ℓ^p} is independent of the choice of Følner sequence.

Proof. It suffices to prove that for any $Y \subset \ell^\infty(\Gamma; V)$ a Γ -invariant set, theorem 5.1 can be invoked, where $a(\varepsilon, \Omega) = \text{wdim}_\varepsilon(B_r^{Y,p}, \text{ev}_{\ell^p(\Omega)})$. Here is why:

- (a) By Γ -invariance of Y .
- (b) As wdim_ε is decreasing in ε (cf. proposition 3.4.a).
- (c) Using proposition 3.1.b $\text{wdim}_\varepsilon(B_r^{\ell^p(\Omega)}, \text{ev}_{\ell^p(\Omega)}) \leq |\Omega|\dim V$. Then proposition 3.4.a allows to conclude as $(B_r^{Y,p}, \text{ev}_{\ell^p(\Omega)})$ can be sent without reducing distance by a continuous map to $(B_r^{\ell^p(\Omega)}, \text{ev}_{\ell^p(\Omega)})$
- (d) Let $\pi_i : \ell^p(\Gamma, V) \rightarrow \ell^p(\Omega_i, V)$, then

$$\pi_1 \times \pi_2 : (B_r^{Y,p}, \text{ev}_{\ell^p(\Omega)}) \rightarrow (B_r^{Y,p}, \text{ev}_{\ell^p(\Omega_1)}) \times_p (B_r^{Y,p}, \text{ev}_{\ell^p(\Omega_2)})$$

is a linear map that does not reduce distances. Applying proposition 3.4.a and 3.4.c, yields

$$\text{wdim}_{2^{1/p}\varepsilon}(B_r^{Y,p}, \text{ev}_{\ell^p(\Omega)}) \leq \text{wdim}_\varepsilon(B_r^{Y,p}, \text{ev}_{\ell^p(\Omega_1)}) + \text{wdim}_\varepsilon(B_r^{Y,p}, \text{ev}_{\ell^p(\Omega_2)}).$$

Thence, we conclude that $a(\varepsilon, \Omega)$ is $2^{-1/p}$ -subadditive. \square

The following notations and definitions will be required in our arguments. The original proof of the Ornstein-Weiss lemma can be found in [13]. The proof that can be better adapted to our case is however that of [6, §1.3.1] (also explained in [8]).

Definition 5.3. Let Γ be a group, let $F \subset \Gamma$ be such that $e_\Gamma \in F$ then the F -boundaries of $\Omega \subset \Gamma$ are defined as

$$\begin{aligned}
\partial_F^+ \Omega &= \{\gamma \notin \Omega \mid \gamma F \cap \Omega \neq \emptyset \text{ and } \gamma F \cap \Omega^c \neq \emptyset\} = F^{-1} \Omega \cap \Omega^c && \text{(outer } F\text{-boundary)} \\
\partial_F^- \Omega &= \{\gamma \in \Omega \mid \gamma F \cap \Omega \neq \emptyset \text{ and } \gamma F \cap \Omega^c \neq \emptyset\} = F^{-1} \Omega^c \cap \Omega && \text{(inner } F\text{-boundary)} \\
\partial_F \Omega &= \{\gamma \in \Gamma \mid \gamma F \cap \Omega \neq \emptyset \text{ and } \gamma F \cap \Omega^c \neq \emptyset\} = \partial_F^+ \Omega \cup \partial_F^- \Omega && \text{(} F\text{-boundary)} \\
\text{int}_F \Omega &= \{\gamma \in \Gamma \mid \gamma F \subset \Omega\} = \Omega \setminus \partial_F \Omega && \text{(} F\text{-interior)} \\
\text{fer}_F \Omega &= \{\gamma \in \Gamma \mid \gamma F \cap \Omega \neq \emptyset\} = \Omega \cup \partial_F^+ \Omega && \text{(} F\text{-closure)}.
\end{aligned}$$

Moreover, let $|\cdot|$ denote a measure on Γ . The relative amenability function will be defined as $\alpha(\Omega; F) = \frac{|\partial_F \Omega|}{|\Omega|}$, given that these numbers are finite.

Before we move on to technical results, observe that the Følner conditions implies that $\alpha(\Omega_i; F) \rightarrow 0$ for any finite set F and any Følner sequence $\{\Omega_i\}$. Another useful property is that if $F' \subset F$, then $\alpha(\Omega; F') \leq \alpha(\Omega; F)$ since $\partial_{F'} \Omega \subset \partial_F \Omega$. We start by showing covering properties of big sets by smaller sets.

Definition 5.4. Let $\varepsilon \in]0, 1[$. Subsets F_i of finite measure of Γ will be said ε -disjoint if there exists $F'_i \subset F_i$ which are disjoint and such that $|F'_i| \geq (1 - \varepsilon)|F_i|$ and $\cup F'_i = \cup F_i$.

A subset of finite measure Ω will be said to admit an ε -quasi-tiling by the subsets F_i if

- (a) $F_i \subset \Omega$,
- (b) the F_i are ε -disjoint,

Here is a first lemma which studies the proportion of a set Ω covered by an ε -quasi-tiling of translates of another set F .

Lemma 5.5. Let Γ be a discrete group endowed with the counting measure, denoted by $|\cdot|$. Let $\Omega \subset \Gamma$ and $e_\Gamma \in F \subset \Gamma$ both finite sets and such that $\alpha(\Omega; F) < 1$. Let $\{\gamma_i\}_{1 \leq i \leq k}$ be a maximal sequence of elements of Γ such that the $\gamma_i F$ form an ε -quasi-tiling of Ω .

Let $U_F^i = \bigcup_{j=1}^i \gamma_j F$, then

$$\frac{|U_F^k|}{|\Omega|} \geq \varepsilon(1 - \alpha(\Omega; F)).$$

Proof. (This proof corresponds to the first part of the proof of the Ornstein-Weiss lemma in [6, §1.3.1].) We shall use this general fact:

$$\begin{aligned}
\int_\Gamma |G_1 \cap \gamma G_2| d\mu(\gamma) &= \int_\Gamma \int_\Gamma \mathbb{1}_{G_1 \cap \gamma G_2}(\gamma') d\mu(\gamma') d\mu(\gamma) \\
&= \int_\Gamma \int_\Gamma \mathbb{1}_{G_1}(\gamma') \mathbb{1}_{G_2}(\gamma' \gamma^{-1}) d\mu(\gamma) d\mu(\gamma') \\
&= \int_\Gamma \mathbb{1}_{G_1}(\gamma') \left(\int_\Gamma \mathbb{1}_{G_2}(\gamma' \gamma^{-1}) d\mu(\gamma) \right) d\mu(\gamma') \\
&= \int_\Gamma \mathbb{1}_{G_1}(\gamma') |G_2| d\mu(\gamma') \\
&= |G_1| |G_2|.
\end{aligned}$$

Thus,

$$|\text{int}_F \Omega|^{-1} \int_{\text{int}_F \Omega} |U_F^k \cap \gamma F| \mathfrak{d}\mu(\gamma) \leq |\text{int}_F \Omega|^{-1} \int_{\Gamma} |U_F^k \cap \gamma F| \mathfrak{d}\mu(\gamma) \leq (1 - \alpha(\Omega; F))^{-1} |\Omega|^{-1} |U_F^k| |F|.$$

Clearly, $|U_F^{i-1} \cap \gamma_i F| \leq \varepsilon |F|$, as the $\gamma_i F$ are ε -disjoint. On the other hand, maximality of k implies that $\forall \gamma \in \text{int}_F \Omega, |U_F^k \cap \gamma F| \geq \varepsilon |F|$. We then observe that

$$|\text{int}_F \Omega|^{-1} \int_{\text{int}_F \Omega} |U_F^k \cap \gamma F| \mathfrak{d}\mu(\gamma) \geq \varepsilon |F|.$$

Consequently, $\varepsilon(1 - \alpha(\Omega; F)) \leq |U_F^k|/|\Omega|$. \square

Note that the quasi-tiling can be empty if $\alpha(\Omega; F) = 1$. More precisely, the proof actually works for $\alpha^-(\Omega; F) = \frac{|\partial_F^- \Omega|}{|\Omega|}$ instead of α . It has the advantage that $\text{int}_F \Omega \neq \emptyset$ implies that $\alpha^-(\Omega; F) < 1$ and the quasi-tiling is non-empty. In any case, in the upcoming applications, F will always be contained in Ω . The three following lemmas are technical ingredients which will be used in the proof of the generalisation of the Ornstein-Weiss lemma.

Lemma 5.6. *Let $\Omega' \subset \Omega \subset \Gamma$ and $F \subset \Gamma$ be finite. Suppose that there exists ε such that $|\Omega \setminus \Omega'| \geq \varepsilon |\Omega|$, then*

$$\alpha(\Omega \setminus \Omega'; F) \leq \frac{\alpha(\Omega'; F) + \alpha(\Omega; F)}{\varepsilon}.$$

Proof. Since $|\partial_F(\Omega \setminus \Omega')| \leq |\partial_F \Omega| + |\partial_F \Omega'| = \alpha(\Omega; F)|\Omega| + \alpha(\Omega'; F)|\Omega'|$, and that $|\Omega \setminus \Omega'| \geq \varepsilon |\Omega| \geq \varepsilon |\Omega'|$, a substitution yields

$$\alpha(\Omega \setminus \Omega'; F) = \frac{|\partial_F(\Omega \setminus \Omega')|}{|\Omega \setminus \Omega'|} \leq \frac{\alpha(\Omega; F)|\Omega|}{\varepsilon |\Omega|} + \frac{\alpha(\Omega'; F)|\Omega'|}{\varepsilon |\Omega'|}. \quad \square$$

Lemma 5.7. *Let $F \subset \Gamma$ be finite, and let $\{D_i\}_{1 \leq i \leq n}$ be an ε -disjoint family of subsets. Then*

$$\alpha(\cup D_i; F) \leq \frac{\max(\alpha(D_i; F))}{1 - \varepsilon}$$

Proof. Since $\partial_F(\cup D_i) \subset \cup \partial_F D_i$, we obtain that

$$|\partial_F(\cup D_i)| \leq \sum |\partial_F D_i| \leq \sum \alpha(D_i; F) |D_i| \leq \max(\alpha(D_i; F)) \sum |D_i|$$

However $(1 - \varepsilon) \sum |D_i| \leq |\cup D_i|$ as they are ε -disjoint. Thus

$$\alpha(\cup D_i; F) = \frac{|\partial_F(\cup D_i)|}{|\cup D_i|} \leq \frac{\max(\alpha(D_i; F))}{1 - \varepsilon}. \quad \square$$

The last lemma is an adaptation of a useful property of \mathbb{Z} to general amenable group: for the typical Følner sequence for \mathbb{Z} $I_i = [-i, i]$, any sufficiently big interval in this family is covered (except for small bits) by translates of the I_i .

Lemma 5.8. *Let $\{F_i\}$ be a Følner sequence, let $\delta \in]0, 1/2[$. Then there exists a subsequence (which depends on δ) $\{F_{n_i}\}$, an integer $N(\delta)$, and a sequence of integers $\{k_i\}_{1 \leq i \leq N}$ such that for all set Ω which contains F_{n_N} and satisfies $\alpha(\Omega, F_{n_N}) \leq 2\delta^{2N}$ there exists a family \mathcal{G} of δ -disjoint sets such that $|\bigcup_{F \in \mathcal{G}} F| \geq (1 - \delta)|\Omega|$ and \mathcal{G} consists in k_i translates of the sets F_{n_i}*

Proof. (We write the argument of [6, §1.3.1], see also [8]; the original result can be found in [13].) In order to better show how the constants enter the proof, we denote $\varepsilon_1 = \delta^{2N}$, $\varepsilon_2 = 2\delta^{2N}$ and $\rho = \delta$. First, $\forall \varepsilon_1 \in]0, 1[$, it is possible to refine the sequence $\{F_i\}$ to have

$$\alpha(F_{i+1}, F_i) \leq \varepsilon_1.$$

Now, let $\Omega^{(1)} = \Omega$ so that $\alpha(\Omega^{(1)}, F_n) \leq \varepsilon_2$, where n will be determined later on. We will cover $\Omega^{(1)}$ to a proportion of $1 - \delta$ by almost disjoint translates of the F_i , where $1 \leq i \leq n$, in n steps (or less). For any $\rho \in]0, \frac{1}{2}[$, lemma 5.5 gives a ρ -quasi-tiling of $\Omega^{(1)}$ by k_n translates of F_n such that $|U_{F_n}^{k_n}| \geq \rho(1 - \varepsilon_2)|\Omega^{(1)}|$. Let $\Omega^{(2)} = \Omega^{(1)} \setminus U_{F_n}^{k_n}$, then $|\Omega^{(2)}| \leq (1 - \rho + \varepsilon_2\rho)|\Omega^{(1)}|$.

If $|\Omega^{(2)}| \leq \delta|\Omega^{(1)}|$ the goal is achieved and there is no need to continue. Otherwise, lemma 5.6 then lemma 5.7 shows that

$$\alpha(\Omega^{(2)}, F_{n-1}) \leq \frac{1}{\delta}(\varepsilon_1 + \alpha(U_{F_n}^{k_n}; F_{n-1})) \leq \frac{1}{\delta}(\varepsilon_1 + \frac{\varepsilon_1}{1 - \rho}) \leq 3\frac{\varepsilon_1}{\delta}.$$

It is now possible to recover $\Omega^{(2)}$ by a ρ -quasi-tiling of k_{n-1} translates of F_{n-1} in such a way that $|U_{F_{n-1}}^{k_{n-1}}| \geq \rho(1 - 3\frac{\varepsilon_1}{\delta})|\Omega^{(2)}|$. We now have a set $\Omega^{(3)}$ such that

$$|\Omega^{(3)}| \leq (1 - \rho - 3\rho\frac{\varepsilon_1}{\delta})|\Omega^{(2)}| \leq (1 - \rho + \varepsilon_2\rho)(1 - \rho + \rho\frac{3\varepsilon_1}{\delta})|\Omega^{(1)}|$$

We will now take $\varepsilon_2 = 2\varepsilon_1$. Proceeding by induction, as long as $|\Omega^{(i-1)}| \geq \varepsilon|\Omega^{(1)}|$, the set $\Omega^{(i)}$ (for $1 \leq i \leq n$) will have the following properties:

1. $\alpha(\Omega^{(i)}, F_{n-i+1}) \leq (1 + i)\varepsilon_1/\delta^{i-1}$
2. $U_{F_{n-i+1}}^{k_{n-i+1}}$ is a ρ -quasi-tiling of $\Omega^{(i)}$ by translates of F_{n-i+1}
3. If $\Omega^{(i+1)} = \Omega^{(i)} \setminus U_{F_{n-i+1}}^{k_{n-i+1}}$ then $|\Omega^{(i+1)}| \leq |\Omega^{(1)}| \prod_{j=1}^i (1 - \rho(1 - (1 + j)\varepsilon_1/\delta^{j-1}))$

Since it is not possible to hope that this process terminates before $i = n$, it remains to be checked that if n is big enough, we still get a quasi-tiling that covers $(1 - \delta)|\Omega^{(1)}|$ elements. To achieve this, observe that the product in the third property above can be bounded if $i = n$ by

$$\prod_{j=1}^n (1 - \rho(1 - (1 + j)\varepsilon_1/\delta^{j-1})) \leq (1 - \rho(1 - (1 + n)\varepsilon_1/\delta^{n-1}))^n.$$

For $\varepsilon_1 = \delta^{2n}$, the right-hand term tends to 0 when n tends to ∞ . Thus, $\exists N(\delta, \rho)$ such that if $\varepsilon_1 = \delta^{2N}$ translates of F_j (where $1 \leq j \leq N$) form a ρ -quasi-tiling of any set $\Omega^{(1)}$ such that $\alpha(\Omega^{(1)}; F_N) \leq \delta^{2N}$.

We substitute as promised $\rho = \delta$ to have: for any fixed δ , we choose a subsequence whose members satisfy $\alpha(F_{n_{i+1}}, F_{n_i}) \leq \delta^{2N}$ where N is such that

$$(1 - \delta(1 - (1 + N)\delta^{N+1}))^N < \delta.$$

Then successive applications of lemma 5.5 give the required translates of F_{n_i} . \square

We are now ready to prove the main result of this section. At a first reading, this proof might be easier to understand with $\Gamma = \mathbb{Z}$ in mind (taking $\Omega_n = [-n, n] \cap \mathbb{Z}$).

Proof of theorem 5.1. Let us first introduce some notations for the functions given by pointwise convergence and their limits. Let $\{\Omega_i^{+, \varepsilon}\}$ and $\{\Omega_i^{-, \varepsilon}\}$ be subsequences of $\{\Omega_i\}$ such that

$$\lim_{i \rightarrow \infty} \frac{a(\varepsilon, \Omega_i^{+, \varepsilon})}{|\Omega_i^{+, \varepsilon}|} = \limsup_{i \rightarrow \infty} \frac{a(\varepsilon, \Omega_i)}{|\Omega_i|} \quad \text{and} \quad \lim_{i \rightarrow \infty} \frac{a(\varepsilon, \Omega_i^{-, \varepsilon})}{|\Omega_i^{-, \varepsilon}|} = \liminf_{i \rightarrow \infty} \frac{a(\varepsilon, \Omega_i)}{|\Omega_i|}.$$

Using (c), these limits are respectively real numbers $l^+(\varepsilon)$ and $l^-(\varepsilon)$ belonging to the interval $[0, K]$. Furthermore, let

$$l^+ := \lim_{\varepsilon \rightarrow 0} l^+(\varepsilon) \quad \text{and} \quad l^- := \lim_{\varepsilon \rightarrow 0} l^-(\varepsilon).$$

Trivially, $l^+(\varepsilon) \geq l^-(\varepsilon)$, but nothing forces $l^\pm(0) = l^\pm$ (in general, equality is not expected). If we try to use the usual argument directly, a problem arises due to the c -subadditivity. Indeed, taking a sequence which converges to $l^+(\varepsilon)$ and decomposing it using another sequence which converges to $l^-(\varepsilon)$ by subadditivity will fail. A factor of c will appear in front of the ε (see (d)), and this would force to pass from the sequence $\Omega_i^{-, \varepsilon}$ to $\Omega_i^{-, c\varepsilon}$ at each step. Diagonal arguments settle this problem. Let

$$b_i(\varepsilon) = \frac{a(\varepsilon, \Omega_i^{-, 1/i})}{|\Omega_i^{-, 1/i}|}.$$

This is a sequence of bounded decreasing functions defined for $\varepsilon \in [0, 1]$ with value in $[0, K]$. Using (one of) Helly's theorem (cf. [7, §36.5 theorem 5, p.372]), there exists a subsequence n_i which possesses a limit at each point. We briefly recall how this subsequence is obtained. First, a sequence $\{r_k\}_{k \geq 1}$ of dense rational number in $[0, 1]$ is taken. Since the $b_i(\varepsilon)$ are bounded, let $n_i^{(j)}$ be the subsequence which converges at r_k for $1 \leq k \leq j$. The diagonal sequence $n_i = n_i^{(i)}$ converges at each r_k , and since the functions $b_i(\varepsilon)$ are decreasing, the function $l^H(\varepsilon) = \lim_{i \rightarrow \infty} b_{n_i}(\varepsilon)$ which is *a priori* only defined for the r_k is also decreasing. It remains to be checked that $l^H(\varepsilon)$ extended at all the points of $[0, 1]$ by approximating by a sequence of increasing r_k is the actual

limit of the subsequence n_i (see the above reference for details). Let us show that $\lim_{\varepsilon \rightarrow 0} l^H(\varepsilon) = l^-$. This follows from

$$\begin{aligned} \forall \delta > 0, \exists N_1(\delta) \text{ such that } N_1(\delta) < i &\Rightarrow |b_{n_i}(\frac{1}{n_i}) - l^H(\frac{1}{n_i})| < \delta; \\ \forall \delta > 0, \exists N_2(\delta) \text{ such that } N_2(\delta) < i &\Rightarrow |b_{n_i}(\frac{1}{n_i}) - l^-(\frac{1}{n_i})| < \delta; \\ \forall \delta > 0, \exists N_3(\delta) \text{ such that } N_3(\delta) < i &\Rightarrow |l^-(\frac{1}{n_i}) - l^-| < \delta. \\ l^H(\varepsilon) \text{ is decreasing in } \varepsilon &\Rightarrow \lim_{\varepsilon \rightarrow 0} l^H(\varepsilon) = \lim_{i \rightarrow \infty} l^H(\frac{1}{n_i}) \end{aligned}$$

These four assertions are respectively consequences of the definition of l^H , the choice of $\Omega_i^{-,1/i}$, the definition of l^- , and the fact that a limit that exists (thanks to monotonicity) is achieved by any sequence. We shall now show that

$$\forall \delta > 0, l^+(\varepsilon) \leq \lim_{\varepsilon' \rightarrow 0} l^H(\varepsilon') + \delta = l^- + \delta.$$

The argument is in essence the same as for subadditive sequences of real numbers: lemma 5.8 plays the role of the decomposition $n = kn' + r$ and c -subadditivity (d) forces $\varepsilon \rightarrow 0$.

Let $\delta \in]0, \frac{1}{2}[$. Denote by $F_i = \Omega_{n_i}^{-,1/n_i}$. It is possible to refine this sequence so that

$$a(\varepsilon, F_i)/|F_i| \leq l^H(\varepsilon) + \delta.$$

Applying lemma 5.8 gives an ε -quasi-tiling (which does not cover a set of proportion δ) of any sufficiently big set by translates of the F_i . Since $\{\Omega_i^{+, \varepsilon}\}$ is also a Følner sequence, for i big enough, lemma 5.8 applies to each element. Take $\Omega = \Omega_i^{+, \varepsilon}$, denote $\gamma_{F_j; m} F_j$ the k_j translates of F_j obtained ($m = 1, \dots, k_j$), and let i_0 be such that $|\Omega^{(i_0)}| \leq \delta |\Omega^{(1)}|$. Thanks to repeated use of c -subadditivity (d), we have that

$$\begin{aligned} a(\varepsilon, \Omega^{(1)}) &\leq \sum_{m=1}^{k_n} a(c^m \varepsilon, \gamma_{F_n; m} F_n) + a(c^{k_n} \varepsilon, \Omega^{(2)}) \\ &\leq \dots \\ &\leq \sum_{i=n-i_0}^n \left(\sum_{m=1}^{k_i} a(c^{\kappa_i + m} \varepsilon, \gamma_{F_i; m} F_i) \right) + a(c^{\kappa_{i_0}} \varepsilon, \Omega^{(i_0)}), \end{aligned}$$

where $\kappa_i = \sum_{j=n-i}^n k_n$. Using Γ -invariance (a), the fact that these functions are decreasing in ε (b), and the K -sublinear property (c), this inequality yields

$$a(\varepsilon, \Omega^{(1)}) \leq \sum_{i=n-i_0}^n \left(\sum_{m=1}^{k_i} a(c^{\kappa_{i_0}} \varepsilon, F_i) \right) + K |\Omega^{(i_0)}|.$$

On one hand, $|\Omega^{(i_0)}| \leq \delta |\Omega^{(1)}|$ and $\frac{a(c^{\kappa_{i_0}} \varepsilon, F_i)}{|F_i|} \leq l^H(c^{\kappa_{i_0}} \varepsilon) + \delta$. Thence,

$$\begin{aligned} \frac{a(\varepsilon, \Omega^{(1)})}{|\Omega^{(1)}|} &\leq \sum_{i,m} \frac{a(c^{\kappa_{i_0}} \varepsilon, F_i)}{|F_i|} \frac{|\gamma_{F_i; m} F_i|}{|\Omega^{(1)}|} + K \frac{|\Omega^{(i_0)}|}{|\Omega^{(1)}|} \\ &\leq (l^H(c^{\kappa_{i_0}} \varepsilon) + \delta) \sum_{i,m} \frac{|\gamma_{F_i; m} F_i|}{|\Omega^{(1)}|} + K \delta \end{aligned}$$

On the other hand, the $\{\gamma_{F_i;m}F_i\}$ are δ -disjoint. Thus $(1 - \delta)\sum |\gamma_{F_i;m}F_i| \leq |\cup \gamma_{F_i;m}F_i| \leq |\Omega^{(1)}|$. This shows that

$$\frac{a(\varepsilon, \Omega_j^{+,\varepsilon})}{|\Omega_j^{+,\varepsilon}|} \leq (l^H(c^{\kappa_{i_0}}) + \delta) \sum_{i,l} \frac{|\gamma_{F_i;l}F_i|}{|\Omega^{(1)}|} + K\delta \leq \frac{l^H(c^{\kappa_{i_0}}) + \delta}{1 - \delta} + K\delta,$$

For all $\Omega_j^{+,\varepsilon}$ big enough, where κ_{i_0} depends on $\Omega_j^{+,\varepsilon}$. Since $l^H(\varepsilon)$ is decreasing and $\lim_{\varepsilon \rightarrow 0} l^H(\varepsilon) = l^-$, taking the limit when j and $\kappa_{i_0} \rightarrow \infty$ is not a problem:

$$l^+(\varepsilon) \leq l^- + \delta(K + l^- + 1).$$

We have shown that $l^+ = l^-$. To deduce the independence on the choice of sequence, notice that given two Følner sequences $\{\Omega_i\}$ and $\{\Omega'_i\}$, the sequence $\{\tilde{\Omega}_i\}$ whose elements alternate between those of the two former sequences will also possess a limit. The limit obtained with $\{\tilde{\Omega}_i\}$ must be equal to the one taken via $\{\Omega_i\}$ or $\{\Omega'_i\}$. \square

Appendix A Von Neumann's dimension and \dim_{ℓ^2}

We recall an argument of Gromov (see [6, §1.12]) that relates von Neumann to the semi-axis of ellipsoids and thus showing that \dim_{ℓ^2} is indeed von Neumann dimension. We briefly review the definition of the latter.

Let $Y \subset \ell^2(\Gamma; \mathbb{R}^s) \subset (\mathbb{R}^s)^\Gamma$ be a Γ -invariant linear subspace, $\forall \Omega \subset \Gamma$ we define the operator $R_\Omega : Y \rightarrow (\mathbb{R}^s)^\Omega_{\ell^2}$ by restriction to Ω : $y \mapsto y|_\Omega$. Its adjoint $R_\Omega^* : (\mathbb{R}^s)^\Omega_{\ell^2} \rightarrow Y$ is the orthogonal projection to Y . To see this, write $R_\Omega(y) = y\mathbb{1}_\Omega$ where $\mathbb{1}_\Omega$ is the characteristic function of Ω , then

$$\langle R_\Omega^*(x), y \rangle := \langle x, R_\Omega y \rangle = \int_\Gamma xy \mathbb{1}_\Omega = \int_\Gamma (\mathbb{1}_\Omega x)y.$$

However this last expression is simply the scalar product of x , extended as a function on all of Γ by 0, with y . Thus, $R_\Omega^*(x)$ is the projection on Y of the extension of x to Γ by 0. In what follows we will omit this inclusion (extension by 0) from $V_{\ell^2}^\Omega$ to $V_{\ell^2}^{\Omega'}$ when $\Omega \subset \Omega'$. Dependence on Ω of R_Ω^* will not be written. A crucial remark is that the invariance of Y by Γ implies that, for $\Omega, \Omega' \subset \Gamma$ finite subsets,

$$\frac{\text{Tr} R_\Omega R^*}{\text{Tr} R_{\Omega'} R^*} = \frac{|\Omega|}{|\Omega'|}.$$

A possible definition of von Neumann dimension (see [11] or [14, §1]) is

$$\dim_{\ell^2}(Y : \Gamma) := |\Omega|^{-1} \text{Tr} R_\Omega R^*$$

for a $\Omega \subset \Gamma$. This quantity is actually independent of the chosen set. The aim of this section is to retrieve this quantity as the wdim of a certain object.

Theorem A.1. (cf. [6, 1.12A]) Let $\Omega_i \subset \Gamma$ be a Følner sequence, let $n_i[a, b]$ be the number of eigenvalues of the operator $R_{\Omega_i}R^*$ (defined relative to Y) contained in the interval $[a, b]$. If $0 < a \leq b < 1$, then

$$\lim_{i \rightarrow \infty} \frac{n_i[a, b]}{|\Omega_i|} = 0$$

Proof. (The proof is with minor differences in notation that of [6].) Since R_{Ω} and R^* are both projections (in ℓ^2), the eigenvalues of $R_{\Omega}R^*$ will be contained in $[0, 1]$. The proof proceeds in three steps.

First, let $x \in \ell^2(\Omega; \mathbb{R}^s)$, it will be called an ε -quasimode of eigenvalue λ for $R_{\Omega}R^*$ if

$$\|R_{\Omega}R^*x - \lambda x\|_{\ell^2} \leq \varepsilon \|x\|_{\ell^2}. \quad (1)$$

If x is such an element, and if its restriction outside Ω is small, more precisely

$$\|R^*x|_{\Gamma \setminus \Omega}\|_{\ell^2} = \|R^*x - R_{\Omega}R^*x\|_{\ell^2} \leq \delta \|x\|_{\ell^2}, \quad (2)$$

then $\lambda(1 - \lambda) \leq 2\varepsilon + \delta$. Indeed, using (1) in (2) yields that $\|R^*x - \lambda x\|_{\ell^2} \leq (\delta + \varepsilon) \|x\|_{\ell^2}$. Since R^* is a projection, $R^*R^* = R^*$ and $\|R^*\| = 1$, whence

$$(1 - \lambda) \|R^*x\|_{\ell^2} = \|R^*x - R^*\lambda x\|_{\ell^2} = \|R^*(R^*x - \lambda x)\|_{\ell^2} \leq \|R^*x - \lambda x\|_{\ell^2} \leq (\delta + \varepsilon) \|x\|_{\ell^2},$$

since the eigenvalues of $R_{\Omega}R^*$ are all contained in $[0, 1]$, $|1 - \lambda| = 1 - \lambda$. Moreover the restriction to Ω can only reduce the norm, $\|R_{\Omega}R^*x\|_{\ell^2} \leq \|R^*x\|_{\ell^2}$. Using (1) anew gives,

$$(1 - \lambda)\lambda \|x\|_{\ell^2} \leq (1 - \lambda) \|R_{\Omega}R^*x\|_{\ell^2} + (1 - \lambda)\varepsilon \|x\|_{\ell^2} \leq (\delta + (2 - \lambda)\varepsilon) \|x\|_{\ell^2}$$

Second, denote by $\Omega^{-\rho} \subset \Omega$ the ρ -interior of Ω , i.e. the set of points with distance at least ρ from a point outside of Ω , where the distance on Γ is the word distance for any fixed generating set. The next argument will consist in showing that most of $x \in \ell^2(\Omega^{-\rho}; V)$ have a small projection to $\Gamma \setminus \Omega$. Precisely, let

$$S_{\rho} = R_{\Gamma \setminus \Omega}R^* = (1 - R_{\Omega})R^* : \ell^2(\Omega^{-\rho}; \mathbb{R}^s) \rightarrow \ell^2(\Gamma \setminus \Omega),$$

then $\text{Tr} S_{\rho}^*S_{\rho} \leq s\beta_0(\rho)|\Omega^{-\rho}|$ where $\beta_0(\rho)$ tends to 0 when $\rho \rightarrow \infty$. The dependence on ρ does not only come from the domain of definition: the operator $S_{\rho}^*S_{\rho}$ is

$$\begin{aligned} S_{\rho}^*S_{\rho} &= R_{\Omega^{-\rho}}(1 - R^*)(1 - R_{\Omega})R^* \\ &= (R_{\Omega^{-\rho}} - R_{\Omega^{-\rho}}R^*)(R^* - R_{\Omega}R^*) \\ &= (R_{\Omega^{-\rho}}R^* - R_{\Omega^{-\rho}}R^*R^* - R_{\Omega^{-\rho}}R_{\Omega}R^* + R_{\Omega^{-\rho}}R^*R_{\Omega}R^*) \\ &= R_{\Omega^{-\rho}}R^*(1 - R_{\Omega}R^*). \end{aligned}$$

Any Dirac mass x_{γ} with support at a point γ satisfies $\|R^*x_{\gamma}\|_{\ell^2} \leq 1$ (R^* is a projection). Thus, if $B_{\rho}(\gamma)$ is the ball of radius ρ , $\|(1 - R_{B_{\rho}(\gamma)})R^*x_{\gamma}\|_{\ell^2} \leq \beta_0(\rho)$, with $\lim_{\rho \rightarrow \infty} \beta_0(\rho) = 0$. Consequently $\|S_{\rho}x_{\gamma}\|_{\ell^2} \leq \beta_0(\rho)$ since $\Gamma \setminus \Omega$ is contained in the complement of the union of the $B_{\rho}(\gamma)$ for $\gamma \in \Omega^{-\rho}$. Since $\|S_{\rho}^*\| \leq 1$, $\|S_{\rho}^*S_{\rho}x_{\gamma}\|_{\ell^2} \leq \beta_0(\rho)$.

The Dirac masses being an orthonormal basis for $\ell^2(\Omega^{-\rho}; \mathbb{R}^s)$, we get that $\text{Tr} S_\rho^* S_\rho \leq s\beta_0(\rho)|\Omega^{-\rho}|$.

Last, we shall evaluate $n_i[a, b]$ for $a, b \in]0, 1[$ and $b - a = \varepsilon \in]0, 1[$. Let X_i be the space generated by eigenvectors of $R_{\Omega_i} R^*$ whose eigenvalue is in $[a, b]$. Then, $\forall \lambda \in [a, b], \forall x \in X_i$, x is an ε -quasimode of eigenvalue λ for $R_{\Omega_i} R^*$. The evaluation of $\dim X_i$ will be done by looking at spaces whose dimension is close. If $X_i^\rho = X_i \cap \ell^2(\Omega_i^{-\rho}, V)$ is the subspace of elements which vanish on the thickened boundary, $\dim X_i - \dim X_i^\rho \leq s|\Omega_i \setminus \Omega_i^{-\rho}|$. The amenability used on Ω_i shows that this difference is negligible, $\lim_{i \rightarrow \infty} (\dim X_i - \dim X_i^\rho)/|\Omega_i| = 0$; it will suffice to evaluate $\dim X_i^\rho$.

Unfortunately, neither X_i^ρ nor X_i is *a priori* invariant by $S_\rho^* S_\rho$. Let's nevertheless look at the intersection of X_i^ρ with the space generated by eigenvectors of $S_\rho^* S_\rho$ of eigenvalue $\leq \beta^2$; we will denote this new intersection by $X_i^{\rho, \beta}$. On this space, $\|S_\rho\| \leq \beta$ since

$$\|S_\rho x\|_{\ell^2}^2 = \langle S_\rho x, S_\rho x \rangle = \langle x, S_\rho^* S_\rho x \rangle \leq \beta^2 \|x\|_{\ell^2}^2$$

Yet again, this space is of dimension close to that of X_i^ρ : if $V^{>\beta^2}$ is the space of eigenvectors of $S_\rho^* S_\rho$ whose eigenvalue is greater than β^2 , then

$$\dim X_i^\rho - \dim X_i^{\rho, \beta} \leq \dim V^{>\beta^2} \leq \beta^{-2} \text{Tr} S_\rho^* S_\rho \leq s|\Omega_i^{-\rho}| \beta_0(\rho) / \beta^2.$$

In other words,

$$\forall \beta > 0, \forall \alpha > 0, \exists \rho \text{ such that } \limsup_{i \rightarrow \infty} \frac{\dim X_i^\rho - \dim X_i^{\rho, \beta}}{|\Omega_i|} \leq \alpha,$$

Thus, it remains to evaluate $\dim X_i^{\rho, \beta}$. To do so, we use the conclusion of the first part for $\lambda = a$ and $\varepsilon = b - a$: this yields $a(1 - a) \leq 2(b - a) + \beta$ given that $\dim X_i^{\rho, \beta} > 0$. Consequently, the inequality $b - a < (a - a^2 - \beta)/2$ implies that $\dim X_i^{\rho, \beta} = 0$. Which means that when $\rho \geq \rho_0(\beta, \alpha)$ is sufficiently big, $\limsup_{i \rightarrow \infty} \dim X_i^\rho / |\Omega_i| \leq \alpha$. It follows that

$$\begin{aligned} \limsup_{i \rightarrow \infty} \frac{\dim X_i}{|\Omega_i|} &\leq \limsup_{i \rightarrow \infty} \frac{\dim X_i - \dim X_i^\rho}{|\Omega_i|} + \limsup_{i \rightarrow \infty} \frac{\dim X_i^\rho}{|\Omega_i|} \\ &\leq 0 + \limsup_{i \rightarrow \infty} \frac{\dim X_i^\rho - \dim X_i^{\rho, \beta}}{|\Omega_i|} + \limsup_{i \rightarrow \infty} \frac{\dim X_i^{\rho, \beta}}{|\Omega_i|} = 0 \end{aligned}$$

since $\alpha \rightarrow 0$ when $\rho \rightarrow \infty$. This proves the theorem for intervals $[a, b]$ satisfying $b - a < a(1 - a)/2$, as the size of β is not constrained. The conclusion is obtained by noticing that any interval strictly contained in $[0, 1]$ can be covered by intervals of this type. \square

This property enables us interpret von Neumann dimension as a wdim for a set with a chosen pseudo-metric.

Corollary A.2. (cf. [6, corollary 1.12.2]) *Let $Y \subset \ell^2(\Gamma; V)$ be an invariant subspace, let $B_1^Y = Y \cap B_1^{\ell^2(\Gamma; V)}$ the intersection of the unit ball with Y . Then, for a given Følner*

sequence $\Omega_i \subset \Gamma$,

$$\forall \varepsilon \in]0, 1[, \quad \lim_{i \rightarrow \infty} \frac{1}{|\Omega_i|} \text{wdim}_\varepsilon(R_{\Omega_i} B_1^Y, \ell^2) = \dim_{\ell^2} Y$$

Proof. (We give the argument of [6] in detail.) To get this result $R_\Omega B_1^Y$ must be seen as an ellipsoid whose semi-axes are related to the eigenvalues of $R_\Omega R^*$. Remark that $B_1^Y = R^* B_1^{\ell^2(\Gamma; V)}$. Then, an ellipsoid can be defined as the image of a ball by a self-adjoint operator, say A ; the semi-axis of this ellipsoid are in correspondence with the eigenvalue of A . It might be worth recalling how this relates to the usual definition of an ellipsoid E (as the set $\{y \mid \langle y, Py \rangle \leq 1\}$ for a positive definite operator P). The semi-axes of E are of the form $\lambda_i(P)^{-1/2}$ for $\lambda_i(P)$ an eigenvalue of P . Indeed let B^V be a ball in a vector space V , and let $A : V \rightarrow V$ be self-adjoint. Restricting to $V' = \text{Im} A = \text{ker} A^\perp \subset V$, it must be shown that for $x \in V'$ such that $\langle x, x \rangle \leq 1$, there exists $P : V' \rightarrow V'$ positive definite such that $\langle Ax, PAx \rangle \leq 1$. Taking $P = A^{-2}$ yields the conclusion: A^{-2} is a positive definite operator on V' whose eigenvalues are $\lambda_i(A)^{-2}$. Thus AB^V is an ellipsoid with semi-axis $\lambda_i(P)^{-1/2} = \lambda_i(A)$.

In our present context, $R_\Omega R^*$ is self-adjoint, thus $R_\Omega R^* B_1^{\ell^2(\Gamma; V)} = R_\Omega B_1^Y$ is an ellipsoid whose semi-axis are the eigenvalues of $R_\Omega R^*$. This ellipsoid contains isometrically the ball obtained by ignoring the semi-axis of length $< \varepsilon$ and replacing the remaining ones by semi-axis of length ε . Thus $\text{wdim}_\varepsilon(R_{\Omega_i} B_1^Y, \ell^2) \geq n_i[\varepsilon, 1]$. On the other hand, $\text{wdim}_\varepsilon(R_{\Omega_i} B_1^Y, \ell^2) \leq n_i[\varepsilon/2, 1]$, as the continuous map obtained by projecting on the sub-ellipsoid formed by the semi-axis of length $> \varepsilon/2$ indicates. When $i \rightarrow \infty$, the eigenvalues of $R_{\Omega_i} R^*$ tend to 0 or 1. In particular, when $i \rightarrow \infty$ the inequality

$$\frac{1}{|\Omega_i|} n_i[\varepsilon, 1] \leq \frac{1}{|\Omega_i|} \text{wdim}_\varepsilon(R_{\Omega_i} B, \ell^2) \leq \frac{1}{|\Omega_i|} n_i[\varepsilon/2, 1]$$

shows that $\lim_{i \rightarrow \infty} \frac{1}{|\Omega_i|} \text{wdim}_\varepsilon(R_{\Omega_i} B_1^Y, \ell^2) = \dim_{\ell^2} Y$, since $n_i[a, 1] \rightarrow \text{Tr} R_{\Omega_i} R^*$. \square

This corollary can be expressed in terms of ℓ^p dimension. Indeed, let $B_1^Y = Y \cap B_1^{\ell^2(\Gamma; V)}$ be endowed with the pseudo-metric of evaluation at $e \in \Gamma$: $ev(x, y) = \|x(e) - y(e)\|_V$. Translation of this pseudo-metric by an element of γ is the evaluation at γ . Thus, $ev_{\ell^2(\Omega)}(x, y) = \|x - y\|_{\ell^2(\Omega)} = \|R_\Omega(x - y)\|_{\ell^2}$. The map $R_\Omega : B_1^Y \rightarrow R_\Omega B$ is continuous for the topology of B_1^Y as a subset of ℓ^p (with τ^* or even with the norm topology). The fibers are of “diameter” 0 given that $\Omega' \subset \Omega$. Thus, corollary A.2 can be expressed as follows:

$$\text{Wgc}_{\ell^2}(B_1^Y, \tau^*, ev, \{\Omega_i\}) = \dim_{\ell^2} Y.$$

Indeed, $R_{\Omega_i} B_1^Y$ injects isometrically in (B_1^Y, ev_{Ω_i}) and (B_1^Y, ev_{Ω_i}) possesses a map to $R_{\Omega_i} B_1^Y$ whose fiber are of “diameter” 0. Thus $\text{wdim}_\varepsilon(B_1^Y, ev_{\Omega_i}) = \text{wdim}_\varepsilon(R_{\Omega_i} B_1^Y, \ell^2)$. This shows that definition 2.3 is equivalent when $p = 2$ to the von Neumann dimension and this for any Følner sequence $\{\Omega_i\}$ chosen.

It would have been surprising that this were not the case in general. An alteration of the Ornstein-Weiss lemma (see section 5) enables to show the independence of the limit on the sequence chosen.

References

- [1] J. Cheeger and M. Gromov. L_2 -cohomology and group cohomology. *Topology*, 25(2):189–215, 1986.
- [2] M. Coornaert. *Dimension topologique et systèmes dynamiques*, volume 14 of *Cours Spécialisés*. Société Mathématique de France, Paris, 2005.
- [3] M. Coornaert and F. Krieger. Mean topological dimension for actions of discrete amenable groups. *Discrete Contin. Dyn. Syst.*, 13(3):779–793, 2005.
- [4] D. L. Donoho. Compressed sensing. *IEEE Trans. Inform. Theory*, 52(4):1289–1306, 2006.
- [5] A. Gournay. Width of ℓ^p balls. arXiv:0711.3081v2 or <http://hal.archives-ouvertes.fr/hal-00189043/fr/>, 2008.
- [6] M. Gromov. Topological invariants of dynamical systems and spaces of holomorphic maps. I. *Math. Phys. Anal. Geom.*, 2(4):323–415, 1999.
- [7] A. N. Kolmogorov and S. V. Fomin. *Introductory real analysis*. Dover Publications Inc., New York, 1975. Translated from the second Russian edition and edited by Richard A. Silverman, Corrected reprinting.
- [8] F. Krieger. Le lemme d’Ornstein-Weiss d’après Gromov. In *Dynamics, ergodic theory, and geometry*, volume 54 of *Math. Sci. Res. Inst. Publ.*, pages 99–111. Cambridge Univ. Press, Cambridge, 2007.
- [9] E. Lindenstrauss and B. Weiss. Mean topological dimension. *Israel J. Math.*, 115:1–24, 2000.
- [10] J. Lindenstrauss and L. Tzafriri. *Classical Banach spaces*. Springer-Verlag, Berlin, 1973. Lecture Notes in Mathematics, Vol. 338.
- [11] W. Lück. L^2 -invariants of regular coverings of compact manifolds and CW-complexes. In *Handbook of geometric topology*, pages 735–817. North-Holland, Amsterdam, 2002.
- [12] F. J. Murray. On complementary manifolds and projections in spaces L_p and l_p . *Trans. Amer. Math. Soc.*, 41(1):138–152, 1937.
- [13] D. S. Ornstein and B. Weiss. Entropy and isomorphism theorems for actions of amenable groups. *J. Analyse Math.*, 48:1–141, 1987.
- [14] P. Pansu. Introduction to L^2 Betti numbers. In *Riemannian geometry (Waterloo, ON, 1993)*, volume 4 of *Fields Inst. Monogr.*, pages 53–86. Amer. Math. Soc., Providence, RI, 1996.
- [15] P. Pansu. L^p -cohomology and pinching. In *Rigidity in dynamics and geometry (Cambridge, 2000)*, pages 379–389. Springer, Berlin, 2002.

- [16] W. Rudin. *Functional analysis*. International Series in Pure and Applied Mathematics. McGraw-Hill Inc., New York, second edition, 1991.
- [17] A. Sobczyk. Projections in Minkowski and Banach spaces. *Duke Math. J.*, 8:78–106, 1941.
- [18] M. Tsukamoto. Macroscopic dimension of the l^p -ball with respect to the l^q -norm. *J. Math. Kyoto Univ.*, 48(2):445–454, 2008.