



HAL
open science

Spectral action for torsion with and without boundaries

Bruno Iochum, Cyril Levy, D. V. Vassilevich

► **To cite this version:**

Bruno Iochum, Cyril Levy, D. V. Vassilevich. Spectral action for torsion with and without boundaries. Communications in Mathematical Physics, 2012, 310 (2), pp.367-382. 10.1007/s00220-011-1406-7. hal-00516327

HAL Id: hal-00516327

<https://hal.science/hal-00516327>

Submitted on 9 Sep 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

CENTRE DE PHYSIQUE THÉORIQUE¹
CNRS–Luminy, Case 907
13288 Marseille Cedex 9
FRANCE

Spectral action for torsion with and without boundaries

B. Iochum^{1,2}, C. Levy³ and D. Vassilevich^{4,5}

Abstract

We derive a commutative spectral triple and study the spectral action for a rather general geometric setting which includes the (skew-symmetric) torsion and the chiral bag conditions on the boundary. The spectral action splits into bulk and boundary parts. In the bulk, we clarify certain issues of the previous calculations, show that many terms in fact cancel out, and demonstrate that this cancellation is a result of the chiral symmetry of spectral action. On the boundary, we calculate several leading terms in the expansion of spectral action in four dimensions for vanishing chiral parameter θ of the boundary conditions, and show that $\theta = 0$ is a critical point of the action in any dimension and at all orders of the expansion.

PACS numbers: 11.10.Nx, 02.30.Sa, 11.15.Kc
MSC–2000 classes: 46H35, 46L52, 58B34
CPT-P050-2010
CPH-SYM-00

¹ UMR 6207

– Unité Mixte de Recherche du CNRS et des Universités Aix-Marseille I, Aix-Marseille II et de l'Université du Sud Toulon-Var (Aix-Marseille Université)

– Laboratoire affilié à la FRUMAM – FR 2291

² Also at Université de Provence, iochum@cpt.univ-mrs.fr

³ University of Copenhagen, Department of Mathematical Sciences, Denmark, levy@math.ku.dk

⁴ CMCC – Universidade Federal do ABC, Santo André, S.P., Brazil, dmitry.vasilevich@ufabc.edu.br

⁵ Also at the Department of Theoretical Physics, St. Petersburg University, Russia

1 Introduction

The central object of the noncommutative geometry approach [11] is given by a spectral triple $(\mathcal{A}, \mathcal{H}, \mathcal{D})$ consisting of an algebra \mathcal{A} , a Hilbert space \mathcal{H} and an unbounded selfadjoint operator \mathcal{D} , like the Dirac operator for manifolds. By making use of the Dirac operator, one can define a spectral action [6] which becomes a natural action for a classical bosonic field theory. Practical calculations of this action are usually done by using the heat kernel expansion, which is a standard instrument of quantum field theory [38]. Although this approach was initially designed for noncommutative spaces, it appeared to be rigid enough to make predictions in the commutative case as well. For example, being applied to Riemannian manifolds with boundary [7], the spectral action reproduces correctly the Einstein–Hilbert action together with the boundary term [25]. It seems also natural to extend this approach to the Riemann–Cartan manifolds and bring into the focus a non-zero torsion.

The spectral action associated to a manifold endowed with a connection with torsion has been computed recently in [23], though technical tools were ready long ago [20, 21, 34]. The extension by torsion [23] of the almost commutative spectral triple of the standard model in particle physics [9] demonstrated that torsion becomes coupled to the Higgs field and, therefore is possibly observable.

The purpose of the present paper is to reconsider the calculations of [23], extend the spectral triple with torsion to manifolds with boundaries, and to compute the corresponding spectral action. We work mostly in four dimensions, although some results are valid for arbitrary even number of dimensions. As in [23] we restrict ourselves to totally skew-symmetric torsion. The presence of a skew-symmetric torsion does not modify the geodesics and is, therefore, a rather minimalist modification of the Riemann geometry. Regarding physical aspects of the space-time torsion, the interested reader can consult [35] (and also [31] for some recent developments, in particular recent limits on torsion given by two different experiments which place careful bounds on the four axial components of torsion [26, 32]).

In an expansion in the large cut-off parameter Λ the spectral action is local, i.e. it is given as a sum of volume and boundary integrals. The volume part of the spectral action with torsion has been calculated in [23], but, as we show below, many terms given in that paper in fact cancel out. In particular, there is no term quadratic in the torsion and linear in the Riemann curvature, no quartic torsion term, etc. These cancellations are not accidental, but they are forced by an important symmetry of the spectral action, which is the chiral symmetry. The importance of chiral symmetry for the spectral action has been stressed recently [1] in a somewhat different context.

This prominent role of the chiral symmetry has motivated us to introduce a chiral phase also on the boundary, i.e. to consider the so called chiral bag boundary conditions. These are local boundary conditions depending on a real parameter θ . They are “mixed oblique” in the sense that in the definition of the domain of squared Dirac operator there is a piece which restricts solely the value of the field at the boundary (as in the Dirichlet case), and a piece which contains the normal derivative at the boundary (as in the Neumann case). This latter piece depends also on the derivatives in the directions tangential to the boundary, and this makes the boundary conditions “oblique”. The case $\theta = 0$ corresponds to the usual local mixed boundary condition.

The bag ($\theta = 0$) and chiral bag ($\theta \neq 0$) boundary conditions appeared first in theoretical physics, in the models of hadrons (see [24] for a review). Properties of the determinants of the Dirac operator for chiral bag conditions and their relations to chiral symmetry breaking were studied in [27, 39]. A more detailed mathematical investigation of the spectral properties of the

Dirac operator for these conditions was performed in [14, 15, 19].

The goal of this paper is to compute the spectral action for spectral triples on compact 4-dimensional manifolds with boundary already considered in [7, 8] but in presence of a totally skew-symmetric torsion T . In order to do so, we exhibit simple spectral triples $(\mathcal{A}, \mathcal{D}, \mathcal{H})$ with vanishing tadpoles (Theorem 3.3) in the case of a chiral bag boundary condition. Essentially, the construction follows the same arguments of [29], with one difference: the boundary operator Π_- , related to the chirality χ_θ , is not selfadjoint. In respect to this, we extend [29, Theorem 4.5] to non-selfadjoint boundary operators in Theorem 3.1.

We generalize [23] for chiral mixed boundary condition (in the case $\theta = 0$) by computing the spectral action in dimension 4 in section 4.2. This restriction to $\theta = 0$ is justified by the fact that this is a critical point for the spectral action, a result obtained in section 5 via the Index Theorem.

2 Notations

Let \widetilde{M} be a smooth compact Riemannian (with the metric g) manifold without boundary of even dimension $n = 2m$. We denote ∇^{LC} the Levi-Civita connection associated to g . We fix a linear connection ∇^g on $T\widetilde{M}$ which is g -compatible and with the same geodesics as ∇^{LC} . Equivalently, ∇^g is a linear connection with a totally skew-symmetric torsion $8T$ (the chosen coefficient 8 is justified below).

It is known in this case that we have $\nabla_X^g Y = \nabla_X^{LC} Y + 4T(X, Y, -)$, or in other words, for any vector fields X, Y, Z ,

$$g(\nabla_X^g Y - \nabla_X^{LC} Y, Z) = 4T(X, Y, Z). \quad (1)$$

Let M be a submanifold of \widetilde{M} of dimension n such that its topological closure \overline{M} is a compact manifold with smooth boundary $\partial M = \overline{M} \setminus M$. This implies that ∂M is a smooth compact submanifold of \widetilde{M} without boundary of dimension $n - 1$. Remark that, for a given manifold M with boundary ∂M , there exists a manifold \widetilde{M} with previous property.

We also assume that \widetilde{M} is endowed with a complex vector bundle \widetilde{V} of dimension 2^m and with a smooth map $\gamma : T\widetilde{M} \rightarrow \text{End}(\widetilde{V})$ giving a structure of Clifford module to \widetilde{V} . This means that for any $x, y \in T\widetilde{M}$, $\{\gamma(x), \gamma(y)\} = 2g(x, y)$.

Let us remark immediately that the convention chosen here $(+2g(x, y))$ for the Clifford anticommutation relations is different from the one $(-2g(x, y))$ used for instance in [2, 4, 17–19]. One can go from one convention to another with a multiplication by i . More precisely, $\gamma' := i\gamma$ satisfies $\{\gamma'(x), \gamma'(y)\} = -2g(x, y)$.

We also fix on \widetilde{V} a Hermitian inner product (\cdot, \cdot) such that $\gamma(x)^* = \gamma(x)$ for any $x \in T\widetilde{M}$. Note that for a given γ , such Hermitian inner product always exists. In the following, we shall say that a given connection $\nabla^{\widetilde{V}}$ on \widetilde{V} is *unitary* if for any $x \in T\widetilde{M}$ and $v, w \in \widetilde{V}$

$$x(v, w) = (\nabla_x^{\widetilde{V}} v, w) + (v, \nabla_x^{\widetilde{V}} w)$$

and *compatible with* $(\nabla^{T\widetilde{M}}, \gamma)$, where $\nabla^{T\widetilde{M}}$ is a given g -compatible connection on $T\widetilde{M}$ with totally skew-symmetric torsion, if for any $x, y \in T\widetilde{M}$ and $v \in \widetilde{V}$,

$$\nabla_x^{\widetilde{V}}(\gamma(y)v) = \gamma(\nabla_x^{T\widetilde{M}}(y))v + \gamma(y)\nabla_x^{\widetilde{V}}v.$$

Our conventions are the following (see [38]): $(e) := \{e_1, \dots, e_n\}$ is a local orthonormal frame of the tangent space where e_n is the inward pointing unit vector field. We assume that Roman

indices a, b, c range from 1 to $n - 1$ and index an orthonormal frame for the tangent bundle of ∂M . Local coordinates charts are denoted (x_μ) , and (∂_μ) is the associated local coordinate frame. We shall use the shorthands: $\gamma_i := \gamma(e_i)$, $\gamma_\mu := \gamma(\partial_\mu)$ and $\gamma^\mu := g^{\mu\nu}\gamma_\nu$. The coordinates of e_i in the frame (∂_μ) are denoted (e_i^μ) and the inverse matrix is denoted (e_i^i) .

By the proof of [18, Lemma 1.1.7], if we fix a given partition of unity on \widetilde{M} and associated local frames for the bundle \widetilde{V} , we can construct a connection on \widetilde{V} which is unitary and compatible with (∇^{LC}, γ) . The idea is to patch together the connections locally defined by the matrices

$$\omega_i^s := -\frac{1}{4}\Gamma_{ijk}\gamma_j^s\gamma_k^s \quad (2)$$

where $s = (s_p)$ is a local frame for \widetilde{V} , (e) is a local orthonormal frame with $e_i\gamma_j^s = 0$, Γ_{ijk} are the Christoffel coefficients of the first kind of the Levi-Civita connection in (e) , and γ_j^s is the matrix of γ_j in the frame s . Note that the minus sign in (2), not present in [18], comes from our different definition of γ .

Actually, we can see from the proof that the same construction using ∇^g instead of ∇^{LC} is still valid, since the fact that ∇^{LC} is torsion-free is not used. The crucial point in the proof is the relation $\Gamma_{ijk} = -\Gamma_{ikj}$, which is true for any g -compatible connection on $T\widetilde{M}$.

As a consequence, we can apply [18, Lemma 1.1.7] to define a ‘‘spin’’ connection $\nabla^{(T)}$ on \widetilde{V} which is unitary and compatible with (∇^g, γ) . The same construction using ∇^{LC} yield to the connection $\nabla^{(0)}$ on \widetilde{V} which is unitary and compatible with (∇^{LC}, γ) . Locally, since T is skew-symmetric, (1) and (2) give $\nabla_j^{(T)} = \nabla_j^{(0)} - T_{jkl}\gamma_k\gamma_l$, where $T_{jkl} := T(e_j, e_k, e_l)$. We define the exterior product \wedge so that $T = 3!\sum_{j<k<l}T_{jkl}\theta_j \wedge \theta_k \wedge \theta_l$, where (θ_j) is the dual coframe of (e_j) . The Clifford action of T (still denoted T) is thus

$$T = 6 \sum_{j<k<l} T_{jkl} (i\gamma_j)(i\gamma_k)(i\gamma_l) = -i \sum_{jkl} T_{jkl} \gamma_j \gamma_k \gamma_l. \quad (3)$$

The Dirac operators respectively associated to $\nabla^{(T)}$ and $\nabla^{(0)}$ are

$$\mathcal{D}^{(T)} := i \sum_{j=1}^n \gamma_j \nabla_j^{(T)}, \quad \mathcal{D}^{(0)} := i \sum_{j=1}^n \gamma_j \nabla_j^{(0)},$$

so we deduce from the preceding equations that

$$\mathcal{D}^{(T)} = \mathcal{D}^{(0)} + T$$

and this relation hence justifies the factor 8 in the definition of the torsion of ∇^g .

The Dirac operators $\mathcal{D}^{(T)}$ and $\mathcal{D}^{(0)}$ are formally selfadjoint, as unbounded operators on $\mathcal{H} := L^2(\widetilde{M}, \widetilde{V})$, at least when \widetilde{M} has no boundary [16]. Of course, since we want to deal with a spectral triple $(\mathcal{A}, \mathcal{D}^{(T)}, \mathcal{H})$, we need a selfadjoint $\mathcal{D}^{(T)}$.

We denote Γ_{ij}^k the Christoffel coefficients of ∇^{LC} . If R_{ijkl} is the component of the Riemann curvature tensor for ∇^{LC} , the Ricci tensor is $R_{jk} := R_{ijki}$ with scalar curvature $R = R_{ii}$ (so $R = +12$ for the 4-sphere).

Moreover, we note $L_{ab} := \Gamma_{ab}^n$ the second fundamental form or extrinsic curvature and $L := L_{aa}$.

In coordinates, we get $(\nabla_\mu^{LC} X)_\nu = \partial_\mu X_\nu + \Gamma_{\mu\rho}^\nu X_\rho$. At the Clifford module level, we get, using previous notations, $(\nabla_\mu^{(0)} v)_p = \partial_\mu v_p + (\omega_\mu^s(v))_p$, where (ω_μ^s) is the connection 1-form associated to $\nabla^{(0)}$ and the frame $s = (s_p)$. A computation shows that $\omega_\mu^s = \frac{1}{4}\sigma_\mu^{\nu\rho}\gamma_\nu^s\gamma_\rho^s = \frac{1}{8}\sigma_\mu^{\nu\rho}[\gamma_\nu^s, \gamma_\rho^s]$ where the spin connection $(\sigma_\mu^{\nu\rho})$ is locally defined by $-e_\mu^j e_k^\nu e_l^\rho \Gamma_{jkl}$.

3 The spectral triple for chiral bag boundary condition

As boundary conditions for the operator $\mathcal{D}^{(T)}$ and its square, we prefer to choose local ones which guarantee its selfadjointness on M , as the chiral bag boundary condition which is based on the existence of the usual chirality γ_{n+1} (which exists on even dimensional manifolds) and a real parameter θ .

It has been shown [2] that such chiral bag boundary conditions yield a strongly elliptic boundary value problem for the Dirac operator and its square. Moreover, the associated heat-kernel asymptotics has been investigated on the Euclidean ball in [15] and in a more general setting in [14]. Some stability properties of heat-kernel coefficients with respect to parameter θ have been established in [19].

We give in this section the construction of a spectral triple based on a Dirac operator with a chiral bag boundary condition.

Recall from (3) that the 3-form T acts on the Clifford module as $-iT_{jkl}\gamma_j\gamma_k\gamma_l$, thus T is formally selfadjoint as an operator (acting by Clifford multiplication) on $L^2(\widetilde{M}, \widetilde{V})$. Recall also that $\mathcal{D}^{(T)} = \mathcal{D}^{(0)} + T$ where $\mathcal{D}^{(0)}$ is the usual Dirac operator based on the spin connection compatible with (∇^{LC}, γ) .

The chirality matrix is defined by

$$\gamma_{n+1} := (-i)^{n/2} \gamma_1 \cdots \gamma_n,$$

so $\gamma_{n+1}^* = \gamma_{n+1}$, $\gamma_{n+1}^2 = \text{Id}_V$ and $\gamma_i \gamma_{n+1} = -\gamma_{n+1} \gamma_i$.

Note that $\{\gamma_{n+1}, \mathcal{D}^{(T)}\} = \{\gamma_{n+1}, \mathcal{D}^{(0)}\} = \{\gamma_{n+1}, T\} = 0$ and if $n = 4$, $\gamma_5 = -\gamma_1 \cdots \gamma_4$.

We also define

$$\chi_\theta := -ie^{\theta\gamma_{n+1}}\gamma_{n+1}\gamma_n.$$

Since γ_n anti-commutes with γ_{n+1} , $\chi_\theta^2 = \text{Id}_V$, so $\Pi_\pm := \frac{1}{2}(\text{Id}_V \pm \chi_\theta)$ are two non-selfadjoint idempotents. A direct computation shows that $\chi_\theta^* = \chi_{-\theta}$ and

$$\chi_{-\theta}\gamma_n + \gamma_n\chi_\theta = 0, \quad \gamma_n + \chi_{-\theta}\gamma_n\chi_\theta = 0. \quad (4)$$

The sub-bundle of \widetilde{V} on \overline{M} (resp. ∂M) is denoted \overline{V} (resp. V) and $H^s(\widetilde{V})$, $H^s(\overline{V})$ are the Sobolev spaces of order $s \in \mathbb{R}$ respectively on \widetilde{M} with bundle \widetilde{V} and \overline{M} with bundle \overline{V} . Recall that by definition

$$H^s(\overline{V}) := H^s(\overline{M}, \overline{V}) := r_+(H^s(\widetilde{V}))$$

where r_+ is the restriction on M . We define

$$C^\infty(\overline{M}, \overline{V}) := r_+(C^\infty(\widetilde{M}, \widetilde{V})), \quad C^\infty(\overline{M}) := r_+(C^\infty(\widetilde{M})).$$

The extension by zero operator e_+ is a linear continuous operator from $H^s(\overline{V})$ into $H^s(\widetilde{V})$ for any $s \in]-\frac{1}{2}, \frac{1}{2}[$ such that $e_+(u) = u$ on M and $e_+(u)(x) = 0$ for any $u \in C^\infty(\overline{M}, \overline{V})$ and $x \in \widetilde{M} \setminus M$.

When P is a pseudodifferential operator on $(\widetilde{M}, \widetilde{V})$, its truncation to M is given by

$$P_+ := r_+ P e_+.$$

Let $\mathcal{D}_{\mathcal{B}_{\chi_\theta}}^{(T)}$ be the realization of $\mathcal{D}^{(T)}$ on $\mathcal{B}_{\chi_\theta}$, that is the operator acting as $\mathcal{D}_+^{(T)}$ on the space of sections ψ in $H^1(\overline{V})$ satisfying the boundary condition $\mathcal{B}_{\chi_\theta}\psi = 0$ with

$$\mathcal{B}_{\chi_\theta}\psi := \Pi_- r(\psi)$$

where we denoted r the restriction operation on ∂M (denoted γ_0 in [22]).

As in [29], define $\mathcal{A}_{\mathcal{D}_{\mathcal{B}_{\chi_\theta}}^{(T)}}$ as the $*$ -algebra of smooth functions $a \in C^\infty(\overline{M})$ such that a and a^* apply $\mathcal{H}_{\mathcal{D}_{\mathcal{B}_{\chi_\theta}}^{(T)}}$ into itself where

$$\mathcal{H}_{\mathcal{D}_{\mathcal{B}_{\chi_\theta}}^{(T)}} := \bigcap_{k \geq 1} \text{Dom } \mathcal{D}_{\mathcal{B}_{\chi_\theta}}^{(T)}.$$

These functions act as multiple of the identity on the Hilbert space $L^2(\overline{V})$.

We refer to [22] for the definition of an elliptic pseudodifferential boundary system and recall the Green formula [22, 1.3.2 Proposition]: if P is a pseudodifferential operator of order $k \in \mathbb{N}$, then for any $u, v \in C^\infty(\overline{M}, \overline{V})$,

$$(P_+ u, v)_M - (u, (P^*)_+ v)_M = (\mathfrak{A}_P \rho u, \rho v)_{\partial M}$$

where $\rho = \{r_0, \dots, r_{k-1}\}$ is the Cauchy boundary operator defined by $r_j u = (-i\partial_n)^j u|_{\partial M}$ with $r_0 := r$ (here, ∂_n is the interior normal derivative) and \mathfrak{A}_P is the Green matrix associated to P . When $k = 1$, \mathfrak{A}_P is an endomorphism on the boundary ∂M . In particular, $\mathfrak{A}_{\mathcal{D}^{(0)}} = -i\gamma_n$.

It is known that $\mathcal{D}_{\mathcal{B}_{\chi_\theta}}^{(T)}$ is selfadjoint [2, Theorem 2.1], but we give here a proof based on another approach more appropriate to noncommutative geometry. We begin with an extension of [29, Theorem 4.5] in which the boundary endomorphism Π is no more supposed to be selfadjoint:

Theorem 3.1. *Let $P \in \Psi^1(\widetilde{V})$ be a symmetric pseudodifferential operator of order one on \widetilde{M} satisfying the transmission property (see [22, 1.2]).*

Let $\Pi \in C^\infty(\partial M, L(V))$ be an idempotent endomorphism on the boundary such that the system $\{P_+, \mathcal{B} := \Pi r\}$ is an elliptic pseudodifferential boundary operator. Then

(i) $P_{\mathcal{B}}$ is selfadjoint if and only if

$$(1 - \Pi^*) \mathfrak{A}_P (1 - \Pi) = 0 \quad \text{and} \quad \Pi \mathfrak{A}_P^{-1} \Pi^* = 0. \quad (5)$$

(ii) When $P_{\mathcal{B}}$ is selfadjoint, $(C^\infty(\overline{M}), L^2(\overline{V}), P_{\mathcal{B}})$ is a spectral triple of dimension $\dim(M)$.

(iii) When P is a differential operator such that P^2 has a scalar principal symbol and $P_{\mathcal{B}}$ is selfadjoint, the spectral triple $(\mathcal{A}_{P_{\mathcal{B}}}, L^2(\overline{V}), P_{\mathcal{B}})$ is regular.

(iv) Under the hypothesis of (iii), $\mathcal{A}_{P_{\mathcal{B}}}$ is the largest algebra \mathcal{A} in $C^\infty(\overline{M})$ such that the triple $(\mathcal{A}, L^2(\overline{V}), P_{\mathcal{B}})$ is regular.

Proof. (i) For any given endomorphism R , we shall denote \widehat{R} the surjective morphism defined as the operator acting as R from the domain of R into the image of R .

Since P is a pseudodifferential operator of order 1 and $\widehat{\Pi}$ is a surjective morphism, we can apply [22, 1.6.11 Theorem] with the following choices: the S of [22] is $\widehat{\Pi}$ and $S' := \widehat{1 - \Pi}$. Note that Π is here not surjective: it is an endomorphism only surjective on $V^+ := \Pi(V)$ with kernel $V^- := (1 - \Pi)(V)$, so V is the direct (not necessarily orthogonal) sum of V^+ and V^- . In the notation of [22], we take $B = P_{\mathcal{B}}$, $G = K = G' = \widetilde{G} = T' = 0$ and $\rho = r$. Remark that $\mathfrak{A}_P^* = -\mathfrak{A}_P$ since $P = P^*$ (viewed as defined on $H^1(\widetilde{M}, \widetilde{V})$). Thus we have (still using the notations of [22]), $T = \mathcal{B}$ and $\widetilde{T} = C'^* \mathfrak{A}_P^* r = -C'^* \mathfrak{A}_P r$ (the matrix I^\times is the number 1 here).

According to [22, 1.6.11 Theorem], since $\{P_+, \mathcal{B}\}$ is elliptic, $P_{\mathcal{B}}$ is selfadjoint if there is a homeomorphism Ψ from $H^s(V^+)$ onto $H^s(V^-)$, such that

$$-C'^* \mathfrak{A}_P r = \Psi \widehat{\Pi} r, \quad (6)$$

with C' satisfying $(\widehat{1 - \Pi})C' = \text{Id}_{V^-}$ and $C'(\widehat{1 - \Pi}) = 1 - \Pi$. In other words, C' is the injection from V^- into V .

By [22, (1.6.52)], when this is the case, Ψ has the form $\Psi = C'^* \mathfrak{A}_P^* C$ with $\widehat{\Pi} C = \text{Id}_{V^+}$ and $C \widehat{\Pi} = \Pi$. Note that \mathfrak{A}_P is invertible as a consequence of the ellipticity of P .

The following computation shows that $C'^* = P_-(1 - \Pi^*) (= P_-)$, where P_- is the orthogonal projection from V onto V^- : for any $u \in V$ and $v \in V^-$,

$$(C'^* u, v)_{V^-} = (u, C' v)_V = (u, v)_V = ((1 - \Pi^*)u, v)_V = (P_-(1 - \Pi^*)u, v)_{V^-}.$$

Now, suppose that $(1 - \Pi^*) \mathfrak{A}_P (1 - \Pi) = 0$ and $\Pi \mathfrak{A}_P^{-1} \Pi^* = 0$. We define $\Psi = -C'^* \mathfrak{A}_P C$ which is a homeomorphism from $H^s(V^+)$ onto $H^s(V^-)$. Indeed, if we set $\Psi^{-1} := -\widehat{\Pi} \mathfrak{A}_P^{-1} (\widehat{1 - \Pi})^*$, we get

$$\begin{aligned} \Psi \circ \Psi^{-1} &= C'^* \mathfrak{A}_P C \widehat{\Pi} \mathfrak{A}_P^{-1} (\widehat{1 - \Pi})^* = P_-(1 - \Pi^*) \mathfrak{A}_P \Pi \mathfrak{A}_P^{-1} (\widehat{1 - \Pi})^* \\ &= P_-(1 - \Pi^*) \mathfrak{A}_P (\Pi + (1 - \Pi)) \mathfrak{A}_P^{-1} (\widehat{1 - \Pi})^* = C'^* (\widehat{1 - \Pi})^* = \text{Id}_{V^-}, \end{aligned}$$

and

$$\begin{aligned} \Psi^{-1} \circ \Psi &= \widehat{\Pi} \mathfrak{A}_P^{-1} (\widehat{1 - \Pi})^* C'^* \mathfrak{A}_P C = \widehat{\Pi} \mathfrak{A}_P^{-1} (1 - \Pi^*) \mathfrak{A}_P C \\ &= \widehat{\Pi} \mathfrak{A}_P^{-1} (1 - \Pi^* + \Pi^*) \mathfrak{A}_P C = \widehat{\Pi} C = \text{Id}_{V^+}. \end{aligned}$$

Moreover,

$$\Psi \widehat{\Pi} = -C'^* \mathfrak{A}_P \Pi = -P_-(1 - \Pi^*) \mathfrak{A}_P (\Pi + (1 - \Pi)) = -C'^* \mathfrak{A}_P.$$

As a consequence, (6) is satisfied and the if part of the assertion follows.

Conversely, suppose that $P_{\mathcal{B}}$ is selfadjoint. Using Green's formula, $(\mathfrak{A}_P r(u), r(v))_{\partial M} = 0$ for any $u, v \in \text{Dom } P_{\mathcal{B}}$. Since $r : H^1(\overline{V}) \rightarrow H^{1/2}(V)$ is surjective, $(\mathfrak{A}_P (1 - \Pi)\psi, (1 - \Pi)\phi)_{\partial M} = 0$ for any $\psi, \phi \in H^{1/2}(V)$ and thus $(1 - \Pi^*) \mathfrak{A}_P (1 - \Pi) = 0$. Again, from [22, Theorem 1.6.11] we get that $\Psi := C'^* \mathfrak{A}_P^* C$ is a homeomorphism from $H^s(V^+)$ onto $H^s(V^-)$ and we check as before that $\Psi^{-1} := -\widehat{\Pi} \mathfrak{A}_P^{-1} (\widehat{1 - \Pi})^*$ is a right-inverse of Ψ , and thus, is the inverse of Ψ .

The equation $\Psi^{-1} \circ \Psi = \text{Id}_{V^+}$ yields $\widehat{\Pi} \mathfrak{A}_P^{-1} \Pi^* \mathfrak{A}_P C = 0$, which gives $\widehat{\Pi} \mathfrak{A}_P^{-1} \Pi^* \mathfrak{A}_P \Pi = 0$. Thus,

$$\widehat{\Pi} \mathfrak{A}_P^{-1} \Pi^* \mathfrak{A}_P = \widehat{\Pi} \mathfrak{A}_P^{-1} \Pi^* \mathfrak{A}_P (1 - \Pi) = \widehat{\Pi} \mathfrak{A}_P^{-1} (\Pi^* + (1 - \Pi^*)) \mathfrak{A}_P (1 - \Pi) = \widehat{\Pi} (1 - \Pi) = 0$$

so $\Pi \mathfrak{A}_P^{-1} \Pi^* = 0$.

(ii, iii, iv) The corresponding proofs of [29, Theorem 4.5] can be applied directly. \square

Corollary 3.2. $\mathcal{D}_{\mathcal{B}_{\chi_\theta}}^{(T)}$ is a selfadjoint operator.

Proof. Since it is a Dirac operator, $\mathcal{D}^{(0)}$ is an elliptic symmetric differential operator of order one such that the principal symbol of its square is scalar, and it is selfadjoint with domain $H^1(\widetilde{V})$.

Since the principal symbol is unchanged by the perturbation $\mathcal{D}^{(0)} \rightarrow \mathcal{D}^{(T)} = \mathcal{D}^{(0)} + T$, and T is selfadjoint, $\mathcal{D}^{(T)}$ is also an elliptic symmetric differential operator of order one such that the principal symbol of its square is scalar, and it is selfadjoint with domain $H^1(\widetilde{V})$.

Moreover, the Green operator $\mathfrak{A}_{\mathcal{D}^{(T)}}$ of $\mathcal{D}^{(T)}$ coincides with $\mathfrak{A}_{\mathcal{D}^{(0)}} = -i\gamma_n = -\mathfrak{A}_{\mathcal{D}^{(0)}}^{-1}$ and we get by (4)

$$\begin{aligned} (1 - \Pi_-^*) \gamma_n (1 - \Pi_-) &= \Pi_+^* \gamma_n \Pi_+ = \frac{1}{4} (\gamma_n + \chi_{-\theta} \gamma_n + \gamma_n \chi_\theta + \chi_{-\theta} \gamma_n \chi_\theta) = 0, \\ \Pi_- \gamma_n \Pi_-^* &= \frac{1}{4} (\gamma_n - \chi_\theta \gamma_n - \gamma_n \chi_{-\theta} + \chi_\theta \gamma_n \chi_{-\theta}) = 0. \end{aligned}$$

Thus, (5) is satisfied and the claim follows from previous theorem (i). \square

If \widetilde{M} (with n even) has a spin structure with Dirac operator $\mathcal{D}^{(0)}$, and J denotes the ordinary charge conjugation on the spin manifold \widetilde{M} , then $\{J, \gamma_i\} = 0$. So $[J, T] = [J, \mathcal{D}^{(0)}] = 0$. Moreover, $J\gamma_{n+1} = \varepsilon' \gamma_{n+1} J$, and thus $J\chi_\theta = \varepsilon' \chi_{\varepsilon'\theta} J$, where $\varepsilon' = -1$ if $n/2$ is odd and $\varepsilon' = 1$ if $n/2$ is even.

If we define $J' := J$ if $n/2$ is even and $J' := J\gamma_{n+1}$ if $n/2$ is odd, then J' is an antilinear isometry satisfying $\mathcal{D}^{(T)} J' = (-1)^{n/2} J' \mathcal{D}^{(T)}$, $J' b J'^{-1} = b^*$ for all $b \in C^\infty(\widetilde{M})$, so is a conjugation operator as defined in [29, Definition 5.2]. Moreover $J' \chi_\theta = \chi_{(-1)^{n/2} \theta} J'$, so if $\theta = 0$ when $n/2$ is odd, $\mathcal{D}_{\mathcal{B}_{x_\theta}}^{(T)} J_+ = (-1)^{n/2} J_+ \mathcal{D}_{\mathcal{B}_{x_\theta}}^{(T)}$.

Thus, we get a spectral triple for a Dirac operator on an even dimensional manifold endowed with a chiral bag boundary condition and a totally skew-symmetric torsion:

Theorem 3.3. *For any $\theta \in \mathbb{R}$, $(\mathcal{A}_{\mathcal{D}_{\mathcal{B}_{x_\theta}}^{(T)}}, L^2(\overline{M}, \overline{V}), \mathcal{D}_{\mathcal{B}_{x_\theta}}^{(T)})$ is a spectral triple which has a simple dimension spectrum and is regular. Moreover, if \widetilde{M} is a spin manifold with Dirac operator $\mathcal{D}^{(0)}$, and if $\theta = 0$ when $n/2$ is odd, this triple has no tadpoles.*

Proof. The proof follows from the same arguments of [29, Theorem 4.8, Theorem 5.8], using Theorem 3.1 instead of [29, Theorem 4.5] and J'_+ being a conjugation operator for the spectral triple $(\mathcal{A}_{\mathcal{D}_{\mathcal{B}_{x_\theta}}^{(T)}}, L^2(\overline{M}, \overline{V}), \mathcal{D}_{\mathcal{B}_{x_\theta}}^{(T)})$ in the sense of [29, Definition 5.2]. \square

4 The spectral action

4.1 Mixed boundary condition: $\theta = 0$

We suppose from now on that the dimension is $n = 4$.

We can rewrite T as a linear combination of $-i\gamma_j \gamma_{n+1}$ matrices. More precisely, we have $T = -i\gamma_j \gamma_{n+1} T_j$ where $T_j := -\varepsilon_{jpk} T_{pkl} \in C^\infty(\widetilde{M}, \mathbb{R})$. In coordinates, $T_\mu = e_\mu^j T_j$. In this respect, the Dirac operator reads

$$\begin{cases} \mathcal{D}^{(T)} = \mathcal{D}^{(0)} + T, \\ \mathcal{D}^{(0)} = i\gamma^\mu (\partial_\mu + \frac{1}{8}[\gamma_\nu, \gamma_\rho] \sigma_\mu^{\nu\rho}), \quad T = i\gamma^\mu (-\gamma_{n+1} T_\mu), \end{cases} \quad (7)$$

where $(\sigma_\mu^{\nu\rho})$ is the spin connection. Remark that $A_\mu = 0$ and $A_\mu^5 = iT_\mu$ in the notation of [38, (3.27)]. Note also that A_μ^5 does not carry any spinor indices, which is a particularity of the chosen dimension $n = 4$.

The square of $\mathcal{D}^{(T)}$ produces a Laplace type operator on $C^\infty(\widetilde{V})$ and by [17, Lemma 4.8.1] there is a unique connection $\nabla^{\mathcal{D}^{(T)}}$ on \widetilde{V} and a unique endomorphism $E \in C^\infty(\text{End}(\widetilde{V}))$ so that

$$\mathcal{D}^{(T)2} = -(g_{\mu\nu} \nabla_\mu^{\mathcal{D}^{(T)}} \nabla_\nu^{\mathcal{D}^{(T)}} + E) \quad (8)$$

where

$$\nabla_\mu^{\mathcal{D}^{(T)}} = \partial_\mu + \omega_\mu, \quad (9)$$

$$\omega_\mu := -\frac{1}{2}[\gamma_\mu, \gamma_\nu] T_\nu \gamma_5 + \frac{1}{8}[\gamma_\rho, \gamma_\sigma] \sigma_\mu^{\rho\sigma}, \quad (10)$$

$$E := 2T^2 - \frac{1}{4}R - \gamma_5 \nabla_\mu T_\mu. \quad (11)$$

We will use the shorthand notations

$$\begin{aligned} T^2 &:= T_\mu T_\mu, \\ \gamma T &:= \gamma_\mu T_\mu, \\ T_{\mu\nu} &:= \nabla_\mu T_\nu - \nabla_\nu T_\mu. \end{aligned}$$

The field strength of the connection ω is

$$\Omega_{\mu\nu} = \gamma^\rho (\nabla_\mu T_\rho \gamma_\nu - \nabla_\nu T_\rho \gamma_\mu) \gamma_5 - T_{\mu\nu} \gamma_5 - \frac{1}{4} \gamma^\rho \gamma^\sigma R_{\rho\sigma\mu\nu} + (\gamma T) \gamma_\mu (\gamma T) \gamma_\nu - (\gamma T) \gamma_\nu (\gamma T) \gamma_\mu.$$

We follow the boundary conditions introduced in [4] and which have been considered in [5, 7, 28, 29, 38]: choose

$$\chi := \chi_0 = i\gamma_1 \gamma_2 \gamma_3 = -i\gamma_5 \gamma_4. \quad (12)$$

Thus $\chi = \chi^*$, $\{\chi, \gamma_n\} = 0$ and $[\chi, \gamma_a] = 0$ for $a = 1, 2, 3$. In particular, $\gamma_5 = i\chi\gamma_4$ is the natural chirality of the even dimensional manifold M .

As above, let $\Pi_\pm := \frac{1}{2}(\text{Id}_V \pm \chi)$ be the projections on the ± 1 eigenvalues of χ and $V_\pm := \Pi_\pm V$ be the sub-bundles of V and fix an auxiliary endomorphism S on V .

We define $\mathcal{D}_{\mathcal{B}}^{(T)}$ as the realization of $\mathcal{D}^{(T)}$ with the boundary condition $\mathcal{B}_\chi \psi = 0$ where \mathcal{B}_χ is defined on $C^\infty(\bar{V})$ by

$$\mathcal{B}_\chi \psi := \Pi_- \psi|_{\partial M}, \quad \psi \in C^\infty(\bar{V}).$$

The mixed boundary conditions $\mathcal{B} = \mathcal{B}(\chi, S)$ are defined on $\mathcal{D}_{\mathcal{B}}^{(T)^2}$ by $\mathcal{B}\psi = 0$ where

$$\mathcal{B}\psi := \Pi_- (\mathcal{D}_+^{(T)} \psi)|_{\partial M} \oplus \Pi_- \psi|_{\partial M}, \quad \psi \in C^\infty(\bar{V}). \quad (13)$$

As in [4], $\mathcal{D}^{(T)^2}$ could also have considered directly with domain given by $\mathcal{B}\psi = 0$ with

$$\mathcal{B}\psi := \Pi_+ (\nabla_n^{\mathcal{D}^{(T)}} + S) \Pi_+ \psi|_{\partial M} \oplus \Pi_- \psi|_{\partial M}, \quad \psi \in C^\infty(\bar{V}). \quad (14)$$

For the choice of $\chi = -i\gamma_5 \gamma_n$, these last two boundary conditions are equivalent [4, Lemma 7] if $S = \frac{1}{2} \Pi_+ (-[i\gamma_n, T] - L_{aa}\chi) \Pi_+$, thus using $\{\gamma_5, \chi\} = 0$,

$$S = -\frac{1}{2} L_{aa} \Pi_+. \quad (15)$$

These boundary conditions generalizes Dirichlet ($\Pi_- = \text{Id}_V$, so $\chi = -\text{id}_V$) and Neumann–Robin ($\Pi_+ = \text{Id}_V$, so $\chi = \text{id}_V$) conditions.

While Corollary 3.2 is the real motivation for the above choice of boundary conditions, it is nevertheless worthwhile to recall as in [7, 8] that signs and ratios in the spectral action given by that choice, are identical to the Euclidean action used in gravitation [25] for a zero torsion.

4.2 Computation of the spectral action for $\theta = 0$

The Chamseddine–Connes spectral action is

$$\mathcal{S}(\mathcal{D}_{\mathcal{B}_\chi}^{(T)}, \Phi, \Lambda) := \text{Tr} (\Phi((\mathcal{D}_{\mathcal{B}_\chi}^{(T)})^2 / \Lambda^2)) \quad (16)$$

where Φ is any positive even function viewed as a cut-off with $\Lambda \in \mathbb{R}$.

As detailed in [12], the spectral action (16) of the above spectral triple (Theorem 3.3, with $\theta = 0$) is related to the asymptotic expansion

$$\mathrm{Tr}(\exp(-t(\mathcal{D}_{\mathcal{B}_\chi}^{(T)})^2)) \underset{t \downarrow 0}{\sim} \sum_{k \in \mathbb{N}} t^{(k-4)/2} a_k(1, \mathcal{D}^{(T)}, \chi) \quad (17)$$

via

$$\mathcal{S}(\mathcal{D}_{\mathcal{B}_\chi}^{(T)}, \Phi, \Lambda) = \Lambda^4 \Phi_4 a_0 + \Lambda^3 \Phi_3 a_1 + \Lambda^2 \Phi_2 a_2 + \Lambda \Phi_1 a_3 + \Phi(0) a_4 + \mathcal{O}(\Lambda^{-1}) \quad (18)$$

with $\Phi_k := \frac{1}{\Gamma(k/2)} \int_0^\infty \Phi(s) s^{k/2-1} ds$. (Note that the decomposition (17) is not the same as in [12, Theorem 1.145] or [29, (1)] so the coefficients Φ_k are different.)

So, using [18, 30, 38] with the decomposition of a_k in $T = 0$ -part and remainder, we get

$$a_0^{(T)} = a_0^{(0)} = \frac{1}{4\pi^2} \mathrm{Vol}(M), \quad (19)$$

$$a_1^{(T)} = a_1^{(0)} = 0, \quad (20)$$

$$a_2^{(T)} = a_2^{(0)} + \frac{1}{6(4\pi)^2} \int_M 48T^2, \quad (21)$$

$$a_2^{(0)} := -\frac{1}{6(4\pi)^2} \left(\int_M 2R + \int_{\partial M} 4L \right).$$

(All integrations are with respect of the volume form on M and of the induced volume form on ∂M .) Using “;” for the tangential covariant differentiation on \widetilde{M} and “:” for the same on the boundary, the computation of

$$\chi_{;a} = iT^c[\gamma_a, \gamma_c]\gamma_n + i\gamma_5 L_{ab}\gamma_b,$$

yields

$$\begin{cases} a_3^{(T)} = a_3^{(0)} - \frac{1}{96(4\pi)^{3/2}} \int_{\partial M} 96T_a T_a, \\ a_3^{(0)} := \frac{1}{96(4\pi)^{3/2}} \int_{\partial M} 3L^2 - 6L_{ab}L_{ab}. \end{cases} \quad (22)$$

For the next coefficient a_4 , we have two contributions [3, 37] (where a common pre-factor of $\frac{4}{360(4\pi)^2}$ is omitted):

- Contributions of individual volume terms in a_4 :

$$\begin{aligned} 60RE & : 120T^2R - 15R^2 \\ 180E^2 & : 720T^4 + \frac{45}{4}R^2 - 180T^2R + 180(\nabla T)^2 \\ 30\Omega^2 & : -60T_{\mu\nu}T^{\mu\nu} - 60(\nabla T)^2 - 120(\nabla_\mu T_\nu)(\nabla_\nu T_\mu) \\ & \quad - \frac{15}{4}R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma} + 60T^2R - 120T^\mu T^\nu R_{\mu\nu} - 720T^4 \\ R^2\text{-terms} & : 5R^2 - 2R_{\mu\nu}R^{\mu\nu} + 2R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma} \end{aligned}$$

- Contributions of individual surface terms in a_4 :

$$\begin{aligned}
180\chi E_{;n} & : 0 \\
30\chi R_{;n} & : 0 \\
24L_{aa:bb} & : 0 \\
120EL_{aa} & : (240T^2 - 30R)L \\
RL & : 20RL + 4R_{anan}L - 12R_{anbn}L_{ab} + 4R_{abcb}L_{ac} \\
L^3\text{-terms} & : \frac{1}{21}(160L^3 - 48L_{ab}L_{ab}L + 272L_{ab}L_{bc}L_{ac}) \\
720SE & : -360LT^2 + 45RL \\
120SR & : -30RL \\
144SL_{aa}L_{bb} & : -36L^3 \\
48SL_{ab}L_{ab} & : -12L_{ab}L_{ab}L \\
480S^2L_{aa} & : 60L^3 \\
480S^3 & : -30L^3 \\
120S_{;aa} & : 0 \\
60\chi\chi_{;a}\Omega_{an} & : -240T_cT_{c;n} + 30L_{ab}R_{anbn} + 120LT_cT_c - 120L_{ab}T_aT_b \\
-12L\chi_{;a}\chi_{;a} & : -12L_{ac}L_{ac}L - 96LT_cT_c \\
-24L_{ab}\chi_{;a}\chi_{;b} & : -24L_{ab}L_{bc}L_{ac} - 96LT_cT_c + 96L_{ab}T_aT_b \\
-120S\chi_{;a}\chi_{;a} & : 30L_{ac}L_{ac}L + 240LT_cT_c
\end{aligned}$$

Remark that

$$\int_M 60 \operatorname{tr}(E_{;\mu\mu}) + \int_{\partial M} \operatorname{tr}((240\Pi_+ - 120\Pi_-)E_{;n}) = 0, \quad (23)$$

$$\int_M 12 \operatorname{tr}(R_{;\mu\mu}) + \int_{\partial M} \operatorname{tr}((42\Pi_+ - 18\Pi_-)R_{;n}) = 0. \quad (24)$$

The following identity will be useful:

$$\int_M (\nabla T)^2 = \int_M \nabla_\mu T_\nu \cdot \nabla_\nu T_\mu + R_{\mu\nu} T^\mu T^\nu + \int_{\partial M} -2T_n T_{a;a} + T_n^2 L + L_{ab} T_a T_b.$$

By adding up all contributions, one obtains

$$\begin{aligned}
a_4^{(T)} = a_4^{(0)} + \frac{4}{360(4\pi)^2} & \left(\int_M -60T_{\mu\nu}T_{\mu\nu} \right. \\
& \left. + \int_{\partial M} 48LT_aT_a + 96L_{ab}T_aT_b - 240T_aT_{a;n} - 240T_nT_{a;a} \right), \quad (25)
\end{aligned}$$

$$\begin{aligned}
a_4^{(0)} := \frac{4}{360(4\pi)^2} & \left(\int_M \frac{5}{4}R^2 - 2R_{\mu\nu}R_{\mu\nu} - \frac{7}{4}R_{\mu\nu\sigma\rho}R_{\mu\nu\sigma\rho} \right. \\
& \left. + \int_{\partial M} 5LR + 4R_{anan}L + 4R_{abcb}L_{ac} + 18L_{ab}R_{anbn} + \frac{34}{21}L^3 + \frac{26}{7}LL_{ab}L_{ab} - \frac{232}{21}L_{ab}L_{bc}L_{ac} \right).
\end{aligned}$$

The spectral action, up to the order Λ^0 , is obtained by the substitution of (21), (22) and (25) in (17).

4.3 Volume part of the spectral action

Let us discuss the volume part of the spectral action, which reads

$$\mathcal{S}(\mathcal{D}^{(T)}, \Phi)_{\text{vol}} = \mathcal{S}(\mathcal{D}^{(0)}, \Phi)_{\text{vol}} + \int_M 8\Lambda^2 \Phi_2 T^2 - \frac{2}{3} \Phi(0) T_{\mu\nu} T_{\mu\nu} + \mathcal{O}(\Lambda^{-1}). \quad (26)$$

This (amazingly simple) result is fully consistent with the previous computations of a_2, a_4 without boundary on Riemann–Cartan manifolds [10, 20, 21, 34]. The paper [23] gives a much longer expression for a_4 containing 6 torsion-dependent terms instead of one in our case. Let us show that the simplification appearing in our calculations is not just a lucky coincidence, but rather reflects an important symmetry which has the spectral action at this order.

Consider the identity

$$\mathcal{D}(\varphi) := e^{\varphi(x)\gamma_5} \mathcal{D}^{(T)} e^{\varphi(x)\gamma_5} = i\gamma^\mu (\nabla_\mu - \gamma_5(T_\mu - \partial_\mu\varphi)) \quad (27)$$

for an arbitrary scalar function $\varphi(x)$, which tells us that the gradient transformation of T_μ is a chiral transformation of the Dirac operator. The corresponding infinitesimal variation of the heat trace can be computed by repeating the same steps as were used to derive conformal variations, see [17, 18]:

$$\frac{d}{d\varepsilon} \Big|_{\varepsilon=0} \text{Tr} \left(e^{-t(\mathcal{D}(\varepsilon\varphi))^2} \right) = - \text{Tr} \left(4t\varphi\gamma_5 (\mathcal{D}(\varphi))^2 e^{-t(\mathcal{D}(\varphi))^2} \right) = 4t \frac{d}{dt} \text{Tr} \left(\varphi\gamma_5 e^{-t(\mathcal{D}(\varphi))^2} \right). \quad (28)$$

Next, we introduce a generalization of the heat trace (17) "smeared" with an arbitrary endomorphism F of the spin bundle, which also has an asymptotic expansion

$$\text{Tr} \left(F e^{-t(\mathcal{D}_{\mathcal{B}\chi}(\varphi))^2} \right) \underset{t \downarrow 0}{\sim} \sum_{k \in \mathbb{N}} a_k(F, \mathcal{D}(\varphi), \chi) t^{(k-4)/2}.$$

By expanding (28) in power series of t , we obtain the equation

$$\frac{d}{d\varepsilon} \Big|_{\varepsilon=0} (a_k)_{\text{vol}}(1, \mathcal{D}(\varepsilon\varphi), \chi) = 2(k-4) (a_k)_{\text{vol}}(\varphi\gamma_5, \mathcal{D}(0), \chi) \quad (29)$$

yielding that the volume part of a_4 is invariant under the transformations (27). Consequently, only the terms which are invariant under $T_\mu \rightarrow T_\mu - \partial_\mu\varphi$ may appear in the volume part of a_4 . The only allowed torsion dependent term is, therefore, $T_{\mu\nu}T^{\mu\nu}$. All couplings of the torsion to the Riemann curvature as well as the T^4 and $(\nabla T)^2$ terms are excluded.

These arguments cannot be used to control the boundary terms since the boundary conditions are not invariant under chiral transformations. The chiral invariance can be restored if one uses the chiral bag boundary condition, but the parameter θ should be made coordinate-dependent, see [33] for a discussion.

When the manifold is non-compact, similar computations of coefficients $a_n^{(T)}$ can be done via a smearing function f as in [18, 38]. However, the cancellations (23), (24) are not satisfied so formulae are more lengthy and, much more important, Theorem 3.3 has to be proved in this context where the spectrum of $\mathcal{D}_{\mathcal{B}}^{(T)}$ is no more discrete.

Moreover, if A is a selfadjoint one-form, i.e. $A = \sum_k a_k[\mathcal{D}_{\mathcal{B}}^{(T)}, b_k]$ with $a_k, b_k \in \mathcal{A}_{\mathcal{D}_{\mathcal{B}}^{(T)}}$, then $A = i\gamma_\mu A_\mu$ with $A_\mu \in C^\infty(\overline{M}, i\mathbb{R})$ and it is possible to compute the fluctuations of the Dirac operator $\mathcal{D}_{\mathcal{B}}^{(T)} + A$ applying [38, eq. (3.26)]. Notice that a real spectral triple has no fluctuations since $\mathcal{D}_{\mathcal{B}}^{(T)} + A + JAJ^{-1} = \mathcal{D}_{\mathcal{B}}^{(T)}$.

5 Stability of spectral action with respect to θ

Let us prove now that $\theta = 0$ is a critical point of the spectral action which justifies the computation of the spectral action in section 4.2.

Proposition 5.1. *In any even dimension, the value $\theta = 0$ is a critical point for the coefficients at any order of the chiral bag spectral action $\mathcal{S}(\mathcal{D}_{\mathcal{B}_{\chi_\theta}}^{(T)}, \Phi, \Lambda)$: precisely, for any $k \in \mathbb{N}$,*

$$\partial_\theta|_{\theta=0} a_k(1, \mathcal{D}^{(T)}, \chi_\theta) = 0. \quad (30)$$

Proof. We shall need the following property

$$\gamma_{n+1} \mathcal{D}^{(T)} + \mathcal{D}^{(T)} \gamma_{n+1} = 0, \quad (31)$$

which is, of course, satisfied by $\mathcal{D}^{(T)}$ given in (7).

If F is an auxiliary endomorphism on \tilde{V} , we define the coefficients $a_k(F, \mathcal{D}^{(T)}, \chi_\theta)$, such that the following complete asymptotic expansion at $t \rightarrow 0$ holds:

$$\mathrm{Tr} \left(F e^{-t(\mathcal{D}_{\mathcal{B}_{\chi_\theta}}^{(T)})^2} \right) \underset{t \downarrow 0}{\sim} \sum_{k \in \mathbb{N}} a_k(F, \mathcal{D}^{(T)}, \chi_\theta) t^{(k-n)/2}.$$

The derivative of the heat kernel coefficients with respect to θ was calculated in [19]:

$$\partial_\theta|_{\theta=0} a_k(1, \mathcal{D}^{(T)}, \chi_\theta) = (n - k) a_k(\gamma_{n+1}, \mathcal{D}^{(T)}, \chi_0). \quad (32)$$

Let V_1 (resp. V_2) be the eigenspace of γ_5 for the eigenvalue 1 (resp. -1) so that $\bar{V} = V_1 \oplus V_2$. By (31), $\mathcal{D}^{(T)} = \mathcal{D}_1^{(T)} \oplus \mathcal{D}_2^{(T)}$ where $\mathcal{D}_1^{(T)}$ is the restriction of $\mathcal{D}^{(T)}$ which is elliptic from $C^\infty(V_1)$ into $C^\infty(V_2)$, and $\mathcal{D}_2^{(T)}$ is the formal adjoint of $\mathcal{D}_1^{(T)}$. The mixed chiral boundary condition \mathcal{B}_{χ_0} can be decomposed along $V_1 \oplus V_2$ as $\mathcal{B}_1 \oplus \mathcal{B}_2$ so that $\mathcal{D}_{\mathcal{B}_{\chi_0}}^{(T)} = (\mathcal{D}_1^{(T)})_{\mathcal{B}_1} \oplus (\mathcal{D}_2^{(T)})_{\mathcal{B}_2}$. By [17, Theorem 1.9.3],

$$\mathrm{Tr}(\gamma_{n+1} e^{-t(\mathcal{D}_{\mathcal{B}_{\chi_0}}^{(T)})^2}) = \mathrm{Index}((\mathcal{D}_1^{(T)})_{\mathcal{B}_1}) \quad \text{for all } t \quad (33)$$

where on the right hand side we have the index of the spin complex, i.e., the difference between the numbers of zero eigenmodes of $\mathcal{D}_{\mathcal{B}_{\chi_0}}^{(T)}$ with positive and negative chirality, or more precisely $\dim \mathrm{Ker}(\mathcal{D}_1^{(T)})_{\mathcal{B}_1} - \dim \mathrm{Ker}(\mathcal{D}_2^{(T)})_{\mathcal{B}_2}$, see [13, 17]. By expanding both sides of this relation in asymptotic series in t , one obtains

$$a_k(\gamma_{n+1}, \mathcal{D}^{(T)}, \chi_0) = 0, \quad \text{for } k \neq n. \quad (34)$$

Combining (34) with (32), one obtains the result (30). \square

Remark 5.2. *More information can be provided about the scale invariant terms $a_n(1, \mathcal{D}^{(T)}, \chi_\theta)$ and $a_n(\gamma_{n+1}, \mathcal{D}^{(T)}, \chi_0)$. First, $a_n(1, \mathcal{D}^{(T)}, \chi_\theta)$ is actually independent of θ as shown in [19]. Moreover, the value of $a_n(\gamma_{n+1}, \mathcal{D}^{(T)}, \chi_0)$ is given by the t -invariant term of (33), that is: $a_n(\gamma_{n+1}, \mathcal{D}^{(T)}, \chi_0) = \mathrm{Index}((\mathcal{D}_1^{(T)})_{\mathcal{B}_1})$.*

Remark 5.3. *In dimension 4, it is actually possible to verify (34) for $k \in \{0, 1, 2, 3\}$ by using the formulae of [14, 33]. Thus, with (32), we can directly get (30) for $k \in \{0, 1, 2, 3, 4\}$. It is for higher coefficient orders that the index formula (33) reveals its power.*

6 Discussion

Let us discuss some physical implications of the spectral action we have just calculated. To achieve positive kinetic energies of the fields of the Standard Model, the coefficient $\Phi(0)$ has to be positive and this is the case. Consequently, the kinetic for T seems to be negative. However, T is an axial vector field rather than a vector field. The standard roles of the Wick rotation for such fields include an additional multiplier of i : an “incorrect” sign of the kinetic energy in Euclidean space corresponds to a positive kinetic energy in the physical Minkowski case. The spectral action also predicts a mass of the torsion, which is restricted from various physical considerations, see [36] for a recent overview, or [35] for a more detailed exposition. It is considerably harder to derive physical consequences of the boundary part of the spectral action. The only thing we can say at the moment is that there are no linear torsion terms, so that it is less probable that certain boundary configurations will decay into torsion.

Our main prediction regarding the chiral bag boundary conditions is that the point $\theta = 0$ is a critical point of the spectral action. This result is based on rather general arguments involving the Index Theorem, and therefore is valid to all orders in Λ and, presumably, for a larger class of spectral triples.

Acknowledgments

We thank Thomas Schücker for useful discussions. This work was started during the visit of the third author (D. V.) to Marseille, which was supported by the CAPES-Mathamsud program and the Université de Provence. D. V. also thanks CNPq and FAPESP for partial support.

References

- [1] A. A. Andrianov and F. Lizzi, “Bosonic spectral action induced from anomaly cancellation,” *JHEP* **1005** (2010), 057, arXiv:1001.2036 [hep-th].
- [2] C. Beneventano, P. Gilkey, K. Kirsten and E. Santangelo, “Strong ellipticity and spectral properties of chiral bag boundary conditions”, *J. Phys. A: Math. Gen.* **36** (2003), 11533–11543, arXiv:hep-th/0306156.
- [3] T. P. Branson and P. B. Gilkey, “The asymptotics of the Laplacian on a manifold with boundary,” *Commun. Part. Diff. Eq.* **15** (1990), 245–272.
- [4] T. Branson and P. Gilkey, “Residues of the eta function for an operator of Dirac type with local boundary conditions”, *Differential Geometry and its Applications* **2** (1992), 249–267.
- [5] T. Branson, P. Gilkey, K. Kirsten and D. Vassilevich, “Heat kernel asymptotics with mixed boundary conditions”, *Nucl. Phys. B* **563** (1999), 603–626, arXiv:hep-th/9906144.
- [6] A. Chamseddine and A. Connes, “The spectral action principle”, *Commun. Math. Phys.* **186** (1997), 731–750, arXiv:hep-th/9606001.
- [7] A. Chamseddine and A. Connes, “Quantum gravity boundary terms from the spectral action on noncommutative space”, *PRL* **99** (2007), 071302, arXiv:0705.1786 [hep-th].

- [8] A. Chamseddine and A. Connes, “Noncommutative geometric spaces with boundary: spectral action”, arXiv:1008.3980v1 [hep-th].
- [9] A. Chamseddine, A. Connes and M. Marcolli, “Gravity and the standard model with neutrino mixing”, *Adv. Theor. Math. Phys.* **11** (2007), 991–1089, arXiv:hep-th/0610241.
- [10] G. Cognola and S. Zerbini, “Heat kernel expansion in geometric fields”, *Phys. Lett. B* **195** (1987), 435–438.
- [11] A. Connes, *Noncommutative Geometry*, Academic Press, London and San Diego, 1994.
- [12] A. Connes and M. Marcolli, *Noncommutative Geometry, Quantum Fields and Motives*, Colloquium Publications, Vol. 55, American Mathematical Society, 2008.
- [13] T. Eguchi, P. B. Gilkey and A. J. Hanson, “Gravitation, gauge theories and differential geometry,” *Phys. Rept.* **66** (1980), 213–393.
- [14] G. Esposito, P. Gilkey and K. Kirsten, “Heat kernel coefficients for chiral bag boundary conditions”, *J. Phys. A: Math. Gen.* **38** (2005), 2259–2276, arXiv:math/0510156.
- [15] G. Esposito and K. Kirsten, “Chiral bag boundary conditions on the ball”, *Phys. Rev. D* **66** (2002), 085014, arXiv:hep-th/0207109.
- [16] Th. Friedrich and S. Sulanke, “Ein kriterium für die formale selbstadjungiertheit des Dirac-operators”, *Coll. Math.* vol XL (1979), 239–247.
- [17] P. B. Gilkey, *Invariance Theory, the heat equation, and the Atiyah–Singer Index theorem*, Mathematics Lecture Series, vol. 11, Publish or Perish, Inc., Wilmington, Delaware, 1984,
- [18] P. B. Gilkey, *Asymptotic Formulae in Spectral Geometry*, Chapman & Hall/CRC, Boca Raton, FL, 2004.
- [19] P. Gilkey and K. Kirsten, “Stability theorems for chiral bag boundary conditions”, *Lett. Math. Phys.* **73** (2005), 147–163, arXiv:math/0510152.
- [20] W. H. Goldthorpe, “Spectral geometry and $SO(4)$ gravity in a Riemann–Cartan spacetime”, *Nucl. Phys. B* **170** (1980), 307–328.
- [21] G. Grensing, “Induced gravity for nonzero torsion”, *Phys. Lett.* **169B** (1986), 333–336.
- [22] G. Grubb, *Functional calculus of pseudodifferential boundary problems*, Second edition, Progress in Mathematics **65**, Birkhäuser, Boston, 1996.
- [23] F. Hanish, F. Pfäffle and C. A. Stephan, “The spectral action for Dirac operators with skew-symmetric torsion”, arXiv:0911.5074 [hep-th].
- [24] P. Hasenfratz and J. Kuti, “The quark bag model,” *Phys. Rept.* **40** (1978), 75–179.
- [25] S. Hawking and G. Horowitz, “The gravitational Hamiltonian, action, entropy and surface terms”, *Class. Quantum Grav.* **13** (1996), 1487–1498.
- [26] B. R. Heckel, E. G. Adelberger, C. E. Cramer, T. S. Cook, S. Schlamminger and U. Schmidt, “Preferred-frame and CP-violation tests with polarized electrons”, *Phys. Rev. D* **78**, 092006 (2008), [arXiv:0808.2673 [hep-ex]].

- [27] P. Hrasco and J. Balog, “The fermion boundary condition and the θ -angle in QED₂’,” Nucl. Phys. B **245** (1984), 118–126.
- [28] B. Iochum and C. Levy, “Tadpoles and commutative spectral triples”, arXiv:0904.0222 [math-ph], to appear in J. Noncommut. Geom.
- [29] B. Iochum and C. Levy, “Spectral triples and manifolds with boundary”, to appear in J. Funct. Anal., arXiv:1001.3927 [math-ph].
- [30] K. Kirsten, *Spectral Functions in Mathematics and Physics*, Chapman & Hall/CRC, Boca Raton, 2002.
- [31] H. Kleinert, “New gauge symmetry in gravity and the evanescent role of torsion”, arXiv:1005.1460 [gr-qc].
- [32] V. A. Kostelecky, N. Russell and J. Tasson, “Constraints on torsion from Lorentz violation”, Phys. Rev. Lett. **100**, 111102 (2008), arXiv:0712.4393 [gr-qc].
- [33] V. N. Marachevsky and D. V. Vassilevich, “Chiral anomaly for local boundary conditions”, Nucl. Phys. B **677** (2004), 535–552, arXiv:hep-th/0309019.
- [34] Y. N. Obukhov, “Spectral geometry of the Riemann–Cartan space-time”, Nucl. Phys. B **212** (1983), 237–254.
- [35] I. L. Shapiro, “Physical aspects of the space-time torsion”, Phys. Rept. **357** (2002), 113–213, arXiv:hep-th/0103093.
- [36] I. L. Shapiro, “Effective QFT and what it tells us about dynamical torsion”, arXiv:1007.5294 [hep-th].
- [37] D. V. Vassilevich, “Vector fields on a disk with mixed boundary conditions”, J. Math. Phys. **36** (1995) 3174–3182, arXiv:gr-qc/9404052.
- [38] D. V. Vassilevich, “Heat kernel expansion: user’s manual”, Phys. Rep. **388** (2003), 279–360, arXiv:hep-th/0306138.
- [39] A. Wipf and S. Dürr, “Gauge theories in a bag”, Nucl. Phys. B **443** (1995), 201–232, arXiv:hep-th/9412018.