



**HAL**  
open science

## Optical precursors in the singular and weak dispersion limits: Reply to reply to comment

Bruno Macke, Bernard Ségard

► **To cite this version:**

Bruno Macke, Bernard Ségard. Optical precursors in the singular and weak dispersion limits: Reply to reply to comment. 2011. hal-00558498v2

**HAL Id: hal-00558498**

**<https://hal.science/hal-00558498v2>**

Preprint submitted on 21 Feb 2011

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Optical precursors in the singular and weak dispersion limits: Reply to reply to comment

Bruno Macke and Bernard Ségard\*

*Laboratoire de Physique des Lasers, Atomes et Molécules ,  
CNRS et Université Lille 1, 59655 Villeneuve d'Ascq, France*

The reply by Oughstun *et al.* [J. Opt. Soc. Am. B **28**, 468-469 (2011)] to our comment [J. Opt. Soc. Am. B **28**, 450-452 (2011)] on their recently published paper [J. Opt. Soc. Am. B **27**, 1664-1670 (2010)] is shown to make no response to the main points raised in our comment and even to bring additional confusion. The present reply provides the necessary clarifications.

OCIS codes: 260.2030, 320.5550, 320.2250.

PACS numbers: 42.25.Bs, 42.50.Md, 41.20.Jb

In a recent comment [1], we pointed out inconsistencies in the paper by Oughstun *et al.* on Sommerfeld and Brillouin precursors [2]. Their reply [3] makes no response to the main points raised in our comment. In addition it is written in such a manner that it may confuse and mislead the unfortunate reader. It thus appears necessary to dot the i's and cross the t's.

The first point concerns the analytical form of the Brillouin precursor. No need to look at the Oughstun's papers abundantly cited in [2, 3] for (possibly) finding the remarkably simple expression  $E_B(z, t') = \frac{b}{\omega_c} \text{Ai}(-bt')$ . As mentioned in our comment [1], this expression was established by Brillouin himself as far back as 1932 [4]. The way by which we have retrieved this result shows that it will be a good approximation of the exact signal when the instantaneous frequency of the precursor is large (small) compared to the damping rate  $\delta$  (the carrier frequency  $\omega_c$ ). This condition is met in the singular dispersion limit, at least for the first oscillations of the precursor. Since  $b \propto z^{-1/3}$  ( $z$  propagation distance), the previous result *analytically* shows that the amplitude of the Brillouin precursor also scales as  $z^{-1/3}$ , a result that was obtained only *numerically* in [2]. The expression of  $E_B(z, t')$  also shows that the instantaneous period of the precursor is proportional to  $1/b$ . In particular, the first half-period  $HP_1$  of the precursor, defined as the time interval between the first maximum and the first minimum of  $E_B(z, t')$ , is equal to  $2.22/b$ . For the parameters used to obtain Fig.3 in [2], we get  $HP_1 = 0.23$  ps in agreement with the exact result (see Fig.1 of our comment [1]). On their side, Oughstun *et al.* give in the equation (15) of [2] an *estimate* of the half-period that, in the same conditions, leads to a value smaller than the right one by about five orders of magnitude!

Contrary to the claim made in [2], we have shown in our comment [1] that the precursors are catastrophically affected by the rise-time  $T_r$  of the incident field even when the latter is considerably faster than the damping time  $\tau = 1/\delta$  of the medium. Our analysis was supported by numerical simulations made for rise times as fast as  $\tau/500$  and  $\tau/100$ . Without disputing the validity of our

simulations, Oughstun *et al.* claim in their reply that our numerical results “do not properly lie in the asymptotic regime of sufficiently large propagation distances” [3]. In fact, the cases considered in our simulations are exactly those considered in their paper [2]. In particular, as noticed both in the text and in the figure caption, the parameters of our Fig.3 are strictly identical to those of their Fig.3. However, as suggested by Oughstun *et al.*, we have made additional simulations for larger propagation distances. These simulations show that the rise-time effects become more and more catastrophic when  $z$  increases. This result was obviously foreseeable. Indeed the effect of the rise-time, *irrespective of the precise shape of the rise*, is to contract the spectrum of the incident field about the carrier frequency  $\omega_c$ . On the other hand, when  $z$  increases, the medium becomes strongly absorbing on a broader and broader spectral region, keeping some transparency only at frequencies  $\omega \gg \omega_c$  and  $\omega \ll \omega_c$ . The precursors, associated with the transparency regions, then cannot be excited by the incident field.

Oughstun *et al.* also invoke in their reply [3] an eventual interaction between the Brillouin precursor and the so-called main signal. Such an interaction, if it exists, is extremely small in the conditions of our figure 3 for which the amplitude  $A_{MS}$  of the main signal is only  $e^{-10}$ , that is about 3700 times smaller than the amplitude  $A_B$  of the Brillouin precursor. The interaction becomes completely negligible when the propagation distance is increased in order to better “lie in the asymptotic regime”. When the optical thickness at  $\omega_c$  is equal to 100 ( $A_{MS} = e^{-100} \approx 3.7 \times 10^{-44}$ ), we have determined the amplitude of the Brillouin precursor for  $T_r = 0$  and  $T_r = \tau/500$  (error-function envelope signal) in the conditions of Fig.3 and Fig.5 in [2]. These figures were intended to respectively illustrate the singular and weak dispersion limits. We find  $A_B(0) \approx 7.9 \times 10^{-3}$  and  $A_B(\tau/500) \approx 6.7 \times 10^{-10}$  in the former case;  $A_B(0) \approx 3.6 \times 10^{-2}$  and  $A_B(\tau/500) \approx 2.2 \times 10^{-8}$  in the latter one. These results confirm *in both cases* the catastrophic effect of the rise time of the incident field even when it is

500 times smaller than the damping time  $\tau = 1/\delta$  of the medium.

In summary, the analysis presented in our comment [1] on the paper [2] is entirely confirmed.



\* Electronic address: [bernard.segard@univ-lille-1.fr](mailto:bernard.segard@univ-lille-1.fr)

- [1] B. Macke and B. Ségard, “Comment on Optical precursor in the singular and weak dispersion limits”, *J. Opt. Soc. Am. B* **28**, 450-452 (2011). See also e-print arXiv: 1009.3146.
- [2] K.E. Oughstun, N.A. Cartwright, D.J. Gauthier, and H. Jeong, “Optical precursor in the singular and weak dispersion limits”, *J. Opt. Soc. Am. B* **27**, 1664-1670 (2010). Most of the content of this paper can also be found in K.E. Oughstun, *Electromagnetic and Optical Pulse Propagation*

*2 : Temporal Pulse Dynamics in Dispersive Attenuative Media*, (Springer, New York, 2009), pp 652-656.

- [3] K.E. Oughstun, N.A. Cartwright, D.J. Gauthier, and H. Jeong, “Optical precursor in the singular and weak dispersion limits: Reply to Comment”, *J. Opt. Soc. Am. B* **28**, 468-469 (2011).
- [4] L. Brillouin, “Propagation des ondes électromagnétiques dans les milieux matériels,” in *Comptes Rendus du Congrès International d’Electricité, Paris 1932* (Gauthier-Villars 1933), Vol.2, pp. 739-788. An English adaptation of this paper can be found in L. Brillouin, *Wave Propagation and Group Velocity* (Academic 1960), Ch. IV and Ch. V (see in particular pp. 108-109 and pp. 127-128). Note that the Airy function  $\mathcal{A}(x)$  used by Brillouin differs from the standard one  $\text{Ai}(x)$ , with  $2\pi\text{Ai}(-x) = 3^{1/3}\mathcal{A}(3^{1/3}x)$ . Peak amplitudes of  $\mathcal{A}(x)$  and  $\text{Ai}(-x)$  are respectively 2.33 and 0.536.