



HAL
open science

Friezes, Strings and Cluster Variables

Ibrahim Assem, Grégoire Dupont, Ralf Schiffler, David Smith

► **To cite this version:**

Ibrahim Assem, Grégoire Dupont, Ralf Schiffler, David Smith. Friezes, Strings and Cluster Variables. Glasgow Mathematical Journal, 2011, 32 p. hal-00518249v2

HAL Id: hal-00518249

<https://hal.science/hal-00518249v2>

Submitted on 7 Mar 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

FRIEZES, STRINGS AND CLUSTER VARIABLES

IBRAHIM ASSEM, GRÉGOIRE DUPONT, RALF SCHIFFLER AND DAVID SMITH

ABSTRACT. To any walk in a quiver, we associate a Laurent polynomial. When the walk is the string of a string module over a 2-Calabi-Yau tilted algebra, we prove that this Laurent polynomial coincides with the corresponding cluster character of the string module, up to an explicit normalising monomial factor.

INTRODUCTION

In the early 2000, S. Fomin and A. Zelevinsky introduced the class of cluster algebras with the purpose of building a combinatorial framework for studying total positivity in algebraic groups and canonical bases in quantum groups, see [27]. Since then, the study of cluster algebras was shown to be connected to several areas of mathematics, notably combinatorics, Lie theory, Poisson geometry, Teichmüller theory, mathematical physics and representation theory of algebras.

A cluster algebra is a commutative algebra generated by a set of variables, called *cluster variables*, obtained recursively by a combinatorial process known as *mutation*, starting from an initial set of cluster variables, the *initial cluster*, and a quiver without cycles of length at most two. One of the most remarkable facts about cluster variables is that they can be expressed as Laurent polynomials in terms of the initial cluster variables [27]; this is the so-called *Laurent phenomenon*. Also, it is conjectured that the coefficients in this expression are always non-negative; this is the *positivity conjecture*. The problem of computing explicitly the cluster variables is a difficult one and has been extensively studied. The most general results known at the present time are in [30, 45, 42].

In order to compute cluster variables, one may use *friezes*. Friezes, which go back to works of Coxeter and Coxeter-Conway [21, 19, 20], are an efficient combinatorial tool which mimics the application of mutations on sinks or sources of the given quiver, hence an obvious combinatorial connection with cluster algebras [17, 14, 46, 40, 5]. It is now known that connections between friezes and cluster algebras are deeper than just combinatorics, see for instance [24, 4, 38, 7, 29]. Our starting point for the present paper was the result in [5] giving an explicit formula as a product of 2×2 matrices for all cluster variables in coefficient-free cluster algebras of type \mathbb{A} and all but finitely many cluster variables in coefficient-free cluster algebras of type $\tilde{\mathbb{A}}$, thus explaining at the same time the Laurent phenomenon and positivity. Our objective here is to show that the same technique can be used for computing the cluster variables associated with the string modules over a 2-Calabi-Yau tilted algebra (in the sense of [48]).

Besides friezes, our second main tool is the notion of a *cluster character*. In [14], Caldero and Chapoton noticed that cluster variables in simply-laced coefficient-free

Date: March 14, 2011.

2010 Mathematics Subject Classification. 13F60, 16G20.

cluster algebras of finite type can be expressed as generating series of Euler-Poincaré characteristics of Grassmannians of submodules. Generalising this work, Caldero-Keller [16], Palu [44] and Fu-Keller [30] introduced the notion of a cluster character associating to each module M over a 2-Calabi-Yau tilted algebra B_T a certain Laurent polynomial X_M^T allowing one to compute a corresponding cluster variable. In general, cluster characters are hard to compute because one first needs to find the Euler characteristics of Grassmannians of submodules, and then the dimensions of certain Hom-spaces in the corresponding 2-Calabi-Yau category.

One class of algebras, however, whose representation theory is reasonably well-understood is the class of string algebras, introduced by Butler and Ringel in [13] (see also [53]). In particular, indecomposable modules over string algebras are partitioned into two sets: string and band modules, and only string modules can be associated with cluster variables. The Euler characteristics of Grassmannians of submodules of string modules were computed by Cerulli and Haupt [18, 33]. Nevertheless, their methods do not allow one to compute explicitly the associated cluster character.

The main result of this paper gives an explicit formula for the cluster character associated with a string module over a 2-Calabi-Yau tilted algebra. This can be stated as follows. To any walk c in a locally finite quiver Q , we associate a Laurent polynomial L_c in the ring of Laurent polynomials in the indeterminates x_i indexed by the set Q_0 of points of Q , which can be expressed as a product of 2×2 matrices as in [5] (see Section 1.3 below). Now, we let T be a tilting object in a Hom-finite triangulated 2-Calabi-Yau category \mathcal{C} and $B_T = \text{End}_{\mathcal{C}}(T)$ be the corresponding 2-Calabi-Yau tilted algebra whose ordinary quiver is denoted by Q . Moreover, to any string B_T -module M , we associate a tuple of integers $\mathbf{n}_M = (n_i)_{i \in Q_0}$ which we call the *normalisation* of M (see Section 5.2), and the Laurent polynomial L_M which is just the Laurent polynomial L_c attached to the string c of M in the quiver Q . Using the notation $\mathbf{x}^{\mathbf{n}_M} = \prod_{i \in Q_0} x_i^{n_i}$, our main result (Theorem 5.11 below) can be stated as saying that :

$$X_M^T = \frac{1}{\mathbf{x}^{\mathbf{n}_M}} L_M.$$

This result entails several interesting consequences. We first obtain the positivity of the Laurent polynomial X_M^T for any string B_T -module M (see Corollary 6.4), thus reproving a result of Cerulli and Haupt [18, 33]. Our results also apply to the cases of string modules over cluster-tilted algebras, string modules over gentle algebras arising from unpunctured surfaces (see [2]) and, more generally, string modules over finite dimensional Jacobian algebras associated with quivers with potentials in the sense of [22]. We also obtain a new proof of the positivity conjecture for cluster algebras arising from surfaces without punctures, see [50, 49, 42].

The paper is organised as follows. Section 1 introduces the basic definitions and presents our combinatorial formula. Section 2 introduces the concept of *realisable quadruples* which is the context in which our formula will actually compute cluster characters. This is closely related to the notion of triangulated 2-Calabi-Yau realisation in the sense of Fu and Keller [30]. Section 3 recalls all the necessary background from [44, 30] concerning cluster characters. Sections 4 and 5 contain the proof of our main result. In Section 4, we actually prove a weaker version of our theorem which will be used in order to prove the general case in Section 5. In Section 6 we present possible applications of the results to the study of positivity in

cluster algebras. In Section 7, we investigate the normalising factor and explicitly compute it for several cases of string modules over cluster-tilted algebras. The last section presents some detailed examples.

1. THE MATRIX FORMULA

1.1. Notations. Throughout the article, \mathbf{k} denotes an algebraically closed field. Given a quiver Q , we denote by Q_0 its set of points and by Q_1 its set of arrows. For any arrow $\alpha \in Q_1$, we denote by $s(\alpha)$ its *source* and by $t(\alpha)$ its *target*. We sometimes simply write $\alpha : s(\alpha) \rightarrow t(\alpha)$ or $s(\alpha) \xrightarrow{\alpha} t(\alpha)$. For any point $i \in Q_0$, we set

$$Q_1(i, -) = \{\alpha \in Q_1 \mid s(\alpha) = i\}, \quad Q_1(-, i) = \{\alpha \in Q_1 \mid t(\alpha) = i\}$$

and if F is a subset of points in Q_0 , we set

$$Q_1(i, F) = \{\alpha \in Q_1 \mid s(\alpha) = i, t(\alpha) \in F\}, \quad Q_1(F, i) = \{\alpha \in Q_1 \mid s(\alpha) \in F, t(\alpha) = i\}$$

and finally, for any $i, j \in Q_0$, we set $Q_1(i, j) = Q_1(i, -) \cap Q_1(-, j)$.

To any quiver Q , we associate a family $\mathbf{x}_Q = \{x_i \mid i \in Q_0\}$ of indeterminates over \mathbb{Z} . We let $\mathcal{L}(\mathbf{x}_Q) = \mathbb{Z}[x_i^{\pm 1} \mid i \in Q_0]$ be the ring of Laurent polynomials in the variables x_i , with $i \in Q_0$ and $\mathcal{F}(\mathbf{x}_Q) = \mathbb{Q}(x_i \mid i \in Q_0)$ be the field of rational functions in the variables x_i , with $i \in Q_0$. For any $\mathbf{d} = (d_i)_{i \in Q_0} \in \mathbb{Z}^{Q_0}$, we set $\mathbf{x}_Q^{\mathbf{d}} = \prod_{i \in Q_0} x_i^{d_i}$.

A *bound quiver* is a pair (Q, I) such that Q is a finite quiver (that is, Q_0 and Q_1 are finite sets) and I is an admissible ideal in the path algebra $\mathbf{k}Q$ of Q . Given a finite dimensional basic \mathbf{k} -algebra B , there exists a bound quiver (Q, I) such that $B \simeq \mathbf{k}Q/I$ and the quiver Q is called the *ordinary quiver* of B (see for instance [6]). We always identify the category $\text{mod-}B$ of finitely generated right B -modules with the category $\text{rep}(Q, I)$ of \mathbf{k} -representations of Q satisfying the relations in I . For a B -module M , we denote by $M(i)$ the \mathbf{k} -vector space at the point $i \in Q_0$ and $M(\alpha)$ the \mathbf{k} -linear map at the arrow $\alpha \in Q_1$.

1.2. Walks and strings. Let Q be a quiver. For any arrow $\beta \in Q_1$, we denote by β^{-1} a *formal inverse* for β , with $s(\beta^{-1}) = t(\beta)$, $t(\beta^{-1}) = s(\beta)$ and we set $(\beta^{-1})^{-1} = \beta$.

A *walk* of length $n \geq 1$ in Q is a sequence $c = c_1 \cdots c_n$ where each c_i is an arrow or a formal inverse of an arrow and such that $t(c_i) = s(c_{i+1})$ for any $i \in \{1, \dots, n-1\}$. The *source of the walk* c is $s(c) = s(c_1)$ and the *target of the walk* c is $t(c) = t(c_n)$. With any point $i \in Q_0$, we associate a walk e_i of length zero which is given by the stationary path at point i . For any walk c on Q , we denote by c_0 the walk of length zero $c_0 = e_{s(c)}$.

If (Q, I) is a bound quiver, a *string* in (Q, I) is either a walk of length zero or a walk $c = c_1 \cdots c_n$ of length $n \geq 1$ such that $c_i \neq c_{i+1}^{-1}$ for any $i \in \{1, \dots, n-1\}$ and such that no walk of the form $c_i c_{i+1} \cdots c_t$ nor its inverse belongs to I for $1 \leq i$ and $t \leq n$. If Q is a quiver, a *string in the quiver* Q is a string in the bound quiver $(Q, (0))$. If $B \simeq \mathbf{k}Q/I$ is a finite dimensional \mathbf{k} -algebra and c is a string in (Q, I) , then we also say that c is a *string in* B .

To any string in a finite dimensional \mathbf{k} -algebra B , we can naturally associate an indecomposable finite dimensional right B -module M_c , called *string module* as in [13, §3]. Namely, if B has ordinary quiver Q and c is a string, we define M_c as follows. If $c = e_v$ has length zero, then M_c is the simple representation S_v

at the point $v \in Q_0$. Otherwise, c is a string of length $n \geq 1$ and we write $c = v_1 \xrightarrow{c_1} \dots \xrightarrow{c_n} v_{n+1}$. For any $v \in Q_0$, we set $I_v = \{i \in [1, n+1] \mid v_i = v\}$ and we define $M_c(v)$ as the $|I_v|$ -dimensional \mathbf{k} -vector space with basis z_i , with $i \in I_v$. For any $1 \leq i \leq n$, if $c_i = \beta \in Q_1$, we set $M_c(\beta)(z_{i-1}) = z_i$ and if $c_i = \beta \in Q_1^{-1}$, we set $M_c(\beta)(z_i) = z_{i-1}$. Finally, if $\alpha : v \rightarrow v'$ is an arrow in Q_1 and if z_j is one of the basis vector of $M_c(v)$ such that $M_c(\alpha)(z_j)$ is not yet defined, we set $M_c(\alpha)(z_j) = 0$.

A string module is also called a *string representation* of the corresponding bound quiver. For any string module M , we denote by $\mathbf{s}(M)$ the corresponding string.

1.3. A formula for walks. For any locally finite quiver Q , we define a family of matrices with coefficients in $\mathbb{Z}[\mathbf{x}_Q] = \mathbb{Z}[x_i \mid i \in Q_0]$ as follows.

For any arrow $\beta \in Q_1$, we set

$$A(\beta) = \begin{bmatrix} x_{t(\beta)} & 0 \\ 1 & x_{s(\beta)} \end{bmatrix} \text{ and } A(\beta^{-1}) = \begin{bmatrix} x_{t(\beta)} & 1 \\ 0 & x_{s(\beta)} \end{bmatrix}.$$

Let $c = c_1 \cdots c_n$ be a walk of length $n \geq 1$ in Q . For any $i \in \{0, \dots, n\}$ we set

$$v_{i+1} = t(c_i)$$

(still with the notation $c_0 = e_{s(c)}$) and

$$V_c(i) = \begin{bmatrix} \prod_{\substack{\alpha \in Q_1(v_i, -) \\ \alpha \neq c_i^{\pm 1}, c_{i-1}^{\pm 1}}} x_{t(\alpha)} & 0 \\ 0 & \prod_{\substack{\alpha \in Q_1(-, v_i) \\ \alpha \neq c_i^{\pm 1}, c_{i-1}^{\pm 1}}} x_{s(\alpha)} \end{bmatrix}.$$

We then set

$$L_c = \frac{1}{x_{v_1} \cdots x_{v_{n+1}}} [1, 1] V_c(1) \left(\prod_{i=1}^n A(c_i) V_c(i+1) \right) \begin{bmatrix} 1 \\ 1 \end{bmatrix} \in \mathcal{L}(\mathbf{x}_Q).$$

If $c = e_i$ is a walk of length 0 at a point i , we similarly set

$$V_{e_i}(1) = \begin{bmatrix} \prod_{\alpha \in Q_1(i, -)} x_{t(\alpha)} & 0 \\ 0 & \prod_{\alpha \in Q_1(-, i)} x_{s(\alpha)} \end{bmatrix}.$$

and


$$L_{e_i} = \frac{1}{x_i} [1, 1] V_{e_i}(1) \begin{bmatrix} 1 \\ 1 \end{bmatrix} \in \mathcal{L}(\mathbf{x}_Q).$$

In other words, if c is any walk, either of length zero, or of the form $c = c_1 \cdots c_n$, we have

$$(1.1) \quad L_c = \frac{1}{\prod_{i=0}^n x_{t(c_i)}} [1, 1] \left(\prod_{i=0}^n A(c_i) V_c(i+1) \right) \begin{bmatrix} 1 \\ 1 \end{bmatrix} \in \mathcal{L}(\mathbf{x}_Q)$$

with the convention that $A(c_0)$ is the identity matrix.

Example 1.1. Consider the quiver



and consider the path $c = \delta^{-1}\beta\gamma$ in Q . Then,

$$\begin{aligned} L_c &= \frac{1}{x_2x_3^2x_4} \left([1, 1] \begin{bmatrix} x_5 & 0 \\ 0 & x_4 \end{bmatrix} \begin{bmatrix} x_3 & 1 \\ 0 & x_2 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & x_1 \end{bmatrix} \right. \\ &\quad \left. \begin{bmatrix} x_4 & 0 \\ 1 & x_2 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x_3 & 0 \\ 1 & x_4 \end{bmatrix} \begin{bmatrix} x_5 & 0 \\ 0 & x_2 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right) \\ &= \frac{x_1x_2^3x_4^2 + 2x_1x_2^2x_4x_5 + x_1x_2x_3x_4x_5 + x_3^2x_4x_5^2 + x_1x_2x_5^2 + x_1x_3x_5^2}{x_2x_3^2x_4} \end{aligned}$$

1.4. A formula for string modules.

Definition 1.2. Let B be a finite dimensional \mathbf{k} -algebra with bound quiver (Q, I) and let M be a string B -module with corresponding string $\mathbf{s}(M)$. We set

$$L_M = L_{\mathbf{s}(M)} \in \mathcal{L}(\mathbf{x}_Q).$$

If Q is a quiver, a *subquiver* R of Q is a quiver R such that $R_0 \subset Q_0$ and such that for any $i, j \in R_0$, the set of arrows from i to j in R_1 is a subset of the set of arrows from i to j in Q_1 . If R is a subquiver of Q , we naturally identify $\mathcal{L}(\mathbf{x}_R)$ with a subring of $\mathcal{L}(\mathbf{x}_Q)$.

A subquiver R of Q is called a *full subquiver* if for all $i, j \in R_0$ the set of arrows from i to j in R_1 equals the set of arrows from i to j in Q_1 . If B is a finite dimensional \mathbf{k} -algebra with bound quiver (Q, I) and M is a B -module, the *support* of M is the full subquiver $\text{supp}(M)$ of Q consisting of the points $i \in Q_0$ such that $M(i) \neq 0$. The *closure of the support of M* is the full subquiver $\overline{\text{supp}(M)}$ of Q formed by the points $i \in Q_0$ which are in the support of M or such that there exists an arrow α such that $s(\alpha) \in \text{supp}(M)_0$ and $t(\alpha) = i$ or $t(\alpha) \in \text{supp}(M)_0$ and $s(\alpha) = i$.

With these identifications, if B is a finite dimensional \mathbf{k} -algebra and M is a string B -module, then

$$L_M \in \mathcal{L}(\overline{\mathbf{x}_{\text{supp}(M)}}).$$

2. REALISABLE QUADRUPLES

In the previous section we associated with any string module M over a finite dimensional algebra B a certain Laurent polynomial L_M . In this section we provide a context in which the algebra B arises in connection with some cluster algebras so that we can compare the Laurent polynomials L_M with cluster variables or, more generally, with cluster characters. The context in which we work is the context of triangulated 2-Calabi-Yau realisations introduced in [30].

2.1. Definitions. An *ice quiver* is a pair (\mathcal{Q}, F) such that \mathcal{Q} is a finite connected quiver without loops and 2-cycles and F is a (possibly empty) subset of points of \mathcal{Q}_0 , called *frozen points*, such that there are no arrows between points in F . The *unfrozen part* of (\mathcal{Q}, F) is the full subquiver of \mathcal{Q} obtained by deleting the points in F .

For any quiver Q , we denote by $B(Q) = (b_{ij}) \in M_{Q_0}(\mathbb{Z})$ the skew-symmetric matrix defined for any $i, j \in Q_0$ by

$$b_{ij} = |Q_1(i, j)| - |Q_1(j, i)|.$$

If (\mathcal{Q}, F) is an ice quiver with unfrozen part Q , then we can fix an ordering of the points in \mathcal{Q}_0 such that

$$B(\mathcal{Q}) = \begin{bmatrix} B(Q) & -C^t \\ C & 0_{F \times F} \end{bmatrix}$$

where C is a matrix in $M_{F \times Q_0}(\mathbb{Z})$.

For any ice quiver (\mathcal{Q}, F) with unfrozen part Q , we denote by $\mathcal{A}(\mathcal{Q}, F)$ the cluster algebra of geometric type with initial seed $(\tilde{B}(\mathcal{Q}), \mathbf{x}, \mathbf{y})$ where

$$\tilde{B}(\mathcal{Q}) = \begin{bmatrix} B(Q) \\ C \end{bmatrix},$$

$\mathbf{x} = (x_i, i \in Q_0)$ and $\mathbf{y} = (x_i, i \in F)$ (see [28]).

A \mathbf{k} -linear category \mathcal{C} is called *Hom-finite* if $\text{Hom}_{\mathcal{C}}(M, N)$ is a finite dimensional \mathbf{k} -vector space for any two objects M, N in \mathcal{C} . A \mathbf{k} -linear triangulated category \mathcal{C} is called *2-Calabi-Yau* if there is a bifunctorial isomorphism

$$\text{Hom}_{\mathcal{C}}(X, Y) \simeq D\text{Hom}_{\mathcal{C}}(Y, X[2])$$

where $D = \text{Hom}_{\mathbf{k}}(-, \mathbf{k})$ is the standard duality and $[1]$ denotes the suspension functor.

For any objects M, N in \mathcal{C} , we denote by $\text{Ext}_{\mathcal{C}}^1(M, N)$ the space $\text{Hom}_{\mathcal{C}}(M, N[1])$. An object M in the category \mathcal{C} is called *rigid* if $\text{Ext}_{\mathcal{C}}^1(M, M) = 0$. An object T in \mathcal{C} is called a *tilting object* in \mathcal{C} if for any object X in \mathcal{C} , the equality $\text{Ext}_{\mathcal{C}}^1(T, X) = 0$ is equivalent to the fact that X belongs to the additive subcategory $\text{add } T$ of \mathcal{C} . Note that these objects are sometimes called *cluster-tilting* in the literature. It is known that the combinatorics of cluster algebras are closely related to the combinatorics of tilting objects in triangulated 2-Calabi-Yau categories [8].

Definition 2.1. A *realisable quadruple* is a quadruple $(\mathcal{Q}, F, \mathcal{C}, T)$ such that :

- (a) (\mathcal{Q}, F) is an ice quiver ;
- (b) \mathcal{C} is a Hom-finite triangulated 2-Calabi-Yau category whose tilting objects form a cluster structure in the sense of [8] ;
- (c) T is a tilting object in \mathcal{C} ;
- (d) the ordinary quiver of $B_T = \text{End}_{\mathcal{C}}(T)$ is \mathcal{Q} .

Following Reiten [48], every algebra of the form $\text{End}_{\mathcal{C}}(T)$ as above is called a *2-Calabi-Yau tilted algebra*. In order to simplify terminology, a category \mathcal{C} satisfying (b) is simply called a *triangulated 2-Calabi-Yau category*.

Definition 2.2. Given a realisable quadruple $(\mathcal{Q}, F, \mathcal{C}, T)$, a B_T -module M is called :

- a) *unfrozen* if $M(i) \neq 0$ implies $i \notin F$;
- b) *unfrozen indecomposable* if it is unfrozen and indecomposable as a B_T -module ;
- c) *unfrozen sincere* if $M(i) \neq 0$ if and only if $i \notin F$.

2.2. Examples. The notion of realisable quadruple covers a lot of situations in the context of cluster algebras. We now list some examples of such situations.

Example 2.3. Let Q be an acyclic quiver and let \mathcal{C} be the *cluster category* of Q , first defined in [10] (see [15] for an alternative description in Dynkin type \mathbb{A}). It is canonically equipped with a structure of triangulated 2-Calabi-Yau category [34] and for any tilting object T in \mathcal{C} , an algebra of the form $\text{End}_{\mathcal{C}}(T)$ is called a *cluster-tilted algebra*, as defined in [11] (see also [15]). Thus, if Q_T is the ordinary quiver of the cluster-tilted algebra $\text{End}_{\mathcal{C}}(T)$, the quadruple $(Q_T, \emptyset, \mathcal{C}, T)$ is realisable and every B_T -module is unfrozen.

Example 2.4. Let (Q, W) be a Jacobi-finite quiver with potential, that is, a quiver with potential in the sense of [22] such that the Jacobian algebra $\mathcal{J}_{(Q,W)}$ is finite dimensional. Let $\mathcal{C}_{(Q,W)}$ be the generalised cluster category constructed by Amiot [1, §3]. Then $\mathcal{C}_{(Q,W)}$ is a triangulated 2-Calabi-Yau category and there exists a tilting object T in $\mathcal{C}_{(Q,W)}$ such that $(Q, \emptyset, \mathcal{C}_{(Q,W)}, T)$ is a realisable quadruple [1, Theorem 3.6].

This example is of particular interest for the study of cluster algebras arising from surfaces in the sense of [26] (see also Section 6.2 for more details). Indeed, Labardini associated a non-degenerate Jacobi-finite quiver with potential (Q, W) to any marked surface (S, M) with non-empty boundary [39]. Thus, the generalised cluster category $\mathcal{C}_{(Q,W)}$ provides a categorification for the cluster algebra $\mathcal{A}(S, M)$ associated with the surface. Moreover, if the surface is *unpunctured*, that is, if there are no marked points in the interior of the surface, it is known that the Jacobian algebra $\mathcal{J}_{(Q,W)}$ is a string algebra (it is in fact a gentle algebra, see [2]). The $\mathcal{J}_{(Q,W)}$ -modules without self-extension are thus string modules and it follows from [30] that the cluster variables in $\mathcal{A}(S, M)$ can be studied via cluster characters associated with string $\mathcal{J}_{(Q,W)}$ -modules (see Section 3 for details).

Example 2.5. Let A be a finite dimensional \mathbf{k} -algebra of global dimension 2 and let \tilde{Q} be the ordinary quiver of the relation extension of A , that is, the trivial extension of A by the A -bimodule $\text{Ext}_A^2(DA, A)$ (see for instance [3]). Then Amiot also associated with A a generalised cluster category \mathcal{C}_A and proved that there exists a tilting object T in \mathcal{C}_A such that $(\tilde{Q}, \emptyset, \mathcal{C}_A, T)$ is a realisable quadruple [1, Theorem 4.10].

Example 2.6. We now give a fundamental example with a non-empty set of frozen points. For any quiver Q , we denote by (Q^{pp}, Q'_0) its *principal extension*. It is defined as follows. We fix a copy $Q'_0 = \{i' \mid i \in Q_0\}$ of Q_0 and we set $Q_0^{\text{pp}} = Q_0 \sqcup Q'_0$. The arrows in Q_1^{pp} between two points in Q_0 are the same as in Q and for any $i \in Q_0$ we add an extra arrow $i' \rightarrow i$ in Q_1^{pp} . Thus, the matrix of Q^{pp} is given by

$$B(Q^{\text{pp}}) = \begin{bmatrix} B(Q) & -I_{Q'_0} \\ I_{Q'_0} & 0 \end{bmatrix}$$

where $I_{Q'_0}$ denotes the identity matrix in $M_{Q'_0}(\mathbb{Z})$.

Now, if Q is the ordinary quiver of a cluster-tilted algebra B_T , it is known that there exists a 2-Calabi-Yau category \mathcal{C}^{pp} , endowed with a tilting object T^{pp} , obtained via a process of *principal gluing*, such that $(Q^{\text{pp}}, Q'_0, \mathcal{C}^{\text{pp}}, T^{\text{pp}})$ is a realisable quadruple for which every B_T -module can naturally be viewed as an unfrozen module (see [30, §6.3] or Corollary 2.12).

2.3. Blown-up ice quivers and their realisations. In the proof of the main theorem of this article, we are interested in a particular family of ice quivers, called *blown-up*. We now give some details concerning these quivers.

Definition 2.7. We say that an ice quiver (\mathcal{Q}, F) is *blown-up* if $|\mathcal{Q}_1(i, -) \cup \mathcal{Q}_1(-, i)| \leq 1$ for every point $i \in F$, that is, if there exists at most one arrow starting or ending at any frozen point.

Example 2.8. a) Any ice quiver with an empty set of frozen points is blown-up.
b) If Q is any quiver, then its principal extension (Q^{pp}, Q'_0) defined in Example 2.6 is blown-up.

Remark 2.9. Any blown-up ice quiver (\mathcal{Q}, F) whose unfrozen part is acyclic can be embedded in a realisable quadruple $(\mathcal{Q}, F, \mathcal{C}, T)$. Indeed, since (\mathcal{Q}, F) is blown-up with an acyclic unfrozen part, \mathcal{Q} is also acyclic. Let thus \mathcal{C} be the cluster category of the quiver \mathcal{Q} . Then the path algebra $\mathbf{k}\mathcal{Q}$ is identified with a tilting object in \mathcal{C} and the corresponding cluster-tilted algebra is isomorphic to $\mathbf{k}\mathcal{Q}$ so that its ordinary quiver is \mathcal{Q} . Thus $(\mathcal{Q}, F, \mathcal{C}, \mathbf{k}\mathcal{Q})$ is a realisable quadruple.

We can construct a wide class of examples of realisable quadruples with the following proposition, due to Amiot :

Proposition 2.10 ([1]). *If \mathcal{C}_1 and \mathcal{C}_2 are two generalised cluster categories associated with Jacobi-finite quivers with potentials (Q_1, W_1) and (Q_2, W_2) , then for any matrix C with non-negative integer entries, there exists a Jacobi-finite quiver with potential (Q', W') whose generalised cluster category $\mathcal{C}_{(Q', W')}$ has a tilting object T' such that the matrix associated with the ordinary quiver of the 2-Calabi-Yau tilted algebra $\text{End}_{\mathcal{C}_{(Q', W')}}(T')$ is*

$$B' = \begin{bmatrix} B(Q_1) & -C^t \\ C & B(Q_2) \end{bmatrix}.$$

Corollary 2.11. *Let (\mathcal{Q}, F) be a blown-up ice quiver with unfrozen part Q . Assume that there exists a potential W on Q such that the Jacobian algebra $\mathcal{J}_{(Q, W)}$ is finite dimensional. Then there exists a triangulated 2-Calabi-Yau category \mathcal{C} and a tilting object T in \mathcal{C} such that $(\mathcal{Q}, F, \mathcal{C}, T)$ is a realisable quadruple.*

Proof. We let $\mathcal{C}^{(0)}$ be the generalised cluster category associated with the quiver with potential (Q, W) . According to Example 2.4, there exists a tilting object $T^{(0)}$ in \mathcal{C} such that the quadruple $(Q, \emptyset, \mathcal{C}^{(0)}, T^{(0)})$ is realisable.

In order to simplify notations, we write $F = \{1, \dots, m\}$. For any $i \in F$, we denote by $\mathcal{C}_{\mathbb{A}_i}$ a copy of the cluster category of type \mathbb{A}_1 , which is a particular case of generalised cluster category in the sense of Amiot. We now construct, by induction on $i \in \{1, \dots, m\}$, a category $\mathcal{C}^{(i)}$ with a tilting object $T^{(i)}$ such that the ordinary quiver of the 2-Calabi-Yau tilted algebra $\text{End}_{\mathcal{C}^{(i)}}(T^{(i)})$ is the full subquiver (\mathcal{Q}, F) formed by points in Q and frozen points in $\{1, \dots, i\}$.

Since (\mathcal{Q}, F) is blown-up, for every $i \in \{1, \dots, m\}$, there exists a unique arrow $\alpha_i \in \mathcal{Q}_1(i, -) \sqcup \mathcal{Q}_1(-, i)$. If $\alpha_i \in \mathcal{Q}_1(i, -)$, we construct the category $\mathcal{C}^{(i)}$ gluing $\mathcal{C}^{(i-1)}$ and $\mathcal{C}_{\mathbb{A}_i}$ as provided by Proposition 2.10 with $\mathcal{C}_1 = \mathcal{C}^{(i-1)}$ and $\mathcal{C}_2 = \mathcal{C}_{\mathbb{A}_i}$ and we denote by $T^{(i)}$ the canonical tilting object in this gluing. If $\alpha_i \in \mathcal{Q}_1(-, i)$, we construct a gluing $\mathcal{C}^{(i)}$ of $\mathcal{C}_{\mathbb{A}_i}$ and $\mathcal{C}^{(i-1)}$ as provided by Proposition 2.10 with $\mathcal{C}_1 = \mathcal{C}_{\mathbb{A}_i}$ and $\mathcal{C}_2 = \mathcal{C}^{(i-1)}$ and we denote by $T^{(i)}$ the canonical tilting object in this gluing. We finally set $\tilde{\mathcal{C}} = \mathcal{C}^{(m)}$ and $\tilde{T} = T^{(m)}$. \square

Corollary 2.12. *Let (\mathcal{Q}, F) be a blown-up ice quiver whose unfrozen part is the ordinary quiver of a cluster-tilted algebra. Then (\mathcal{Q}, F) can be embedded in a realisable quadruple $(\mathcal{Q}, F, \mathcal{C}, T)$.*

Proof. It is proved in [35] and [9] that any cluster-tilted algebra is the Jacobian algebra of a Jacobi-finite quiver with potential. The result thus follows from Corollary 2.11. \square

3. CLUSTER CHARACTERS WITH COEFFICIENTS

In this section, we collect some background concerning Fu-Keller's cluster characters [30]. These characters allow one to give an explicit realisation of the elements in a cluster algebra in terms of the geometry and the homology underlying a 2-Calabi-Yau category.

We fix a triangulated 2-Calabi-Yau category \mathcal{C} with suspension functor [1] and we fix a tilting object T in \mathcal{C} .

We let \mathcal{Q} be the ordinary quiver of the 2-Calabi-Yau tilted algebra $B_T = \text{End}_{\mathcal{C}}(T)$. Indecomposable direct summands of T label the points in \mathcal{Q}_0 . Let F be a set of frozen points in \mathcal{Q}_0 . We set $T_F = \bigoplus_{i \in F} T_i$. Consider the full subcategory \mathcal{U} of \mathcal{C} formed by the objects X such that $\text{Hom}_{\mathcal{C}}(T_F, X) = 0$. Let $(\text{add } T[1])$ be the ideal consisting of those morphisms factoring through objects of $\text{add } T[1]$.

Theorem 3.1 ([11, 36]). *The functor $\text{Hom}_{\mathcal{C}}(T, -)$ induces an equivalence*

$$\text{Hom}_{\mathcal{C}}(T, -) : \mathcal{C}/(\text{add } T[1]) \xrightarrow{\sim} \text{mod-}B_T.$$

The equivalence $\text{Hom}_{\mathcal{C}}(T, -)$ induces an equivalence between $\mathcal{U}/(\text{add } T[1])$ and the subcategory of $\text{mod-}B_T$ consisting of B_T -modules supported on the unfrozen part of (\mathcal{Q}, F) , which is denoted by Q . Slightly abusing notations, an object M in $\mathcal{U}/(\text{add } T[1])$ is identified with its image $\text{Hom}_{\mathcal{C}}(T, M)$ in $\text{mod-}B_T$. Conversely, any B_T -module supported on Q is viewed as an object in $\mathcal{U}/(\text{add } T[1])$.

For any $i \in Q_0$, we denote by S_i the simple B_T -module corresponding to the point i . We denote by $\langle -, - \rangle$ the *truncated Euler form* on $\text{mod-}B_T$ defined by

$$\langle M, N \rangle = \dim \text{Hom}_{B_T}(M, N) - \dim \text{Ext}_{B_T}^1(M, N)$$

for any B_T -modules M and N . We denote by $\langle -, - \rangle_a$ the *anti-symmetrised Euler form* on $\text{mod-}B_T$ defined by

$$\langle M, N \rangle_a = \langle M, N \rangle - \langle N, M \rangle$$

for any B_T -modules M and N .

Lemma 3.2 ([44]). *For any $i \in \mathcal{Q}_0$, the form $M \mapsto \langle S_i, M \rangle$ on $\text{mod-}B_T$ only depends on the class $[M]$ of M in the Grothendieck group $K_0(\text{mod-}B_T)$ of $\text{mod-}B_T$.*

For any B_T -module M and any $\mathbf{e} \in K_0(\text{mod-}B_T)$, we let $\text{Gr}_{\mathbf{e}}(M)$ denote the set of submodules N of M whose class $[N]$ in $K_0(\text{mod-}B_T)$ equals \mathbf{e} . This set is called the *Grassmannian of submodules of M of dimension \mathbf{e}* . It is a projective variety and we denote by $\chi(\text{Gr}_{\mathbf{e}}(M))$ its Euler-Poincaré characteristic (with respect to the singular cohomology if \mathbf{k} is the field of complex numbers, and to the étale cohomology with compact support if \mathbf{k} is arbitrary).

Definition 3.3 ([44]). The *cluster character* associated with (\mathcal{C}, T) is the unique map

$$X_{\mathcal{C}}^T : \text{Ob}(\mathcal{C}) \longrightarrow \mathcal{L}(\mathbf{x}_{\mathcal{Q}})$$

such that

- a) $X_{T_i[1]}^T = x_i$ for any $i \in \mathcal{Q}_0$;
- b) If M is indecomposable and not isomorphic to any $T_i[1]$, then

$$X_M^T = \sum_{\mathbf{e} \in \mathbb{N}^{\mathcal{Q}_0}} \chi(\mathrm{Gr}_{\mathbf{e}}(\mathrm{Hom}_{\mathcal{C}}(T, M))) \prod_{i \in \mathcal{Q}_0} x_i^{\langle S_i, \mathbf{e} \rangle_a - \langle S_i, \mathrm{Hom}_{\mathcal{C}}(T, M) \rangle} ;$$

- c) For any two objects M, N in \mathcal{C} ,

$$X_{M \oplus N}^T = X_M^T X_N^T.$$

We recall that an indecomposable object X in \mathcal{C} is called *reachable from T* if it is a direct summand of a tilting object which can be obtained from T by a finite number of mutations, see [8]. In particular, any reachable object is rigid.

Theorem 3.4 ([30]). *The map X_T^T induces a surjection from the set of indecomposable objects in \mathcal{C} which are reachable from T to the set of cluster variables in the cluster algebra $\mathcal{A}(\mathcal{Q}, F)$.*

Note that identifying $\mathcal{C}/(\mathrm{add} T[1])$ with $\mathrm{mod}\text{-}B_T$ using the functor $\mathrm{Hom}_{\mathcal{C}}(T, -)$, we associate to any indecomposable B_T -module M the cluster character of an indecomposable lifting \overline{M} of M in $\mathcal{C}/(\mathrm{add} T[1])$. Thus, we set

$$X_M^T = X_{\overline{M}}^T = \sum_{\mathbf{e} \in \mathbb{N}^{\mathcal{Q}_0}} \chi(\mathrm{Gr}_{\mathbf{e}}(M)) \prod_{i \in \mathcal{Q}_0} x_i^{\langle S_i, \mathbf{e} \rangle_a - \langle S_i, M \rangle}$$

and this way, we may view X_T^T as a map on the set of objects in $\mathrm{mod}\text{-}B_T$.

4. A FORMULA FOR DYNKIN TYPE \mathbb{A} WITH COEFFICIENTS

We now start the proof of our main result, which will be stated in Theorem 5.11. This section is devoted to the first step of the proof in which we establish the theorem for specific modules in the particular case of blown-up quivers with unfrozen part of Dynkin type \mathbb{A} . The result we prove in this section is the following :

Theorem 4.1. *Let (\mathcal{Q}, F) be a blown-up ice quiver with unfrozen part Q of Dynkin type \mathbb{A} and let $(\mathcal{Q}, F, \mathcal{C}, T)$ be a realisable quadruple. Let $B_T = \mathrm{End}_{\mathcal{C}}(T)$. Then for any indecomposable B_T -module M , the following hold :*

- a) *If M is unfrozen, then it is a string module ;*
- b) *If M is a submodule of the unique unfrozen sincere module, then the corresponding cluster character is given by*

$$X_M^T = L_M.$$

Every B_T -module M supported on Q has the structure of a module over the path algebra $H = \mathbf{k}Q$ of the quiver Q . Since Q is of Dynkin type \mathbb{A} every such module M is a string module if it is indecomposable. This proves the first point of the theorem.

To prove the second point, we need to collect the necessary background concerning the matrix product formula from [5], which we do in Section 4.1. Section 4.2 is devoted to the proof when (\mathcal{Q}, F) is the principal extension of a Dynkin quiver of type \mathbb{A} . In Section 4.3, we use the Fomin-Zelevinsky separation formula in order to deduce the general case.

observe that in $\text{mod-}\mathbf{k}Q$, this part of the grid contains exactly all the indecomposable submodules of the unique indecomposable sincere $\mathbf{k}Q$ -module, which is exactly positioned at the intersection of the vertical and the horizontal lines of asterisks (see for instance [32]). This observation will be crucial in Lemma 4.2 below and in Section 5.

We give more details on this region. For any cell (u, v) located in the north-west component of the grid (and not on its south-east frontier of asterisks), its horizontal and vertical projections onto the initial variables x_i determine a word $x_k x_{k+1} \dots x_{k+l+1}$. Following [5], for each $j = 1, 2, \dots, l-1$, let

$$M(x_j, x_{j+1}) = \begin{cases} \begin{bmatrix} x_j & 1 \\ 0 & x_{j+1} \end{bmatrix} & \text{if } x_j \text{ is to the left of } x_{j+1}, \\ \begin{bmatrix} x_{j+1} & 0 \\ 1 & x_j \end{bmatrix} & \text{if } x_j \text{ is below } x_{j+1}. \end{cases}$$

Then

$$t_{Q^{\text{op}}}(u, v) = \frac{1}{x_{k+1} \dots x_{k+l}} [1, x_k] \left(\prod_{j=1}^{l-1} M(x_{k+j}, x_{k+j+1}) \right) \begin{bmatrix} 1 \\ x_{k+l+1} \end{bmatrix}.$$

To determine the corresponding indecomposable $\mathbf{k}Q$ -module $M_{(u,v)}$ for which $t_{Q^{\text{op}}}(u, v) = X_{M_{(u,v)}}$, the horizontal and vertical projections onto the cluster variables X_{P_i} gives the word $X_{P_{k+1}} X_{P_{k+2}} \dots X_{P_{k+l}}$, meaning that $M_{(u,v)}$ is the string module corresponding to the unique string from the point $k+1$ to the point $k+l$ in Q . In our example, the cell (u, v) gives rise to the word $x_2 x_3 \dots x_7$, leading to

$$t_{Q^{\text{op}}}(u, v) = \frac{1}{x_3 x_4 x_5 x_6} [1, x_2] \begin{bmatrix} x_3 & 1 \\ 0 & x_4 \end{bmatrix} \begin{bmatrix} x_5 & 0 \\ 1 & x_4 \end{bmatrix} \begin{bmatrix} x_5 & 1 \\ 0 & x_6 \end{bmatrix} \begin{bmatrix} 1 \\ x_7 \end{bmatrix}$$

and $M_{(u,v)}$ corresponds to the string module whose corresponding string is $\gamma^{-1} \delta \epsilon^{-1}$ in Q .

Now, a close inspection of the formulae (in the coefficient-free situation) gives $t_{Q^{\text{op}}}(u, v) = L_{M_{(u,v)}}$ whenever (u, v) lies in the north-west region.

As mentioned above, we also need to consider the south-east frontier of asterisks of the north-west region. Observe that on this frontier, $L_{M_{(u,v)}}$ is defined, while $t_{Q^{\text{op}}}(u, v)$ is not. To fix this, one can augment $\overline{Q^{\text{op}}}$ with two new sinks, labeled 0 and $n+1$, in order to obtain a Dynkin quiver $\overline{Q^{\text{op}}}$ of type \mathbb{A}_{n+2} in such a way the cells which were on the frontier of asterisks now lie in the north-west region in the grid corresponding to $\overline{Q^{\text{op}}}$; thus $t_{\overline{Q^{\text{op}}}}(u, v)$ can be defined. In our example, it suffices to let $\overline{Q^{\text{op}}}$ be given by

$$0 \longleftarrow 1 \xleftarrow{\alpha} 2 \xleftarrow{\beta} 3 \xrightarrow{\gamma} 4 \xleftarrow{\delta} 5 \xrightarrow{\epsilon} 6 \xrightarrow{\zeta} 7 \xrightarrow{\eta} 8 \xleftarrow{\theta} 9 \xleftarrow{\iota} 10 \xrightarrow{\kappa} 11 \longrightarrow 12.$$

It is then easily checked that $L_{M_{(u,v)}} = t_{\overline{Q^{\text{op}}}}(u, v)|_{\substack{x_0=1 \\ x_{n+1}=1}}$, that is, $L_{M_{(u,v)}}$ is obtained from $t_{\overline{Q^{\text{op}}}}(u, v)$ by specialising the initial cluster variables x_0 and x_{n+1} to 1. Observe that this relation also holds true for any cell located in the north-west region. So, in general, one can write

$$L_{M_{(u,v)}} = t_{\overline{Q^{\text{op}}}}(u, v)|_{\substack{x_0=1 \\ x_{n+1}=1}}$$

whenever (u, v) lies in the north-west region or on its south-east frontier of asterisks.

In the situation where we deal with arbitrary coefficients, let (\mathcal{Q}, F) be an ice quiver with unfrozen part Q of Dynkin type A . Then, generalising the coefficient-free situation, we let

$$t_{Q^{\text{op}}}(u, v) = L_{M_{(u, v)}}.$$

For each $i \in Q_0$, let

$$y_i = \prod_{\alpha \in \mathcal{Q}_1(F, i)} x_{s(\alpha)} \quad \text{and} \quad z_i = \prod_{\alpha \in \mathcal{Q}_1(i, F)} x_{t(\alpha)}.$$

and for any $\mathbf{d} = (d_i)_{i \in Q_0} \in \mathbb{N}^{Q_0}$, we set $\mathbf{y}^{\mathbf{d}} = \prod_{i \in Q_0} y_i^{d_i}$ and $\mathbf{z}^{\mathbf{d}} = \prod_{i \in Q_0} z_i^{d_i}$.

Then, a tedious and combinatorial adaptation of Lemmata 5, 6 and Theorem 4 in [5], in which one needs to embed Q^{op} in $\overline{Q^{\text{op}}}$ as above, allows us to obtain the following recurrence relations, whose verification is left to the reader.

Lemma 4.2. *a) For any $i \in Q_0$, we have*

$$(4.2) \quad x_i L_{P_i} - y_i \left(\prod_{\alpha \in Q_1(i, -)} L_{P_{t(\alpha)}} \right) \left(\prod_{\alpha \in Q_1(-, i)} x_{s(\alpha)} \right) = \mathbf{z}^{\dim P_i}.$$

b) For any non-projective indecomposable submodule M of the unique indecomposable unfrozen sincere $\mathbf{k}Q$ -module, we have

$$(4.3) \quad L_{\tau M} L_M - L_E = \mathbf{y}^{\dim \tau M} \mathbf{z}^{\dim M}$$

where E is the middle term of the almost split exact sequence $0 \rightarrow \tau M \xrightarrow{i} E \xrightarrow{p} M \rightarrow 0$ in $\text{mod-}\mathbf{k}Q$.

4.2. Proof of Theorem 4.1 - first step : Principal coefficients. We start with two lemmata concerning cluster characters associated with principal extensions of acyclic quivers. These are analogues to [14, Lemma 3.9 and Proposition 3.10] (see also [25, Lemma 2.2]) which provide representation-theoretical interpretations of certain exchange relations in a cluster algebra with principal coefficients at an acyclic seed.

Let Q be an acyclic quiver and let $H = \mathbf{k}Q$ be the path algebra of Q , which is finite-dimensional and hereditary. The map sending an H -module to its dimension vector allows us to identify the Grothendieck group $K_0(\text{mod-}H)$ with \mathbb{Z}^{Q_0} . If (\mathcal{Q}, F) is the principal extension (Q^{pp}, Q'_0) of Q defined in Example 2.6, we denote by X_j^{pp} the corresponding cluster character and by L^{pp} the matrix formula of equation (1.1). For every $i' \in Q'_0$, we set $y_i = x_{i'}$. It is known that

$$X_M^{\text{pp}} = \sum_{\mathbf{e} \in \mathbb{N}^{Q_0}} \chi(\text{Gr}_{\mathbf{e}}(M)) \prod_{i \in Q_0} x_i^{-\langle \mathbf{e}, [S_i] \rangle - \langle [S_i], [M] - \mathbf{e} \rangle} y_i^{m_i - e_i}$$

where the Euler forms and the Grassmannian are considered in $\text{mod-}H$ and where $[M] = (m_i)_{i \in Q_0}$ (see for instance [25, Remark 2.4]).

Lemma 4.3. *Let Q be an acyclic quiver. Then for any $i \in Q_0$, we have*

$$(4.4) \quad x_i X_{P_i}^{\text{pp}} - y_i \left(\prod_{\alpha \in Q_1(i, -)} X_{P_{t(\alpha)}}^{\text{pp}} \right) \left(\prod_{\alpha \in Q_1(-, i)} x_{s(\alpha)} \right) = 1.$$

where P_j is the indecomposable projective $\mathbf{k}Q$ -module associated with the point $j \in Q_0$.

Proof. The proof is a straightforward adaptation of the proof of [14, Lemma 3.9], we give it for completeness. We recall that for any $i \in Q_0$, we have

$$\text{Rad } P_i = \bigoplus_{\alpha \in Q_1(i, -)} P_{t(\alpha)} \text{ and thus } X_{\text{Rad } P_i}^{\text{PP}} = \prod_{\alpha \in Q_1(i, -)} X_{P_{t(\alpha)}}^{\text{PP}}.$$

We also recall that $P_i/\text{Rad } P_i \simeq S_i$ and that a submodule M of P_i either equals P_i or is a submodule of $\text{Rad } P_i$. We set $\mathbf{dim } P_i = \mathbf{m} = (m_j)_{j \in Q_0}$ and let δ be such that $\delta_{ij} = 1$ if $i = j$ and $\delta_{ij} = 0$ otherwise. Thus

$$\begin{aligned} X_{\text{Rad } P_i}^{\text{PP}} &= \sum_{\mathbf{e}} \chi(\text{Gr}_{\mathbf{e}}(\text{Rad } P_i)) \prod_{l \in Q_0} x_l^{-\langle \mathbf{e}, [S_l] \rangle - \langle [S_l], \mathbf{m} - [S_l] - \mathbf{e} \rangle} y_l^{m_l - \delta_{il} - e_l} \\ &= \sum_{\mathbf{e}} \chi(\text{Gr}_{\mathbf{e}}(\text{Rad } P_i)) \prod_{l \in Q_0} \left(x_l^{-\langle \mathbf{e}, [S_l] \rangle - \langle [S_l], \mathbf{m} - \mathbf{e} \rangle} y_l^{m_l - e_l} \right) x_l^{\langle [S_l], [S_l] \rangle} y_l^{-\delta_{il}} \\ &= y_i^{-1} \left(\prod_{\alpha \in Q_1(-, i)} x_{s(\alpha)}^{-1} \right) x_i \sum_{\mathbf{e}} \chi(\text{Gr}_{\mathbf{e}}(\text{Rad } P_i)) \prod_{l \in Q_0} x_l^{-\langle \mathbf{e}, [S_l] \rangle - \langle [S_l], \mathbf{m} - \mathbf{e} \rangle} y_l^{m_l - e_l} \end{aligned}$$

but

$$X_{P_i}^{\text{PP}} = x_i^{-1} + \sum_{\mathbf{e}} \chi(\text{Gr}_{\mathbf{e}}(\text{Rad } P_i)) \prod_{l \in Q_0} x_l^{-\langle \mathbf{e}, [S_l] \rangle - \langle [S_l], \mathbf{m} - \mathbf{e} \rangle} y_l^{m_l - e_l}.$$

Thus,

$$x_i X_{P_i}^{\text{PP}} = y_i \left(\prod_{\alpha \in Q_1(-, i)} x_{s(\alpha)} \right) X_{\text{Rad } P_i}^{\text{PP}} + 1$$

from which we deduce (4.4). \square

Lemma 4.4. *Let Q be an acyclic quiver. Then for any non-projective $\mathbf{k}Q$ -module M , we have*

$$(4.5) \quad X_{\tau M}^{\text{PP}} X_M^{\text{PP}} - X_E^{\text{PP}} = \mathbf{y}^{\mathbf{dim } \tau M}$$

where E is the central term of the almost split exact sequence $0 \rightarrow \tau M \xrightarrow{i} E \xrightarrow{p} M \rightarrow 0$.

Proof. This result is an analogue of [14, Proposition 3.10]. We sketch the proof for the convenience of the reader.

We write $N = \tau M$, $\mathbf{dim } N = \mathbf{n} = (n_l)_{l \in Q_0}$ and $\mathbf{dim } M = \mathbf{m} = (m_l)_{l \in Q_0}$. It follows from the definition of the character that we have

$$X_N^{\text{PP}} X_M^{\text{PP}} = X_{N \oplus M}^{\text{PP}} = \sum_{\mathbf{e} \in \mathbb{N}^{Q_0}} \chi(\text{Gr}_{\mathbf{e}}(N \oplus M)) \prod_{l \in Q_0} x_l^{-\langle \mathbf{e}, [S_l] \rangle - \langle [S_l], \mathbf{n} + \mathbf{m} - \mathbf{e} \rangle} y_l^{m_l + n_l - e_l}.$$

but there is a surjective map with affine fibres

$$\text{Gr}_{\mathbf{e}}(N \oplus M) \longrightarrow \bigsqcup_{\mathbf{f} + \mathbf{g} = \mathbf{e}} \text{Gr}_{\mathbf{f}}(N) \times \text{Gr}_{\mathbf{g}}(M)$$

so that $\chi(\text{Gr}_{\mathbf{e}}(N \oplus M)) = \sum_{\mathbf{f} + \mathbf{g} = \mathbf{e}} \chi(\text{Gr}_{\mathbf{f}}(N)) \chi(\text{Gr}_{\mathbf{g}}(M))$ and thus

$$X_{N \oplus M}^{\text{PP}} = \sum_{\mathbf{f}, \mathbf{g}} \chi(\text{Gr}_{\mathbf{f}}(N)) \chi(\text{Gr}_{\mathbf{g}}(M)) \prod_{l \in Q_0} x_l^{-\langle \mathbf{f} + \mathbf{g}, [S_l] \rangle - \langle [S_l], \mathbf{n} + \mathbf{m} - \mathbf{f} - \mathbf{g} \rangle} y_l^{m_l + n_l - f_l - g_l}.$$

Now, it follows from [14, Lemma 3.11] that every fibre of the map

$$\zeta : \begin{cases} \mathrm{Gr}_{\mathbf{e}}(N \oplus M) & \longrightarrow & \bigsqcup_{\mathbf{f}+\mathbf{g}=\mathbf{e}} \mathrm{Gr}_{\mathbf{f}}(N) \times \mathrm{Gr}_{\mathbf{g}}(M) \\ U & \longmapsto & (i^{-1}(U), p(U)) \end{cases}$$

is an affine space except over the point $(0, M)$ where it is empty. It thus follows that

$$X_{N \oplus M}^{\mathrm{PP}} = X_E^{\mathrm{PP}} + \prod_{l \in Q_0} y_l^{n_l}$$

which establishes (4.5). \square

We now prove the second point of Theorem 4.1 for principal coefficients :

Proposition 4.5. *Let Q be a Dynkin quiver of type \mathbb{A} . Then for any indecomposable submodule M of the unique unfrozen sincere $\mathbf{k}Q$ -module, we have*

$$X_M^{\mathrm{PP}} = L_M^{\mathrm{PP}}.$$

Proof. The proof directly follows from Lemmata 4.2, 4.3 and 4.4, keeping in mind that for principal coefficients we have $z_i = 1$ for each i in Lemma 4.2. \square

4.3. Proof of Theorem 4.1 - second step. We now finish the proof of Theorem 4.1. For this, the strategy is to apply the Fomin-Zelevinsky separation formula to the equality established in Proposition 4.5. We let (\mathcal{Q}, F) be a blown-up ice quiver with unfrozen part Q of Dynkin type \mathbb{A} and we fix a realisable quadruple $(\mathcal{Q}, F, \mathcal{C}, T)$ and an unfrozen indecomposable B_T -module M which has a natural structure of H -module where H is the path algebra of Q . We denote by (Q^{PP}, Q'_0) the principal extension of Q . Every H -module can naturally be viewed as a B_T -module or as a $\mathbf{k}Q^{\mathrm{PP}}$ -module. According to Proposition 4.5, we know that $L_M^{\mathrm{PP}} = X_M^{\mathrm{PP}}$. We now want to prove that $L_M = X_M^T$ where L_M denotes the matrix formula associated with (\mathcal{Q}, F) in equation (1.1).

In order to simplify the notations, we identify Q_0 with $\{1, \dots, n\}$ and for every $i \in Q_0$, we denote i' by $n + i$. Let \mathbb{P} be the tropical semifield generated by the x_i , with $i \in F$, and endowed with the auxiliary addition

$$\prod_i x_i^{a_i} \oplus \prod_i x_i^{b_i} = \prod_i x_i^{\min\{a_i, b_i\}}.$$

For any $i \in Q_0$, we set

$$w_i = \left(\prod_{\alpha \in \mathcal{Q}_1(F, i)} x_{s(\alpha)} \right) \left(\prod_{\alpha \in \mathcal{Q}_1(i, F)} x_{t(\alpha)}^{-1} \right).$$

With the notations of Section 4.1, we can thus write $w_i = y_i z_i^{-1}$. Following [28], for every subtraction-free rational expression f in the variables x_1, \dots, x_{2n} , we define the *separation of f* as

$$\sigma(f) = \frac{f(x_1, \dots, x_n, w_1, \dots, w_n)}{f|_{\mathbb{P}}(1, \dots, 1, w_1, \dots, w_n)}$$

where the $f|_{\mathbb{P}}$ means that we have replaced the ordinary addition in $\mathcal{F}(\mathbf{x}_F)$ by the auxiliary addition \oplus of the semifield \mathbb{P} . This can be done since f is subtraction-free.

Now, we note that, for every H -module, L_M^{PP} is a subtraction-free rational expression by definition. Also, for every indecomposable H -module M , X_M^{PP} is a

cluster variable so that it is defined with a finite number of mutations, which are all subtraction-free. Thus, X_M^{pp} is also a subtraction-free rational expression (see [28, §3]) and we can apply σ to both X_M^{pp} and L_M^{pp} . Since $L_M^{\text{pp}} = X_M^{\text{pp}}$ by Proposition 4.5, we have $\sigma(L_M^{\text{pp}}) = \sigma(X_M^{\text{pp}})$. Thus, we only need to prove that $\sigma(X_M^{\text{pp}}) = X_M^T$ and $\sigma(L_M^{\text{pp}}) = L_M$.

The equality $\sigma(X_M^{\text{pp}}) = X_M^T$ follows directly from [28, Theorem 3.7] since X_7^{pp} and X_7^T induce bijections from the set of indecomposable H -modules to the set of cluster variables in the cluster algebras $\mathcal{A}(Q^{\text{pp}}, Q_0^{\text{pp}})$ and $\mathcal{A}(\mathcal{Q}, F)$ respectively and these bijections respect denominator vectors.

The equality $\sigma(L_M^{\text{pp}}) = L_M$ is obtained from the following observation. We write $c = \mathbf{s}(M)$. For each $j \in F$, x_j appears in exactly one w_i , since there is exactly one $i \in Q_0$ which is adjacent to j in \mathcal{Q} . On the other hand, in the product of matrices L_M^{pp} , x_{n+i} appears in at most one matrix, namely in the matrix $V_c^{\text{pp}}(i)$ (where $V_c^{\text{pp}}(i)$ denotes the matrices arising in the matrix product L_c^{pp}). Thus, using the definition of the addition in \mathbb{P} and the fact that $L_M^{\text{pp}}(1, \dots, 1, x_{n+1}, \dots, x_{2n})$ has constant term 1, it follows that

$$L_M^{\text{pp}}|_{\mathbb{P}}(1, \dots, 1, w_1, \dots, w_n) = \prod_{i \in (\text{supp}(M))_0} \prod_{\alpha \in Q_1^{\text{pp}}(i, F)} x_{t(\alpha)}^{-1}.$$

Now, since

$$\left(\prod_{\alpha \in \mathcal{Q}_1(i, F)} x_{t(\alpha)} \right) (V_c^{\text{pp}}(i)(x_1, \dots, x_n, w_1, \dots, w_n)) = V_c(i),$$

we get $\sigma(L_M^{\text{pp}}) = L_M$. This ends the proof of Theorem 4.1. \square

5. THE GENERAL CASE

We now deduce the main theorem from Theorem 4.1. For this, we use some “blow-up” techniques which are described in Section 5.1. These techniques introduce some “error” in the computation of the characters but this can be controlled with a normalising factor that we introduce in Section 5.2. The main result (Theorem 5.11) is stated in Section 5.4.

5.1. Blowing up quivers along string modules. In this section, we fix a quiver Q , we denote by M a string representation of Q and we write $c = \mathbf{s}(M)$. If c is of positive length $n \geq 1$, we write $c = c_1 \cdots c_n$.

The *blow-up* \widetilde{Q}_M of Q along M is the quiver constructed as follows. Let $\{v_1, \dots, v_{n+1}\}$ be a set. For any $i \in \{1, \dots, n\}$, we set β_i from v_i to v_{i+1} which is an arrow (or a formal inverse of an arrow, respectively) in $(\widetilde{Q}_M)_1$ if c_i is an arrow (or a formal inverse of an arrow, respectively) in Q_1 . For any $i \in \{1, \dots, n\}$ and for any arrow $\alpha \in Q_1$ such that $\alpha \neq c_i^{\pm 1}, c_{i-1}^{\pm 1}$, if $s(\alpha) = t(c_i)$ (or $t(\alpha) = s(c_i)$, respectively), we create a new point, denoted by $t(\alpha)^{\alpha; i}$ (or $s(\alpha)^{\alpha; i}$, respectively) and an arrow $\alpha_{v_i} : v_i \rightarrow t(\alpha)^{\alpha; i}$ (or $\alpha_{v_i} : s(\alpha)^{\alpha; i} \rightarrow v_i$, respectively).

Example 5.1. Consider the quiver

$$Q : \quad 1 \xrightarrow{\alpha} 2 \begin{array}{c} \xrightarrow{\epsilon} \\ \xrightarrow{\gamma} \end{array} 3 \xrightarrow{\delta} 4$$

and let M be the string representation corresponding to the walk $c = \epsilon^{-1}\gamma$, that is

$$M : \quad 0 \longrightarrow \mathbf{k} \xrightarrow{\begin{matrix} [1,0]^t \\ \cong \\ [0,1]^t \end{matrix}} \mathbf{k}^2 \longrightarrow 0.$$

Then, the quiver \widetilde{Q}_M is

$$\begin{array}{ccccccc} \widetilde{Q}_M & & 2\gamma;1 & \xrightarrow{\gamma_{v_1}} & v_1 & \xleftarrow{\beta_1} & v_2 & \xrightarrow{\beta_2} & v_3 & \xleftarrow{\epsilon_{v_3}} & 2\epsilon;3 \\ & & & & \downarrow \delta_{v_1} & & \uparrow \alpha_{v_1} & & \downarrow \delta_{v_3} & & \\ & & & & 4\delta;1 & & 1\alpha;2 & & 4\delta;3 & & \end{array}$$

We recall the following definition from [33] :

Definition 5.2. Let Q and S be two quivers. A *winding of quivers* $\Phi : Q \rightarrow S$ is a pair $\Phi = (\Phi_0, \Phi_1)$ where $\Phi_0 : Q_0 \rightarrow S_0$ and $\Phi_1 : Q_1 \rightarrow S_1$ are such that :

- a) Φ is a morphism of quivers, that is $s \circ \Phi_1 = \Phi_0 \circ s$ and $t \circ \Phi_1 = \Phi_0 \circ t$;
- b) If $\alpha, \alpha' \in Q_1$ with $\alpha \neq \alpha'$ and $s(\alpha) = s(\alpha')$, then $\Phi_1(\alpha) \neq \Phi_1(\alpha')$;
- c) If $\alpha, \alpha' \in Q_1$ with $\alpha \neq \alpha'$ and $t(\alpha) = t(\alpha')$, then $\Phi_1(\alpha) \neq \Phi_1(\alpha')$;

With the above notations, the maps

$$\Phi_0 : \begin{cases} v_i & \mapsto t(c_{i-1}) & \text{for any } i \in \{1, \dots, n+1\} \\ v^{\alpha; i} & \mapsto v & \text{for any } v \in \overline{\text{supp}(M)} \text{ and any } \alpha, i \end{cases}$$

and

$$\Phi_1 : \begin{cases} \beta_i & \mapsto c_i & \text{for any } i \in \{1, \dots, n\} \\ \alpha_{v_i} & \mapsto \alpha & \text{for any arrow of the form } \alpha_{v_i} \end{cases}$$

induce a winding of quivers $\Phi : \widetilde{Q}_M \rightarrow \overline{\text{supp}(M)}$.

Let Φ_* be the map from the set of objects in $\text{rep}(\widetilde{Q}_M)$ to the set of objects in $\text{rep}(\overline{\text{supp}(M)})$ which associates to a representation \widetilde{V} of \widetilde{Q}_M the representation V given by

$$V(i) = \bigoplus_{j \in \Phi_0^{-1}(i)} \widetilde{V}(j) \text{ and } V(\alpha) = \bigoplus_{\beta \in \Phi_1^{-1}(\alpha)} \widetilde{V}(\beta)$$

for any $i \in (\overline{\text{supp}(M)})_0$ and any $\alpha \in (\overline{\text{supp}(M)})_1$. At the level of dimension vectors, Φ_* induces a natural map $\phi : \mathbb{N}^{(\widetilde{Q}_M)_0} \rightarrow \mathbb{N}^{(\overline{\text{supp}(M)})_0}$.

We define a representation \widetilde{M} of \widetilde{Q}_M by setting for any point $v \in (\widetilde{Q}_M)_0$

$$\widetilde{M}(v) = \begin{cases} \mathbf{k} & \text{if } v = v_i \text{ for some } i \in \{0, \dots, n\}, \\ 0 & \text{otherwise} \end{cases}$$

and for any arrow $\alpha \in (\widetilde{Q}_M)_1$

$$\widetilde{M}(\alpha) = \begin{cases} 1_{\mathbf{k}} & \text{if } \alpha = \beta_i^{\pm 1} \text{ for some } i \in \{1, \dots, n\}, \\ 0 & \text{otherwise.} \end{cases}$$

Lemma 5.3. *Let B be a finite dimensional \mathbf{k} -algebra. Then for any string B -module M , we have $\Phi_*(\widetilde{M}) \simeq M$.*

Proof. This follows from the construction. \square

Definition 5.4. If B is a finite dimensional \mathbf{k} -algebra with bound quiver (Q, I) and M is a B -module, the *border* ∂M of M is the set of points in the closure $\text{supp}(M)$ of the support of M in Q and which do not lie in the support $\text{supp}(M)$ of M .

Thus, with the above notations, the border $\partial \widetilde{M}$ consists of all the points in $(\widetilde{Q}_M)_0$ which do not lie in the support of \widetilde{M} .

Lemma 5.5. *Let B be a finite dimensional \mathbf{k} -algebra. Then for any string B -module M , the pair $(\widetilde{Q}_M, \partial \widetilde{M})$ is a blown-up ice quiver whose unfrozen part is of Dynkin type \mathbb{A} . Moreover, the representation \widetilde{M} is a sincere unfrozen string representation of \widetilde{Q}_M .*

Proof. The first assertion follows from the construction of \widetilde{Q}_M and \widetilde{M} . For the second assertion, we observe that \widetilde{M} is supported on the unfrozen part of \widetilde{Q}_M which is of Dynkin type \mathbb{A} . It is unfrozen sincere and indecomposable by construction so that it is a string representation. \square

Example 5.6. Let B be the path algebra of the quiver Q considered in Example 5.1 and let M be the string module considered in that same example. Then, the representation \widetilde{M} of the quiver \widetilde{Q}_M is

$$\widetilde{M} : \quad \begin{array}{ccccccc} 0 & \longrightarrow & \mathbf{k} & \xleftarrow{1_{\mathbf{k}}} & \mathbf{k} & \xrightarrow{1_{\mathbf{k}}} & \mathbf{k} \longleftarrow 0 \\ & & \downarrow & & \uparrow & & \downarrow \\ & & 0 & & 0 & & 0 \end{array}$$

so that $(\widetilde{Q}_M, \partial \widetilde{M})$ is indeed a blown-up ice quiver with unfrozen part of Dynkin type \mathbb{A}_3 .

5.2. Normalisation. In this section M denotes a string module over a finite dimensional algebra B with bound quiver (Q, I) .

Definition 5.7. The *normalising vector* of M is $\mathbf{n}_M = (n_i)_{i \in \overline{\text{supp}(M)}_0} \in \overline{\mathbb{N}^{\text{supp}(M)}_0}$ given by

$$n_i = \langle S_i, M \rangle - \sum_{j \in \Phi_0^{-1}(i)} \langle S_j, \widetilde{M} \rangle$$


for any $i \in \overline{\text{supp}(M)}_0$ where the first truncated Euler form is considered in $\text{mod-}B$ and the second truncated Euler form is considered in $\text{mod-}\mathbf{k}\widetilde{Q}_M$.

The *normalising factor* of M is

$$\mathbf{x}^{\mathbf{n}_M} = \prod_{i \in \overline{\text{supp}(M)}_0} x_i^{n_i}.$$

This normalisation is actually easy to compute in several usual situations as it is explained in Section 7.

Example 5.8. Consider the finite dimensional algebra B whose ordinary quiver is



and whose relations are given by the vanishing of all paths of length two, that is $\alpha\beta = \beta\gamma = \gamma\alpha = 0$. It is a cluster-tilted algebra of type \mathbb{A}_3 . Consider the projective module M associated with the point 1. It is a string B -module with string α . The associated blown-up quiver \widetilde{Q}_M is $3^{\gamma;1} \rightarrow v_1 \rightarrow v_2 \rightarrow 3^{\beta;2}$ and the representation \widetilde{M} is $0 \rightarrow \mathbf{k} \xrightarrow{1_{\mathbf{k}}} \mathbf{k} \rightarrow 0$. Then, one has $\langle S_1, M \rangle = \langle S_{v_1}, \widetilde{M} \rangle = 0$, $\langle S_2, M \rangle = \langle S_{v_2}, \widetilde{M} \rangle = 1$ and $\langle S_3, M \rangle = 0$ whereas $\langle S_{3^{\beta;2}}, \widetilde{M} \rangle + \langle S_{3^{\gamma;1}}, \widetilde{M} \rangle = 0 - 1 = -1$. Thus, the normalisation is $\mathbf{n}_M = (0, 0, 1)$.

5.3. Blow-ups and cluster characters. We keep the notations of Section 5.1. As usual, we naturally identify $\mathcal{L}(\overline{\mathbf{x}_{\text{supp}(M)}})$ to a subring of $\mathcal{L}(\mathbf{x}_{\mathcal{Q}})$. We consider the following surjective morphism of \mathbb{Z} -algebras :

$$\pi : \mathcal{L}(\mathbf{x}_{\widetilde{Q}_M}) \rightarrow \mathcal{L}(\overline{\mathbf{x}_{\text{supp}(M)}})$$

defined by

$$\pi(x_{v_i}) = x_{t(c_{i-1})}$$

for any $1 \leq i \leq n+1$ and

$$\pi(x_{v_{\alpha;i}}) = x_v$$

for any $v \in \overline{\text{supp}(M)}$ and any α, i .

We now observe that for any unfrozen string module M with respect to a realisable quadruple $(\mathcal{Q}, F, \mathcal{C}, T)$, the Laurent polynomial X_M^T is in the image of the function π .

Lemma 5.9. *Let $(\mathcal{Q}, F, \mathcal{C}, T)$ be a realisable quadruple and let M be an unfrozen string module. Then*

$$X_M^T \in \mathcal{L}(\overline{\mathbf{x}_{\text{supp}(M)}}).$$

Proof. We recall that unfrozen B_T -modules are identified via $\text{Hom}_{\mathcal{C}}(T, -)$ with the objects M in $\mathcal{U}/(\text{add } T[1])$. We have

$$X_M^T = \sum_{\mathbf{e} \in \mathbb{N}^{\mathcal{Q}_0}} \chi(\text{Gr}_{\mathbf{e}}(M)) \prod_{i \in \mathcal{Q}_0} x_i^{\langle S_i, \mathbf{e} \rangle_a - \langle S_i, M \rangle}.$$

Any dimension vector \mathbf{e} such that $\text{Gr}_{\mathbf{e}}(M) \neq \emptyset$ is supported on $\text{supp}(M)$. Thus, if $i \in \mathcal{Q}_0$ is not in $\overline{\text{supp}(M)}$, then $\langle S_i, \mathbf{e} \rangle_a = 0$ and $\langle S_i, M \rangle = 0$. In particular,

$$X_M^T = \sum_{\mathbf{e} \in \mathbb{N}^{\mathcal{Q}_0}} \chi(\text{Gr}_{\mathbf{e}}(M)) \prod_{i \in \overline{\text{supp}(M)}} x_i^{\langle S_i, \mathbf{e} \rangle_a - \langle S_i, M \rangle} \in \mathcal{L}(\overline{\mathbf{x}_{\text{supp}(M)}}).$$

□

Let $\widetilde{\mathcal{C}}$ be the cluster category of the quiver \widetilde{Q}_M and let $\widetilde{T} = \mathbf{k}\widetilde{Q}_M$ be the path algebra of \widetilde{Q}_M , which is identified with a tilting object in $\widetilde{\mathcal{C}}$. We denote by $\widetilde{X}_?$ the corresponding cluster character with values in $\mathcal{L}(\mathbf{x}_{\widetilde{Q}_M})$.

Proposition 5.10. *Let $(\mathcal{Q}, F, \mathcal{C}, T)$ be a realisable quadruple and let M be an unfrozen string module. Then,*

$$\pi(\widetilde{X}_{\widetilde{M}}) = \mathbf{x}^{\mathbf{n}_M} X_M^T.$$

Proof. We have

$$\begin{aligned} X_M^T &= \sum_{\mathbf{e} \in \mathbb{N}^{\mathcal{Q}_0}} \chi(\mathrm{Gr}_{\mathbf{e}}(M)) \prod_{i \in \mathcal{Q}_0} x_i^{\langle S_i, \mathbf{e} \rangle_a - \langle S_i, M \rangle} \\ &= \sum_{\mathbf{e} \in \mathbb{N}^{\mathrm{supp}(M)_0}} \chi(\mathrm{Gr}_{\mathbf{e}}(M)) \prod_{i \in \overline{\mathrm{supp}(M)}_0} x_i^{\langle S_i, \mathbf{e} \rangle_a - \langle S_i, M \rangle} \end{aligned}$$

Now, for any $\mathbf{e} \in \mathbb{N}^{\mathrm{supp}(M)_0}$, since Φ is a winding of quivers and $\Phi_*(\widetilde{M}) = M$, it follows from [33, Theorem 1.2 (a)] that

$$\chi(\mathrm{Gr}_{\mathbf{e}}(M)) = \sum_{\mathbf{f} \in \phi^{-1}(\mathbf{e})} \chi(\mathrm{Gr}_{\mathbf{f}}(\widetilde{M})).$$

Fix $\mathbf{e} \in \mathbb{N}^{\mathrm{supp}(M)_0}$ and $\mathbf{f} \in \phi^{-1}(\mathbf{e})$. We now prove that for any $i \in \overline{\mathrm{supp}(M)}$, we have

$$(5.6) \quad x_i^{\langle S_i, \mathbf{e} \rangle_a} = \pi \left(\prod_{j \in \Phi_0^{-1}(i)} x_j^{\langle S_j, \mathbf{f} \rangle_a} \right).$$

where the anti-symmetrised Euler form in the left-hand side is taken in $\mathrm{mod}\text{-}\mathbf{k}\widetilde{Q}_M$ and the anti-symmetrised Euler forms in the right-hand side are taken in $\mathrm{mod}\text{-}B$.

First note that

$$\pi \left(\prod_{j \in \Phi_0^{-1}(i)} x_j^{\langle S_j, \mathbf{f} \rangle_a} \right) = x_i^{\sum_{j \in \Phi_0^{-1}(i)} \langle S_j, \mathbf{f} \rangle_a}$$

so that it is enough to prove that

$$\langle S_i, \mathbf{e} \rangle_a = \sum_{j \in \Phi_0^{-1}(i)} \langle S_j, \mathbf{f} \rangle_a.$$

For any module U and any integer $n \geq 1$, we denote by nU the direct sum of n copies of U . Now, since $\langle S_i, - \rangle_a$ is well-defined on the Grothendieck group $K_0(\mathrm{mod}\text{-}B_T)$ (Lemma 3.2), we have

$$\begin{aligned} \langle S_i, \mathbf{e} \rangle_a &= \sum_{k \in \mathrm{supp}(M)_0} \langle S_i, e_k S_k \rangle_a \\ &= \sum_{k \in \mathrm{supp}(M)_0} \left\langle S_i, \left(\sum_{l \in \Phi_0^{-1}(k)} f_l \right) S_k \right\rangle_a \\ &= \sum_{k \in \mathrm{supp}(M)_0} \left(\sum_{l \in \Phi_0^{-1}(k)} f_l \right) \langle S_i, S_k \rangle_a \end{aligned}$$

and

$$\begin{aligned} \sum_{j \in \Phi_0^{-1}(i)} \langle S_j, \mathbf{f} \rangle_a &= \sum_{k \in \text{supp}(M)_0} \sum_{l \in \Phi_0^{-1}(k)} \sum_{j \in \Phi_0^{-1}(i)} f_l \langle S_j, S_l \rangle_a \\ &= \sum_{k \in \text{supp}(M)_0} \sum_{l \in \Phi_0^{-1}(k)} f_l \left(\sum_{j \in \Phi_0^{-1}(i)} \langle S_j, S_l \rangle_a \right). \end{aligned}$$

But for any $k \in \text{supp}(M)_0$ and any $l \in \Phi_0^{-1}(k)$, we have

$$\sum_{j \in \Phi_0^{-1}(i)} \langle S_j, S_l \rangle_a = \langle S_i, S_k \rangle_a$$

which proves (5.6).

Now,

$$\tilde{X}_{\tilde{M}} = \sum_{\mathbf{f} \in \mathbb{N}^{(\tilde{Q}_M)_0}} \chi(\text{Gr}_{\mathbf{f}}(\tilde{M})) \prod_{i \in (\tilde{Q}_M)_0} x_i^{\langle S_i, \mathbf{f} \rangle_a - \langle S_i, \tilde{M} \rangle}$$

so that,

$$\begin{aligned} \pi(\tilde{X}_{\tilde{M}}) &= \sum_{\mathbf{e} \in \mathbb{N}^{\text{supp}(M)_0}} \sum_{\mathbf{f} \in \phi^{-1}(\mathbf{e})} \chi(\text{Gr}_{\mathbf{f}}(\tilde{M})) \prod_{i \in \text{supp}(M)_0} \pi \left(\prod_{j \in \Phi_0^{-1}(i)} x_j^{\langle S_j, \mathbf{f} \rangle_a - \langle S_j, \tilde{M} \rangle} \right) \\ &= \sum_{\mathbf{e} \in \mathbb{N}^{\text{supp}(M)_0}} \left(\sum_{\mathbf{f} \in \phi^{-1}(\mathbf{e})} \chi(\text{Gr}_{\mathbf{f}}(\tilde{M})) \right) \prod_{i \in \text{supp}(M)_0} x_i^{\langle S_i, \mathbf{e} \rangle_a} \pi \left(\prod_{j \in \Phi_0^{-1}(i)} x_j^{-\langle S_j, \tilde{M} \rangle} \right) \\ &= \sum_{\mathbf{e} \in \mathbb{N}^{\text{supp}(M)_0}} \chi(\text{Gr}_{\mathbf{e}}(M)) \prod_{i \in \text{supp}(M)_0} x_i^{\langle S_i, \mathbf{e} \rangle_a - \sum_{j \in \Phi_0^{-1}(i)} \langle S_j, \tilde{M} \rangle} \\ &= \mathbf{x}^{\mathbf{n}_M} X_M^T. \end{aligned}$$

This finishes the proof. \square

5.4. The main theorem. We can now prove the main theorem of the article :

Theorem 5.11. *Let $(\mathcal{Q}, F, \mathcal{C}, T)$ be a realisable quadruple and let M be an unfrozen string module with respect to this quadruple. Then*

$$X_M^T = \frac{1}{\mathbf{x}^{\mathbf{n}_M}} L_M.$$

Proof. We first notice that with the above notations, it follows directly from the definitions that

$$\pi(\tilde{L}_{\tilde{M}}) = L_M.$$

Thus,

$$X_M^T = \frac{1}{\mathbf{x}^{\mathbf{n}_M}} \pi(\tilde{X}_{\tilde{M}}) = \frac{1}{\mathbf{x}^{\mathbf{n}_M}} \pi(\tilde{L}_{\tilde{M}}) = \frac{1}{\mathbf{x}^{\mathbf{n}_M}} L_M$$

where the first equality follows from Proposition 5.10, the second equality from Theorem 4.1 and Lemma 5.5 and the third from the above observation. \square

6. APPLICATIONS

In this section, we give several applications of Theorem 5.11.

6.1. Computing cluster variables. We now prove that the formula given in equation (1.1) allows one to compute the cluster variables in several situations.

Corollary 6.1. *Let (\mathcal{Q}, F) be an ice quiver. Assume that there exists a realisable quadruple $(\mathcal{Q}, F, \mathcal{C}, T)$ such that every rigid object in \mathcal{C} is reachable from T . Then for any unfrozen $\text{End}_{\mathcal{C}}(T)$ -module M which is rigid and indecomposable, the Laurent polynomial $\frac{1}{\mathbf{x}^{\mathbf{n}_M}} L_M$ is a cluster variable in $\mathcal{A}(\mathcal{Q}, F)$.*

Proof. We identify M with an indecomposable lifting in \mathcal{C} for the functor $\text{Hom}_{\mathcal{C}}(T, -)$, which is also rigid in \mathcal{C} by [52, 31]. It follows from Theorem 3.4 that the map $M \mapsto X_M^T$ induces a surjection from the set of reachable rigid objects in to the set of cluster variables in $\mathcal{A}(\mathcal{Q}, F)$. It is also rigid in \mathcal{C} , hence, by hypothesis, it is reachable and thus X_M^T is a cluster variable in $\mathcal{A}(\mathcal{Q}, F)$. \square

Corollary 6.2. *Let B_T be a cluster-tilted algebra. Then for any rigid string B_T -module M , the Laurent polynomial $\frac{1}{\mathbf{x}^{\mathbf{n}_M}} L_M$ is a cluster variable in the coefficient-free cluster algebra associated with the ordinary quiver of B_T .*

Proof. Let \mathcal{C} be a cluster category, T be a tilting object in \mathcal{C} and Q_T be the ordinary quiver of B_T . Then the quadruple $(Q_T, \emptyset, \mathcal{C}, T)$ is realisable (see Example 2.3). Moreover, it follows from [10] that every indecomposable rigid object in \mathcal{C} is reachable from T . The result thus follows from Corollary 6.1. \square

Corollary 6.3. *Let B_T be a cluster-tilted algebra of Dynkin type \mathbb{A} or euclidean type $\tilde{\mathbb{A}}$. Then, the map*

$$M \mapsto \frac{1}{\mathbf{x}^{\mathbf{n}_M}} L_M$$

induces a bijection from the set of indecomposable rigid B_T -modules to the set of cluster variables in the coefficient-free cluster algebra associated with the ordinary quiver of B_T which do not belong to the initial cluster.

Proof. Cluster-tilted algebras of Dynkin type \mathbb{A} or euclidean type $\tilde{\mathbb{A}}$ are string algebras [2]. Thus, each indecomposable rigid B_T -module is a string module. Now if \mathcal{C} is a cluster category and T is a tilting object in \mathcal{C} such that $B_T = \text{End}_{\mathcal{C}}(T)$, it follows in particular from Theorem 5.11 that for every indecomposable rigid B_T -module M , we have $X_M^T = \frac{1}{\mathbf{x}^{\mathbf{n}_M}} L_M$. But then it follows from [44] that $M \mapsto \frac{1}{\mathbf{x}^{\mathbf{n}_M}} L_M$ is a bijection from the set of indecomposable rigid B_T -modules to the set of cluster variables in the coefficient-free cluster algebra $\mathcal{A}(Q_T, \emptyset)$ which do not belong to the initial cluster \mathbf{x}_{Q_T} where Q_T denotes the ordinary quiver of B_T . \square

6.2. Positivity in cluster algebras. We now prove positivity results in cluster algebras using formula (1.1). We first recall the basic notions concerning positivity in cluster algebras.

If \mathbb{P} is an abelian group, we denote by $\mathbb{Z}\mathbb{P}$ its group ring. Let $n \geq 1$ be an integer and $\mathbf{x} = (x_1, \dots, x_n)$ be an n -tuple of indeterminates. We denote by $\mathbb{Z}_{\geq 0}\mathbb{P}[\mathbf{x}^{\pm 1}]$ the semiring of subtraction-free Laurent polynomials in the variables in \mathbf{x} with coefficients in $\mathbb{Z}\mathbb{P}$, that is, the set of elements of the form

$$\frac{f(x_1, \dots, x_n)}{x_1^{d_1} \cdots x_n^{d_n}}$$

where f is a polynomial in n variables whose coefficients are non-negative linear combinations of elements of \mathbb{P} and $d_1, \dots, d_n \in \mathbb{Z}$. In particular, in the case when

$\mathbb{P} = \{1\}$, the notation $\mathbb{Z}_{\geq 0}[\mathbf{x}^{\pm 1}]$ denotes the semiring of Laurent polynomials in variables x_1, \dots, x_n with non-negative coefficients.

If Q is a quiver without loops and 2-cycles, if \mathbb{P} is any semifield and if \mathbf{y} is a Q_0 -tuple of elements of \mathbb{P} , it follows from the Laurent phenomenon that every cluster variable in the cluster algebra $\mathcal{A}(Q, \mathbf{x}_Q, \mathbf{y})$ is an element of $\mathbb{Z}\mathbb{P}[\mathbf{c}^{\pm 1}]$ when expressed in any cluster \mathbf{c} of $\mathcal{A}(Q, \mathbf{x}_Q, \mathbf{y})$ [27]. The Fomin and Zelevinsky *positivity conjecture*, stated in [27], asserts that every cluster variable is actually an element of $\mathbb{Z}_{\geq 0}\mathbb{P}[\mathbf{c}^{\pm 1}]$. This conjecture was proved for rank two cluster algebras [51, 41, 23], cluster algebras arising from surfaces with or without punctures [50, 49, 42], cluster algebras with bipartite seeds [43] and weaker versions were proved for acyclic cluster algebras [5, 47, 43].

Corollary 6.4. *Let $(\mathcal{Q}, F, \mathcal{C}, T)$ be a realisable quadruple and let M be an unfrozen string module with respect to this quadruple. Then*

$$X_M^T \in \mathbb{Z}_{\geq 0}[\mathbf{x}_{\text{supp}(M)}^{\pm 1}].$$

Proof. The set $\mathbb{Z}_{\geq 0}[\mathbf{x}_{\text{supp}(M)}]$ is a semiring for the usual sum and product. Thus, the set $M_2(\mathbb{Z}_{\geq 0}[\mathbf{x}_{\text{supp}(M)}])$ is also a semiring for the usual sum and product of matrices. It follows that $L_M \in \mathbb{Z}_{\geq 0}[\mathbf{x}_{\text{supp}(M)}^{\pm 1}]$ and by Theorem 5.11, $X_M^T = L_M$. This proves the corollary. \square

This corollary was also obtained by Cerulli and Haupt [33, 18].

Corollary 6.5. *Let B_T be a cluster-tilted algebra. Then $X_M^T \in \mathbb{Z}_{\geq 0}[\mathbf{x}_{\text{supp}(M)}^{\pm 1}]$ for any string B_T -module M .* \square

We now provide a new proof of the Fomin and Zelevinsky positivity conjecture for cluster algebras arising from unpunctured surfaces. We first set some notations. If (S, M) is an oriented bordered marked surface, Fomin, Shapiro and Thurston associated with any triangulation Γ of (S, M) a quiver Q_Γ [26]. For any semifield \mathbb{P} and any $(Q_\Gamma)_0$ -tuple \mathbf{y} of elements of \mathbb{P} , we denote by $\mathcal{A}(Q_\Gamma, \mathbf{x}_{Q_\Gamma}, \mathbf{y})$ the cluster algebra with initial seed $(Q_\Gamma, \mathbf{x}_{Q_\Gamma}, \mathbf{y})$ (see [28]). Such a cluster algebra does not depend on the choice of the triangulation Γ of (S, M) , up to an isomorphism of $\mathbb{Z}\mathbb{P}$ -algebras [26]. It is called the *cluster algebra arising from the surface (S, M)* and is denoted by $\mathcal{A}(S, M)$.

For this class of cluster algebras, Fomin and Zelevinsky positivity conjecture amounts to saying that any cluster variable z in $\mathcal{A}(S, M)$ belongs to $\mathbb{Z}_{\geq 0}\mathbb{P}[\mathbf{x}_{Q_\Gamma}^{\pm 1}]$ for any choice of triangulation Γ [27, §3]. This conjecture was proved in [50, 49] for unpunctured surfaces and in [42] for arbitrary surfaces. We now provide a new independent representation-theoretical proof of the positivity conjecture for cluster algebras arising from unpunctured surfaces.

Corollary 6.6. *Let (S, M) be an unpunctured surface. Then the positivity conjecture holds for $\mathcal{A}(S, M)$ equipped with an arbitrary choice of coefficients.*

Proof. Using the Fomin-Zelevinsky separation formula [28, Theorem 3.7], it is enough to prove the result for principal coefficients. Let Γ be any triangulation of (S, M) and let (Q_Γ, W_Γ) be the quiver with potential associated with this triangulation as in [2] (see also [39]). Let $\mathcal{J}_\Gamma = \mathcal{J}_{(Q_\Gamma, W_\Gamma)}$ be the corresponding Jacobian algebra which is finite dimensional [2]. It follows from Corollary 2.11 that the principal extension $(Q_\Gamma^{\text{pp}}, (Q_\Gamma)_0')$ can be embedded in a realisable quadruple

$(Q_\Gamma^{\text{pp}}, (Q_\Gamma)_0', \mathcal{C}, T)$. Now, let z be a cluster variable in $\mathcal{A}(S, M)$, it follows from Theorem 3.4 that either z belongs to the initial cluster \mathbf{x}_{Q_Γ} or $z = X_M^T$ for some indecomposable rigid \mathcal{J}_Γ -module M . Since the algebra \mathcal{J}_Γ is a string algebra, any indecomposable rigid \mathcal{J}_Γ -module is a string module. Thus, if we set $y_i = x_{i'}$ for any $i \in (Q_T)_0$, it follows from Corollary 6.4 that

$$z = X_M^T \in \mathbb{Z}_{\geq 0}[\mathbf{x}_{\text{supp}(M)}^{\pm 1}] \subset \mathbb{Z}_{\geq 0}[\mathbf{y}^{\pm 1}][\mathbf{x}_{Q_\Gamma}^{\pm 1}] = \mathbb{Z}_{\geq 0}\mathbb{P}[\mathbf{x}_{Q_\Gamma}^{\pm 1}].$$

□

6.3. $SL(2, \mathbb{Z})$ and total Grassmannians of submodules.

Definition 6.7. Given a finite dimensional \mathbf{k} -algebra B and a finitely generated B -module M , the *total Grassmannian of submodules of M* is the set

$$\text{Gr}(M) = \bigsqcup_{\mathbf{e} \in K_0(\text{mod-}B)} \text{Gr}_{\mathbf{e}}(M).$$

Since M is finite dimensional, all but finitely many $\text{Gr}_{\mathbf{e}}(M)$ are empty so that $\text{Gr}(M)$ is a finite disjoint union of projective varieties. It is thus endowed with a structure of projective variety and we can consider its Euler-Poincaré characteristic $\chi(\text{Gr}(M)) = \sum_{\mathbf{e}} \chi(\text{Gr}_{\mathbf{e}}(M)) \in \mathbb{Z}$.

For modules over hereditary algebras, these numbers can be computed with friezes, as shown in [4, §4]. Here we prove that, for string modules over 2-Calabi-Yau tilted algebras, these numbers can be computed with products of matrices in $SL(2, \mathbb{Z})$.

If Q is any quiver, for any $\beta \in Q_1$, we define two matrices in $SL(2, \mathbb{Z})$:

$$a(\beta) = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} \text{ and } a(\beta^{-1}) = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$$

and for any walk $c = c_1 \cdots c_n$ in Q , we set

$$l_c = [1, 1] \left(\prod_{i=1}^n a(c_i) \right) \begin{bmatrix} 1 \\ 1 \end{bmatrix} \in \mathbb{N}.$$

Finally, if c is a walk of length 0, we set $l_c = 2$.

Proposition 6.8. *For any string module M over a finite dimensional algebra, we have*

$$\chi(\text{Gr}(M)) = l_{\mathbf{s}(M)}.$$

Proof. Let B be a finite-dimensional algebra with ordinary quiver Q and let M be a string B -module. We use the notations of Section 5.1. Let \widetilde{M} be the representation of the blow-up \widetilde{Q}_M of Q along M . Then \widetilde{M} is a string representation of \widetilde{Q}_M and we have $l_{\mathbf{s}(M)} = l_{\mathbf{s}(\widetilde{M})}$. Moreover, it follows from [33, Theorem 1.2 (a)] that

$$\sum_{\mathbf{e} \in \mathbb{N}^{Q_0}} \chi(\text{Gr}_{\mathbf{e}}(M)) = \sum_{\mathbf{e} \in \mathbb{N}^{Q_0}} \sum_{\mathbf{f} \in \phi^{-1}(\mathbf{e})} \chi(\text{Gr}_{\mathbf{f}}(\widetilde{M})) = \sum_{\mathbf{f} \in \mathbb{N}^{(\widetilde{Q}_M)_0}} \chi(\text{Gr}_{\mathbf{f}}(\widetilde{M}))$$

so that the characteristic of the total Grassmannian of M equals the characteristic of the total Grassmannian of \widetilde{M} . Since Grassmannians of submodules of \widetilde{M} only depends on the support of \widetilde{M} , it is enough to prove the proposition for indecomposable modules over path algebras of Dynkin quivers of type \mathbb{A} .

Let thus B be the path algebra of a Dynkin quiver Q of type \mathbb{A} , let \mathcal{C} be the cluster category of Q and let $T = \mathbf{k}Q$. Then $(Q, \emptyset, \mathcal{C}, T)$ is a realisable quadruple such that $B = \text{End}_{\mathcal{C}}(T)$. According to Theorem 5.11, we have

$$X_M^T = \frac{1}{\mathbf{x}_Q^{\mathbf{n}_M}} L_M.$$

Consider the surjective morphism of \mathbb{Z} -algebras.

$$p: \begin{cases} \mathcal{L}(\mathbf{x}_Q) & \longrightarrow & \mathbb{Z} \\ x_i & \mapsto & 1 \end{cases} \quad \text{for any } i \in Q_0.$$

Then it follows from the definition of the cluster character that $p(X_M^T) = \chi(\text{Gr}(M))$.

We now prove that $p(L_M) = l_{\mathbf{s}(M)}$. If the string $\mathbf{s}(M)$ of M is of length zero, M is simple and thus

$$\chi(\text{Gr}(M)) = \chi(\text{Gr}_{[0]}(M)) + \chi(\text{Gr}_{[M]}(M)) = 1 + 1 = 2 = l_{\mathbf{s}(M)}.$$

Otherwise, we write $\mathbf{s}(M) = c_1 \cdots c_n$ with $n \geq 1$. Then, for any $i \in \{1, \dots, n\}$, we have $p(A(c_i)) = a(c_i)$ and $p(V_c(s(c_i))) = p(V_c(t(c_i)))$ is the identity matrix. Since p is a ring homomorphism, it induces a ring homomorphism at the level of matrices and thus $p(L_M) = p(L_c) = l_c$. Since $p(\frac{1}{\mathbf{x}_Q^{\mathbf{n}_M}}) = 1$, it follows that

$$\chi(\text{Gr}(M)) = p(X_M^T) = p(L_M) = l_{\mathbf{s}(M)}$$

and the proposition is proved. \square

Example 6.9. Let (S, M) be an unpunctured surface. Let Γ be a triangulation of (S, M) , let (Q_Γ, W_Γ) be the quiver with potential associated with this triangulation and let $\mathcal{J}_\Gamma = \mathcal{J}_{(Q_\Gamma, W_\Gamma)}$ be the corresponding Jacobian algebra. Then for any string module M over \mathcal{J}_Γ , we have $\chi(\text{Gr}(M)) = l_{\mathbf{s}(M)}$.

7. MORE ABOUT THE NORMALISATION

We now describe some situations in which the normalisation can be omitted or computed combinatorially.

7.1. The hereditary case. We first observe that the normalisation can be omitted in several cases.

Lemma 7.1. *Let $(\mathcal{Q}, F, \mathcal{C}, T)$ be a realisable quadruple and let M be an unfrozen string module. If the full subcategory of $\text{mod-}B_T$ formed by modules which are supported on $\overline{\text{supp}(M)}$ is hereditary, then $\mathbf{n}_M = 0$.*

Proof. Let $i \in \overline{\text{supp}(M)}_0$, since M and S_i are supported on $\overline{\text{supp}(M)}$, the truncated Euler form $\langle S_i, M \rangle$ only depends on the dimension vectors. But since \widetilde{Q}_M is acyclic, $\sum_{j \in \Phi_0^{-1}(i)} \langle S_j, \widetilde{M} \rangle$ also only depends on the dimension vectors, it thus follows that $\langle S_i, M \rangle = \sum_{j \in \Phi_0^{-1}(i)} \langle S_j, \widetilde{M} \rangle$ and thus $\mathbf{n}_M = 0$. \square

Corollary 7.2. *Let $(\mathcal{Q}, F, \mathcal{C}, T)$ be a realisable quadruple such that B_T is hereditary. Then $\mathbf{n}_M = 0$ for every unfrozen string module M .* \square

7.2. The maximal factor method for modules over cluster-tilted algebras.

For any walk c in a locally finite quiver Q we set

$$N_c = [1, 1] \left(\prod_{i=0}^n A(c_i) V_c(i+1) \right) \begin{bmatrix} 1 \\ 1 \end{bmatrix} \in \mathcal{L}(\mathbf{x}_Q)$$

so that (1.1) becomes

$$L_c = \frac{1}{\prod_{i=0}^n x_{t(c_i)}} N_c.$$

Let $\eta_c \in \mathbb{N}^{Q_0}$ be such that

$$N_c = \mathbf{x}_Q^{\eta_c} P_c(\mathbf{x}_Q)$$

with $P_c(\mathbf{x}_Q)$ not divisible by any x_i with $i \in Q_0$.

Let \mathcal{C} be a cluster category, $T = T_1 \oplus \dots \oplus T_n$ be a tilting object in \mathcal{C} , then $B_T = \text{End}_{\mathcal{C}}(T)$ is a cluster-tilted algebra and we denote by Q its ordinary quiver. Then $(Q, \emptyset, \mathcal{C}, T)$ is a realisable quadruple and every B_T -module is unfrozen.

We now prove that for such modules, we can compute the normalisation combinatorially :

Proposition 7.3. *Consider the above notations and assume moreover that $\text{End}_{\mathcal{C}}(T_i) \simeq \mathbf{k}$ for any $i \in \{1, \dots, n\}$. Then for every rigid string B_T -module M we have $\mathbf{n}_M = \eta_{\mathbf{s}(M)}$.*

Proof. For any Laurent polynomial $L(x_1, \dots, x_n) \in \mathbb{Z}[x_1^{\pm 1}, \dots, x_n^{\pm 1}]$, we denote by $\delta(L) \in \mathbb{Z}^n$ its *denominator vector*, that is, the unique vector (d_1, \dots, d_n) such that $L(x_1, \dots, x_n) = f(x_1, \dots, x_n) / \prod_{i=1}^n x_i^{d_i}$ where $f(x_1, \dots, x_n)$ is a polynomial not divisible by any x_i .

Since $\text{End}_{\mathcal{C}}(T_i) \simeq \mathbf{k}$ for any $i \in \{1, \dots, n\}$, it follows from [12] that for any indecomposable rigid B_T -module

$$\delta(X_M^T) = (\dim \text{Hom}_{\mathcal{C}}(T_i, \overline{M}))_{1 \leq i \leq n}$$

where \overline{M} is an indecomposable rigid object in \mathcal{C} such that $\text{Hom}_{\mathcal{C}}(T, \overline{M}) = M$.

Since $\text{Hom}_{\mathcal{C}}(T, -)$ induces an equivalence of categories $\mathcal{C}/(\text{add } T[1]) \simeq \text{mod-}B_T$, it follows that

$$\delta(X_M^T) = (\dim \text{Hom}_{B_T}(P_i, M))_{1 \leq i \leq n}$$


where P_i is the indecomposable projective B_T -module at the point i . For any $i \in \{1, \dots, n\}$, the dimension of $\text{Hom}_{B_T}(P_i, M)$ is the multiplicity of the simple B_T -module S_i as a composition factor of M . If M is a string module, this multiplicity is also equal to the number of occurrences of i along the string $\mathbf{s}(M)$. Thus, if $\mathbf{s}(M) = c_1 \dots c_n$, the denominator of X_M^T in its irreducible form is equal to $\prod_{i=0}^n x_{t(c_i)}$.

On the other hand the denominator of L_M (in its irreducible form) is $\mathbf{x}^{-\eta_{\mathbf{s}(M)}} \prod_{i=0}^n x_{t(c_i)}$. Now it follows from Theorem 5.11 that $X_M^T = \frac{1}{\mathbf{x}^{\mathbf{n}_M}} L_M$ and thus the denominators coincide, that is to say $\mathbf{n}_M = \eta_{\mathbf{s}(M)}$. \square

8. EXAMPLES

8.1. A first example in type A_2 . We start with a very simple and detailed example which is nevertheless instructive in order to understand the behaviour of the normalisation.

We consider the frozen quiver (\mathcal{Q}, F) where



and $F = \{3\}$.

The corresponding cluster algebra is of geometric type with initial seed $(\mathbf{x}, \mathbf{y}, B)$ where $\mathbf{x} = (x_1, x_2)$, $\mathbf{y} = (x_3)$ and

$$B = \begin{bmatrix} 0 & 1 \\ -1 & 0 \\ 1 & -1 \end{bmatrix}$$

Its unfrozen part Q is of Dynkin type A_2 . Note that \mathcal{Q} is the ordinary quiver of a cluster-tilted algebra of type A_3 whose relations are $\alpha\beta = \beta\gamma = \gamma\alpha = 0$. Thus (\mathcal{Q}, F) can be embedded in a realisable quadruple $(\mathcal{Q}, F, \mathcal{C}, T)$ where \mathcal{C} is a cluster category of type A_3 . There are three indecomposable modules supported on Q which are the simple S_1 , the simple S_2 and the projective $\mathbf{k}Q$ -module P_1 and all these modules are string modules.

Since (\mathcal{Q}, F) is not blown-up (there are two arrows entering or leaving the frozen point 3), we cannot simply apply Theorem 4.1. So we apply Theorem 5.11, which means that we need to consider normalising factors.

The string of S_1 is e_1 and thus, applying formula (1.1) to S_1 , we get $L_{S_1} = \frac{x_2+x_3}{x_1}$. The quiver \widetilde{Q}_{S_1} is $3^{\gamma:1} \rightarrow v_1 \rightarrow 2^{\alpha:1}$. It is easily computed that $\mathbf{n}_{S_1} = 0$ so that $X_{S_1} = L_{S_1}$ which is indeed the cluster variable in $\mathcal{A}(\mathcal{Q}, F)$ corresponding to the simple module S_1 . We similarly prove that $X_{S_2} = L_{S_2} = \frac{x_1+x_3}{x_2}$.

The case of P_1 is slightly more instructive. We apply the matrix formula and we get $L_{P_1} = \frac{x_3(x_1+x_2+x_3)}{x_1x_2}$. We can now use Proposition 7.3 in order to conclude that the normalising factor is x_3 but we do the computation in order to see the blow-up technique working. The quiver $\text{supp}(\widetilde{P}_1)$ is \mathcal{Q} itself. Its blow-up along P_1 is thus $\widetilde{Q}_{P_1} : 3^{\gamma:1} \rightarrow v_1 \rightarrow v_2 \rightarrow 3^{\beta:2}$ and \widetilde{M} is the representation $\widetilde{P}_1 : 0 \rightarrow \mathbf{k} \xrightarrow{1_{\mathbf{k}}} \mathbf{k} \rightarrow 0$. We compute that

$$\widetilde{L}_{\widetilde{P}_1} = \frac{x_{v_1}x_{3^{\gamma:1}} + x_{v_2}x_{3^{\beta:2}} + x_{3^{\gamma:1}}x_{3^{\beta:2}}}{x_{v_1}x_{v_2}}$$

so that the morphism π sending x_{v_1} to x_1 , x_{v_2} to x_2 and $x_{3^{\gamma:1}}, x_{3^{\beta:2}}$ to x_3 satisfies $\pi(\widetilde{L}_{\widetilde{M}}) = L_M$. And computing directly, or else applying Theorem 4.1, we get $\widetilde{X}_{\widetilde{M}} = \widetilde{L}_{\widetilde{P}_1}$.

The normalisation of M is $\mathbf{n}_M = (0, 0, 1)$ (see Example 5.8) so that we get $X_{P_1} = \frac{1}{x_3}L_{P_1} = \frac{x_1+x_2+x_3}{x_1x_2}$ which is indeed the cluster variable in $\mathcal{A}(\mathcal{Q}, F)$ corresponding to this module.

8.2. Regular cluster variables in type \widetilde{A} . Let Q be a euclidean quiver of type \widetilde{A} equipped with an acyclic orientation. In [5, Theorem 4], the authors provided a combinatorial formula for expressing all but finitely many cluster variables in the coefficient-free cluster algebra associated with Q . The cluster variables they computed correspond in fact to the cluster characters associated with indecomposable postprojective $\mathbf{k}Q$ -modules. Using their methods, it is possible to compute the cluster characters associated with indecomposable postprojective and preinjective

$\mathbf{k}Q$ -modules but not the remaining cluster variables, namely those which correspond to indecomposable regular rigid $\mathbf{k}Q$ -modules. We now use Theorem 4.1 to complete the formula and compute cluster variables associated with regular modules.

It is known that the Auslander-Reiten quiver of $\text{mod-}\mathbf{k}Q$ contains at most two exceptional tubes (that is, tubes of rank ≥ 2) and all the rigid indecomposable regular modules belong to these tubes. Every indecomposable regular module (rigid or not) in such a tube is a string module and its string may be completely described. For simplicity, we only compute the cluster variables associated with quasi-simple regular modules in such tubes, this can easily be extended to any module in one of the exceptional tubes. It is well-known (see for instance [4, §5.2]) that any such module is of the form

$$\cdots 0 \xleftarrow{0} \mathbf{k} \xrightarrow{1_k} \mathbf{k} \xrightarrow{1_k} \cdots \xrightarrow{1_k} \mathbf{k} \xrightarrow{1_k} \mathbf{k} \xleftarrow{0} 0 \cdots$$

so that, if we depict locally the quiver Q as

$$\cdots 0 \longleftarrow 1 \xrightarrow{\alpha_1} 2 \xrightarrow{\alpha_2} \cdots \xrightarrow{\alpha_{n-2}} n-1 \xrightarrow{\alpha_{n-1}} n \longleftarrow n+1 \cdots,$$

the string is $c = \alpha_1 \cdots \alpha_{n-1}$ and we compute

$$L_c = \frac{1}{\prod_{i=1}^{n+1} x_i} [1, 1] \begin{bmatrix} x_0 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} \prod_{i=1}^{n-1} x_{i+1} & 0 \\ \sum_{j=1}^{n-1} \frac{\prod_{i=1}^n x_i}{x_j x_{j+1}} & \prod_{i=1}^{n-1} x_i \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & x_{n+1} \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix}.$$

Now, if M_c denotes the quasi-simple regular module associated with the string c , then it follows from Lemma 7.1 that L_c is the cluster character corresponding to the module M_c and thus, is a cluster variable in the cluster algebra $\mathcal{A}(Q, \emptyset)$.

8.3. An example for the n -Kronecker quiver with principal coefficients.

Let $n \geq 2$ and let K_n^{pp} be the quiver

$$K_n^{\text{pp}} : \begin{array}{ccc} & 1' & 2' \\ & \downarrow & \downarrow \\ & 1 & 2 \\ & \xrightarrow{\alpha_1} & \xrightarrow{\alpha_n} \end{array}$$

with n arrows $\alpha_1, \dots, \alpha_n$ from 1 to 2. We set $F = \{1', 2'\}$ so that K_n^{pp} is the principal extension of the n -Kronecker quiver K_n . We fix two distinct arrows among $\alpha_1, \dots, \alpha_n$ which we denote by α and β .

For any $p \geq 1$, we define a string representation M^p of K_n by setting $M^p(1) = \mathbf{k}^p$, $M^p(2) = \mathbf{k}^{p+1}$, $M^p(\alpha) = 1_{\mathbf{k}^p} \oplus 0$ and $M^p(\beta) = 0 \oplus 1_{\mathbf{k}^p}$ and we view M^p as a representation of K_n^{pp} which is supported on the unfrozen part K_n . For any $p \geq 1$, the string $\mathbf{s}(M^p)$ is $(\alpha^{-1}\beta)^p$ so that it has length $2p$.

We write $y_1 = x_{1'}$ and $y_2 = x_{2'}$. Then, a direct computation shows that

$$(8.7) \quad L_{M^p} = \frac{1}{x_1^p x_2^{p+1}} [1, 1] \begin{bmatrix} 1 & 0 \\ 0 & y_2 x_1^{n-1} \end{bmatrix} \begin{bmatrix} y_1 + x_2^n & y_1 y_2 x_1^{n-1} \\ y_1 x_1 & y_1 y_2 x_1^n \end{bmatrix}^p \begin{bmatrix} 1 \\ x_1 \end{bmatrix}.$$

Applying Corollary 2.11, we see that there exists a realisable quadruple $(K_n^{\text{pp}}, (K_n)'_0, \mathcal{C}, T)$. Since K_n^{pp} is acyclic, it follows from [37] that B_T is hereditary and thus, Lemma 7.1 implies that the normalisation vanishes. Thus, $X_{M^p}^T = L_{M^p}$ is given by the formula (8.7) for every $p \geq 1$.

8.4. **Cyclic cluster-tilted algebras of Dynkin type \mathbb{D} .** Let B be the quotient of the path algebra of the quiver

$$Q : \quad 1 \begin{array}{c} \xrightarrow{\quad} \\ \xrightarrow{\quad} \end{array} 2 \xrightarrow{\quad} 3 \xrightarrow{\quad} \cdots \xrightarrow{\quad} n$$

by the ideal generated by all paths of length $n - 1$. Here we suppose that $n \geq 3$. B is a cluster-tilted algebra of type \mathbb{D}_n if $n \geq 4$ and a cluster-tilted algebra of type

\mathbb{A}_3 if $n = 3$. Let M be the indecomposable module given by the Loewy series $\begin{matrix} 2 \\ 3 \\ \vdots \\ m \end{matrix}$, where $2 \leq m \leq n$.

Proposition 8.1. *Let Q denote the above quiver and let $(\mathcal{Q}, F, \mathcal{C}, T)$ be a realisable quadruple with unfrozen part Q . For any $i \in Q_0$, let, as in Section 4.1,*

$$y_i = \prod_{\alpha \in \mathcal{Q}_1(F, i)} x_{s(\alpha)} \quad \text{and} \quad z_i = \prod_{\alpha \in \mathcal{Q}_1(i, F)} x_{t(\alpha)}.$$

Let M be as above. Then

$$(8.8) \quad L_M = \sum_{\ell=1}^m \frac{x_1 x_2 \cdots x_{m+1}}{x_\ell x_{\ell+1}} \left(\prod_{i=2}^{\ell} y_i \right) \left(\prod_{i=\ell+1}^m z_i \right) \frac{1}{\prod_{j=2}^m x_j}$$

if $m < n$; and if $n = m$ then L_M is given by (8.8) divided by x_1 .

Proof. We proceed by induction on m . If $m = 2$, then M is simple and L_M is given by the exchange relation

$$(8.9) \quad L_M = (x_1 y_2 + x_3 z_2) \frac{1}{x_2}.$$

On the other hand, (8.8) gives

$$\left(\frac{x_1 x_2 x_3}{x_1 x_2} z_2 + \frac{x_1 x_2 x_3}{x_2 x_3} y_2 \right) \frac{1}{x_2},$$

which is equal to the right-hand side of equation (8.9).

Suppose now that $m > 2$. Let N_M be the numerator of the Laurent polynomial L_M . Our formula gives N_M as the product of $2m - 1$ matrices :

$$(8.10) \quad \begin{bmatrix} 1 & x_1 \end{bmatrix} \begin{bmatrix} z_2 & 0 \\ 0 & y_2 \end{bmatrix} \cdots \begin{bmatrix} x_m & 0 \\ 1 & x_{m-1} \end{bmatrix} \begin{bmatrix} z_m & 0 \\ 0 & y_m \end{bmatrix} \begin{bmatrix} x_{m+1} \\ 1 \end{bmatrix}.$$

The product of the last three matrices in (8.10) is equal to

$$\begin{bmatrix} x_m x_{m+1} z_m \\ x_{m+1} z_m + x_{m-1} y_m \end{bmatrix} = \begin{bmatrix} x_m \\ 1 \end{bmatrix} x_{m+1} z_m + \begin{bmatrix} 0 \\ 1 \end{bmatrix} x_{m-1} y_m.$$

Let us denote the product of the first $2m - 4$ matrices by $\begin{bmatrix} a & b \end{bmatrix}$, so

$$N_M = \begin{bmatrix} a & b \end{bmatrix} \left(\begin{bmatrix} x_m \\ 1 \end{bmatrix} x_{m+1} z_m + \begin{bmatrix} 0 \\ 1 \end{bmatrix} x_{m-1} y_m \right).$$

Now let M' be the indecomposable module given by the Loewy series $\begin{matrix} 2 \\ 3 \\ \vdots \\ m-1 \end{matrix}$.

Then, our formula gives $N_{M'}$ as a product of $2m - 3$ matrices, and the first $2m - 4$

matrices are just the same as the first $2m - 4$ matrices in (8.10), and therefore

$$N_{M'} = \begin{bmatrix} a & b \end{bmatrix} \begin{bmatrix} x_m \\ 1 \end{bmatrix}.$$

Thus

$$N_M = N_{M'} x_{m+1} z_m + \begin{bmatrix} a & b \end{bmatrix} \begin{bmatrix} 0 \\ 1 \end{bmatrix} x_{m-1} y_m,$$

which, by induction, is equal to

$$\sum_{\ell=1}^{m-1} \frac{x_1 x_2 \cdots x_m}{x_\ell x_{\ell+1}} \left(\prod_{i=2}^{\ell} y_i \right) \left(\prod_{i=\ell+1}^{m-1} z_i \right) x_{m+1} z_m + \frac{x_1 x_2 \cdots x_m}{x_{m-1} x_m} \left(\prod_{i=2}^{m-1} y_i \right) x_{m-1} y_m,$$

which in turn is equal to

$$\sum_{\ell=1}^{m-1} \frac{x_1 x_2 \cdots x_{m+1}}{x_\ell x_{\ell+1}} \left(\prod_{i=2}^{\ell} y_i \right) \left(\prod_{i=\ell+1}^m z_i \right) + \frac{x_1 x_2 \cdots x_{m+1}}{x_m x_{m+1}} \left(\prod_{i=2}^m y_i \right),$$

and this shows the formula (8.8). The statement for $m = n$ follows from the normalising factor. \square

ACKNOWLEDGMENTS

Work on this problem was started during the *2010 South American Meeting on Representations of Algebras and Related Topics* in Mar del Plata (Argentina); the authors wish to thank the organisers for their kind invitation.

The first author gratefully acknowledges partial support from the NSERC of Canada, the FQRNT of Québec and the Université de Sherbrooke.

This paper was written while the second author was at the Université de Sherbrooke as a CRM-ISM postdoctoral fellow under the supervision of the first author, Thomas Brüstle and Virginie Charette.

The third author is supported by the NSF grants DMS-0908765 and DMS-1001637 and by the University of Connecticut.

The fourth author gratefully acknowledges support from the NSERC of Canada and Bishop's University.

REFERENCES

- [1] C. Amiot. Cluster categories for algebras of global dimension 2 and quivers with potential. *Ann. Inst. Fourier (Grenoble)*, 59(6):2525–2590, 2009.
- [2] I. Assem, T. Brüstle, G. Charbonneau-Jodoin, and P. Plamondon. Gentle algebras arising from surface triangulations. *Algebra and Number Theory*, 4(2):201–229, 2010.
- [3] I. Assem, T. Brüstle, and R. Schiffler. Cluster-tilted algebras as trivial extensions. *Bull. Lond. Math. Soc.*, 40(1):151–162, 2008.
- [4] I. Assem and G. Dupont. Friezes and a construction of the euclidean cluster variables. *Journal of Pure and Applied Algebra*, 2011.
- [5] I. Assem, C. Reutenauer, and D. Smith. Friezes. *Adv. Math.*, 225:3134–3165, 2010.
- [6] I. Assem, D. Simson, and A. Skowroński. *Elements of representation theory of Associative Algebras, Volume 1: Techniques of Representation Theory*, volume 65 of *London Mathematical Society Student Texts*. Cambridge University Press, 2005. MR2197389 (2006j:16020).
- [7] K. Baur and R. Marsh. Categorification of a frieze pattern determinant. *arXiv:1008.5329v1 [math.CO]*, 2010.
- [8] A. Buan, O. Iyama, I. Reiten, and J. Scott. Cluster structures for 2-Calabi-Yau categories and unipotent groups. *Compos. Math.*, 145(4):1035–1079, 2009.
- [9] A. Buan, O. Iyama, I. Reiten, and D. Smith. Mutation of cluster-tilting objects and potentials. *arXiv:0804.3813v3 [math.RT]*, 2009. to appear in *American J. Math.*

- [10] A. Buan, R. Marsh, M. Reineke, I. Reiten, and G. Todorov. Tilting theory and cluster combinatorics. *Adv. Math.*, 204(2):572–618, 2006. MR2249625 (2007f:16033).
- [11] A. Buan, R. Marsh, and I. Reiten. Cluster-tilted algebras. *Trans. Amer. Math. Soc.*, 359(1):323–332, 2007. MR2247893 (2007f:16035).
- [12] A. Buan, R. Marsh, and I. Reiten. Denominators of cluster variables. *J. Lond. Math. Soc.* (2), 79(3):589–611, 2009.
- [13] M. Butler and C. M. Ringel. Auslander-Reiten sequences with few middle terms and applications to string algebras. *Communications in Algebra*, 15(1-2):145–179, 1987.
- [14] P. Caldero and F. Chapoton. Cluster algebras as Hall algebras of quiver representations. *Commentarii Mathematici Helvetici*, 81:596–616, 2006. MR2250855 (2008b:16015).
- [15] P. Caldero, F. Chapoton, and R. Schiffler. Quivers with relations arising from clusters (A_n case). *Transactions of the AMS*, 358:1347–1354, 2006. MR2187656 (2007a:16025).
- [16] P. Caldero and B. Keller. From triangulated categories to cluster algebras II. *Annales Scientifiques de l'École Normale Supérieure*, 39(4):83–100, 2006. MR2316979 (2008m:16031).
- [17] G. Carroll and G. Price. The new combinatorial models for the Ptolemy recurrence. *unpublished*, 2003.
- [18] G. Cerulli Irelli. Quiver Grassmannians associated with string modules. *Journal of Algebraic Combinatorics*, 33(2):259–276, 2011.
- [19] J. Conway and H. Coxeter. Triangulated polygons and frieze patterns. *Math. Gaz.*, 57(401):175–183, 1973.
- [20] J. Conway and H. Coxeter. Triangulated polygons and frieze patterns. *Math. Gaz.*, 57(400):87–94, 1973.
- [21] H. Coxeter. Frieze patterns. *Acta Arith.*, 18:297–310, 1971.
- [22] H. Derksen, J. Weyman, and A. Zelevinsky. Quivers with potentials and their representations. I. Mutations. *Selecta Math. (N.S.)*, 14(1):59–119, 2008.
- [23] G. Dupont. Positivity in coefficient-free rank two cluster algebras. *Electronic Journal of Combinatorics*, 16(1):R98, 2009.
- [24] G. Dupont. Cluster multiplication in regular components via generalized Chebyshev polynomials. *Algebras and Representation Theory*, 2010. To appear.
- [25] G. Dupont. Quantized Chebyshev polynomials and cluster characters with coefficients. *Journal of Algebraic Combinatorics*, 31(4):501–532, june 2010.
- [26] S. Fomin, M. Shapiro, and D. Thurston. Cluster algebras and triangulated surfaces. I. Cluster complexes. *Acta Math.*, 201(1):83–146, 2008.
- [27] S. Fomin and A. Zelevinsky. Cluster algebras I: Foundations. *J. Amer. Math. Soc.*, 15:497–529, 2002. MR1887642 (2003f:16050).
- [28] S. Fomin and A. Zelevinsky. Cluster algebras IV: Coefficients. *Compositio Mathematica*, 143(1):112–164, 2007. MR2295199 (2008d:16049).
- [29] A. Fordy and R. Marsh. Cluster mutation-periodic quivers and associated Laurent sequences. *Journal of Algebraic Combinatorics*, 2011.
- [30] C. Fu and B. Keller. On cluster algebras with coefficients and 2-Calabi-Yau categories. *Trans. Amer. Math. Soc.*, 362:859–895, 2010.
- [31] C. Fu and P. Liu. Lifting to cluster-tilting objects in 2-Calabi-Yau triangulated categories. *Comm. Algebra*, 37(7):2410–2418, 2009.
- [32] P. Gabriel. Auslander-Reiten sequences and representation-finite algebras. In *Representation theory, I (Proc. Workshop, Carleton Univ., Ottawa, Ont., 1979)*, volume 831 of *Lecture Notes in Math.*, pages 1–71. Springer, Berlin, 1980.
- [33] N. Haupt. Euler characteristics of quiver Grassmannians and Ringel-Hall algebras of string algebras. *Algebras and Representation Theory*, pages 1–39, 2010.
- [34] B. Keller. On triangulated orbit categories. *Documenta Mathematica*, 10:551–581, 2005. MR2184464 (2007c:18006).
- [35] B. Keller. Deformed Calabi-Yau completions. *Journal für die reine und angewandte Mathematik*, 2011.
- [36] B. Keller and I. Reiten. Cluster-tilted algebras are Gorenstein and stably Calabi-Yau. *Adv. Math.*, 211(1):123–151, 2007.
- [37] B. Keller and I. Reiten. Acyclic Calabi-Yau categories. *Compositio Mathematicae*, 144(5):1332–1348., 2008. MR2457529.
- [38] B. Keller and S. Scherotzke. Linear recurrence relations for cluster variables of affine quivers. *arXiv:1004.0613v2 [math.RT]*, 2010.

- [39] D. Labardini-Fragoso. Quivers with potentials associated to triangulated surfaces. *Proc. London Math. Soc.*, 98(3):787–839, 2009.
- [40] G. Musiker. A graph theoretic expansion formula for cluster algebras of classical type. *arXiv:0710.3574v2 [math.CO]*, 2007.
- [41] G. Musiker and J. Propp. Combinatorial interpretations for rank-two cluster algebras of affine type. *Electron. J. Combin.*, 14, 2006. MR2285819 (2008j:05374).
- [42] G. Musiker, R. Schiffler, and L. Williams. Positivity for cluster algebras from surfaces. *arXiv:0906.0748v1 [math.CO]*, 2009.
- [43] H. Nakajima. Quiver varieties and cluster algebras. *arXiv:0905.0002v5 [math.QA]*, 2010.
- [44] Y. Palu. Cluster characters for 2-Calabi-Yau triangulated categories. *Ann. Inst. Fourier (Grenoble)*, 58(6):2221–2248, 2008.
- [45] P. Plamondon. Cluster algebras via cluster categories with infinite-dimensional morphism spaces. *arXiv:1004.0830v2 [math.RT]*, 2010.
- [46] J. Propp. The combinatorics of frieze patterns and Markoff numbers. *arXiv:math/0511633v4 [math.CO]*, 2008.
- [47] F. Qin. Quantum cluster variables via Serre polynomials. *arXiv:1004.4171v2 [math.QA]*, 2010.
- [48] I. Reiten. Calabi-Yau categories. Talk at the meeting : Calabi-Yau algebras and N-Koszul algebras (CIRM), 2007.
- [49] R. Schiffler. On cluster algebras arising from unpunctured surfaces II. *Adv. Math.*, 223:1885–1923, 2010.
- [50] R. Schiffler and H. Thomas. On cluster algebras arising from unpunctured surfaces. *Int. Math. Res. Not.*, (17):3160–3189, 2009.
- [51] P. Sherman and A. Zelevinsky. Positivity and canonical bases in rank 2 cluster algebras of finite and affine types. *Mosc. Math. J.*, 4:947–974, 2004. MR2124174 (2006c:16052).
- [52] D. Smith. On tilting modules over cluster-tilted algebras. *Illinois J. Math.*, 52:1223–1247, 2008.
- [53] B. Wald and J. Waschbüsch. Tame biserial algebras. *J. Algebra*, 95(2):480–500, 1985.

UNIVERSITÉ DE SHERBROOKE, SHERBROOKE QC, CANADA

E-mail address: `ibrahim.assem@usherbrooke.ca`

E-mail address: `gregoire.dupont@usherbrooke.ca`

UNIVERSITY OF CONNECTICUT, STORRS CT, USA

E-mail address: `schiffler@math.uconn.edu`

BISHOP'S UNIVERSITY, SHERBROOKE QC, CANADA

E-mail address: `dsmith@ubishops.ca`