



HAL
open science

Absence of traveling wave solutions of conductivity type for the Novikov-Veselov equations at zero energy

Anna Kazeykina

► **To cite this version:**

Anna Kazeykina. Absence of traveling wave solutions of conductivity type for the Novikov-Veselov equations at zero energy. 2011. hal-00603403v2

HAL Id: hal-00603403

<https://hal.science/hal-00603403v2>

Preprint submitted on 15 Jul 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Absence of traveling wave solutions of conductivity type for the Novikov-Veselov equation at zero energy

A.V. Kazeykina ¹

Abstract. We prove that the Novikov-Veselov equation (an analog of KdV in dimension $2 + 1$) at zero energy does not have sufficiently localized soliton solutions of conductivity type.

1 Introduction

In this note we are concerned with the Novikov-Veselov equation at zero energy

$$\begin{aligned} \partial_t v &= 4\operatorname{Re}(4\partial_z^3 v + \partial_z(vw)), \\ \partial_{\bar{z}} w &= -3\partial_z v, \quad v = \bar{v}, \\ v &= v(x, t), \quad w = w(x, t), \quad x = (x_1, x_2) \in \mathbb{R}^2, \quad t \in \mathbb{R}, \end{aligned} \tag{1}$$

where

$$\partial_t = \frac{\partial}{\partial t}, \quad \partial_z = \frac{1}{2} \left(\frac{\partial}{\partial x_1} - i \frac{\partial}{\partial x_2} \right), \quad \partial_{\bar{z}} = \frac{1}{2} \left(\frac{\partial}{\partial x_1} + i \frac{\partial}{\partial x_2} \right).$$

Definition 1. A pair (v, w) is a sufficiently localized solution of equation (1) if

- $v, w \in C(\mathbb{R}^2 \times \mathbb{R})$, $v(\cdot, t) \in C^3(\mathbb{R}^3)$,
- $|\partial_x^j v(x, t)| \leq \frac{q(t)}{(1 + |x|)^{2+\varepsilon}}$, $|j| \leq 3$, for some $\varepsilon > 0$, $w(x, t) \rightarrow 0$, $|x| \rightarrow \infty$,
- (v, w) satisfies (1).

Definition 2. A solution (v, w) of (1) is a soliton (a traveling wave) if $v(x, t) = V(x - ct)$, $c \in \mathbb{R}^2$.

Equation (1) is an analog of the classic KdV equation. When $v = v(x_1, t)$, $w = w(x_1, t)$, then equation (1) is reduced to KdV. Besides, equation (1) is integrable via the scattering transform for the 2-dimensional Schrödinger equation

$$\begin{aligned} L\psi &= 0, \\ L &= -\Delta + v(x, t), \quad \Delta = 4\partial_z\partial_{\bar{z}}, \quad x \in \mathbb{R}^2. \end{aligned} \tag{2}$$

Equation (1) is contained implicitly in [M] as an equation possessing the following representation

$$\frac{\partial(L - E)}{\partial t} = [L - E, A] + B(L - E), \tag{3}$$

where L is defined in (2), A and B are suitable differential operators of the third and zero order respectively and $[\cdot, \cdot]$ denotes the commutator. In the explicit form equation (1) was written in [NV1], [NV2], where it was also studied in the periodic setting. For the rapidly decaying potentials the studies of equation (1) and the scattering problem for (2) were carried out in [BLMP], [GN] [T], [LMS]. In [LMS] the relation with the Calderón conductivity problem was discussed in detail.

¹CMAP, Ecole Polytechnique, Palaiseau, 91128, France; email: kazeykina@cmap.polytechnique.fr

Definition 3. A potential $v \in L^p(\mathbb{R}^2)$, $1 < p < 2$, is of conductivity type if $v = \gamma^{-1/2} \Delta \gamma^{1/2}$ for some real-valued positive $\gamma \in L^\infty(\mathbb{R}^2)$, such that $\gamma \geq \delta_0 > 0$ and $\nabla \gamma^{1/2} \in L^p(\mathbb{R}^2)$.

The potentials of conductivity type arise naturally when the Calderón conductivity problem is studied in the setting of the boundary value problem for the 2-dimensional Schrödinger equation at zero energy (see [Nov1], [N], [LMS]); in addition, in [N] it was shown that for this type of potentials the scattering data for (2) are well-defined everywhere.

The main result of the present note consists in the following: there are no solitons of conductivity type for equation (1). The proof is based on the ideas proposed in [Nov2].

This work was fulfilled in the framework of research carried out under the supervision of R.G. Novikov.

2 Scattering data for the 2-dimensional Schrödinger equation at zero energy with a potential of conductivity type

Consider the Schrödinger equation (2) on the plane with the potential $v(z)$, $z = x_1 + ix_2$, satisfying

$$\begin{aligned} v(z) &= \overline{v(\bar{z})}, \quad v(z) \in L^\infty(\mathbb{C}), \\ |v(z)| &< q(1 + |z|)^{-2-\varepsilon} \text{ for some } q > 0, \varepsilon > 0. \end{aligned} \quad (4)$$

For $k \in \mathbb{C}$ we consider solutions $\psi(z, k)$ of (2) having the following asymptotics

$$\psi(z, k) = e^{ikz} \mu(z, k), \quad \mu(z, k) = 1 + o(1), \text{ as } |z| \rightarrow \infty, \quad (5)$$

i.e. Faddeev's exponentially growing solutions for the two-dimensional Schrödinger equation (2) at zero energy, see [F], [GN], [Nov1].

It was shown that if v satisfies (4) and is of conductivity type, then $\forall k \in \mathbb{C} \setminus 0$ there exists a unique continuous solution of (2) satisfying (5) (see [N]). Thus the scattering data b for the potential v of conductivity type are well-defined and continuous:

$$b(k) = \iint_{\mathbb{C}} e^{i(ky + \bar{k}\bar{y})} v(y) \mu(y, k) d\text{Re}y d\text{Im}y, \quad k \in \mathbb{C} \setminus 0. \quad (6)$$

In addition (see [N]), the function $\mu(z, k)$ from (5) satisfies the following $\bar{\partial}$ -equation

$$\frac{\partial \mu(z, k)}{\partial \bar{k}} = \frac{1}{4\pi \bar{k}} e^{-i(kz + \bar{k}\bar{z})} b(k) \overline{\mu(z, k)}, \quad z \in \mathbb{C}, \quad k \in \mathbb{C} \setminus 0 \quad (7)$$

and the following limit properties:

$$\mu(z, k) \rightarrow 1, \text{ as } |k| \rightarrow \infty, \quad (8)$$

$$\mu(z, k) \text{ is bounded in the neighborhood of } k = 0. \quad (9)$$

The following lemma describes the scattering data corresponding to a shifted potential.

Lemma 1. Let $v(z)$ be a potential satisfying (4) with the scattering data $b(k)$. The scattering data $b_y(k)$ for the potential $v_y(z) = v(z - y)$ are related to $b(k)$ by the following formula

$$b_y(k) = e^{i(ky + \bar{k}\bar{y})} b(k), \quad k \in \mathbb{C} \setminus 0, \quad y \in \mathbb{C}. \quad (10)$$

Proof. We note that $\psi(z - y, k)$ satisfies (2) with $v_y(z)$ and has the asymptotics $\psi(z - y, k) = e^{ik(z-y)}(1 + o(1))$ as $|z| \rightarrow \infty$. Thus $\psi_y(z, k) = e^{iky}\psi(z - y, k)$ and $\mu_y(z, k) = \mu(z - y, k)$. Finally, we have

$$\begin{aligned} b_y(k) &= \iint_{\mathbb{C}} e^{i(k\zeta + \bar{k}\bar{\zeta})} v_y(\zeta) \mu_y(\zeta, k) d\operatorname{Re}\zeta d\operatorname{Im}\zeta = \\ &= \iint_{\mathbb{C}} e^{i(k\zeta + \bar{k}\bar{\zeta})} v(\zeta - y) \mu(\zeta - y, k) d\operatorname{Re}\zeta d\operatorname{Im}\zeta = e^{i(ky + \bar{k}\bar{y})} b(k). \end{aligned}$$

□

As for the time dynamics of the scattering data, in [BLMP], [GN] it was shown that if the solution (v, w) of (1) exists and the scattering data for this solution are well-defined, then the time evolution of these scattering data is described as follows:

$$b(k, t) = e^{i(k^3 + \bar{k}^3)t} b(k, 0), \quad k \in \mathbb{C} \setminus 0, \quad t \in \mathbb{R}. \quad (11)$$

3 Absence of solitons of conductivity type

Theorem 1. *Let (v, w) be a sufficiently localized traveling wave solution of (1) of conductivity type. Then $v \equiv 0$, $w \equiv 0$.*

Scheme of proof. From (10), (11), continuity of $b(k)$ on $\mathbb{C} \setminus 0$ and the fact that the functions k , \bar{k} , k^3 , \bar{k}^3 , 1 are linearly independent in the neighborhood of any point, it follows that $b \equiv 0$. Equation (7) implies that in this case the function $\mu(z, k)$ is holomorphic on k , $k \in \mathbb{C} \setminus 0$. Using properties (8) and (9) we apply Liouville theorem to obtain that $\mu \equiv 1$. Then $\psi(z, k) = e^{ikz}$ and from (2) it follows that $v \equiv 0$. □

References

- [BLMP] Boiti M., Leon J.J.P., Manna M., Pempinelli F.: On a spectral transform of a KdV-like equation related to the Schrodinger operator in the plane. *Inverse Problems*. 3, 25–36 (1987)
- [F] Faddeev L.D. Growing solutions of the Schrödinger equation. *Dokl. Akad. Nauk SSSR*. 165, 514-517 (1965); translation in *Sov. Phys. Dokl.* 10, 1033-1035 (1966)
- [GN] Grinevich P.G., Novikov S.P.: Two-dimensional “inverse scattering problem” for negative energies and generalized-analytic functions. I. Energies below the ground state. *Funct. Anal. Appl.* 22(1), 19–27 (1988)
- [LMS] Lassas M., Mueller J.L., Siltanen S.: Mapping properties of the nonlinear Fourier transform in dimension two. *Communications in Partial Differential Equations*. 32, 591–610 (2007)
- [M] Manakov S.V.: The inverse scattering method and two-dimensional evolution equations. *Uspekhi Mat. Nauk.* 31(5), 245–246 (1976) (in Russian)
- [N] Nachman A.I.: Global uniqueness for a two-dimensional inverse boundary value problem. *Annals of Mathematics*. 143, 71–96 (1995)

- [Nov1] Novikov R.G.: Multidimensional inverse spectral problem for the equation $-\Delta\psi+(v(x)-Eu(x))\psi=0$. *Funkt. Anal. i Pril.* 22(4), 11-22 (1988); translation in *Funct. Anal. Appl.* 22, 263-272 (1988)
- [Nov2] Novikov R.G.: Absence of exponentially localized solitons for the Novikov–Veselov equation at positive energy. *Physics Letters A.* 375, 1233–1235 (2011)
- [NV1] Novikov S.P., Veselov A.P.: Finite-zone, two-dimensional, potential Schrödinger operators. Explicit formula and evolutions equations. *Dokl. Akad. Nauk SSSR.* 279, 20–24 (1984), translation in *Sov. Math. Dokl.* 30, 588-591 (1984)
- [NV2] Novikov S.P., Veselov A.P.: Finite-zone, two-dimensional Schrödinger operators. Potential operators. *Dokl. Akad. Nauk SSSR.* 279, 784–788 (1984), translation in *Sov. Math. Dokl.* 30, 705–708 (1984)
- [T] Tsai T.-Y. The Schrödinger operator in the plane. *Inverse Problems.* 9, 763–787 (1993)