Shoreline instability under low-angle wave incidence

Déborah Idier, Albert Falqués, Gerben Ruessink, Roland Garnier

► To cite this version:

Déborah Idier, Albert Falqués, Gerben Ruessink, Roland Garnier. Shoreline instability under low-angle wave incidence. *Journal of Geophysical Research: Earth Surface*, 2011, 111 (F4), 12 p. 10.1029/2010JF001894 . hal-00635211

HAL Id: hal-00635211

<https://hal-brgm.archives-ouvertes.fr/hal-00635211>

Submitted on 24 Oct 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Shoreline instability under low-angle wave incidence

D. Idier

BRGM, 3 Avenue C. Guillemin, BP 6009, 45060 Orléans Cedex 2, France

A. Falqués

UPC, C/ Jordi Girona 1-3, Modul B4/B5, despatx 103 - E-08034, Barcelona,

Catalonia, Spain

B.G. Ruessink

Utrecht University, Institute for Marine and Atmospheric Research Utrecht,

Faculty of Geosciences, Department of Physical Geography, P.O. box 80115,

3508 TC Utrecht, The Netherlands

R. Garnier

Instituto de Hidráulica Ambiental IH Cantabria, Universidad de Cantabria,

ETSI Caminos Canales y Puertos, Avda. los Castros s/n, 39005 Santander,

Cantabria, Spain

D. Idier, BRGM, 3 Avenue C. Guillemin, BP 6009, 45060 Orléans Cedex 2, France.

(d.idier@brgm.fr)

A. Falqués, UPC, C/ Jordi Girona 1-3, Modul B4/B5, despatx 103 - E-08034, Barcelona,

Catalonia, Spain. (falques@fa.upc.edu)

Abstract.

The growth of megacusps as shoreline instabilities is investigated by examining the coupling between wave transformation in the shoaling zone, long-shore transport in the surf zone, cross-shore transport, and morphological evolution. This coupling is known to drive a potential positive feedback in case of very oblique wave incidence, leading to an unstable shoreline and the consequent formation of shoreline sandwaves. Here, using a linear stability model based on the one-line concept, we demonstrate that such instabilities can also develop in case of low-angle or shore-normal incidence, under certain conditions (small enough wave height and/or large enough beach slope). The wavelength and growth time scales are much smaller than those of high-angle wave instabilities and are nearly in the range of those of surf zone rhythmic bars, $O(10^2 - 10^3 \text{ m})$ and $O(1 - 10 \text{ days})$, respectively. The feedback mechanism is based on: (1) wave refraction by a shoal (defined as a cross-shore extension of the shoreline perturbation) leading to wave convergence shoreward of it, (2) longshore sediment flux convergence between the shoal and the shoreline, resulting in megacusp formation, and (3) cross-shore sed-

B.G. Ruessink, Utrecht University, Institute for Marine and Atmospheric Research Utrecht, Faculty of Geosciences, Department of Physical Geography, P.O. box 80115, 3508 TC Utrecht, The Netherlands. (b.g.ruessink@uu.nl)

R. Garnier, IH Cantabria, Universidad de Cantabria, Avda. los Castros s/n, 39005 Santander, Cantabria, Spain. (garnierr@unican.es)

28 iment flux from the surf to the shoaling zone, feeding the shoal. Even though
29 the present model is based on a crude representation of nearshore dynam-
30 ics, a comparison of model results with existing 2DH model output and lab-
31 oratory experiments suggests that the instability mechanism is plausible. Ad-
32 ditional work is required to fully assess whether and under which conditions
33 this mechanism exists in nature.

1. Introduction

Rhythmic shorelines featuring planview undulations with a relatively regular spacing or wavelength are quite common on sandy coasts. In the present study, we focus on undulations that are linked to submerged bars or shoals and are generally known as shoreline sandwaves [Komar, 1998; Bruun, 1954]. These sandwaves can be classified according to their length scale as short and long sandwaves (see, e.g., Stewart and Davidson-Arnott [1988]).

The spacing of short sandwaves ranges from several tens to several hundreds of meters and their seaward perturbations are known as megacusps. Observations show that these megacusps can develop shoreward of crescentic bar systems during the typical “Rhythmic Bar and Beach” morphological beach state or can develop from the shore attachment of transverse bars that characterise the “Transverse Bar and Beach” state [Wright and Short, 1984]. These transverse bars can appear where the horns of a previous crescentic bar approach the shoreline [Wright and Short, 1984; Sonu, 1973; Ranasinghe et al., 2004; Lafon et al., 2004; Castelle et al., 2007]. On the other hand, transverse bars can also develop freely, independently of any offshore rhythmic system (e.g. the “transverse finger bars” [Sonu, 1968, 1973; Ribas and Kroon, 2007]). The formation of rhythmic surf zone bars and associated megacusps is believed to be due primarily to an instability of the coupling between the evolving bathymetry and the distribution of wave breaking (bed-surf coupling) [Falqués et al., 2000]. The developing shoals and channels cause changes in wave breaking, which in turn cause gradients in radiation stresses and thereby horizontal circulation with rip currents. If the sediment fluxes carried by this circulation converge

55 over the shoals and diverge over the channels, a positive feedback arises and the coupled
56 system self-organizes to produce certain patterns, both morphological and hydrodynamic
57 (see, e.g., *Reniers et al.* [2004]; *Garnier et al.* [2008]). In the case of oblique wave inci-
58 dence, a meandering of the longshore current is also essential to the instability process
59 [*Garnier et al.*, 2006]. Two important characteristics of all available models of the self-
60 organized formation of rhythmic surf zone bars are that they (1) are essentially based
61 on sediment transport driven by the longshore current and rip currents only, i.e. ignore
62 cross-shore transport induced by undertow and wave non-linearity and (2) do not consider
63 morphological changes beyond the offshore reach of the rip-current circulation.

64 Rhythmic shorelines can also develop as a result of an instability not related to bed-
65 surf coupling. *Ashton et al.* [2001] and *Ashton and Murray* [2006a, b] have shown that
66 sandy shorelines are unstable for wave angles (angle between wave fronts and the local
67 shoreline orientation) larger than about 42° in deep water, leading to the formation of
68 shoreline sandwaves, cusped features and spits. *Falqués and Calvete* [2005] have found
69 that the initial characteristic wavelength of the emerging sandwaves is in the range of 3 to
70 15 km, i.e., much larger than that of surf zone rhythmic bars. This instability caused by
71 high-angle waves will henceforth be referred to as HAWI (High-Angle Wave Instability).
72 The physical mechanism can be explained as follows. For oblique wave incidence, there
73 are essentially two counteracting effects. On one hand, the angle relative to the local
74 shoreline is larger on the lee of a cusped feature than on the updrift side. This tends to
75 cause larger alongshore sediment flux at the lee and thereby divergence of sediment flux
76 along the bump, which therefore tends to erode. On the other hand, since the refractive
77 wave ray turning is stronger at the lee than at the updrift flank, there is more wave energy

78 spreading due to crest stretching at the lee causing smaller waves and smaller alongshore
79 sediment flux. This produces convergence of sediment flux at the bump, which therefore
80 tends to grow. For high angle waves the latter effect dominates and, if the bathymetric
81 perturbation associated with the shoreline feature extends far enough offshore, it leads to
82 the development of the cusped feature. In contrast to the bed-surf instability for rhythmic
83 surf zone bars, this instability mechanism depends essentially on the coupling between the
84 surf and shoaling zones. Indeed, the gradients in alongshore sediment flux that induce
85 bathymetric changes in the surf zone occur because of wave field perturbations induced by
86 bathymetric features in the shoaling zone. Thus, in order to achieve a positive feedback,
87 the shoals (or the bed depressions) in the surf zone must extend into the shoaling zone.
88 This is achieved by the cross-shore sediment fluxes induced, for instance, by wave non-
89 linearity, gravity and undertow, which force the cross-shore shoreface profile to reach an
90 equilibrium profile. HAWI may provide an explanation for the self-organized formation
91 of some long shoreline sand waves which are reported in the literature [*Verhagen*, 1989;
92 *Inman et al.*, 1992; *Thevenot and Kraus*, 1995; *Ruessink and Jeuken*, 2002; *Davidson-*
93 *Arnott and van Heyningen*, 2003].

94 On the other hand, some observations for low incidence angles show that longshore
95 currents can converge on megacusps because of wave refraction [*Komar*, 1998]. Such cur-
96 rent convergence may lead to longshore sediment flux convergence and hence to megacusp
97 growth. If the submerged part of the megacusp grows and extends far enough into the
98 shoaling zone (due to the cross-shore transport leading to an equilibrium profile), wave
99 refraction would be enhanced and a positive feedback would arise. This might provide a
100 mechanism for shoreline instability formation under low-angle wave incidence that bears

some similitude with HAWI in the sense that the coupling between the surf and shoaling zones is essential, in contrast to the bed-surf instability. We will refer to this potential mechanism as Low-Angle Wave Instability (LAWI). The aim of the present contribution is to investigate this new morphodynamic instability mechanism and discuss whether the resulting shoreline instability may be found in nature.

The lay-out of this paper is as follows. First (Section 2), we introduce the 1D-morfo model [Falqués and Calvete, 2005] that we used to investigate LAWI. Numerical experiments of idealized cases are presented and analyzed in Section 3. We find that shoreline sandwaves can indeed develop because of LAWI, with a length scale comparable to those of megacusps and rhythmic surf zone bars. In Section 4, we analyze the physics of the instability mechanism. We conclude our paper with a discussion and a summary of the main results.

2. The shoreline stability model: 1D-morfo

Owing to the similitude with HAWI [Ashton *et al.*, 2001; Falqués and Calvete, 2005], the LAWI mechanism is assumed to be un-related to surf zone processes like rip or longshore meandering currents. Thus, the engineering simplification of one-line modeling [Dean and Dalrymple, 2002] is used, where the shoreline dynamics are based on alongshore gradients in the total alongshore transport rate, Q (i.e., the total volume of sand carried by the wave-driven longshore current that crosses a cross-section of the surf zone area for unit of time (m^3/s)). Using such a simple model, which neglects numerous aspects of surf zone dynamics, including rip current circulation, longshore current meandering, and thus the bed-surf coupling phenomena, allows to investigate properly whether the LAWI mechanism is supported by the governing equations. Furthermore, the consistency

123 of the sediment transport patterns from the one-line modelling has been confirmed by
 124 using a 2DH model (delft3D) [*List and Ashton*, 2007], at least in case of HAWI. These
 125 reasons, in addition to the fact that HAWI has been studied and reproduced with a linear
 126 stability model called 1D-morfo based on the one-line concept, lead us to use the same
 127 model to investigate LAWI and its differences with HAWI. A very brief description of the
 128 1D-morfo model is given here. The reader is referred to *Falqués and Calvete* [2005] for
 129 further details.

130 The model describes the dynamics of small amplitude perturbations of an otherwise rec-
 131 tilinear coastline. Following the one-line concept, the dynamics are governed by gradients
 132 in the total alongshore wave-driven transport rate Q :

$$\bar{D} \frac{\partial x_s}{\partial t} = - \frac{\partial Q}{\partial y} \quad (1)$$

133 A Cartesian coordinate system is used, with x increasing seaward in the unperturbed
 134 cross-shore direction and y running alongshore (Figure 1). The position of the shoreline
 135 is given by $x = x_s(y, t)$, where t is time and $\bar{D}$ is the active water depth (as defined by
 136 *Falqués and Calvete* [2005]), which is of the order of the depth of closure. This active
 137 water depth is directly related to the one-line model concept, for more details, see *Falqués*
 138 *and Calvete* [2005]. The transport rate Q is computed according to the longshore sediment
 139 transport equation of *Ozasa and Brampton* [1980]. In this formulation, Q is the sum of
 140 two terms: the first one (Q_1) is driven by waves approaching the shore at an angle and
 141 is equivalent to the CERC formula [*USACE*, 1984], and the second one (Q_2) takes into
 142 account the influence of wave set-up induced currents related to the alongshore gradient
 143 in the wave height.

Figure 1

144 The equation of the sediment transport rate Q can be written as follows:

$$Q = Q_1 + Q_2 \quad Q = \mu H_b^{5/2} \left(\sin(2(\theta_b - \phi)) - r \frac{2}{\beta} \cos(\theta_b - \phi) \frac{\partial H_b}{\partial y} \right) \quad (2)$$

145 where H_b is the (rms) wave height at breaking (index b), θ_b is the angle between wave
146 fronts and the unperturbed coastline at breaking, and $\phi = \tan^{-1}(\partial x_s / \partial y)$ is the local
147 orientation of the perturbed shoreline. β is the beach slope at the instantaneous shoreline
148 (i.e. the waterline). The constant μ is proportional to the empirical parameter K_1 of
149 the original CERC formula and is $\sim 0.1 - 0.2 \text{ m}^{1/2}\text{s}^{-1}$. For the reference case of this
150 paper, a value, $\mu = 0.15 \text{ m}^{1/2}\text{s}^{-1}$, was chosen, which corresponds to $K_1 = 0.525$. The
151 nondimensional parameter r is equal to K_2/K_1 , where K_2 is the empirical parameter of
152 *Ozasa and Brampton* [1980]. According to *Horikawa* [1988], r ranges between 0.5 and 1.5,
153 whereas *Ozasa and Brampton* [1980] suggest a value of 1.62. The value $r = 1$ is used for
154 the present reference case, which is equivalent to $K_2 = K_1$ and has been used in several
155 earlier studies on shoreline instabilities [*Bender and Dean*, 2004; *List et al.*, 2008; *van den*
156 *Berg et al.*, 2011]. However, the term Q_2 is not always taken into account, and its validity
157 and application range are uncertain. In section 3.2, we will study the sensitivity of our
158 results to the value of r .

159 Some discussion exists about the capacity of the CERC formula (Q_1) to predict correctly
160 gradients in alongshore sediment transport in the presence of bathymetric perturbations
161 [*List et al.*, 2006, 2008; *van den Berg et al.*, 2011]. The results of *List and Ashton* [2007]
162 suggest that the CERC formula predicts qualitatively correct transport gradients for large
163 scale shoreline undulations (alongshore lengths of 1-8 km). The term Q_2 was introduced to
164 describe the sediment transport resulting from alongshore variations in the breaker wave
165 height induced by diffraction near coastal structures. These breaker-height variations in-

166 duce alongshore gradients in set-up, which drive longshore currents and hence sediment
 167 transport. In our work, alongshore variability in breaker heights is related to wave re-
 168 fraction rather than to wave diffraction, but the subsequent mechanism for alongshore
 169 sediment transport remains the same.

170 To compute the sediment transport rate according to eq. 2, $H_b(y, t)$ and $\theta_b(y, t)$ are
 171 needed. The procedure to determine them is as follows. It is assumed that the wave
 172 height and wave angle are alongshore uniform in deep water. Then, wave transformation,
 173 including refraction and shoaling, is performed from deep water up to the breaking point
 174 so that $H_b(y, t)$ and $\theta_b(y, t)$ are determined and Q can be computed. To do the wave
 175 transformation, a perturbed nearshore bathymetry coupled to the shoreline changes is
 176 assumed:

$$D(x, y, t) = D_0(x) - \beta f(x) x_s(y, t) \quad (3)$$

177 where $D(x, y, t)$ and $D_0(x)$ are the perturbed and unperturbed water depth, respectively,
 178 and $f(x)$ is a shape function. Figure 2a shows some examples of possible perturbation
 179 profiles: constant bed perturbation in the surf zone and decreasing exponentially in the
 180 offshore direction (P1), bed perturbation decreasing exponentially in the offshore direction
 181 from the coast (P2), and bed perturbation similar to a shoal (P3 and P4).

Figure 2

182 The offshore extension of the bathymetric perturbation is controlled by its “charac-
 183 teristic” length, xl , which is a free parameter in the model. It was shown by *Falqués*
 184 *and Calvete* [2005] that the coupling between the surf and shoaling zones is crucial for
 185 HAWI. This is accomplished only if xl is at least a couple of times larger than the surf
 186 zone width. The parameter xl can be seen as a way to parameterise cross-shore sediment
 187 transport, especially between the surf and shoaling zones. This makes HAWI essentially

different from the surf zone morphodynamic instabilities that lead to rhythmic bars and
 rip channels. The changes in the shoreline cause changes in the bathymetry (both in the
 surf and shoaling zones), which in turn cause changes in the wave field. The changes in
 the wave field affect the sediment transport that drives shoreline evolution. Therefore,
 the shoreline, the bathymetry and the wave field are fully coupled.

Following the linear stability concept, the perturbation of the shoreline is assumed to
 be

$$x_s(y, t) = ae^{\sigma t + iKy} + c.c. \quad (4)$$

where a is a small amplitude. For each given (real) wavenumber, K , this expression is
 inserted into the governing equation (eq. 1), and into the perturbed bathymetry equation
 (eq. 3). By computing the perturbed wave field and inserting H_b and θ_b in eq. 1, the
 complex growth rate, $\sigma(K) = \sigma_r + i\sigma_i$, is determined. All of the equations are linearized
 with respect to the amplitude, a . Then, for those K such that $\sigma_r(K) > 0$ a sandwave with
 wavelength $\lambda = 2\pi/K$ tends to emerge from a positive feedback between the morphology
 and the wave field. The pattern that has the maximum growth rate is called the Linearly
 Most Amplified mode (LMA mode).

3. Numerical experiments on idealised cases

To investigate the possible mechanism causing shoreline instabilities under low wave
 incidence angles, numerical experiments on idealised cases are performed. First, numerical
 experiments and results are given. Then, a sensitivity study is done to assess better the
 results.

3.1. Instabilities versus beach slope and wave incidence angle

3.1.1. Configuration

A Dean profile (Figure 2b) is chosen as the equilibrium profile, using various beach slopes (Figures 2b and c). The adopted profile is given by:

$$D_0(x) = b \left((x + x_0)^{2/3} - x_0^{2/3} \right) \quad (5)$$

which has been modified from the original Dean profile to avoid an infinite slope at the shoreline. The constants b and x_0 are determined from the prescribed slope β at the coastline and the prescribed distance x_c from the coastline to the location of the closure depth D_c (see *Falqués and Calvete* [2005] for details). The forcing conditions are waves with $H_{rms} = 1.5$ m and $Tp = 8$ s at a water depth of 25 m. The wave direction ranges from 0° to 85° in increments of 5° . A closure depth $\bar{D} = 20$ m is chosen. To perform the linear stability analysis, the shape function for the bathymetric perturbations is assumed to be constant in the surf zone and to decrease exponentially seaward. Its cross-shore extent is given by the characteristic distance xl corresponding to the closure depth of 20 m, i.e. $xl = 1410$ m for the present case (P1-perturbation in Figure 2a).

3.1.2. Results

Figure 3a shows the growth rate of the LMA mode versus the wave angle and the beach slope. The wave angle is given for a water depth of 25 m. The beach slope is defined as the beach slope at the shoreline. For small beach slopes (< 0.04) the coast behaves as expected: it is unstable only if the wave incidence angle is large enough. In this case, the shoreline instabilities clearly correspond to HAWI. For instance, for a beach slope of 0.02,

Figure 3

a wave incidence of 70° leads to the largest growth rate ($3.13 \times 10^{-9} \text{ s}^{-1}$), corresponding to a wavelength of 7000 m (Figure 3b).

However, for larger beach slopes, instabilities occur for all wave directions, and especially for low wave incidence angles. These instabilities correspond to what will be called LAWI in the present paper. Furthermore, among all the wave incidence angles, the most amplified mode is for shore-normal wave incidence. For a beach slope of 0.1, an angle of 0° leads to the largest growth rate ($1.82 \times 10^{-6} \text{ s}^{-1}$), corresponding to a wavelength of 571 m (Figure 3b).

3.2. Sensitivity analysis

The sensitivity analysis is performed using a planar beach, as the aim is to focus on the mechanisms. However, simulations with other (e.g. barred) profiles also lead to LAWI (not shown).

3.2.1. Wave height and beach slope

Keeping the same reference configuration as above and focusing on shore-normal wave incidence, we investigate the sensitivity of the instability to the wave height for a range of beach slopes. For normally incident waves, instabilities develop only for a beach slope that exceeds 0.04 (Figure 4). In this case, there is an optimum in the wave height for which the growth rate is largest. This wave height increases with the beach slope. For instance, for a beach slope of 0.1 and 0.18), the optimal wave height H_{rms} is 1.75 m and 2.5 m, respectively. A wave height increase also leads to an increase in the LMA mode wavelength, which is due to the corresponding increase in surf zone width.

Figure 4

3.2.2. Bathymetric perturbation length

The influence of the parameter xl was previously investigated by *Falqués and Calvete* [2005], who showed that xl must exceed a threshold to initiate HAWI (the perturbation must extend across both the surf and shoaling zones). Similar behaviour is found here by exploring the range between $xl = 10$ and $xl = 10^4$ m. Figure 5 shows that there is a threshold, $xl > 100$ m, above which shoreline instabilities may occur. This value appears physically reasonable since the width of the surf zone in the reference case is about 73 m. For $125 \leq xl \leq 250$ m, the shoreline instability wavelength decreases significantly (from 1040 to 530 m) with increasing perturbation length, whereas for $xl \geq 250$ m, the wavelength increases only slightly until reaching a nearly constant value of 574 m. The growth rate increases with increasing xl for values below 500 m but decreases for xl exceeding 500 m, reaching a nearly constant value for large perturbation length scales. The main conclusion is that instabilities occur only if the perturbation extends far enough into the shoaling zone. When the perturbation length is about the width of the surf zone, it influences the wavelength and growth rate of the LMA mode strongly, whereas for larger perturbation length values, this influence is negligible.

3.2.3. Initial perturbation shape

To evaluate the influence of the bathymetric perturbation shape function on the results we have presented so far, several shapes were investigated using the same wave-boundary conditions and a perturbation length xl of 2000 m. The shape functions we consider are (Figure 2a):

- Perturbation P1: bed perturbation constant in the surf zone and decreasing exponentially in the offshore direction

269 • Perturbation P2: bed perturbation decreasing exponentially from the coast in the
270 offshore direction

271 • Perturbation P3: P2 perturbation with a shoal located only in the shoaling zone,
272 from 400 to 800 m, with a maximum height at $x_1 = 600$ m.

273 • Perturbation P4: P2 perturbation with a shoal located in both the surf and shoaling
274 zones, from 0 to 1400 m, with a maximum height at $x_1 = 600$ m.

275 The reference configuration is still the same ($H_{rms}=1.5$ m, $T = 8$ s, $\theta = 0^\circ$). The four
276 shape functions result in LMA mode wavelengths of 571, 571, 608 and 608 m, respectively,
277 and growth rates of 1.7, 1.5, 1.7, $1.7 \times 10^{-6} \text{ s}^{-1}$, respectively. Thus, the results are
278 slightly sensitive (mean wavelength of 598 m and a standard deviation of 18 m) to the
279 bed perturbation type, but all perturbation types cause LAWI with similar growth rates.

280 3.2.4. Sediment transport equation

281 To investigate the sensitivity of the results to the sediment transport equation, compu-
282 tations were carried out with $r = 0$, which reduces eq. 2 to the CERC equation. This
283 sensitivity study is done in beach slope - wave angle space. The LMA characteristics are
284 quite similar with $r = 1$ (Figure 3) and $r = 0$ (Figure 6), except for the case of normal
285 wave incidence. In this case ($r = 0$), for small beach slopes, all of the perturbations are
286 damped, as for $r = 1$. For larger beach slopes, the growth rate increases with decreasing
287 perturbation wavelength without reaching a local maximum (Figure 7). Thus there is no
288 LMA mode for shore-normal wave incidence (X symbol on Figure 6). This specific case for
289 shore-normal wave incidence will be discussed in section 4. To summarize, the previous
290 results are not highly sensitive to the second term of the sediment transport equation,
291 except for the case of shore-normal wave incidence.

Figure 6

Figure 7

4. The physical mechanism

Here we investigate the physics behind the model prediction of shoreline instabilities caused by low wave incidence angles. The physical processes are analysed based on the study of the growth rate components, and the hydrodynamic and sediment transport patterns, before identifying the main mechanisms.

4.1. Growth rate analysis

In the perturbed situation where the shoreline position is given by eq. 4, the wave height and wave angle at breaking are given by:

$$\begin{aligned} H_b(y, t) &= H_b^0 + (\hat{H}'_{br} + i\hat{H}'_{bi})e^{\sigma t + iKy} + c.c. \\ \theta_b(y, t) &= \theta_b^0 + (\hat{\theta}'_{br} + i\hat{\theta}'_{bi})e^{\sigma t + iKy} + c.c. \end{aligned} \quad (6)$$

where H_b^0, θ_b^0 are the wave height and angle for the unperturbed situation. Then, according to *Falqués and Calvete* [2005], the growth rate (the real part of the complex growth rate) is:

$$\sigma_r = \underbrace{2\frac{\mu}{D}H_b^0K^2\cos(2\theta_b^0)}_{e_0} \left(\underbrace{-1}_{e_1} + \underbrace{\frac{\hat{\theta}'_{bi}}{aK}}_{e_2} + \underbrace{\frac{5\hat{H}'_{bi}}{4aKH_b^0}\tan(2\theta_b^0)}_{e_3} - r \underbrace{\frac{\hat{H}'_{br}}{a\beta}\frac{\cos(\theta_b^0)}{\cos(2\theta_b^0)}}_{e_4} \right) \quad (7)$$

A clue to the physical mechanism is provided by a careful analysis of the meaning and behaviour of each term:

- e_0 : common to all terms. It does not contribute to the stability/instability since it is positive. This is because we can assume that $\theta_b^0 < 45^\circ$ due to wave refraction. It is the magnitude of the growth rate.

- e_1 : always negative. It represents the contribution due to the changes in shoreline orientation when there is no perturbation in the wave field. This is the only term arising

in case of the classical analytical one-line modeling (Pelnard-Considère equation) [Dean and Dalrymple, 2002]. It is a damping term describing the shoreline diffusivity in that approach.

- e_2 : its sign depends on $\hat{\theta}'_{bi}$. Numerical computations [Falqués and Calvete, 2005] demonstrates that it is always positive. This results from the fact that refracted wave rays tend to rotate in the same direction as the shoreline. Thus e_2 is a growing term.

- e_3 : its sign depends on $\hat{H}'_{bi}$. Numerical computations [Falqués and Calvete, 2005] show that $\hat{H}'_{bi} > 0$ for long sandwaves and < 0 for short sandwaves. This term is related to energy spreading due to wave crest stretching as waves refract. Thus e_3 is a growing or damping term, depending on the sandwave wavelength, $2\pi/K$. Moreover, its magnitude increases with an increasing incident wave angle. These two properties explain that e_3 is an essential growing term for HAWI formation [Falqués and Calvete, 2005], whereas, for LAWI, its magnitude is smaller and it is, most of the time, negative.

- e_4 : this term stems from the alongshore gradients in H_b in the sediment transport equation (eq. 2). Its sign is the opposite to that of $\hat{H}'_{br}$, which is numerically found to be always positive. This is related to the fact that the maximum in wave energy is always located close to the sandwave crest (wave focusing). Thus e_4 is a damping term.

The corresponding growth rate contributions, $\sigma_1 = e_0 e_1$, $\sigma_2 = e_0 e_2$, ... are plotted in Figure 8. It can be seen that σ_2 is always positive leading to the development of shoreline sandwaves, whereas σ_1 and σ_4 are always negative, leading to the damping of the sandwaves. The term σ_3 can be either positive, for small beach slope (eg smaller than 0.05 to 0.08), or negative, for larger beach slopes. Even if the behaviour of this term is not monotonous, σ_3 generally increases with the wave angle. It is remarkable that σ_2 becomes

Figure 8

331 very large for large beach slopes and for small wave angles. For normal wave incidence
332 ($\theta_b^0 = 0$), $\sigma_3 = 0$ and σ_2 is the only contribution leading to the instability. We therefore
333 conclude that wave refraction is responsible for LAWI in the case of very steep beach
334 slopes.

335 Based on the growth rate results, it is also possible to analyse the relative influence
336 of the wave incidence induced Q_1 and wave set-up induced Q_2 sediment transport. The
337 analysis of the growth rate versus the perturbation wavelength for $r = 0$ or $r = 1$ and
338 several wave incidence angles θ (Figure 9) shows that the relative influence of Q_2 decreases
339 with increasing the wave incidence angle. In other words, wave height gradients (wave set-
340 up induced sediment fluxes) largely influence (damp) the shoreline instability for low wave
341 angle incidence (LAWI), whereas their impact is almost negligible for large wave incidence
342 angles (HAWI). The main driving term of the LAWI is due to the wave incidence induced
343 sediment transport flux Q_1 . This means that the use of the CERC equation alone, taking
344 into account wave refraction in the shoaling zone, can cause LAWI. The Q_2 term influences
345 this instability by changing the growth rate and the favored wavelength. This influence
346 increases with decreasing wave incidence until the case of perfectly shore-normal waves,
347 for which there is a preferred wavelength (LMA mode) only if the Q_2 term is taken into
348 account (Figure 6).

Figure 9

4.2. Model results analysis: Hydrodynamic and sediment transport

349 To understand better how wave refraction causes shoreline instabilities, hydrodynamic
350 and sediment transport model results are analysed next. Here, we focus on the case
351 of shore-normal wave incidence, considering the LMA mode obtained for a beach slope
352 equal to 0.1 (case a). The model results are compared with the same wave incidence, but

a smaller beach slope (0.02) for which the shoreline is stable (case b). The perturbation wavelength (571 m) is the same for both cases, corresponding to the LMA mode of case (a). Although the linear stability analysis is strictly valid only in the limit $a \rightarrow 0$ we choose a shoreline sandwave amplitude $a = 10$ m (for visualization and comparison of the different sources of sediment transport). In addition, to make the analysis simpler, we assume shore-normal waves and $r = 0$. Figures 10a and b show planviews of the wave angle, as well as a longshore cross-section of several quantities along the breaking line. First, it can be noted that the breaking line is much farther offshore in case (b) because of the shallower bathymetry. This is directly linked with the bathymetry. These planviews illustrate the wave focussing on the cusp, which implies increasing wave height and converging waves at the cusps.

Looking at the alongshore cross-section (Figure 10c), the wave angle amplitude is much larger for case (a) than for case (b), about 14° versus 3.2° , indicating a stronger refraction up to the breaking line in case (a). The corresponding amplitude of the oscillation in the shoreline angle is about 6.3° , and is thus within the range of those two values. This means that the angle of the wave fronts with respect to the local shoreline reverses when passing from case (a) to (b), implying a reversal in the direction of sediment transport. This can be traced back to equation 6. The wave angle amplitude is much larger for case (a). Coming back to equation 6, in the case of shore-normal waves and $r = 0$, there are only two terms left: e_1 , which is the contribution due to shoreline change only, and e_2 , which represents the wave refraction-induced sediment flux. The analytical computation for the present case leads to: $e_1 = -1$ for both cases, whereas $e_2 = 2.23$ for case (a) and $e_2 = 0.507$ (case b), consistent with the different amplitudes of alongshore wave angle

Figures 10

376 oscillation. This clearly shows that the growth rate is positive for case (a) and negative
 377 for case (b). The alongshore cross-section of the resulting sediment flux Q (Figure 10c)
 378 illustrates the opposite behaviour for the two cases. Our sign convention is that positive
 379 Q represents sediment transport directed in the direction of the increasing y coordinate
 380 (i.e. to the right on the cross-sections). A positive (negative) longshore gradient indicates
 381 a convergence (divergence), assumed to cause shoreline accretion (erosion). Thus Figure
 382 10c shows that Q for case (a) has a spatial phase-lag compared to the shoreline such that
 383 the shoreline perturbation should be amplified, whereas Q for case (b) has an opposite
 384 phase-lag, leading to the damping of the perturbation. This spatial phase-lag change
 385 results from the continuous amplitude changes of the terms e_1 and e_2 : the phase-lag
 386 between shoreline and longshore sediment fluxes is either 90° or -90° , implying that
 387 there is no migration and either amplification or damping of the shoreline perturbation.
 388 Thus, for shore-normal waves and neglecting the damping term related with wave set-
 389 up induced sediment flux (second term in eq. 2), the instability of the shoreline results
 390 from an alongshore oscillation in the angle of wave refraction, which is stronger than the
 391 oscillation in the angle of shoreline orientation.

4.3. Mechanism

392 From the above, we can draw the following conclusion: the main growing term is re-
 393 lated to the wave refraction toward the cusp, leading to wave incidence induced sediment
 394 transport converging at the cusp. This term strongly increases with beach slope. The
 395 damping is due to three components: (1) longshore sediment transport due to the shore-
 396 line orientation only (and not refraction, term e_1), (2) wave energy spreading (term e_3),

and (3) wave height gradients (set-up) (term e_4), the second component having a smaller damping effect than the two others.

Figure 11

Now we can figure out how LAWI works (Figure 11). Let us consider shore-normal wave incidence. In this case, the wave energy spreading has no influence on the instability ($e_3 = 0$). If a cusped feature with an associated shoal develops on a coastline, wave refraction bends wave rays towards the tip of the feature. Depending on the orientation of the refracted wave fronts with respect to the local shoreline along the cusped feature, the alongshore sediment flux can be directed towards the tip, reinforcing it and leading to a positive feedback between flow and morphology. Whether the transport is directed to the tip, depends on the bathymetry and wave conditions. For a given offshore extent, xl , of the associated shoal and a given wave height, the surf zone will become narrower if the beach slope increases. Then, the shoal will extend a longer distance beyond the surf zone, and the waves will be refracted strongly when they reach the breaking point, increasing the wave incidence related sediment flux (e_2) convergence whereas the divergence term (e_1) is constant. As shown by the model results (Figure 10), the contribution of the refracted wave angle (e_2) can exceed the contribution of the shoreline orientation to the sediment flux (e_1), such that Q_1 (resulting from e_1 and e_2) converges near the cusp. This leads to the development of the cusp. If the beach slope is mild, the surf zone will be wider, and wave refraction over the shoal before breaking will be less intense, leading to smaller wave incidence angle induced sediment fluxes, which are dominated instead by the diverging sediment flux induced by shoreline orientation changes. In this case, as shown in the model results (Figure 10c), the sediment flux is directed away from the tip of the cusp.

5. Discussion

5.1. Linear stability analysis validity

The 1D-morfo model has been applied to investigate HAWI and to study the sandwaves generation along the Dutch coast [Falqués, 2006], El Puntal beach - Spain [Medellín et al., 2008, 2009]. The results indicated similarities with the wavelengths observed in nature. This supports the use of this linear stability model to investigate the mechanisms of shore-line sandwave formation. In the present paper, it is clear that LAWI is a robust output of the 1D-morfo model, and the physical mechanism causing instabilities is wave refraction induced by an offshore shoal associated with a cusped feature. This wave refraction leads to two counter-acting phenomena: sediment transport induced by converging waves counteracted by diverging wave height gradients. The present paper investigates the linear generation only. The pros and cons of linear stability analysis have been discussed extensively in [Blondeaux, 2001; Dodd et al., 2003; Falqués et al., 2008; Tiessen et al., 2010]. In any case, the fundamental assumption of infinitesimal amplitude growth makes comparisons to field data questionable. Nonlinear model studies for other rhythmic features, such as crescentic sandbars and sand ridges [Calvete, 1999; Damgaard et al., 2002], have sometimes shown the finite-amplitude dynamics to be dominated by the LMA mode ; in other cases, modes other than the LMA became dominant. We leave the nonlinear modeling of LAWI, including the study on cessation of the growth, to future work.

5.2. Analogy with megacusps: growth rates and circulation patterns

Although the model results are given for a planar beach, LAWI is also found in the presence of a shore-parallel bar (not shown). Thus, for intermediate morphological beach state where crescentic bars and associated megacusp systems usually develop, the model

440 predicts LAWI. To survive in the finite amplitude domain, the LAWI must grow at a
441 rate comparable to that of co-existing instabilities. Our sensitivity studies indicate LAWI
442 growth rates to range from 10^{-6}s^{-1} (for a beach slope of 0.05) to 10^{-5}s^{-1} (for a beach
443 slope of 0.2). The typical generation time scale thus ranges from 1.5 to 11.5 days. These
444 time scales were obtained for shore-normal waves having a moderate wave height of 1.5
445 m and wave period of 8 s. A typical time scale for the LMA mode of crescentic bars is
446 several days [Damgaard *et al.*, 2002; Garnier *et al.*, 2010]. Thus, for specific beach slope
447 and wave conditions, the LMA shoreline instabilities have comparable initial growth rates
448 as those of crescentic bar patterns.

449 Computations for the idealized cases gives LAWI wavelengths of the same order of
450 magnitude as the observed spacing of crescentic bars and associated megacusps. The
451 distinction between these two kinds of instabilities is therefore difficult and the validation
452 of the presence of LAWI in a Rhythmic Bar and Beach morphological environment is not
453 straightforward. More generally, a proper validation of the present results would need
454 dataset of shoreline evolution, together with bathymetric, wave and current data, starting
455 from an initial longshore uniform beach. To our knowledge, such data do not exist.

456 Although we cannot validate the model results with wavelengths observed in the field,
457 it is possible to discuss whether the type of nearshore circulation linked to LAWI, that is,
458 a longshore sediment flux pointing toward the cusped feature at both sides, is realistic or
459 not in nature and in the framework of 2DH modeling. According to Komar [1998], both
460 types of longshore current patterns, either converging or diverging at a megacusp, are
461 observed in nature. Another example showing that this type of circulation is realistic is
462 the case of a submerged breakwater. Both observations and numerical modelling indicate

463 that if the breakwater is beyond the breaker line, the waves drive longshore currents that
464 converge at the lee of the breakwater to build a salient [*Ranasinghe et al.*, 2006]. This
465 converging type of circulation at a megacusp was also observed by *Haller et al.* [2002] in
466 laboratory experiments on barred beaches with rip channels. One of their six experimental
467 configurations may be quite close to a LAWI configuration. This configuration had the
468 largest average water depth at the bar crest and the smallest rip velocity at the rip neck,
469 such that, in addition to the rip current circulation, they found a secondary circulation
470 system near the shoreline, likely forced by the breaking of the larger waves that propagated
471 through the channel. As these waves are breaking close to the shoreline, they drove
472 longshore currents away from the rip channels into the shallowest area. This experiment
473 shows that breaking close to the shoreline counteracts the rip-induced circulation, leading
474 to current convergence in the shallowest area.

475 The studies of *Calvete et al.* [2005] and *Orzech et al.* [2011] give other elements to
476 investigate the plausibility of the LAWI mechanism, in rip channels configurations. For
477 the case of a barred-beach, *Calvete et al.* [2005] developed a 2DH linear stability model,
478 having a fixed shoreline, that describes the formation of rip channels from an initially
479 straight shore-parallel bar. For shore-normal waves, the circulation linked to rip channel
480 formation is offshore through the channels and onshore over the shoals or horns of the
481 developing crescentic bar as is clearly observed in nature (e.g. [*MacMahan et al.*, 2006]).
482 However, they also noticed small secondary circulation cells near the shoreline flowing in
483 the opposite direction, leading to presence of megacusp formed in front of the horns of
484 the crescentic bar; therefore, the shoreline undulations were out of phase (spatial phase-
485 lag of 180°) with the crescentic bars, meaning that the amplitude of the wave-refracted

486 terms should dominate the amplitude of the wave set-up terms (Eq. 2). The formation of
487 those megacusps was not part of the instability leading to the crescentic bar development
488 but was forced by the hydrodynamics associated with it. More importantly, the small
489 secondary circulation cells were essentially related to wave refraction: if wave refraction
490 from the model was eliminated, they did not develop. Thus, wave refraction by offshore
491 shoals (those of the crescentic bar in this case) can induce a circulation that may move
492 sediment toward a developing cusped feature. A recent study, based on both observation
493 (video images) and non-linear morphodynamic modeling [Orzech *et al.*, 2011] also showed
494 the occurrence of two types of megacusp (shoreward of the shoal or shoreward of the
495 rip), and the associated converging sediment fluxes toward the megacusps. This tends to
496 support our mechanism analysis of LAWI formation.

497 The similarity between LAWI and megacusps in both wavelength and growth time
498 is certainly intriguing given the fact that 1D-morfo is mainly based on the gradients
499 in longshore sediment transport and wave set-up induced sediment transport (damping
500 term), but neglects many surf zone processes like rip current circulation, which is known
501 to be essential to crescentic bar dynamics [Calvete *et al.*, 2005; Garnier *et al.*, 2008].
502 However, the analysis above tends to show that there could be configurations, where the
503 processes taken into account in 1D-morfo are dominant in the system, at the initial stage.
504 This could explain why similarities are observed with the various numerical experiments
505 done with more sophisticated models. Furthermore, we should keep in mind the fact that
506 the LAWI mechanism is not related to any longshore bar, and thus that short shoreline
507 sandwaves such as megacusps could develop without a bar, whereas it was thought that,
508 for barred beaches, short shoreline sandwaves develop due to surf zone sand bar variability.

6. Conclusions

A one-line linear stability model, which was initially created to describe the formation of shoreline sandwaves under high-angle wave incidence, has revealed shoreline instabilities for low to shore-normal wave incidence (LAWI). The most amplified mode has wavelengths of ~ 500 m and characteristic growth time scales of a few days, which are smaller than those of the high angle wave instabilities. Sensitivity analyses focusing on wave height, wave incidence angle, beach slope, beach profile, model free parameters and the sediment transport equation show that, for low to shore-normal wave incidence, instabilities develop for sufficiently large beach slopes (e.g. 0.06) and for sufficiently small wave heights (smaller than 2 m for a beach slope of 0.06).

The main process causing the instabilities for low to shore-normal wave incidence is wave refraction on a shoal in the shoaling zone, which focuses wave fronts onshore of it, leading to wave incidence induced sediment transport converging at the cusp. This effect strongly increases with beach slope. The damping is due to three longshore transport components: (1) that caused by shoreline orientation only (and not refraction), (2) that caused by wave energy spreading (minor effect for low-angle wave incidence), (3) that caused by wave height gradients (set-up). Whether LAWI develops or not depends on the balance between these growing and damping terms. If this shoreline sand accumulation can feed the initial shoal through cross-shore sediment transport, a positive feedback arises.

Acknowledgments. The authors thank the reviewers (including A. Ashton) for their comments and suggestions, D. Calvete for fruitful discussions, J. Thiébot for his comments on this paper. M. Yates-Michelin is also acknowledged for her careful English corrections. Funding from the ANR VMC 2006 - project VULSACO n°ANR-06-VULN-009, the Span-

ish Government (CTM2009-11892/IMNOBE project and TM2006-08875/MAR project),
the Netherlands Organisation for Scientific Research (NWO) project 818.01.009, the Uni-
versity of Nottingham and the Juan de la Cierva program are also acknowledged.

References

- Ashton, A., and A. B. Murray, High-angle wave instability and emergent shoreline
shapes: 1. modeling of sand waves, flying spits, and capes, *J.Geophys.Res.*, *111*,
F04,011,doi:10.1029/2005JF000,422, 2006a.
- Ashton, A., and A. B. Murray, High-angle wave instability and emergent shoreline
shapes: 2. wave climate analysis and comparisons to nature, *J.Geophys.Res.*, *111*,
F04,012,doi:10.1029/2005JF000,423, 2006b.
- Ashton, A., A. B. Murray, and O. Arnault, Formation of coastline features by large-scale
instabilities induced by high-angle waves, *Nature*, *414*, 296–300, 2001.
- Bender, C., and R. Dean, Potential shoreline changes induced by three-dimensional bathy-
metric anomalies with gradual transitions in depth, *Coast. Eng.*, *51*, 1143–1161, 2004.
- Blondeaux, P., Mechanics of coastal forms, *Ann. Rev. Fluid Mech.*, *33*, 339–370, 2001.
- Bruun, P., Migrating sand waves or sand humps, with special reference to investigations
carried out on the Danish North Sea Coast, in *Coastal Eng. 1954*, pp. 269–295, Am.
Soc. of Civ. Eng., 1954.
- Calvete, D., Morphological stability models: Shoreface-connected sand ridges, Ph.D. the-
sis, Appl. Physics Dept., Univ. Politècnica de Catalunya, Barcelona, Spain, 1999.
- Calvete, D., N. Dodd, A. Falqués, and S. M. van Leeuwen, Morphological development
of rip channel systems: Normal and near normal wave incidence, *J. Geophys. Res.*,

110(C10006), doi:10.1029/2004JC002803, 2005.

Castelle, B., P. Bonneton, H. Dupuis, and N. Snchal, Double bar beach dynamics on the high-energy meso-macrotidal french aquitanian coast : a review, *Mar. Geol.*, *245*, 141–159, 2007.

Damgaard, J., N. Dodd, L. Hall, and T. Chesher, Morphodynamic modelling of rip channel growth, *Coastal Eng.*, *45*, 199–221, 2002.

Davidson-Arnott, R. G. D., and A. van Heyningen, Migration and sedimentology of long-shore sandwaves, Long Point, Lake Erie, Canada, *Sedimentology*, *50*, 1123–1137, 2003.

Dean, R. G., and R. A. Dalrymple, *Coastal Processes*, Cambridge University Press, Cambridge, 2002.

Dodd, N., P. Blondeaux, D. Calvete, H. E. de Swart, A. Falqus, S. J. M. H. Hulscher, G. Rzynski, and G. Vittori, The use of stability methods in understanding the morphodynamical behavior of coastal systems, *J. Coastal Res.*, *19*(4), 849–865, 2003.

Falqus, A., Wave driven alongshore sediment transport and stability of the Dutch coastline, *Coastal Eng.*, *53*, 243–254, 2006.

Falqus, A., and D. Calvete, Large scale dynamics of sandy coastlines. Diffusivity and instability, *J. Geophys. Res.*, *110*(C03007), doi:10.1029/2004JC002587, 2005.

Falqus, A., G. Coco, and D. A. Huntley, A mechanism for the generation of wave-driven rhythmic patterns in the surf zone, *J. Geophys. Res.*, *105*(C10), 24,071–24,088, 2000.

Falqus, A., N. Dodd, R. Garnier, F. Ribas, L. MacHardy, P. Larroud, D. Calvete, and F. Sancho, Rhythmic surf zone bars and morphodynamic self-organization, *Coastal Eng.*, *55*, 622–641, doi:10.1016/j.coastaleng.2007.11.012, 2008.

574 Garnier, R., D. Calvete, A. Falqués, and M. Caballeria, Generation and nonlinear evolu-
575 tion of shore-oblique/transverse sand bars, *J. Fluid Mech.*, 567, 327–360, 2006.

576 Garnier, R., D. Calvete, A. Falqués, and N. Dodd, Modelling the formation and the
577 long-term behavior of rip channel systems from the deformation of a longshore bar, *J.*
578 *Geophys. Res.*, 113(C07053), doi:10.1029/2007JC004632, 2008.

579 Garnier, R., N. Dodd, A. Falqués, and D. Calvete, Mechanisms controlling crescentic bar
580 amplitude, *J. Geophys. Res.*, 115, doi:10.1029/2009JF001407, 2010.

581 Haller, M. C., R. A. Dalrymple, and I. A. Svendsen, Experimental study of
582 nearshore dynamics on a barred beach with rip channels, *J. Geophys. Res.*, 107(C6),
583 10.1029/2001JC000,955, 2002.

584 Horikawa, K., *Nearshore Dynamics and Coastal Processes*, University of Tokio Press,
585 Tokio, Japan, 1988.

586 Inman, D. L., M. H. S. Elwany, A. A. Khafagy, and A. Golik, Nile delta profiles and
587 migrating sand blankets, in *Coastal Eng. 1992*, pp. 3273–3284, Am. Soc. of Civ. Eng.,
588 1992.

589 Komar, P. D., *Beach Processes and Sedimentation*, second ed., Prentice Hall, Englewood
590 Cliffs, N.J., 1998.

591 Lafon, V., D. D. M. Apoluceno, H. Dupuis, D. Michel, H. Howa, and J. M. Froidefond,
592 Morphodynamics of nearshore rhythmic sandbars in a mixed-energy environment (SW
593 France): I. Mapping beach changes using visible satellite imagery, *Estuarine, Coastal*
594 *and Shelf Science*, 61, 289–299, 2004.

595 List, J. H., and A. D. Ashton, A circulation modeling approach for evaluating the condi-
596 tions for shoreline instabilities, in *Coastal Sediments 2007*, pp. 327–340, ASCE, 2007.

597 List, J. H., D. M. Hanes, and P. Ruggiero, Predicting longshore gradients in alongshore
598 transport: comparing the cerc formula to delft3d, in *Coastal Eng. 2006*, pp. 3370–3380,
599 World Scientific, 2006.

600 List, J. H., L. Benedet, D. M. Hanes, and P. Ruggiero, Understanding differences between
601 delft3d and emperical predictions of alongshore sediment transport gradients, in *Coastal*
602 *Eng. 2008*, pp. 1864–1875, World Scientific, 2008.

603 MacMahan, J. H., E. B. Thornton, and A. J. H. M. Reniers, Rip current review, *Coastal*
604 *Eng.*, *53*, 191–208, 2006.

605 Medellín, G., R. Medina, A. Falqués, and M. González, Coastline sand waves on a low
606 energy beach at 'El Puntal' spit, Spain, *Mar. Geol.*, *250*, 143–156, 2008.

607 Medellín, G., A. Falqués, R. Medina, and M. González, Sand waves on a low-energy beach
608 at el 'puntal' spit, Spain: Linear Stability Analysis, *J. Geophys. Res.*, *114*(C03022),
609 doi:10.1029/2007JC004426, 2009.

610 Orzech, M., A. Reniers, E. Thornton, and J. MacMahan, Megacusps on
611 rip channel bathymetry: Observations and modeling, *Coastal Eng.*, p.
612 doi:10.1016/j.coastaleng.2011.05.001, 2011.

613 Ozasa, H., and A. H. Brampton, Mathematical modelling of beaches backed by seawalls,
614 *Coastal Eng.*, *4*, 47–63, 1980.

615 Ranasinghe, R., G. Symonds, K. Black, and R. Holman, Morphodynamics of intermediate
616 beaches: A video imaging and numerical modelling study, *Coastal Eng.*, *51*, 629–655,
617 2004.

618 Ranasinghe, R., I. L. Turner, and G. Symonds, Shoreline response to multi-functional
619 artificial surfing reefs: A numerical and physical modelling study, *Coastal Eng.*, *53*,

589–611, 2006.

Reniers, A. J. H. M., J. A. Roelvink, and E. B. Thornton, Morphodynamic modeling of an embayed beach under wave group forcing, *J. Geophys. Res.*, *109*(C01030), doi:10.1029/2002JC001586, 2004.

Ribas, F., and A. Kroon, Characteristics and dynamics of surfzone transverse finger bars, *J. Geophys. Res.*, *112*(F03028), doi:10.1029/2006JF000685, 2007.

Ruessink, B. G., and M. C. J. L. Jeuken, Dunefoot dynamics along the dutch coast, *Earth Surface Processes and Landforms*, *27*, 1043–1056, 2002.

Sonu, C. J., Collective movement of sediment in littoral environment, in *Coastal Eng. 1968*, pp. 373–400, Am. Soc. of Civ. Eng., 1968.

Sonu, C. J., Three-dimensional beach changes, *J. Geology*, *81*, 42–64, 1973.

Stewart, C. J., and R. G. D. Davidson-Arnott, Morphology, formation and migration of longshore sandwaves; long point, lake erie, canada, *Mar. Geol.*, *81*, 63–77, 1988.

Thevenot, M. M., and N. C. Kraus, Longshore sandwaves at Southampton Beach, New York: observations and numerical simulation of their movement, *Mar. Geology*, *126*, 249–269, 1995.

Tiessen, M. C. H., S. M. V. Leeuwen, D. Calvete, and N. Dodd, A field test of a linear stability model for crescentic bars, *Coastal Engineering*, *57*, 41–51, 2010.

USACE, Shore protection manual, *Tech. Rep. 4th ed., 2 Vol.*, USACE, U.S. Government Printing Office, Washington, D.C., 1984.

van den Berg, N., A. Falqués, and R. Ribas, Long-term evolution of nourished beaches under high angle wave conditions, *Journal of Marine Systems*, *to appear*, 2011.

Verhagen, H. J., Sand waves along the dutch coast, *Coastal Eng.*, *13*, 129–147, 1989.

643 Wright, L. D., and A. D. Short, Morphodynamic variability of surf zones and beaches: A
644 synthesis, *Mar. Geol.*, 56, 93–118, 1984.

Figure 1. Sketch of the geometry and the variables. The angle between the wave fronts and the local shoreline is $\alpha = \theta - \phi$.

Figure 2. Bathymetric perturbations (a) and cross-shore Dean beach profile for a various beach slopes (b,c). P1-perturbation: constant in the surf zone and exponential decrease, P2-perturbation: exponential decrease, P3-perturbation: exponential decrease with a shoal in the shoaling zone, P4-perturbation: exponential decrease with a shoal in the surf and shoaling zones.

Figure 3. LMA (a) growth rate and (b) wavelength as a function of beach slope and angle of incidence. Circles indicate the 1D-morfo computations. In white: no growing perturbation. The model parameters are : $\mu = 0.15$, $r = 1$, P1-Perturbation and $xl=1410$ m for Dean profiles with $H_{rms} = 1.5$ m and $T_p = 8$ s at a water depth of 25 m.

Figure 4. LMA (a) growth rate and (b) wavelength as a function of beach slope and wave height, for shore-normal waves. Circles indicate the 1D-morfo computations. In white: no growing perturbation. The model parameters are : $\mu = 0.15$, $r = 1$, P1-Perturbation and $xl=1410$ m for Dean profiles with $\theta = 0^\circ$ and $T_p = 8$ s at a water depth of 25 m.

Figure 5. LMA mode (a) growth rate and (b) wavelength as a function of the shoreline perturbation length xl for a Dean profile with a beach slope of 0.1. Crosses on the xl axis ($10 \leq xl \leq 100$) indicate that there is no LMA mode for the given shoreline perturbation length.

Figure 6. LMA (a) growth rate and (b) wavelength as function of beach slope and wave angle, for $r = 0$. Circles indicate the 1D-morfo computations. In white: no LMA mode. The symbol x indicates that, even if the growth rate was positive, there was a singularity at $\theta = 0^\circ$ (growth rate continuously increasing with decreasing wavelength), and therefore no LMA mode (see Figure 7).

Figure 7. LMA growth rate versus wavelength for various beach slopes, using $r = 0$ and shore-normal waves. The shoreline is stable for a beach slope $\beta < 0.05$. The model parameters are : $\mu = 0.15$, P1-Perturbation and $xl=1410$ m for the Dean profile with $H_{rms} = 1.5$ m and $T_p = 8$ s at a water depth of 25 m.

Figure 8. Growth rate components for the LMA mode as a function of beach slope and angle of incidence. Model parameters are: $H_{rms} = 1.5$ m, $T_p = 8$ s and cross-shore perturbation of type P1 with $x_l = 1410$ m. (a) σ_1 , (b) σ_2 , (c) σ_3 , (d) σ_4 . Circles indicate the 1D-morfo computations. In white: no LMA mode.

Figure 9. Growth rate versus wavelength for several combinations of r and wave incidence angle θ . The beach slope is 0.05.

Figure 10. Model results for a perturbation wavelength of 571 m (LMA mode for a beach slope of 0.1), shore-normal wave incidence, and a shoreline wave amplitude of $a = 10$ m. (a) and (b) show the topographic contours and the refracted wave angle for beach slopes of 0.1 (a) and 0.02 (b). (c) shows the longshore profiles of (Top) the shoreline position, (Middle) the shoreline angle (solid line) and refracted wave angles and (Bottom) the sediment fluxes. Dashed-dotted lines, and dotted lines represent a beach slope of 0.1 and 0.02, respectively.

Figure 11. Sketch of the physical mechanisms causing LAWI. Sediment transport components induced by a shoal for unstable (a) and stable (b) situations, corresponding to a narrow and wide surf zone, respectively.