



HAL
open science

The closure of a sheet is not always a union of sheets

Michael Bulois

► **To cite this version:**

| Michael Bulois. The closure of a sheet is not always a union of sheets. 2010. hal-00660584v1

HAL Id: hal-00660584

<https://hal.science/hal-00660584v1>

Preprint submitted on 17 Jan 2012 (v1), last revised 29 Mar 2019 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The closure of a sheet is not always a union of sheets, a short note

MICHAËL BULOIS *

Abstract

In this note we answer to a frequently asked question. If G is an algebraic group acting on a variety V , a G -sheet of V is an irreducible component of $V^{(m)}$, the set of elements of V whose G -orbit has dimension m . We focus on the case of the adjoint action of a semisimple group on its Lie algebra. We give two families of examples of sheets whose closure is not a union of sheets in this setting.

Let \mathfrak{g} be a semisimple Lie algebra defined over an algebraically closed field \mathbb{k} of characteristic zero. Let G be the adjoint group of \mathfrak{g} . For any integer m , one defines

$$\mathfrak{g}^{(m)} = \{x \in \mathfrak{g} \mid \dim G.x = m\}.$$

A G -sheet (or simply *sheet*) is an irreducible component of $\mathfrak{g}^{(m)}$ for some $m \in \mathbb{N}$. We refer to [TY, §39] for elementary properties of sheets. The most important one is that each sheet contains a unique nilpotent orbit.

There exists a well known subdivision of sheets which forms a stratification. The objects considered in this subdivision are Jordan classes and generalize the classical Jordan's block decomposition in \mathfrak{gl}_n . Those classes and their closures are widely studied in [Bo] (cf. also [TY, §39] for a more elementary viewpoint). Since sheets are locally closed, a natural question is then the following.

If S is a sheet, is \overline{S} is a union of sheets?

The answer is negative in general. We give two families of counterexamples below.

1. A nilpotent orbit \mathcal{O} of \mathfrak{g} is said to be rigid if it is a sheet of \mathfrak{g} . Rigid orbits are key objects in the description of sheets given in [Bo]. They were classified for the first time in [Sp, §II.7&II.10]. The closure ordering of nilpotent orbits (or *Hasse diagram*) can be found in [Sp, §II.8&IV.2]. In the classical cases, a more recent reference for these lists is [CM]. One easily checks from these classifications that there may exists some rigid nilpotent orbit \mathcal{O}_1 that contains a non-rigid nilpotent orbit \mathcal{O}_2 in its closure. Then, we set $S = \mathcal{O}_1$ and we get $\mathcal{O}_2 \subset \overline{S} \subset \mathcal{N}(\mathfrak{g})$ where $\mathcal{N}(\mathfrak{g})$ is the set of nilpotent elements of \mathfrak{g} . Since \mathcal{O}_2 is not rigid, the sheets containing

*michael.bulois@univ-angers.fr

Université d'Angers, Laboratoire de Mathématiques d'Angers, UMR 6093, 2 boulevard Lavoisier 49045 Angers CEDEX 1, France

\mathcal{O}_2 are not wholly included in $\mathcal{N}(\mathfrak{g})$. Therefore, the closure of S is not a union of sheets.

Here are some examples of such nilpotent orbits. In the classical cases, we embed \mathfrak{g} in \mathfrak{gl}_n in the natural way. Then, we can assign to each nilpotent orbit \mathcal{O} , a partition of n , denoted by $\Gamma(\mathcal{O})$. This partition defines the orbit \mathcal{O} , sometimes up to an element of $\text{Aut}(\mathfrak{g})$. In the case $\mathfrak{g} = \mathfrak{so}_8$ (type D_4), there is exactly one rigid orbit \mathcal{O}_1 , such that $\Gamma(\mathcal{O}_1) = [3, 2^2, 1]$. It contains in its closure the non-rigid orbit \mathcal{O}_2 such that $\Gamma(\mathcal{O}_2) = [3, 1^5]$ (cf. [Mo, Table2, p.15]). Very similar examples can be found in types C and B.

In the exceptional cases, we denote nilpotent orbits by their Bala-Carter symbol as in [Sp]. Let us give some examples of the above described phenomenon.

- in type E_6 ($\mathcal{O}_1 = 3A_1$ and $\mathcal{O}_2 = 2A_1$),
 - in type E_7 ($\mathcal{O}_1 = A_2 + 2A_1$ and $\mathcal{O}_2 = A_2 + A_1$),
 - in type E_8 ($\mathcal{O}_1 = A_2 + A_1$ and $\mathcal{O}_2 = A_2$)
 - and in type F_4 ($\mathcal{O}_1 = A_2 + A_1$ and A_2).
2. In the case $\mathfrak{g} = \mathfrak{sl}_n$ of type A , there is only one rigid nilpotent orbit, the null one. Hence the phenomenon depicted in 1 can not arise in this case. Let S be a sheet and let $\lambda_S = (\lambda_1 \geq \dots \geq \lambda_{k(\lambda_S)})$ be the partition of n associated to the nilpotent orbit \mathcal{O}_S of S . As a consequence of the theory of induction of orbits, cf. [Bo], we have

$$\overline{S} = \overline{G \cdot \mathfrak{h}_S}^{reg} \quad (1)$$

where \mathfrak{h}_S is the centre of a Levi subalgebra \mathfrak{l}_S . The size of the blocks of \mathfrak{l}_S yield a partition of n , which we denote by $\tilde{\lambda}_S = (\tilde{\lambda}_1 \geq \dots \geq \tilde{\lambda}_{p(\lambda_S)})$. In fact $\tilde{\lambda}$ is the dual partition of λ , i.e. $\tilde{\lambda}_i = \#\{j \mid \lambda_j \geq i\}$ (see, e.g., [Kr, §2.2]). In particular, the map sending a sheet S to its nilpotent orbit \mathcal{O}_S is a bijection.

An easy consequence of (1) is the following (see [Kr, Satz 1.4]). Given any two sheets S and S' of \mathfrak{g} , we have $S \subset \overline{S'}$ if and only if \mathfrak{h}_S is G -conjugate to a subspace of $\mathfrak{h}_{S'}$ or, equivalently, $\mathfrak{l}_{S'}$ is conjugate to a subspace of \mathfrak{l}_S . This can be translated in terms of partitions by defining a partial ordering on the set of partitions of n as follows. We say that $\lambda \preceq \lambda'$ if there exists a partition $(J_i)_{i \in [1, p(\lambda)]}$ of $[1, p(\lambda)]$ such that $\tilde{\lambda}_i = \sum_{j \in J_i} \tilde{\lambda}'_j$. Hence, we have the following characterization.

Lemma 1. $S \subset \overline{S'}$ if and only if $\lambda_S \preceq \lambda_{S'}$.

One sees that this criterion is strictly stronger than the characterization of inclusion relations of closures of nilpotent orbits (see, e.g., [CM, §6.2]). More precisely, one easily finds two sheets S and S' such that $\mathcal{O}_S \subset \overline{\mathcal{O}_{S'}}$ while $\lambda_S \not\preceq \lambda_{S'}$. Then, $\mathcal{O}_S \subset \overline{S'}$, S is the only sheet containing \mathcal{O}_S and $S \not\subset \overline{S'}$. For instance, take $\lambda_{S'} = [3, 2]$, $\lambda_S = [3, 1, 1]$. Their respective dual partitions being $[2, 2, 1]$ and $[3, 1, 1]$, we have $\lambda_S \not\preceq \lambda_{S'}$.

References

- [Bo] W. Borho, Über schichten halbeinfacher Lie-algebren, *Invent. Math.*, **65** (1981), 283-317.
- [CM] D. H. Collingwood and W. M. McGovern, *Nilpotent orbits in semisimple Lie algebras*, Van Nostrand Reinhold Mathematics Series, New York, (1993).
- [Kr] H. Kraft, Parametrisierung von Konjugationklassen in \mathfrak{sl}_n , *Math. Ann.*, **234** (1978), 209-220.
- [Mo] A. Moreau, On the dimension of the sheets of a reductive Lie algebra, *J. Lie Theory*, **18** (2008), 671-696.
- [Sp] N. Spaltenstein, Classes unipotentes et sous-groupes de Borel, *Lecture Notes in Math.* **946**, Springer-Verlag, New York, 1982.
- [TY] P. Tauvel and R. W. T. Yu, *Lie algebras and algebraic groups*, Springer Monographs in Mathematics, Springer-Verlag, 2005.