



HAL
open science

Eigenvarieties for classical groups and complex conjugations in Galois representations

Olivier Taïbi

► **To cite this version:**

Olivier Taïbi. Eigenvarieties for classical groups and complex conjugations in Galois representations. 2012. hal-00675682

HAL Id: hal-00675682

<https://hal.science/hal-00675682>

Preprint submitted on 1 Mar 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Eigenvarieties for classical groups and complex conjugations in Galois representations

Olivier Taïbi

Abstract

The goal of this paper is to remove the irreducibility hypothesis in a theorem of Richard Taylor describing the image of complex conjugations by p -adic Galois representations associated with regular, algebraic, essentially self-dual, cuspidal automorphic representations of GL_{2n+1} over a totally real number field F . We also extend it to the case of representations of GL_{2n}/F whose multiplicative character is “odd”. We use a p -adic deformation argument, more precisely we prove that on the eigenvarieties for symplectic and even orthogonal groups, there are “many” points corresponding to (quasi-)irreducible Galois representations. The recent work of James Arthur describing the automorphic spectrum for these groups is used to define these Galois representations, and also to transfer self-dual automorphic representations of the general linear group to these classical groups.

Contents

1	Introduction	3
2	Assumptions on the forthcoming chapter of Arthur’s book	6
3	The eigenvariety for definite symplectic groups	7
3.1	The eigenvariety	7
3.1.1	Symplectic groups compact at the archimedean places	7
3.1.2	The Atkin-Lehner algebra	7
3.1.3	p -adic automorphic forms	8
3.1.4	Existence and properties of the eigenvariety	9
3.2	Unramified and “completely refinable” points	11
3.2.1	Small slope p -adic eigenforms are classical	11
3.2.2	Representations having Iwahori-invariants and unramified principal series	13
3.2.3	Most points of the eigenvariety arise from unramified, completely refinable representations	
4	Galois representations associated with automorphic representations of symplectic groups	
4.1	A consequence of Arthur’s description of the discrete spectrum for classical groups	15
4.1.1	Automorphic self-dual representations of GL_{2n+1} of orthogonal type	15
4.1.2	p -adic Galois representations associated with RLASDC representations of GL_N	17
4.1.3	The Galois pseudocharacter on the eigenvariety	18
4.2	Galois representations stemming from symplectic forms are generically almost irreducible	19
4.2.1	Crystalline representations over $\overline{\mathbb{Q}}_p$	19

4.2.2	Variation of the crystalline Frobenius on the eigenvariety	20
4.2.3	Main result	21
5	Similar results for even orthogonal groups	24
6	The image of complex conjugation: relaxing hypotheses in Taylor's theorem	25
6.1	Regular, L-algebraic, self-dual, cuspidal representations of $\mathrm{GL}_{2n}(\mathbb{A}_F)$ having Iwahori-invariants	26
6.2	Regular, L-algebraic, self-dual, cuspidal representations of $\mathrm{GL}_{2n+1}(\mathbb{A}_F)$ having Iwahori-invariants	27
6.3	Almost general case	33
6.4	A supplementary, non-regular case	35

1 Introduction

Let p be a prime. Let us choose once and for all algebraic closures $\overline{\mathbb{Q}}, \overline{\mathbb{Q}}_p, \mathbb{C}$ and embeddings $\iota_p : \overline{\mathbb{Q}} \hookrightarrow \overline{\mathbb{Q}}_p, \iota_\infty : \overline{\mathbb{Q}} \hookrightarrow \mathbb{C}$. Let F be a totally real number field. A regular, L-algebraic, essentially self-dual, cuspidal (RLAESDC) representation of $\mathrm{GL}_n(\mathbb{A}_F)$ is a cuspidal automorphic representation π together with an algebraic character $\eta|\cdot|^q$ of $\mathbb{A}_F^\times/F^\times$ (η being an Artin character, and q an integer) such that

- $\pi^\vee \simeq \eta|\det|^q \otimes \pi$,
- $\eta_v(-1)$ does not depend on the real place v of F (the common value will be denoted $\eta_\infty(-1)$),
- For any real place v of F , $\mathcal{LL}(\pi_v)|_{W_{\mathbb{C}}} \simeq \bigoplus_i (z \mapsto z^{a_{v,i}} \bar{z}^{b_{v,i}})$ where \mathcal{LL} is the local Langlands correspondence, $W_{\mathbb{C}} \simeq \mathbb{C}^\times$ is the Weil group of \mathbb{C} , and $a_{v,i}, b_{v,i}$ are integers and $a_{v,i} \neq a_{v,j}$ if $i \neq j$.

By definition, π is regular, L-algebraic, essentially self-dual, cuspidal (RLAESDC) if and only if $\pi \otimes |\det|^{(n-1)/2}$ is regular, *algebraic* (in the sense of Clozel), essentially self-dual, cuspidal (RAESDC). The latter is the notion of “algebraic” usually found in the literature, and is called “C-algebraic” in [BG10]. Given a RLAESDC representation π of $\mathrm{GL}_n(\mathbb{A}_F)$, there is (Theorem 4.1.2) a unique continuous, semisimple Galois representation $\rho_{\iota_p, \iota_\infty}(\pi) : G_F \rightarrow \mathrm{GL}_n(\overline{\mathbb{Q}}_p)$ such that $\rho_{\iota_p, \iota_\infty}(\pi)$ is unramified at any finite place v of F not lying above p for which π_v is unramified, and $\iota_\infty \iota_p^{-1} \mathrm{Tr}(\rho_{\iota_p, \iota_\infty}(\pi)(\mathrm{Frob}_v))$ is equal to the trace of the Satake parameter of π_v (contained in this assertion is the fact that this trace is algebraic over \mathbb{Q}). It is conjectured that for any real place v of F , if $c_v \in G_F$ is the conjugacy class of complex conjugations associated with v , the conjugacy class of $\rho_{\iota_p, \iota_\infty}(\pi)(c_v)$ is determined by $\mathcal{LL}(\pi_v)$ (see [BG10][Lemma 2.3.2] for the case of an arbitrary reductive group). In the present case, by Clozel’s purity lemma and by regularity, $\mathcal{LL}(\pi_v)$ is completely determined by its restriction to $W_{\mathbb{C}}$, and since $\det(\rho_{\iota_p, \iota_\infty}(\pi))$ is known, the determination of $\rho_{\iota_p, \iota_\infty}(\pi)(c_v)$ amounts to the following

Conjecture. *Under the above hypotheses, $|\mathrm{Tr}(\rho_{\iota_p, \iota_\infty}(\pi)(c_v))| \leq 1$.*

There are several cases for which this is known. When $\eta_\infty(-1)(-1)^q = -1$ (this happens only if n is even, and by [BC11] this means that $\rho_{\iota_p, \iota_\infty}(\pi)$ together with the character $\rho_{\iota_p, \iota_\infty}(\eta|\cdot|^q) = (\eta \circ \mathrm{rec})\mathrm{cyclo}^q$, is “symplectic”), $\rho_{\iota_p, \iota_\infty}(\pi)(c_v)$ is conjugate to $-\rho_{\iota_p, \iota_\infty}(\pi)(c_v)$, so the trace is obviously zero.

In [Tay], Richard Taylor proves the following

Theorem (Taylor). *Let F be a totally real number field, $n \geq 1$ an integer. Let π be a regular, L-algebraic, essentially self-dual, cuspidal automorphic representation of GL_{2n+1}/F . Assume that the attached Galois representation $\rho_{\iota_p, \iota_\infty}(\pi) : G_F \rightarrow \mathrm{GL}_{2n+1}(\overline{\mathbb{Q}}_p)$ is irreducible. Then for any real place v of F ,*

$$\mathrm{Tr}(\rho_{\iota_p, \iota_\infty}(\pi)(c_v)) = \pm 1.$$

Although one expects $\rho_{\iota_p, \iota_\infty}(\pi)$ to be always irreducible, this is not known in general. However it is known when $n \leq 5$ by [CG], and for a set of primes p of density one when n is arbitrary but the weights are “extremely regular” by [BLGGT, Theorem D].

In this paper, the following cases are proved:

Theorem A (Theorem 6.3.4). *Let $n \geq 2$, F a totally real number field, π a regular, L -algebraic, essentially self-dual, cuspidal representation of $\mathrm{GL}_n(\mathbb{A}_F)$, such that $\pi^\vee \simeq ((\eta) \cdot |^q) \circ \det \otimes \pi$, where η is an Artin character and q an integer. Suppose that one of the following conditions holds*

1. n is odd.
2. n is even, q is even, and $\eta_\infty(-1) = 1$.

Then for any complex conjugation $c \in G_F$, $|\mathrm{Tr}(\rho_{\iota_p, \iota_\infty}(\pi)(c))| \leq 1$.

This is achieved thanks to the result of Taylor, Arthur’s endoscopic transfer between twisted general linear groups and symplectic or orthogonal groups, and using eigenvarieties for these groups. Let us describe the natural strategy that one might consider to prove the odd-dimensional case using these tools, to explain why it fails and how a detour through the even-dimensional case allows to conclude.

Let π be a RLAESDC representation of $\mathrm{GL}_{2n+1}(\mathbb{A}_F)$. Up to a twist by an algebraic character π is self-dual and has trivial central character. Conjecturally, there should be an associated self-dual Langlands parameter $\phi_\pi : L_F \rightarrow \mathrm{GL}_{2n+1}(\mathbb{C})$ where L_F is the conjectural Langlands group. Up to conjugation, ϕ_π takes values in $\mathrm{SO}_{2n+1}(\mathbb{C})$, and by functoriality there should be a discrete automorphic representation Π of $\mathrm{Sp}_{2n}(\mathbb{A}_F)$ such that $\mathcal{LL}(\Pi_v)$ is equal to $\mathcal{LL}(\pi_v)$ via the inclusion $\mathrm{SO}_{2n+1}(\mathbb{C})$ for any place of F which is either archimedean or such that π_v is unramified. Arthur’s results in his upcoming book [Art11] imply that this (in fact, much more) holds. To construct p -adic families of automorphic representations (i.e. eigenvarieties) containing Π , it is preferable to work with a group which is *compact* at the real places of F , and work with representations having Iwahori-invariants at the p -adic places. A suitable solvable base change allows to assume that $[F : \mathbb{Q}]$ is even and that π_v has Iwahori-invariants for $v|p$. The last chapter of [Art11] will allow to “transfer” π to an automorphic representation Π of \mathbf{G} , the inner form of Sp_{2n} which is split at the finite places and compact at the real places of F . By [Loe11] (which generalizes [Che04]), the eigenvariety \mathcal{X} for \mathbf{G} is available. Thanks to [Art11], one can associate p -adic Galois representations $\rho_{\iota_p, \iota_\infty}(\cdot)$ to automorphic representations of \mathbf{G} , yielding a family of Galois representations on \mathcal{X} , that is to say a continuous map $T : G_F \rightarrow \mathcal{O}(\mathcal{X})$ which specializes to $\mathrm{Tr}(\rho_{\iota_p, \iota_\infty}(\cdot))$ at the points of \mathcal{X} corresponding to automorphic representations of $\mathbf{G}(\mathbb{A}_F)$. One can then hope to prove a result similar to [BC11, Lemma 3.3], i.e. show that one can “deform” Π (on \mathcal{X}) to reach a point corresponding to an automorphic representation Π' whose Galois representation is irreducible (even when restricted to the decomposition group of a p -adic place of F). Since $\rho_{\iota_p, \iota_\infty}(\Pi')$ comes from an automorphic representation π' of GL_{2n+1} , π' is necessarily cuspidal and satisfies

the hypotheses of Taylor’s theorem. Since $T(c_v)$ is locally constant on \mathcal{X} , we would be done.

Unfortunately, it does not appear to be possible to reach a representation Π' whose Galois representation is irreducible by using local arguments on the eigenvariety. However we will prove the following, which includes the case of some even-dimensional special orthogonal groups as it will be needed later:

Theorem B (Theorem 4.2.2, Theorem 5.0.3). *Let \mathbf{G} be an inner form of Sp_{2n} or SO_{4n} over a totally real number field, compact at the real places and split at the p -adic ones. Let Π be an irreducible automorphic representation of $\mathbf{G}(\mathbb{A}_F)$ having Iwahori invariants at all the places of F above p , and having invariants under an open subgroup U of $\mathbf{G}(\mathbb{A}_{F,f}^{(p)})$. Let $\rho_{\iota_p, \iota_\infty}(\Pi)$ denote the p -adic representation of the absolute Galois group G_F of F associated with Π and embeddings $\iota_p : \overline{\mathbb{Q}} \hookrightarrow \overline{\mathbb{Q}}_p$, $\iota_\infty : \overline{\mathbb{Q}} \hookrightarrow \mathbb{C}$. Let N be an integer. There exists an automorphic representation Π' of $\mathbf{G}(\mathbb{A}_F)$ such that:*

- Π' is unramified at the places above p , and has invariants under U ;
- The restriction of $\rho_{\iota_p, \iota_\infty}(\Pi')$ to the decomposition group at any place above p is either irreducible or the sum of an Artin character and an irreducible representation of dimension $2n$ (the latter occurring only in the symplectic case);
- For all g in G_F , $\mathrm{Tr}(\rho_{\iota_p, \iota_\infty}(\Pi')(g)) \equiv \mathrm{Tr}(\rho_{\iota_p, \iota_\infty}(\Pi)(g)) \pmod{p^N}$.

The possible presence of an Artin character (in the case of inner forms of Sp_{2n}) comes from the fact that the “standard” representation of $\mathrm{SO}_{2n+1}(\mathbb{C})$ in $\mathrm{GL}_{2n+1}(\mathbb{C})$ is not minuscule: the set of characters of a torus $T(\mathbb{C})$ of $\mathrm{SO}_{2n+1}(\mathbb{C})$ in this representation has two orbits under the Weyl group, one of which contains only the trivial character. The key fact allowing to prove the above theorem is that classical points on the eigenvariety for \mathbf{G} correspond to automorphic representations Π of $\mathbf{G}(\mathbb{A}_F)$ (say, unramified at the p -adic places) and a refinement of each Π_v , $v|p$, that is to say a particular element in $T(\mathbb{C})$ in the conjugacy class of the Satake parameter of Π_v . The variation of the crystalline Frobenius of $\rho_{\iota_p, \iota_\infty}(\cdot)$ on the eigenvariety with respect to the weight and the freedom to change the refinement (by the action of the Weyl group) are at the heart of the proof of Theorem B.

Although the strategy outlined above fails, Theorem A can be deduced from Theorem B. Indeed the precise description of the discrete automorphic spectrum of symplectic and orthogonal groups by Arthur shows that formal sums of distinct cuspidal self-dual representations of general linear groups “contribute” to this spectrum. The even-dimensional case in Theorem A will be proved by transferring $\pi \boxplus \pi_0$, where π, π_0 are regular, L -algebraic, self-dual, cuspidal representations of $\mathrm{GL}_{2n}(\mathbb{A}_F)$ (resp. $\mathrm{GL}_3(\mathbb{A}_F)$) with distinct weights at any real place of F , to an automorphic representation Π of an inner form \mathbf{G} of Sp_{2n+2}/F . Since $\rho_{\iota_p, \iota_\infty}(\pi) \oplus \rho_{\iota_p, \iota_\infty}(\pi_0)$ does not contain any Artin character (the zero Hodge-Tate weights come from $\rho_{\iota_p, \iota_\infty}(\pi_0)$, which is known to be irreducible), for big enough N any representation Π' as in B has an irreducible Galois representation.

To treat the original case of a regular, L -algebraic, self-dual, cuspidal representation of $\mathrm{GL}_{2n+1}(\mathbb{A}_F)$ having trivial central character, we appeal to Theorem B for special orthog-

onal groups. For example, if n is odd, $\pi \boxplus \pi_0$, where π_0 is the trivial character of $\mathbb{A}_F^\times/F^\times$, contributes to the automorphic spectrum of \mathbf{G} , which is now the special orthogonal group of a quadratic form on F^{2n+2} which is definite at the real places and split at the finite places of F . Note that $\pi \boxplus \pi_0$ is not regular: the zero weight appears twice at each real place of F . However the Langlands parameters of representations of the compact group $\mathrm{SO}_{2n+2}(\mathbb{R})$ are of the form

$$\bigoplus_{i=1}^{n+1} \mathrm{Ind}_{W_{\mathbb{R}}}^{W_{\mathbb{C}}} \left(z \mapsto (z/\bar{z})^{k_i} \right)$$

when composed with $\mathrm{SO}_{2n+2}(\mathbb{C}) \hookrightarrow \mathrm{GL}_{2n+2}(\mathbb{C})$, with $k_1 > \dots > k_{n+1} \geq 0$. Moreover $\mathcal{LL}((\pi \boxplus \pi_0)_v)$ is of the above form, with $k_{n+1} = 0$. The rest of the proof is identical to the odd-dimensional case.

This fact also shows that some *non-regular*, L-algebraic, self-dual, cuspidal representations of $\mathrm{GL}_{2n}(\mathbb{A}_F)$ contribute to the automorphic spectrum of \mathbf{G} . Consequently we can also extend Taylor’s result to the Galois representations associated with these slightly non-regular automorphic representations. These Galois representations were shown to exist by Wushi Goldring [Gol].

We now fix some notations for the rest of the article. The valuation v_p of $\overline{\mathbb{Q}}_p$ is the one sending p to 1, and $|\cdot|$ will denote the norm $p^{-v_p(\cdot)}$. All the number fields in the paper will sit inside $\overline{\mathbb{Q}}$. We have chosen arbitrary embeddings $\iota_p : \overline{\mathbb{Q}} \hookrightarrow \overline{\mathbb{Q}}_p$, $\iota_\infty : \overline{\mathbb{Q}} \hookrightarrow \mathbb{C}$. In fact, the constructions will only depend on the identification between the algebraic closures of \mathbb{Q} in $\overline{\mathbb{Q}}_p$ and \mathbb{C} (informally, $\iota_p \iota_\infty^{-1}$). Observe that the choice of a p -adic place v of a number field F and of an embedding $F_v \hookrightarrow \overline{\mathbb{Q}}_p$ is equivalent, via ι_p , to the choice of an embedding $F \hookrightarrow \overline{\mathbb{Q}}$. The same holds for the infinite places and ι_∞ . Thus if F is totally real, $\iota_p \iota_\infty^{-1}$ defines a bijection between the set of infinite places of F and the set of p -adic places v of F together with an embedding $F_v \hookrightarrow \overline{\mathbb{Q}}_p$. The eigenvarieties will be rigid analytic spaces (in the sense of Tate). If \mathcal{X} is a rigid analytic space over a finite extension E of \mathbb{Q}_p , $|\mathcal{X}|$ will denote its points.

I would like to thank Gaëtan Chenevier for introducing me to this fascinating subject, Colette Moeglin and Jean-Loup Waldspurger for Lemma 6.1.1, and James Arthur for his help.

2 Assumptions on the forthcoming chapter of Arthur’s book

As the results of this paper rely on the main theorem of chapter 9 of [Art11] (the case of inner forms of quasi-split classical groups), which is not yet available, we have stated some properties as assumptions: Assumptions 4.1.1, 6.1.2, 6.2.2 and 6.4.1. These will all be consequences of the main theorem of [Art11, Chapter 9], very similar to [Art11, Theorem 1.5.2]. In fact the difference will only be of local nature (see [Art06] and [Kot86]). Since the inner forms of classical groups we will consider are split at all the finite places, to state the analog of [Art11, Theorem 1.5.2] in these cases one would only need to know how to normalize the local A-packets, i.e. fix a Whittaker functional at each finite place and specify a “strong form” above the relevant inner form at the archimedean places. We are

unable to guess what global choice could determine these compatible local normalizations in general, except for the case of *pure* inner forms. This is the case for the orthogonal groups of section 5, but not for the “symplectic” groups of section 3.

A subsequent version of this paper will have the assumptions replaced by actual propositions or lemmas.

3 The eigenvariety for definite symplectic groups

In this section we recall the main result of [Loe11] in our particular case (existence of the eigenvariety for symplectic groups), and show that the points corresponding to unramified, “completely refinable” automorphic forms, with weight far from the walls, are “dense” in this eigenvariety.

3.1 The eigenvariety

3.1.1 Symplectic groups compact at the archimedean places

Let F be a totally real number field of even degree over \mathbb{Q} , and let D be a quaternion algebra over F , unramified at all the finite places of F ($F_v \otimes_F D \simeq M_2(F_v)$), and definite at all the real places of F . Such a D exists thanks to the exact sequence relation the Brauer groups of F and the F_v . Let n be a positive integer, and let \mathbf{G} be the algebraic group over F defined by the equation $M^*M = I_n$ for $M \in M_n(D)$, where $(M^*)_{i,j} = M_{j,i}^*$, and \cdot^* denotes conjugation in D .

Then $\mathbf{G}(F \otimes_{\mathbb{Q}} \mathbb{R})$ is a compact Lie group, and for all finite places v of F , $\mathbf{G} \times_F F_v \simeq \mathbf{Sp}_{2n}/F_v$.

Fix a prime p . We will apply the results of [Loe11] to the group $\mathbf{G}' = \text{Res}_{\mathbb{Q}}^F \mathbf{G}$. Let E be a finite and Galois extension of \mathbb{Q}_p , containing all the F_v (v over p).

3.1.2 The Atkin-Lehner algebra

The algebraic group $\mathbf{G}' \times_{\mathbb{Q}} \mathbb{Q}_p = \prod_{v|p} \mathbf{G} \times_{\mathbb{Q}} F_v$ (where v runs over the places of F) is isomorphic to $\prod_{v|p} \text{Res}_{\mathbb{Q}_p}^{F_v} \mathbf{Sp}_{2n}/F_v$, which is quasi-split but not split in general. The algebraic group \mathbf{Sp}_{2n} is defined over \mathbb{Z} by the equation ${}^t M J M = J$ in M_{2n} , where $J =$

$$\begin{pmatrix} 0 & J_n \\ -J_n & 0 \end{pmatrix} \text{ and } J_n = \begin{pmatrix} 0 & & & 1 \\ & \ddots & & \\ & & \ddots & \\ 1 & & & 0 \end{pmatrix}. \text{ We define its algebraic subgroups } \mathbf{T}_v, \mathbf{B}_v, \bar{\mathbf{B}}_v,$$

$\mathbf{N}_v, \bar{\mathbf{N}}_v$ of diagonal, upper triangular, lower triangular, unipotent upper triangular, and unipotent lower triangular matrices of $\text{Res}_{\mathbb{Q}_p}^{F_v} \mathbf{Sp}_{2n}/F_v$, and let $\mathbf{T} = \prod_{v|p} \mathbf{T}_v$, $\mathbf{B} = \prod_{v|p} \mathbf{B}_v$, and so on. In [Loe11, 2.4], only the action of the maximal split torus of $\mathbf{G}' \times_{\mathbb{Q}} \mathbb{Q}_p$ is considered. For our purpose, we will need to extend this and consider the action of a maximal (non-split in general) torus, that is \mathbf{T} , instead of a maximal split torus $\mathbf{S} \subset \mathbf{T}$. The results in [Loe11] are easily extended to this bigger torus, essentially because $\mathbf{T}(\mathbb{Q}_p)/\mathbf{S}(\mathbb{Q}_p)$ is compact. Moreover, we let I_v be the compact subgroup of $\mathbf{Sp}_{2n}(\mathcal{O}_v)$ consisting of matrices with invertible diagonal elements and elements of positive valuation below

$\mathcal{C}(\mathcal{U}, k)$ (k big enough such that χ is k -analytic on the open affinoid \mathcal{U}), which interpolates p -adically the restriction to $\mathbf{G}'(\mathbb{Q}_p)$ of algebraic representations of $\mathbf{G}'(\overline{\mathbb{Q}}_p)$. From there one can define the spaces $M(e, \mathcal{U}, k)$ ([Loe11, Definition 3.7.1]) of p -adic automorphic forms (or overconvergent automorphic forms, by analogy with the rigid-geometric case of modular forms) above an open affinoid or a point \mathcal{U} of \mathcal{W} which are k -analytic and fixed by the idempotent e . This space has an action of \mathcal{H}^+ . By [Loe11, Corollary 3.7.3], when considering p -adic automorphic forms which are eigenvectors for $[G_0 u G_0]$ for some $u \in \Sigma^{++}$ and for a non-zero eigenvalue (“finite slope” p -adic eigenforms), one can forget about k , and we will do so in the sequel.

3.1.4 Existence and properties of the eigenvariety

We choose the element

$$\eta = \left(\left(\begin{array}{ccccccc} \varpi_v^{-n} & & & & & & \\ & \ddots & & & & & \\ & & \varpi_v^{-1} & & & & \\ & & & \varpi_v & & & \\ & & & & \ddots & & \\ & & & & & \varpi_v^n & \\ & & & & & & \varpi_v^n \end{array} \right) \right)_v \in \Sigma^{++}$$

Theorem 3.1.1. *There exists a reduced rigid space \mathcal{X} over E , together with an E -algebra morphism $\Psi : \mathcal{H}^+ \rightarrow \mathcal{O}(\mathcal{X})^\times$ and a morphism of rigid spaces $w : \mathcal{X} \rightarrow \mathcal{W}$ such that:*

1. *The morphism $(w, \Psi([G_0 \eta G_0])^{-1}) : \mathcal{X} \rightarrow \mathcal{W} \times \mathbb{G}_m$ is finite*
2. *For each point x of \mathcal{X} , $\Psi \otimes w^\sharp : \mathcal{H}^+ \otimes_E \mathcal{O}_{w(x)} \rightarrow \mathcal{O}_x$ is surjective*
3. *For every finite extension E'/E , $\mathcal{X}(E')$ is in bijection with the finite slope systems of eigenvalues of \mathcal{H}^+ acting on the space of “overconvergent” automorphic forms, via evaluation of the image of Ψ at a given point.*

Moreover, for any point $x \in |\mathcal{X}|$, there is an arbitrarily small open affinoid \mathcal{V} containing x and an open affinoid \mathcal{U} of \mathcal{W} such that $\mathcal{V} \subset w^{-1}(\mathcal{U})$, the morphism $w|_{\mathcal{V}} : \mathcal{V} \rightarrow \mathcal{U}$ is finite, and surjective when restricted to any irreducible component of \mathcal{V} .

Proof. This is [Loe11, Theorems 3.11.2 and 3.12.3], except for the last assertion. To prove it, we need to go back to the construction of the eigenvariety in [Buz07]. Buzzard begins by constructing the Fredholm hypersurface \mathcal{Z} (encoding only the value of $\Psi([G_0 \eta G_0])$), together with a flat morphism $\mathcal{Z} \rightarrow \mathcal{W}$, before defining the finite morphism $\mathcal{X} \rightarrow \mathcal{Z}$. By [Buz07, Theorem 4.6], \mathcal{Z} can be admissibly covered by its open affinoids \mathcal{V}_0 such that w restricted to \mathcal{V}_0 induces a finite, surjective morphism to an open affinoid \mathcal{U} of \mathcal{W} , and \mathcal{V}_0 is a connected component of the pullback of \mathcal{U} . We can assume that \mathcal{U} is connected, and hence irreducible, since \mathcal{W} is normal. The morphism $\mathcal{V}_0 \rightarrow \mathcal{U}$ is both open (since it is flat: [Bos09, Corollary 7.2]) and closed (since it is finite), so that any irreducible component of \mathcal{V}_0 is mapped onto \mathcal{U} . This can be seen more naturally by observing that the irreducible components of \mathcal{V}_0 are also Fredholm hypersurfaces, by [Con99, Theorem 4.3.2].

By [Che04, Proposition 6.4.2], if \mathcal{V} denotes the pullback to \mathcal{X} of \mathcal{V}_0 , each irreducible component of \mathcal{V} is mapped onto an irreducible component of \mathcal{V}_0 (more precisely, this is a consequence of [Che04, Lemme 6.2.10]). To conclude, we only need to show that if $x \in \mathcal{V}$, up to restricting \mathcal{U} , the connected component of \mathcal{V} containing x can be arbitrarily small. This is a consequence of the following lemma. \square

Lemma 3.1.2. *Let $f : \mathcal{X}_1 \rightarrow \mathcal{X}_2$ be a finite morphism of rigid analytic spaces. Then the connected components of $f^{-1}(U)$, for U admissible open of \mathcal{X}_2 , form a basis for the canonical topology on \mathcal{X}_1 .*

Proof. It is enough to consider the case $\mathcal{X}_1 = \mathrm{Sp}A_1$, $\mathcal{X}_2 = \mathrm{Sp}A_2$. Let x_1 be a maximal ideal of A_1 . Then $f^{-1}(\{f(x_1)\}) = \{x_1, \dots, x_m\}$. We choose generators t_1, \dots, t_n of $f(x_1)$, and $r_1^{(i)}, \dots, r_{k_i}^{(i)}$ of x_i . Using the maximum modulus principle, it is easily seen that $\Omega_{j,N} := \{y \in \mathcal{X}_2 \mid |t_j(y)| \geq p^{-N}\}_{j,N}$ is an admissible covering of the admissible open $\mathcal{X}_2 \setminus \{f(x)\}$ of \mathcal{X}_2 . Let V_M be the admissible open $\{x \in \mathcal{X}_1 \mid \forall i, \exists k, |r_k^{(i)}(x)| \geq p^{-M}\}$, which is a finite union of open affinoids, hence quasi-compact. Consequently, the admissible open sets

$$\begin{aligned} U_{j,N} &:= V_M \cap f^{-1}(\Omega_{j,N}) \\ &= \left\{ x \in \mathcal{X}_1 \mid \forall i, \exists k, |r_k^{(i)}(x)| \geq p^{-M} \text{ and } |f^{\sharp}(t_j)(x)| \geq p^{-N} \right\}_{j,N} \end{aligned}$$

form an admissible covering of V_M . Therefore there is an N big enough so that

$$V_M = \bigcup_{j=1}^r U_{j,N}$$

which implies that

$$f^{-1}(\{y \in \mathcal{X}_2 \mid |t_j(y)| \leq p^{-N-1}\}) \subset \bigcup_i \left\{ x \in \mathcal{X}_1 \mid \forall k, |r_k^{(i)}(x)| \leq p^{-M} \right\}$$

and when M goes to infinity, the right hand side is the disjoint union of arbitrarily small affinoid neighbourhoods of the x_i . \square

We define the algebraic points of $\mathcal{W}(E)$ to be the ones of the form

$$(x_{v,i})_{v,i} \mapsto \prod_{v,\sigma} \sigma \left(\prod_{i=1}^n x_{v,i}^{k_{v,\sigma,i}} \right)$$

where $k_{v,\sigma,i}$ are integers, and such a point is called dominant if $k_{v,\sigma,1} \geq k_{v,\sigma,2} \geq \dots \geq k_{v,\sigma,n} \geq 0$.

Recall that a set $S \subset |\mathcal{X}|$ is said to *accumulate* at a point $x \in |\mathcal{X}|$ if x has a basis of affinoid neighbourhoods in which S is Zariski dense.

Proposition 3.1.3. *Let $(\phi_r)_r$ be a finite family of linear forms on \mathbb{R}^A where A is the set of triples (v, σ, i) for v a place of F above p , $\sigma : F_v \rightarrow E$ and $1 \leq i \leq n$, and let $(c_r)_r$ be a family of elements in $\mathbb{R}_{\geq 0}$. Assume that the open affine cone $C = \{y \in \mathbb{R}^A \mid \forall r, \phi_r(y) > c_r\}$ is nonempty. Then the set of algebraic characters in C yields a Zariski dense set in the weight space \mathcal{W} , which accumulates at all the algebraic points.*

Proof. [Che09, Lemma 2.7]. □

In particular the property of being dominant or “very regular” can be expressed in this way.

By finiteness of $\mathbf{G}(F)\backslash\mathbf{G}(\mathbb{A}_{F,f})/U$ for any open subgroup U of $\mathbf{G}(\mathbb{A}_{F,f})$, if Π is an automorphic representation of $\mathbf{G}(\mathbb{A}_F)$, the representation Π_f is defined over $\iota_\infty(\overline{\mathbb{Q}})$. Loeffler defines ([Loe11, Definition 3.9.1]) the classical subspace of the space of p -adic automorphic forms above an algebraic and dominant point w of the weight space. This subspace is isomorphic to $\iota_p \iota_\infty^{-1} \left(e(\mathcal{C}^\infty(\mathbf{G}(F)\backslash\mathbf{G}(\mathbb{A}_F)) \otimes W^*)^{\mathbf{G}(F \otimes_{\mathbb{Q}} \mathbb{R})} \right)$ as \mathcal{H}^+ -module, with W the representation of $\mathbf{G}'(F \otimes_{\mathbb{Q}} \mathbb{R})$ which is the restriction of the algebraic representation of $\mathbf{G}' \times_{\mathbb{Q}} \mathbb{C}$ having highest weight $\iota_\infty^{-1} \iota_p(w)$. The classical points of the eigenvariety are the ones having eigenvectors in the classical subspace.

We need to give an interpretation of classical points on the eigenvariety \mathcal{X} , in terms of automorphic representations of $\mathbf{G}(\mathbb{A}_F)$. Namely, there is a classical point $x \in \mathcal{X}(E')$ defining a character $\Psi_x : \mathcal{H} \rightarrow E'$ (here $E \subset E' \subset \overline{\mathbb{Q}}_p$) if and only if there is an automorphic representation $\Pi = \otimes'_v \Pi_v = \Pi_\infty \otimes \Pi_p \otimes \Pi_f^{(p)}$ of $\mathbf{G}(\mathbb{A}_F)$ such that:

- $\iota_p \iota_\infty^{-1} (\otimes_{v|\infty} \Pi_v)$ is the algebraic representation having highest weight $w(x)$;
- $\iota_p \left((e^S \otimes e_S) \Pi_f^{(p)} \right)$ contains a non-zero vector on which $\mathcal{H}^S \otimes \mathcal{H}_S$ acts according to Ψ_x ;
- $\iota_p(e_{G_0} \Pi_p)$ contains a non-zero vector on which \mathcal{H}_p acts according to $\mu_{w(x)} \Psi_x$, where $\mu_{w(x)}([G_0 \xi G_0]) = w(x)(\xi)$ if $\xi \in \Sigma^+$.

The twist by the character $\mu_{w(x)}$ is explained by the fact that the classical overconvergent automorphic forms are constructed by induction of characters of the torus extended from T_0 (on which they are defined by w) to T trivially on Σ .

3.2 Unramified and “completely refinable” points

3.2.1 Small slope p -adic eigenforms are classical

The algebraic and dominant points of \mathcal{W} are the ones of the form

$$(x_{v,i})_{v,i} \mapsto \prod_{v,\sigma} \sigma \left(\prod_{i=1}^n x_{v,i}^{k_{v,\sigma,i}} \right)$$

where $k_{v,\sigma,1} \geq k_{v,\sigma,2} \geq \dots \geq k_{v,\sigma,n} \geq 0$ are integers. The proof of the criterion given in [Loe11, Theorem 3.9.6] contains a minor error, because it “sees” only the restriction of these characters to the maximal split torus \mathbf{S} (over \mathbb{Q}_p), and the BGG resolution has to be applied to *split* semi-simple Lie algebras.

We correct it in the case of quasi-split reductive groups (in particular the restriction to a subfield of a quasi-split group remains quasi-split), and give a stronger criterion. This criterion could be used on an eigenvariety for which only the weights corresponding to a given p -adic place of F vary. For this purpose we use the “dual BGG resolution” given in [Jon]. The proof will be very close to that of [Loe11, Propositions 2.6.3-2.6.4]. In the

following \mathbf{G}' could be any quasi-split reductive group over \mathbb{Q}_p , and we could replace E/\mathbb{Q}_p by any extension splitting \mathbf{G}' .

Let \mathbf{B} be a Borel subgroup of \mathbf{G}' , \mathbf{S} a maximal split torus in \mathbf{B} , \mathbf{T} the centralizer of \mathbf{S} , a maximal torus. This determines an opposite Borel subgroup $\bar{\mathbf{B}}$ such that $\bar{\mathbf{B}} \cap \mathbf{B} = \mathbf{T}$. Let Φ^+ (resp. Δ) be the set of positive (resp. simple) roots of $\mathbf{G}' \times_{\mathbb{Q}_p} E$, with respect to the maximal torus \mathbf{T} of the Borel subgroup \mathbf{B} . One can split $\Delta = \sqcup_i \Delta_i$ where α, β belong to the same Δ_i if and only if $\alpha|_{\mathbf{S}} = \beta|_{\mathbf{S}}$ (equivalently, the Δ_i are the Galois orbits of Δ). Let Σ be a subgroup of $\mathbf{T}(\mathbb{Q}_p)$ supplementary to its maximal compact subgroup, and Σ^+ the submonoid consisting of the $z \in \mathbf{T}(\mathbb{Q}_p)$ such that $|\alpha(z)| \geq 1$ for all $\alpha \in \Delta$. For each i , define η_i to be the element of $\Sigma^+ / (Z(\mathbf{G}')(\mathbb{Q}_p) \cap \Sigma)$ generating $\cap_{j \neq i} \ker |\alpha_j(\cdot)|$ (here α_j denotes any element of Δ_j , and $|\alpha_j(\cdot)|$ does not depend on this choice).

Assume that G_0 is a compact open subgroup of $\mathbf{G}'(\mathbb{Q}_p)$ having an Iwahori factorization $\bar{N}_0 T_0 N_0$. Using a lattice in the Lie algebra of N and the exponential map, it is easily seen that N_0 admits a decreasing, exhaustive filtration by open subgroups $(N_k)_{k \geq 1}$ having a canonical rigid-analytic structure. Moreover any ordering of Φ^+ endows the Banach space of \mathbb{Q}_p -analytic functions on N_k taking values in E with an orthonormal basis consisting of monomials on the weight spaces.

Let λ be an algebraic and dominant weight of $\mathbf{T} \times_{\mathbb{Q}_p} E$. By [Jon], there is an exact sequence of $E[\mathbb{I}]$ -modules, where $\mathbb{I} = G_0 \Sigma^+ G_0 = \bar{B}_0 \Sigma^+ N_0$ is the monoid generated by G_0 and Σ^+ :

$$0 \rightarrow \text{Ind}_{\mathbf{B}}^{\mathbf{G}}(\lambda) \otimes \text{sm-Ind}_{\bar{B}_0}^{\bar{B}_0 N_0} 1 \rightarrow \text{la-Ind}_{\bar{B}}^{\bar{B} N_0}(\lambda) \rightarrow \bigoplus_{\alpha \in \Delta} \text{la-Ind}_{\bar{B}}^{\bar{B} N_0}(s_\alpha(\lambda + \rho) - \rho) \quad (3.2.1)$$

where $2\rho = \sum_{\alpha \in \Phi^+} \alpha$, “sm” stands for “smooth” and “la” for “locally analytic”. The relation with Loeffler’s $\text{Ind}(V)_k$ is $\text{la-Ind}_{\bar{B}}^{\bar{B} N_0}(\lambda) \otimes \lambda_{\text{sm}}^{-1} = \varinjlim_k \text{Ind}(E_\lambda)_k$, where λ_{sm} is the character on T which is trivial on its maximal compact subgroup and agrees with λ on Σ . Naturally $\text{Ind}_{\mathbf{B}}^{\mathbf{G}}(\lambda) \otimes \text{sm-Ind}_{\bar{B}_0}^{\bar{B}_0 N_0} 1 \otimes \lambda_{\text{sm}}^{-1} = \varinjlim_k \text{Ind}(E_\lambda)_k^{\text{cl}}$.

To prove a classicity criterion, we need to bound the action of η_i on the factors of the RHS of (3.2.1) twisted by λ_{sm}^{-1} . Let $n_\alpha = \alpha^\vee(\lambda) \in \mathbb{N}$ for $\alpha \in \Delta$, then $s_\alpha(\lambda + \rho) - \lambda - \rho = -(1+n_\alpha)\alpha$. The Banach space of k -analytic functions on N_0 is the direct sum of the spaces of analytic functions on xN_k , $x \in N_0/N_k$, and each of these spaces has an orthonormal (with respect to the supremum norm) basis $(v_{j,x})_{j \in J}$ where $J = \mathbb{N}^{\Phi^+}$ (monomials on the weights spaces). This basis depends on the choice of a representative x , but if we fix i and $x_0 \in N_0$, we can choose $\eta_i^{-1} x_0 \eta_i$ as a representative of its class. Then if $\phi = \sum_j a_j v_{j, \eta_i^{-1} x_0 \eta_i}$ (with $a_j \rightarrow 0$) is an element of $\text{la-Ind}_{\bar{B}}^{\bar{B} N_0}(s_\alpha(\lambda + \rho) - \rho) \otimes \lambda_{\text{sm}}^{-1}$, and $\xi \in N_k$,

$$\begin{aligned} (\eta_i \cdot \phi)(x_0 \xi) &= \eta_i^{-(1+n_\alpha)\alpha} \sum_{j \in J} a_j v_{j, \eta_i^{-1} x_0 \eta_i}(\eta_i^{-1} x_0 \xi \eta_i) \\ &= \sum_{j \in J} a_j \eta_i^{-(1+n_\alpha)\alpha - s(j)} v_{j, x_0}(x_0 \xi) \end{aligned}$$

where $s(j) = \sum_{\beta \in \Phi^+} j(\beta)\beta$. This shows that $|\eta_i \cdot \phi| \leq |\alpha(\eta_i)|^{-(1+n_\alpha)} |\phi|$, and so the operator η_i has norm less than or equal to $|\alpha(\eta_i)|^{-(1+n_\alpha)}$ on $\text{la-Ind}_{\bar{B}}^{\bar{B} N_0}(s_\alpha(\lambda + \rho) - \rho) \otimes \lambda_{\text{sm}}^{-1}$.

We can then apply the exact functor which to an $E[\mathbb{I}]$ -module W associates the automorphic forms taking values in W , and take the invariants under the idempotent e (this functor is left exact). We obtain that $M(e, E_\lambda)/M(e, E_\lambda)_{\text{cl}}$ (the space of p -adic automorphic forms modulo the classical automorphic forms) embeds in $\bigoplus_{\alpha \in \Delta} M_\alpha$ where each M_α is a Banach space on which the operator $[G_0 \eta_i G_0]$ has norm $\leq |\alpha(\eta_i)|^{-(1+n_\alpha)}$. The following criterion follows:

Lemma 3.2.1. *If an overconvergent eigenform $f \in M(e, E_\lambda)$ satisfies $[G_0 \eta_i G_0] f = \mu_i f$ with $\mu_i \neq 0$ and*

$$v_p(\mu_i) < \inf_{\alpha \in \Delta_i} -(1+n_\alpha)v_p(\alpha(\eta_i))$$

for all i , then f is classical.

In the case of the symplectic group \mathbf{G}' , the family $(\eta_i)_i$ can be indexed by the couples (v, i) where v is a place of F above p and $1 \leq i \leq n$, and $\Delta_{v,i}$ is indexed by the embeddings $F_v \hookrightarrow E$. Specifically, $\eta_{v,i}$ is trivial at all the places except for v , where it equals

$$\text{Diag}(x_1, \dots, x_n, x_n^{-1}, \dots, x_1^{-1})$$

$$\text{with } x_j = \begin{cases} \varpi_v^{-1} & \text{if } j \leq i \\ 1 & \text{if } j > i \end{cases}$$

The conditions in the previous lemma can be written

$$\begin{cases} v_p(\mu_{v,i}) < \frac{1}{e_v} \inf_\sigma (1 + k_{v,\sigma,i} - k_{v,\sigma,i+1}) & \text{for } i < n \\ v_p(\mu_{v,n}) < \frac{1}{e_v} \inf_\sigma (2 + 2k_{v,\sigma,n}). \end{cases}$$

3.2.2 Representations having Iwahori-invariants and unramified principal series

We recall results of Casselman showing that irreducible representations having Iwahori-invariants appear in unramified principal series, and giving the Atkin-Lehner eigenvalues in terms of the unramified character being induced.

In this subsection, we fix a place v of F above p . Recall I_v has an Iwahori decomposition $I_v = N_{v,0} T_{v,0} \bar{N}_{v,0}$. As in [Cas], if (Π, V) is a smooth representation of $\mathbf{G}(F_v)$, $V(\bar{N}_v)$ is the subspace of V spanned by the $\Pi(\bar{n})(x) - x$, $\bar{n} \in \bar{N}_v$, $V_{\bar{N}_v} = V/V(\bar{N}_v)$ and if $\bar{N}_{v,i}$ is a compact subgroup of \bar{N}_v , $V(\bar{N}_{v,i}) = \left\{ v \in V \mid \int_{\bar{N}_{v,i}} \Pi(\bar{n})(v) d\bar{n} = 0 \right\}$.

Lemma 3.2.2. *Let (Π, V) be an admissible representation of $\mathbf{G}(F_v)$ over \mathbb{C} . Then the natural (vector space) morphism from V^{I_v} to $(V_{\bar{N}_v})^{T_{v,0}}$ is an isomorphism, inducing a Σ_v^+ -equivariant isomorphism*

$$\Pi^{I_v} \xrightarrow{\sim} (\Pi_{\bar{N}_v})^{T_{v,0}} \otimes \delta_{\bar{B}_v}^{-1}$$

where $\delta_{\bar{B}_v}$ denotes the modulus morphism of \bar{B}_v , and $u \in \Sigma_v^+$ acts on Π^{I_v} by $[I_v u I_v]$.

Proof. Let $\bar{N}_{v,1}$ be a compact subgroup of \bar{N}_v such that $V^{I_v} \cap V(\bar{N}_v) \subset V(\bar{N}_{v,1})$. There is a $u \in \Sigma_v^+$ such that $u \bar{N}_{v,1} u^{-1} \subset \bar{N}_{v,0}$. By [Cas, Prop. 4.1.4], and using the fact that

Definition 3.2.5. An irreducible representation Π_v of $\mathbf{G}(F_v)$ is completely refinable if it is isomorphic to $\text{Ind}_{\bar{B}_v}^{\text{Sp}_{2n}(F_v)} \chi$ for some unramified character χ .

An automorphic representation Π of $\mathbf{G}(\mathbb{A}_F)$ is completely refinable if Π_v is completely refinable for any $v|p$.

Note that completely refinable representations are unramified (for any choice of hyperspecial subgroup). A representation Π_v is completely refinable if and only if $(\Pi_v)_{\bar{N}_v}^{\text{ss}}$ is the sum of $|W(T_v, \mathbf{G}(F_v))|$ unramified characters.

Recall that classical points on the eigenvariety are determined by an automorphic representation Π together with a refinement of each Π_v , $v|p$. Completely refinable automorphic representations are the ones giving the greatest number of points on the eigenvariety. When one can associate Galois representations to automorphic representations, each refinement of Π comes with a “ p -adic family” of Galois representations going through the same one.

Proposition 3.2.6. Let $f_1, \dots, f_r \in \mathcal{O}(\mathcal{X})^\times$. The set \mathcal{S} of points corresponding to completely refinable, unramified classical points at which

$$\min_{v,\sigma} \min\{k_{v,\sigma,1} - k_{v,\sigma,2}, \dots, k_{v,\sigma,n-1} - k_{v,\sigma,n}, k_{v,\sigma,n}\} \geq \max\{v_p(f_1), \dots, v_p(f_r)\} \quad (3.2.2)$$

is Zariski dense and accumulates at all the algebraic points.

Compare [Che04, Proposition 6.4.7], [Loe11, Corollary 3.13.3].

Proof. The hypotheses in the classicality criterion 3.2.1 and the ones in Theorem 3.2.4 are implied by inequalities of the form 3.2.2. First we prove the accumulation property. We can restrict to open affinoids \mathcal{V} of the eigenvariety, and hence assume that the right hand side of 3.2.2 is replaced by a constant. By Theorem 3.1.1, \mathcal{V} can be an arbitrarily small open affinoid containing an algebraic point x of \mathcal{X} , such that there is open affinoid \mathcal{U} of \mathcal{W} such that $\mathcal{V} \subset w^{-1}(\mathcal{U})$, the morphism $w|_{\mathcal{V}} : \mathcal{V} \rightarrow \mathcal{U}$ is finite, and surjective when restricted to any irreducible component of \mathcal{V} . By Proposition 3.1.3, the algebraic weights satisfying 3.2.2 are Zariski dense in the weight space \mathcal{W} and accumulate at all the algebraic points of \mathcal{W} . [Che04, Lemme 6.2.8] shows that $\mathcal{S} \cap \mathcal{V}$ is Zariski-dense in \mathcal{V} .

Each irreducible component \mathcal{X}' of \mathcal{X} is mapped onto a Zariski-open subset of a connected component of \mathcal{W} , by [Che04, Corollaire 6.4.4] (which is a consequence of the decomposition of a Fredholm series into a product of prime Fredholm series, [Con99, Corollary 4.2.3]), so \mathcal{X}' contains at least one algebraic point (the algebraic weights intersect all the connected components of \mathcal{W}), and hence the Zariski closure of $\mathcal{S} \cap \mathcal{X}'$ contains an open affinoid of \mathcal{X}' , which is Zariski dense in \mathcal{X}' . \square

4 Galois representations associated with automorphic representations of symplectic groups

4.1 A consequence of Arthur's description of the discrete spectrum for classical groups

4.1.1 Automorphic self-dual representations of GL_{2n+1} of orthogonal type

According to Arthur's conjectural parametrization of discrete automorphic representations, each such representation of $\mathbf{G}(\mathbb{A}_F)$ should be part of an A-packet corresponding to a discrete parameter, which is a representation

$$\mathcal{L}_F \times \mathrm{SL}_2(\mathbb{C}) \rightarrow \mathrm{SO}_{2n+1}(\mathbb{C})$$

such that (among other conditions) the commutant of the image is finite.

The standard embedding $\mathrm{SO}_{2n+1}(\mathbb{C}) \hookrightarrow \mathrm{GL}_{2n+1}(\mathbb{C})$ “transfers” this parameter to a parameter of GL_{2n+1}/F , which is not discrete in general, and thus it corresponds to an automorphic representation of $\mathrm{GL}_{2n+1}(\mathbb{A}_F)$. Here we define an automorphic representation π of $\mathrm{GL}_N(\mathbb{A}_F)$ as a formal sum of discrete automorphic representations π_i of GL_{n_i} such that $\sum_i n_i = N$. We will write $\pi = \boxplus_i \pi_i$. By [MW89], each π_i is the Langlands quotient of the parabolic induction of twists of a single cuspidal representation by powers of $|\det|$. We will not need this generality, as we will force the representations π_i to be cuspidal in the sequel.

Since π comes from a self-dual parameter, it is self-dual: $\pi^\vee \simeq \pi$. Even though π is not discrete in general, the discreteness of the parameter which takes values in SO_{2n+1} implies that the π_i 's are self-dual.

If $\Pi = \otimes_v \Pi_v$ is an automorphic representation of $\mathbf{G}(\mathbb{A}_F)$, then for any archimedean place v of F , the local Langlands parameter of Π_v (composed with $\mathrm{SO}_{2n+1}(\mathbb{C}) \hookrightarrow \mathrm{GL}_{2n+1}(\mathbb{C})$) is of the form:

$$\mathcal{L}\mathcal{L}(\Pi_v) \simeq \epsilon^n \oplus \bigoplus_{i=1}^n \mathrm{Ind}_{W_{\mathbb{C}}}^{W_{\mathbb{R}}} (z \mapsto (z/\bar{z})^{r_i})$$

where ϵ is the only non-trivial character of $W_{\mathbb{C}}/W_{\mathbb{R}}$, and the r_i are integers, with $r_n > r_{n-1} > \dots > r_1 > 0$. We define $A_{\mathrm{Sp}_{2n}}$ to be the set of automorphic representations such that for each infinite place v of F , $r_1 \geq 2$ and $r_{i+1} \geq r_i + 2$. The equivalence above is meant as representations of $W_{\mathbb{R}}$, although $\mathcal{L}\mathcal{L}(\Pi_v)$ is a parameter taking values in $\mathrm{SO}_{2n+1}(\mathbb{C})$ (the two notions coincide).

Similarly, let $A_{\mathrm{GL}_{2n+1}}$ be the set of formal sums of self-dual cuspidal representations $\pi = \boxplus_i \pi_i = \otimes_v \pi_v$ of $\mathrm{GL}_{2n+1}(\mathbb{A}_F)$ such that for each infinite place v of F ,

$$\mathcal{L}\mathcal{L}(\pi_v) \simeq \epsilon^n \oplus \bigoplus_{i=1}^n \mathrm{Ind}_{W_{\mathbb{C}}}^{W_{\mathbb{R}}} (z \mapsto (z/\bar{z})^{r_i})$$

where the r_i 's are integers, such that $r_1 \geq 2$, $r_{i+1} \geq r_i + 2$, and such that the product of the central characters of the π_i 's is trivial.

These inequalities are imposed to ensure that the corresponding global parameters are trivial on Arthur's $\mathrm{SL}_2(\mathbb{C})$, to simplify the statements. That is why we take formal sums of *cuspidal* (not discrete) representations.

Note that there is no non-zero alternate bilinear form preserved by such a parameter (one could say that the parameter is “completely orthogonal”).

Assumption 4.1.1. *For any $\Pi \in A_{\mathrm{Sp}_{2n}}$, there is a $\pi \in A_{\mathrm{GL}_{2n+1}}$, such that the local Langlands parameters match at the infinite places, and for any finite place v of F , π_v is unramified if Π_v is unramified, and in that case the local parameters match, by means of the inclusion $\mathrm{SO}_{2n+1}(\mathbb{C}) \subset \mathrm{GL}_{2n+1}(\mathbb{C})$.*

4.1.2 p -adic Galois representations associated with RLASDC representations of GL_N

An automorphic cuspidal representation π of $\mathrm{GL}_N(\mathbb{A}_F)$ is said to be L -algebraic if for any infinite place v of F , the restriction of $\mathcal{LL}(\pi_v)$ to \mathbb{C}^\times is of the form

$$z \mapsto \mathrm{Diag} \left(\left(z^{a_{v,i}} \bar{z}^{b_{v,i}} \right)_i \right)$$

where $a_i, b_i \in \mathbb{Z}$. By the “purity lemma” [Clo88, Lemme 4.9], $a_{v,i} + b_{v,i}$ does not depend on v, i . We will say that π is L -algebraic *regular* if for any v as above, the $a_{v,i}$ are distinct. By purity, this implies that if v is real,

$$\mathcal{LL}(\pi_v) | \cdot |^{-s} = \begin{cases} \epsilon^e \oplus_i \mathrm{Ind}_{W_{\mathbb{C}}}^{W_{\mathbb{R}}} \left(z \mapsto (z/\bar{z})^{a'_{v,i}} \right) & \text{if } N \text{ is odd, with } e = 0, 1 \\ \oplus_i \mathrm{Ind}_{W_{\mathbb{C}}}^{W_{\mathbb{R}}} \left(z \mapsto (z/\bar{z})^{a'_{v,i}} \right) & \text{if } N \text{ is even} \end{cases}$$

for some integer s , and integers $0 < a'_{v,1} < \dots < a'_{v, \lfloor N/2 \rfloor}$.

As a special case of [CH, Theorem 4.2] (which builds on previous work of Clozel, Harris, Kottwitz, Labesse, Shin, Taylor), we have the following theorem

Theorem 4.1.2. *Let π be a regular L -algebraic, self-dual, cuspidal (RLASDC) representation of $\mathrm{GL}_{2n+1}(\mathbb{A}_F)$. Then π is L -arithmetic, and there is a continuous Galois representation*

$$\rho_{\iota_p, \iota_\infty}(\pi) : G_F \longrightarrow \mathrm{GL}_{2n+1}(\overline{\mathbb{Q}}_p)$$

such that if v is a finite place of F and π_v is unramified,

1. if v is coprime to p , then $\rho_{\iota_p, \iota_\infty}(\pi)|_{G_{F_v}}$ is unramified, and

$$\det(T\mathrm{Id} - \rho_{\iota_p, \iota_\infty}(\pi)(\mathrm{Frob}_v)) = \iota_p \iota_\infty^{-1} \det(T\mathrm{Id} - A)$$

where $A \in \mathrm{GL}_N(\mathbb{C})$ is associated with π_v via the Satake isomorphism.

2. if v lies above p , $\rho_{\iota_p, \iota_\infty}(\pi)|_{G_{F_v}}$ is crystalline. The associated filtered φ -module (over $F_{v,0} \otimes_{\mathbb{Q}_p} \overline{\mathbb{Q}}_p$) is such that

$$\det_{\overline{\mathbb{Q}}_p} \left(T\mathrm{Id} - \varphi^{f_v} \right) = \iota_p \iota_\infty^{-1} \det(T\mathrm{Id} - A)^{f_v}$$

where $A \in \mathrm{GL}_N(\mathbb{C})$ is associated with π_v via the Satake isomorphism. For any $\sigma : F_v \rightarrow \overline{\mathbb{Q}}_p$, the σ -Hodge-Tate weights are the $a_{w,i}$, where w is the real place of F defined by σ , ι_p and ι_∞ .

The power f_v appearing at places above p may seem more natural to the reader (and will actually disappear) after reading subsection 4.2.1.

Combining this theorem with the transfer detailed in the last section, we obtain

Corollary 4.1.3. *Let Π be an automorphic representation of $\mathbf{G}(\mathbb{A}_F)$, whose weights $k_{w,1} \geq k_{w,2} \geq \dots k_{w,n} \geq 0$ at the real places w are far from the walls ($\Pi \in A_{\mathrm{Sp}_{2n}}$ is enough), and unramified at the places above p . There exists a continuous semisimple Galois representation*

$$\rho_{\iota_p, \iota_\infty}(\Pi) : G_F \longrightarrow \mathrm{GL}_{2n+1}(\overline{\mathbb{Q}}_p)$$

such that for any finite place v of F such that Π_v is unramified

1. if v is coprime to p , then $\rho_{\iota_p, \iota_\infty}(\Pi)|_{G_{F_v}}$ is unramified, and

$$\det(T\mathrm{Id} - \rho_{\iota_p, \iota_\infty}(\Pi)(\mathrm{Frob}_v)) = \iota_p \iota_\infty^{-1} \det(T\mathrm{Id} - A)$$

where $A \in \mathrm{GL}_N(\mathbb{C})$ is associated with Π_v via the Satake isomorphism.

2. if v lies above p , $\rho_{\iota_p, \iota_\infty}(\Pi)|_{G_{F_v}}$ is crystalline. The associated filtered φ -module is such that

$$\det_{\overline{\mathbb{Q}}_p}(T\mathrm{Id} - \varphi^{f_v}) = \iota_p \iota_\infty^{-1} \det(T\mathrm{Id} - A)^{f_v}$$

where $A \in \mathrm{SO}_{2n+1}(\mathbb{C}) \subset \mathrm{GL}_{2n+1}(\mathbb{C})$ is associated with Π_v via the Satake isomorphism. For any $\sigma : F_v \rightarrow \overline{\mathbb{Q}}_p$, the σ -Hodge-Tate weights are $k_{w,1} + n > k_{w,2} + n - 1 > \dots > k_{w,1} + 1 > 0 > -k_{w,1} - 1 > \dots > -k_{w,1} - n$, where w is the real place of F defined by σ , ι_p and ι_∞ .

Proof. There is an automorphic representation $\pi = \boxplus_i \pi_i$ of $\mathrm{GL}_{2n+1}(\mathbb{A}_F)$ corresponding to Π by Assumption 4.1.1, obtained by induction from distinct cuspidal representations π_i . Let $\rho_{\iota_p, \iota_\infty}(\Pi) = \oplus_i \rho_{\iota_p, \iota_\infty}(\pi_i)$. □

Note that in that case, since Π_∞ is \mathbb{C} -algebraic, Π is obviously \mathbb{C} -arithmetic (which is equivalent to \mathbb{L} -arithmetic in the case of Sp_{2n}), and thus the coefficients of the polynomials appearing in the corollary lie in a finite extension of \mathbb{Q} .

4.1.3 The Galois pseudocharacter on the eigenvariety

To study families of representations, it is convenient to use *pseudorepresentations* (or *pseudocharacters*), which are simply the traces of semi-simple representations when the coefficient ring is an algebraically closed field of characteristic zero. We refer to [Tay91] for the definition, and [Tay91, Theorem 1] is the ‘‘converse theorem’’ we will need.

On $\mathcal{O}(\mathcal{X})$, we put the topology of uniform convergence on open affinoids.

The Zariski-density of the classical points at which we can define an attached Galois representation implies the following

Proposition 4.1.4. *There is a continuous pseudocharacter $T : G_F \rightarrow \mathcal{O}(\mathcal{X})$, such that at every classical unramified point of the eigenvariety having weight far from the walls, T specializes to the character of the Galois representation associated with the automorphic representation by Corollary 4.1.3.*

Proof. This is identical to the unitary case, and thus is a consequence of [Che04, Proposition 7.1.1], by Proposition 3.2.6. \square

Thus at any (classical or not) point of the eigenvariety, there is an attached Galois representation.

4.2 Galois representations stemming from symplectic forms are generically almost irreducible

4.2.1 Crystalline representations over $\overline{\mathbb{Q}_p}$

We fix a finite extension K of \mathbb{Q}_p , and denote K_0 the maximal unramified subextension, $e = [K : K_0]$, $f = [K_0 : \mathbb{Q}_p]$. Let $\rho : G_K \rightarrow \mathrm{GL}(V)$ be a continuous representation of the absolute Galois group of K , where V is a finite dimensional vector space over L , a finite Galois extension of \mathbb{Q}_p . We will take L to be big enough so as to be able to assume in many situations that $L = \overline{\mathbb{Q}_p}$. For example, we can assume that L is an extension of K , and that ρ has a composition series $0 = V_1 \subset \dots \subset V_r = V$ such that each quotient V_{i+1}/V_i is absolutely irreducible.

For any such ρ , we denote $D_{\mathrm{cris}}(V) = (B_{\mathrm{cris}} \otimes_{\mathbb{Q}_p} V)^{G_K}$. From now on we assume that ρ is a crystalline representation, which means that $\dim_{K_0} D_{\mathrm{cris}}(V) = \dim_{\mathbb{Q}_p} V$. It is well-known that $D_{\mathrm{cris}}(V)$ is a filtered φ -module over K , and since V is a vector space over L , $D_{\mathrm{cris}}(V)$ is a φ -module over $K_0 \otimes_{\mathbb{Q}_p} L$, and $D_{\mathrm{dR}}(V) = K \otimes_{K_0} D_{\mathrm{cris}}(V)$ is a module over $K \otimes_{\mathbb{Q}_p} L$ with a filtration by projective submodules.

We have a natural decomposition $K_0 \otimes_{\mathbb{Q}_p} L \simeq \prod_{\sigma_0 \in \Upsilon_0} L_{\sigma_0}$ with $\Upsilon_0 = \mathrm{Hom}_{\mathbb{Q}_p\text{-alg.}}(K_0, L)$ and $L_{\sigma_0} \simeq L$, given by the morphisms $\sigma_0 \otimes \mathrm{Id}_L$. Similarly, $K \otimes_{\mathbb{Q}_p} L \simeq \prod_{\sigma \in \Upsilon} L_{\sigma}$ with $\Upsilon = \mathrm{Hom}_{\mathbb{Q}_p\text{-alg.}}(K, L)$.

Hence we have decompositions

$$D_{\mathrm{cris}}(V) = \prod_{\sigma_0 \in \Upsilon_0} D_{\mathrm{cris}}(V)_{\sigma_0}, \quad D_{\mathrm{dR}}(V) = \prod_{\sigma \in \Upsilon} D_{\mathrm{dR}}(V)_{\sigma}.$$

The operator φ restricts as linear isomorphisms from $D_{\mathrm{cris}}(V)_{\sigma_0}$ to $D_{\mathrm{cris}}(V)_{\sigma_0 \circ \varphi^{-1}}$, and so φ^f is a L_{σ_0} -linear automorphism on each $D_{\mathrm{cris}}(V)_{\sigma_0}$, which are isomorphic as vector spaces over L equipped with the linear automorphism φ^f .

Each $D_{\mathrm{dR}}(V)_{\sigma}$ comes with a filtration, and hence defines $\dim_L V = N$ Hodge-Tate weights $k_{\sigma,1} \leq \dots \leq k_{\sigma,N}$ (the jumps of the filtration).

Although we will not use it, it should be noted that by [BM02, Proposition 3.1.1.5], to verify the weak admissibility of a filtered φ -module D over K with an action of L commuting with φ and leaving the filtration stable, it is enough to check the inequality $t_N(D') \geq t_H(D')$ for sub- $K_0 \otimes L$ -modules stable under φ .

If φ^f has eigenvalues $\varphi_1, \dots, \varphi_N$, with $v_p(\varphi_1) \leq \dots \leq v_p(\varphi_N)$, we can in particular choose $D' = \bigoplus_{i \leq j} \ker(\varphi^f - \varphi_i)$ (if the eigenvalues are distinct, but even if they are not, we can choose D' such that $\varphi^f|_{D'}$ has eigenvalues $\varphi_1, \dots, \varphi_j$, counted with multiplicities).

The worst case for the filtration yields the inequalities

$$\begin{aligned} v_p(\varphi_1) &\geq \frac{1}{e} \sum_{\sigma} k_{\sigma,1} \\ v_p(\varphi_1\varphi_2) &\geq \frac{1}{e} \sum_{\sigma} k_{\sigma,1} + k_{\sigma,2} \\ &\vdots \end{aligned}$$

In the sequel, we will only use these inequalities, and we will not be concerned with the subtleties of the filtrations.

4.2.2 Variation of the crystalline Frobenius on the eigenvariety

In this section we explicit the formulas relating the eigenvalues of the crystalline Frobenius at classical, unramified points of the eigenvariety and the eigenvalues of the Hecke-Iwahori operators acting on p -adic automorphic forms. Let x be a classical point on the eigenvariety. There is an automorphic representation Π of $\mathbf{G}(\mathbb{A}_F)$ such that $\iota_p \iota_{\infty}^{-1}(\Pi_{\infty})$ is the representation having highest weight $w(x)$. Assume that Π_p is unramified. The point x defines a refinement of Π_p , that is an unramified character $\chi_x : T_0 \rightarrow \mathbb{C}^{\times}$ such that $\Pi_p \hookrightarrow \text{Ind}_{\bar{B}}^{\mathbf{G}'(\mathbb{Q}_p)} \chi_x$, or equivalently the character $\delta_{\bar{B}}^{1/2} \chi_x$ appearing in $(\Pi_p)_{\bar{N}}$. By 3.2.2, for any $u \in \Sigma^+$, $\mu_{w(x)} \Psi_x |_{\mathcal{H}_p} = (\iota_p \circ \iota_{\infty}^{-1} \circ \chi_x) \delta_{\bar{B}}^{1/2}$.

The diagonal torus in $\text{SO}_{2n+1}(\mathbb{C})$ and the identification of it with the dual of the diagonal torus of Sp_{2n}/F_v being fixed, the character χ_x is mapped by the unramified Langlands correspondence for tori to $y = (y_v)_{v|p}$ with $y_v = \text{Diag}(y_{1,v}, \dots, y_{n,v}, 1, y_{n,v}^{-1}, \dots, y_{1,v}^{-1})$, and $y_{v,i} = \chi_x(\text{Diag}(1, \dots, \varpi_v, \dots, 1, 1, \dots, \varpi_v^{-1}, \dots, 1))$ (ϖ_v being the i -th element). Thus the linearization of the crystalline Frobenius φ^{f_v} on $D_{\text{cris}}(\rho_{\iota_p, \iota_{\infty}}(\pi)|_{G_{F_v}})_{\sigma_0}$ (for any choice of $\sigma_0 : F_v \rightarrow E$ in $\Upsilon_{0,v}$) has eigenvalues

$$\iota_p \iota_{\infty}^{-1}(y_{v,i}) = q_v^{n+1-i} \phi_{v,n+1-i}(x) \prod_{\sigma \in \Upsilon_v} \sigma(\varpi_v)^{k_{v,\sigma,i}}$$

and their inverses, together with the eigenvalue 1. Here $\phi_{v,n+1-i} \in \mathcal{O}(\mathcal{X})$ is defined by

$$\phi_{v,n+1-i} = \frac{\Psi([G_0 u_{i-1} G_0])}{\Psi([G_0 u_i G_0])}$$

with $u_i = \text{Diag}(\varpi_v^{-1}, \dots, \varpi_v^{-1}, 1, \dots, 1, \varpi_v, \dots, \varpi_v)$ (the last ϖ_v^{-1} is the i -th element), and $k_{v,\sigma,i}$ the integers defining the weight $w(x)$.

Assume furthermore that Π_p admits another refinement $\chi_{x'} = \chi_x^a$ for some $a = (a_v)_{v|p}$ in the Weyl group $W(\mathbf{G}'(\mathbb{Q}_p), \mathbf{T}(\mathbb{Q}_p)) = \prod_v W(\mathbf{G}(F_v), T_v)$. Each $W(\mathbf{G}(F_v), T_v)$ can be identified with the group of permutations $a_v : \{-n, \dots, n\} \rightarrow \{-n, \dots, n\}$ such that $a_v(-i) = -a_v(i)$ for all i , acting by

$$a_v(\text{Diag}(x_1, \dots, x_n, x_n^{-1}, \dots, x_1^{-1})) = \text{Diag}(x_{a_v^{-1}(1)}, \dots, x_{a_v^{-1}(n)}, x_{a_v^{-1}(-n)}, \dots, x_{a_v^{-1}(1)})$$

on T_v , where for commodity we set $x_{-i} = x_i^{-1}$ for $i < 0$. Similarly we define $k_{v,\sigma,-i} = -k_{v,\sigma,i}$ and $\phi_{v,-i} = \phi_{v,i}^{-1}$. We also set $k_{v,\sigma,0} = 0$, $\phi_{v,0} = 1$. The equality $\chi_{x'} = \chi_x^a$ can also be

written

$$q_v^{(n+1)\text{sign}(w(i))-w(i)} \phi_{v,n+1-w(i)}(x) \prod_{\sigma \in \Upsilon_v} \sigma(\varpi_v)^{k_{v,\sigma,w(i)}} = q_v^{n+1-i} \phi_{v,n+1-i}(x') \prod_{\sigma \in \Upsilon_v} \sigma(\varpi_v)^{k_{v,\sigma,i}}$$

which is valid for any $-n \leq i \leq n$ if we set $\text{sign}(i) = -1$ (resp. 0, 1) if i is negative (resp. zero, positive), and equivalent to

$$\phi_{v,n+1-i}(x') = \phi_{v,n+1-w(i)}(x) q_v^{i-w(i)+(n+1)(\text{sign}(i)-\text{sign}(w(i)))} \prod_{\sigma \in \Upsilon_v} \sigma(\varpi_v)^{k_{v,\sigma,w(i)}-k_{v,\sigma,i}}.$$

This last formula will be useful in the proof of the main result.

4.2.3 Main result

Lemma 4.2.1. *Let K be a finite extension of \mathbb{Q}_p , and let $\rho : G_K \rightarrow \text{GL}_N(\overline{\mathbb{Q}}_p)$ be a crystalline representation. Let $(D, \varphi, \text{Fil}^i D \otimes_{K_0} K)$ be the associated filtered φ -module. Let $\kappa_{\sigma,1} \leq \dots \leq \kappa_{\sigma,N}$ be the Hodge-Tate weights associated with the embedding $\sigma : K \hookrightarrow \overline{\mathbb{Q}}_p$. Let $\varphi_1, \dots, \varphi_N$ be the eigenvalues of the linear operator φ^f (on any of the D_{σ_0} , $\sigma_0 \in \Upsilon_0$), and suppose they are distinct. Finally, assume that for some $\tau \in \Upsilon$, for all i ,*

$$\left| v_p(\varphi_i) - \frac{1}{e} \sum_{\sigma \in \Upsilon} \kappa_{\sigma,i} \right| \leq \frac{1}{eN} \min_{1 \leq j \leq N-1} \kappa_{\tau,j+1} - \kappa_{\tau,j}.$$

Then if $D' \subset D$ is an admissible sub- φ -module over $K_0 \otimes_{\mathbb{Q}_p} \overline{\mathbb{Q}}_p$ (corresponding to a sub-representation), there is a subset I of $\{1, \dots, N\}$ such that D' has φ^f -eigenvalues $(\varphi_i)_{i \in I}$ and τ -Hodge-Tate weights $(\kappa_{\sigma,i})_{i \in I}$.

Proof. Since the eigenvalues of φ^f are distinct, and D' is stable under φ , there is a subset I of $\{1, \dots, N\}$ such that $D' = \ker \prod_{i \in I} (\varphi^f - \varphi_i)$. There are unique increasing functions $\theta_{1,\sigma} : I \rightarrow \{1, \dots, N\}$ such that the σ -weights of D' are the $\kappa_{\sigma,\theta_{1,\sigma}(i)}$, for $i \in I$. By ordering similarly the weights of D/D' , we define increasing functions $\theta_{2,\sigma} : \{1, \dots, N\} \setminus I \rightarrow \{1, \dots, N\}$, and we can glue the $\theta_{\cdot,\sigma}$ to get bijective maps $\theta_\sigma : \{1, \dots, N\} \rightarrow \{1, \dots, N\}$. We will show that $\theta_\tau = \text{Id}$

We now write the admissibility condition for D' and D/D' . Let i_1 be the smallest element of I . Then $\ker(\varphi^f - \varphi_{i_1})$ is a sub- φ -module of D' . Its induced σ -weight is one of the $\kappa_{\sigma,\theta_\sigma(i)}$ for $i \in I$, thus it is greater than or equal to $\kappa_{\sigma,\theta_\sigma(i_1)}$. This implies that $v_p(\varphi_{i_1}) \geq 1/e \sum_{\sigma \in \Upsilon} \kappa_{\sigma,\theta_\sigma(i_1)}$. We can proceed similarly for the submodules

$$\ker \left((\varphi^f - \varphi_{i_1}) \dots (\varphi^f - \varphi_{i_r}) \right)$$

(where the i_\cdot are the ordered elements of I), to get the inequality

$$\sum_{1 \leq x \leq r} v_p(\varphi_{i_x}) \geq \frac{1}{e} \sum_{1 \leq x \leq r} \sum_{\sigma \in \Upsilon} \kappa_{\sigma,\theta_\sigma(i_x)}$$

The same applies to D/D' , and by adding both inequalities, we finally get

$$\sum_{1 \leq i \leq s} v_p(\varphi_i) \geq \frac{1}{e} \sum_{1 \leq i \leq s} \sum_{\sigma \in \Upsilon} \kappa_{\sigma,\theta_\sigma(i)}$$

We now isolate τ , using the fact that $\sum_{1 \leq i \leq s} \kappa_{\sigma, \theta_\sigma(i)} \geq \sum_{1 \leq i \leq s} \kappa_{\sigma, i}$ for $\sigma \neq \tau$, and obtain the inequality

$$\sum_{1 \leq i \leq s} v_p(\varphi_i) - \frac{1}{e} \sum_{1 \leq i \leq s} \sum_{\sigma \in \Upsilon} \kappa_{\sigma, i} \geq \frac{1}{e} \sum_{1 \leq i \leq s} \kappa_{\tau, \theta_\tau(i)} - \kappa_{\tau, i}$$

Let r be minimal such that $\theta_\tau(s) \neq s$ (if no such s exists, we are done). In that case, we necessarily have $\theta_\tau(s) \geq s + 1$, and the previous inequality yields

$$\sum_{1 \leq i \leq s} v_p(\varphi_i) - \frac{1}{e} \sum_{1 \leq i \leq s} \sum_{\sigma \in \Upsilon} \kappa_{\sigma, i} \geq \frac{\kappa_{\tau, s+1} - \kappa_{\tau, s}}{e}$$

but the hypothesis implies that the left hand side is less than $\min_j (\kappa_{\tau, j+1} - \kappa_{\tau, j}) / e$, and we get a contradiction. \square

Theorem 4.2.2. *Let Π be an irreducible automorphic representation of $\mathbf{G}(\mathbb{A}_F)$ having Iwahori invariants at all the places of F above p , and having invariants under an open subgroup U of $\mathbf{G}(\mathbb{A}_{F, f}^{(p)})$. Let N be an integer. There exists an automorphic representation Π' of $\mathbf{G}(\mathbb{A}_F)$ such that:*

- Π' is unramified at the places above p , and has invariants under U ;
- The restriction of $\rho_{\iota_p, \iota_\infty}(\Pi')$ to the decomposition group at any place above p is either irreducible or the sum of an Artin character and an irreducible representation of dimension $2n$;
- For all g in G_F , $\text{Tr}(\rho_{\iota_p, \iota_\infty}(\Pi')(g)) \equiv \text{Tr}(\rho_{\iota_p, \iota_\infty}(\Pi)(g)) \pmod{p^N}$.

Proof. We will write $\Pi' \equiv \Pi \pmod{p^N}$ for the last property.

Recall that for v a place of F above p , there are elements $\phi_{v,1}, \dots, \phi_{v,n} \in \mathcal{O}(\mathcal{X})^\times$ such that for any unramified classical point $x \in \mathcal{X}(\overline{\mathbb{Q}}_p)$ refining an automorphic representation Π , the filtered φ -module associated with the crystalline representation $\rho_{\iota_p, \iota_\infty}(\Pi)|_{G_{F_v}}$ has φ^{f_v} -eigenvalues

$$\left(\phi_{v,-n}(x) q_v^{-n} \prod_{\sigma} \sigma(\varpi_v)^{k_{v,\sigma,-1}}, \dots, \phi_{v,-1}(x) q_v^{-1} \prod_{\sigma} \sigma(\varpi_v)^{k_{v,\sigma,-n}}, 1, \right. \\ \left. \phi_{v,1}(x) q_v \prod_{\sigma} \sigma(\varpi_v)^{k_{v,\sigma,n}}, \dots, \phi_{v,n}(x) q_v^n \prod_{\sigma} \sigma(\varpi_v)^{k_{v,\sigma,1}} \right)$$

and σ -Hodge-Tate weights

$$k_{v,\sigma,-1} - n, \dots, k_{v,\sigma,-n} - 1, 0, k_{v,\sigma,n} + 1, \dots, k_{v,\sigma,1} + n$$

In the following if x_b or x'_b is a classical point, $k_{v,\sigma,i}^{(b)}$ will be the weights defining $w(x_b)$. The representation Π corresponds to at least one point x of the eigenvariety \mathcal{X} for \mathbf{G}' and the idempotent $e_U \otimes e_{G_0}$. By Proposition 3.2.6, and since G_F is compact, there exists a point $x_1 \in \mathcal{X}(E')$ (near x , and for some finite extension E' of E) corresponding to an

unramified, completely refinable automorphic representation Π_1 and a refinement χ , such that for any v ,

$$\frac{2}{e_v} \sum_{i=1}^n \sum_{\sigma} k_{v,\sigma,i}^{(1)} > -v_p(\phi_{v,1}(x_1) \dots \phi_{v,n}(x_1)) + 3n(n+1)f_v$$

and $\Pi_1 \equiv \Pi \pmod{p^N}$. Since Π_1 is completely refinable, there is a point $x'_1 \in \mathcal{X}(E')$ associated with the representation Π_1 and the character χ^a , where a is the element of the Weyl group acting as $-\text{Id}$ on the roots. Specifically, $\Psi_{x_1}|_{\mathcal{H}^S \otimes \mathcal{H}_S \otimes e_{G_0}} = \Psi_{x'_1}|_{\mathcal{H}^S \otimes \mathcal{H}_S \otimes e_{G_0}}$, but

$$\phi_{v,n+1-i}(x'_1) = \phi_{v,-n-1+i}(x_1) q_v^{2i+(2n+2)} \prod_{\sigma} \sigma(\varpi_v)^{-2k_{v,\sigma,i}^{(1)}}$$

for $i = 1, \dots, n$, and all places v . There exists a point $x_2 \in \mathcal{X}(E')$ (near x'_1 , and up to enlarging E') corresponding to an unramified, completely refinable automorphic representation Π_2 and a refinement, such that for any v and any $j < 0$,

$$\frac{1}{e_v} \sum_{\sigma} k_{v,\sigma,n+j}^{(2)} - k_{v,\sigma,n+j+1}^{(2)} > -v_p(\phi_{v,-j+1}(x_2)) - f_v$$

and $\Pi_2 \equiv \Pi_1 \equiv \Pi \pmod{p^N}$. Like before, since Π_2 is completely refinable, there is a point $x'_2 \in \mathcal{X}(E')$ such that $\Psi_{x_2}|_{\mathcal{H}^S \otimes \mathcal{H}_S \otimes e_{G_0}} = \Psi_{x'_2}|_{\mathcal{H}^S \otimes \mathcal{H}_S \otimes e_{G_0}}$, and

$$\begin{aligned} \phi_{v,n}(x'_2) &= \phi_{v,1}(x_2) q_v^{1-n} \prod_{\sigma} \sigma(\varpi_v)^{k_{v,\sigma,n}^{(2)} - k_{v,\sigma,1}^{(2)}} \\ \phi_{v,i}(x'_2) &= \phi_{v,i+1}(x_2) q_v \prod_{\sigma} \sigma(\varpi_v)^{k_{v,\sigma,n-i}^{(2)} - k_{v,\sigma,n-i+1}^{(2)}} \text{ for } i = 1, \dots, n-1. \end{aligned}$$

Here we used the element of the Weyl group corresponding (at each v) to the permutation

$$\begin{pmatrix} -n & -n+1 & \dots & -2 & -1 & 1 & \dots & n \\ -n+1 & -n+2 & \dots & -1 & -n & n & \dots & n-1 \end{pmatrix}.$$

Again, we can choose a point $x_3 \in \mathcal{X}(E')$ (near x'_1 , and up to enlarging E') corresponding to an unramified automorphic representation Π_3 and a refinement, such that for any v and any $\tau \in \Upsilon$,

$$\frac{1}{e_v(2n+1)} \min \left\{ k_{v,\tau,1}^{(3)} - k_{v,\tau,2}^{(3)}, \dots, k_{v,\tau,n-1}^{(3)} - k_{v,\tau,n}^{(3)}, k_{v,\tau,n}^{(3)} \right\} > \max \{0, |v_p(\phi_{v,\tau,1}(x_3))|, \dots, |v_p(\phi_{v,\tau,n}(x_3))|\}$$

and $\Pi_3 \equiv \Pi \pmod{p^N}$. We show that Π_3 has the desired properties. First we apply the previous lemma to the local Galois representations associated with Π_3 , at the places above p , which are crystalline. Since the differences $v_p(\varphi_i) - \frac{1}{e} \sum_{\sigma \in \Upsilon} \kappa_{\sigma,i}$ in the hypotheses of the lemma are equal in our case to

$$-v_p(\phi_{v,n}(x_3)), \dots, -v_p(\phi_{v,1}(x_3)), 0, v_p(\phi_{v,1}(x_3)), \dots, v_p(\phi_{v,n}(x_3)),$$

the hypotheses of the lemma are satisfied for all $\tau \in \Upsilon$. Thus if $\rho_{\nu_p, \nu_\infty}(\pi_3)|_{G_{F_v}}$ is not irreducible, there is a subset $\emptyset \subsetneq I \subsetneq \{-n, \dots, n\}$ such that if $i_1 < \dots < i_r$ are the elements of I and $j_1 < \dots < j_{2n+1-r}$ those of $J = \{-n, \dots, n\} \setminus I$,

$$\begin{aligned}
v_p(\phi_{v, i_1}(x_3)) &\geq 0 \\
v_p(\phi_{v, i_1}(x_3)) + v_p(\phi_{v, i_2}(x_3)) &\geq 0 \\
&\vdots \\
v_p(\phi_{v, i_1}(x_3)) + \dots + v_p(\phi_{v, i_r}(x_3)) &= 0 \\
v_p(\phi_{v, j_1}(x_3)) &\geq 0 \\
v_p(\phi_{v, j_1}(x_3)) + v_p(\phi_{v, j_2}(x_3)) &\geq 0 \\
&\vdots \\
v_p(\phi_{v, j_1}(x_3)) + \dots + v_p(\phi_{v, j_{2n+1-r}}(x_3)) &= 0
\end{aligned}$$

by the admissibility of the corresponding filtered φ -modules. For all i , $v_p(\phi_{v, i}(x'_2)) = v_p(\phi_{v, i}(x_3))$, so all these conditions hold also at x'_2 . Up to exchanging I and J , we can assume that $i_1 = -n$. If $j_1 < 0$,

$$v_p(\phi_{v, j_1}(x'_2)) = -v_p(\phi_{v, -j_1}(x'_2)) = -v_p(\phi_{v, -j_1+1}(x_2)) - f_v - \frac{1}{e_v} \sum_{\sigma} k_{v, \sigma, n+j_1}^{(2)} - k_{v, \sigma, n+j_1+1}^{(2)}$$

and x_2 was chosen to ensure that this quantity is negative, so we are facing a contradiction. Thus J has only nonnegative elements, and $\{-n, \dots, -1\} \subset I$. If we do not assume that $i_1 = -n$, we have in general that $\{-n, \dots, -1\}$ is contained in I or J . Similarly, suppose $i_r = n$. If $j_{2n+1-r} > 0$,

$$\begin{aligned}
v_p(\phi_{v, j_{2n+1-r}}(x'_2)) &= v_p(\phi_{v, j_{2n+1-r}}(x_2)) \\
&= v_p(\phi_{v, j_{2n+1-r}}(x_2)) + f_v + \frac{1}{e_v} \sum_{\sigma} k_{v, \sigma, n-j_{2n+1-r}}^{(2)} - k_{v, \sigma, n-j_{2n+1-r}+1}^{(2)}
\end{aligned}$$

is positive, another contradiction. Therefore $\{1, \dots, n\}$ is contained in I or J .

Assume for example that $\{-n, \dots, -1\} \subset I$ and $\{1, \dots, n\} \subset J$. In that case

$$\begin{aligned}
v_p(\phi_{v, j_1}(x_3) \dots \phi_{v, j_{2n+1-r}}(x_3)) &= v_p(\phi_{v, 1}(x_2) \dots \phi_{v, n}(x_2)) \\
&= v_p(\phi_{v, 1}(x'_1) \dots \phi_{v, n}(x'_1)) \\
&= -v_p(\phi_{v, 1}(x_1) \dots \phi_{v, n}(x_1)) + 3n(n+1)f_v \\
&\quad - \frac{2}{e_v} \sum_{i=1}^n \sum_{\sigma} k_{v, \sigma, i}^{(1)}
\end{aligned}$$

is negative, which is yet another contradiction.

As a consequence, we can conclude that I or J is equal to $\{0\}$, and this shows that at each place v of F above p , the semisimplification of $\rho_{\nu_p, \nu_\infty}(\Pi_3)|_{G_{F_v}}$ is either irreducible or the sum of an Artin character and an irreducible representation of dimension $2n$. Consequently Π_3 has the required properties. \square

5 Similar results for even orthogonal groups

In this section we explain (very) briefly how the same method as in the previous sections applies to orthogonal groups.

Let F be a totally real number field of even degree over \mathbb{Q} . Then F has an even number of 2-adic places of odd degree over \mathbb{Q}_2 , and since these are the only finite places of F at which $(-1, -1)_v = -1$ (where $(\cdot, \cdot)_v$ denotes the Hilbert symbol), $\prod_v (-1, -1)_v = 1$ where the product ranges over the finite places of F . Consequently, there is a unique quadratic form on F^4 which is positive definite at the real places of F , and split (isomorphic to $(x, y, z, t) \mapsto xy + zt$) at the finite places. It has Hasse invariant $(-1, -1)_v$ at each finite place v of F , and its discriminant is 1. As a consequence, for any integer $n \geq 1$, there is a connected reductive group \mathbf{G} over F which is compact (and connected) at the real places (isomorphic to $\mathrm{SO}_{4n}/\mathbb{R}$) and split at all the finite places (isomorphic to the split SO_{4n}). As before, we let $\mathbf{G}' = \mathrm{Res}_{\mathbb{Q}}^F \mathbf{G}$. The proofs of the existence and properties of the attached eigenvariety $\mathcal{X} \rightarrow \mathcal{W}$ are identical to the symplectic case. We could not find a result as precise as Theorem 3.2.4 in the literature, however by [Cas80, Proposition 3.5] unramified principal series are irreducible on an explicit Zariski-open subset of the unramified characters. Specifically, if $\mathrm{SO}_{4n}(F_v) = \{M \in \mathrm{M}_{4n}(F_v) \mid {}^t M J_{4n} M = J_{4n}\}$,

$$T = \left\{ \left(\begin{array}{ccccccc} x_1 & & & & & & \\ & \ddots & & & & & \\ & & x_n & & & & \\ & & & x_n^{-1} & & & \\ & & & & \ddots & & \\ & & & & & & x_1^{-1} \end{array} \right) \mid x_i \in F_v^\times \right\}$$

and P is any parabolic subgroup containing T , then for an unramified character $\chi = (\chi_1, \dots, \chi_n)$ of T (χ_i is a character of the variable x_i), $\mathrm{Ind}_P^{\mathrm{SO}_{4n}(F_v)} \chi$ is irreducible if $\chi_i(\varpi_v)^2 \neq 1$ for all i and $\chi_i(\varpi_v) \chi_j(\varpi_v)^{\pm 1} \neq 1, q_v, q_v^{-1}$ for all $i < j$. Note that this is not an equivalence.

The existence of Galois representations $\rho_{\iota_p, \iota_\infty}(\Pi)$ attached to automorphic representations Π of $\mathbf{G}(\mathbb{A}_F)$ is identical to Assumption 4.1.1. We now state the main result for orthogonal groups.

Theorem 5.0.3. *Let Π be an irreducible automorphic representation of $\mathbf{G}(\mathbb{A}_F)$ having Iwahori invariants at all the places of F above p , and having invariants under an open subgroup U of $\mathbf{G}(\mathbb{A}_{F,f}^{(p)})$. Let N be an integer. There exists an automorphic representation Π' of $\mathbf{G}(\mathbb{A}_F)$ such that:*

- Π' is unramified at the places above p , and has invariants under U ;
- The restriction of $\rho_{\iota_p, \iota_\infty}(\Pi')$ to the decomposition group at any place above p is irreducible;
- For all g in G_F , $\mathrm{Tr}(\rho_{\iota_p, \iota_\infty}(\Pi')(g)) \equiv \mathrm{Tr}(\rho_{\iota_p, \iota_\infty}(\Pi)(g)) \pmod{p^N}$.

Proof. The proof is nearly identical to that of Theorem 4.2.2. In the orthogonal case the Weyl group is a bit smaller: it is the semi-direct product of S_{2n} and a hyperplane of $(\mathbb{Z}/2\mathbb{Z})^{2n}$. Alternatively, it is the group of permutations w of $\{-2n, \dots, -1, 1, \dots, 2n\}$ such that $w(-i) = -w(i)$ for all i and $\prod_{i=1}^{2n} w(i) > 0$. The two elements of the Weyl group used in the proof of Theorem 4.2.2 have natural counterparts in this Weyl group. The only difference lies in the fact that there is no Hodge-Tate weight equal to 0 in the orthogonal case, hence the simpler conclusion “ $\rho_{\iota_p, \iota_\infty}(\Pi')|_{G_{F_v}}$ is irreducible for $v|p$ ”. \square

6 The image of complex conjugation: relaxing hypotheses in Taylor’s theorem

Let us apply the previous results to the determination of the image of the complex conjugations under the p -adic Galois representations associated with regular, algebraic, essentially self-dual, cuspidal automorphic representations of $\mathrm{GL}_n(\mathbb{A}_F)$, F totally real. Recall that these representations are constructed by “patching” representations of Galois groups of CM extensions of F , on Shimura varieties for unitary groups. The complex conjugations are lost when we restrict to CM fields. In [Tay], Taylor proves that the image of any complex conjugation is given by (the “discrete” part of) the local Langlands parameter at the corresponding real place, assuming n is odd and the Galois representation is irreducible, by constructing the complex conjugation on the Shimura datum. Of course the Galois representation associated with a cuspidal representation of GL_n is conjectured to be irreducible, but unfortunately this is (at the time of writing) still out of reach in the general case (however, see [CG] for $n \leq 5$ and [BLGGT, Theorem D] for a “density one” result).

The results of the first part of this paper allow to remove the irreducibility hypothesis in Taylor’s theorem, and to extend it to some (“half”) cases of even n , using Arthur’s endoscopic transfer. Unfortunately some even-dimensional cases are out of reach using this method, because odd-dimensional essentially self-dual cuspidal representations are (up to a twist) self-dual, whereas some even-dimensional ones are not.

Since the proof is not direct, let us outline the strategy. First we deduce the even-dimensional self-dual case from Taylor’s theorem by adding a cuspidal self-dual (with appropriate weights) representation of GL_3 , we get an automorphic self-dual representation of GL_{2n+3} which (up to base change) can be “transferred” to a discrete representation of the symplectic group in dimension $2n$. Since the associated Galois representation contains no Artin character, it can be deformed irreducibly, and Taylor’s theorem applies. Then the general odd-dimensional case is deduced from the even-dimensional one, by essentially the same method, using the eigenvariety for orthogonal groups.

Finally we prove a supplementary, non-regular case, thanks to the fact that discrete Langlands parameters for the group $\mathrm{SO}_{2n}/\mathbb{R}$ are not always discrete when seen as parameters for GL_{2n} , i.e. can correspond to a non-regular representation of $\mathrm{GL}_{2n}/\mathbb{R}$.

6.1 Regular, L-algebraic, self-dual, cuspidal representations of $\mathrm{GL}_{2n}(\mathbb{A}_F)$ having Iwahori-invariants

In this subsection \mathbf{G} will denote the symplectic group in dimension $2n+2$ defined in section 3.

The following is due to C. Mœglin and J.-L. Waldspurger.

Lemma 6.1.1. *Let K be a finite extension of \mathbb{Q}_p . Let $\phi : W_K \times \mathrm{SU}(2) \rightarrow \mathrm{SO}_{2n+3}(\mathbb{C})$ be a Langlands parameter (equivalently, a generic Arthur parameter). Assume that the subgroup $I \times \{1\}$ (I being the inertia subgroup of W_K) is contained in the kernel of ϕ .*

Then the A-packet associated with ϕ contains a representation having a non-zero vector fixed under the Iwahori subgroup of $\mathrm{Sp}_{2n+2}(K)$.

Proof. Let $\{\Pi_1, \dots, \Pi_k\}$ denote the A-packet. Since Arthur's construction of the Π_i 's is inductive for parameters trivial on the supplementary $\mathrm{SL}_2(\mathbb{C})$, and subquotients of parabolic inductions of representations having Iwahori-invariants have too, it is enough to prove the result when ϕ is discrete. Let τ be the irreducible smooth representation of $\mathrm{GL}_{2n+3}(K)$ having parameter ϕ , then $\tau \simeq \mathrm{Ind}_L^{\mathrm{GL}_{2n+3}} \sigma$, where σ is the tensor product of (square-integrable) Steinberg representations $\mathrm{St}(\chi_i, n_i)$ of $\mathrm{GL}_{n_i}(K)$ ($i \in \{1, \dots, r\}$), χ_i are unramified, auto-dual characters of K^\times (thus $\chi_i = 1$ or $(-1)^{v(\cdot)}$), and the couples (χ_i, n_i) are distinct. Here L denotes the standard parabolic associated with the decomposition $2n+3 = \sum_i n_i$. Since ϕ is self-dual, τ can be extended (not uniquely, but this will not matter for our purpose) to a representation of $\widetilde{\mathrm{GL}}_{2n+3}^+ = \mathrm{GL}_{2n+3} \rtimes \{1, \theta\}$, where

$$\theta(g) = \begin{pmatrix} & & & 1 \\ & & -1 & \\ & \ddots & & \\ 1 & & & \end{pmatrix} {}_t g^{-1} \begin{pmatrix} & & & 1 \\ & & -1 & \\ & \ddots & & \\ 1 & & & \end{pmatrix}$$

Let also $\widetilde{\mathrm{GL}}_{2n+3} = \mathrm{GL}_{2n+3} \rtimes \theta$.

Let N_0 be the number of i such that n_i is odd, and for $j \geq 1$ let N_j be the number of i such that $n_i \geq 2j$. Then $N_0 + 2 \sum_{j \geq 1} N_j = 2n+3$, and if s is maximal such that $N_s > 0$, we let

$$M = \mathrm{GL}_{N_s} \times \dots \times \mathrm{GL}_{N_1} \times \mathrm{GL}_{N_0} \times \mathrm{GL}_{N_1} \times \dots \times \mathrm{GL}_{N_s}$$

which is a θ -stable Levi subgroup of GL_{2n+3} , allowing us to define \widetilde{M}^+ and \widetilde{M} . Since the standard (block upper triangular) parabolic containing M is also stable under θ , τ_M is naturally a representation of \widetilde{M}^+ , denoted by $\tau_{\widetilde{M}}$. The constituents of the semi-simplification of $\tau_{\widetilde{M}}$ either stay irreducible when restricted to M , in which case they are of the form $\sigma_1 \otimes \sigma_0 \otimes \theta(\sigma_1)$ where σ_1 is a representation of $\mathrm{GL}_{N_s} \times \dots \times \mathrm{GL}_{N_1}$ and σ_0 is a representation of $\widetilde{\mathrm{GL}}_{N_0}$; or they are induced from M to \widetilde{M}^+ , and the restriction of their character to \widetilde{M} is zero. Since we are precisely interested in that character, we can forget about the second case. By the geometrical lemma,

$$\tau_M^{\mathrm{ss}} \simeq \bigoplus_{w \in W^{L,M}} \mathrm{Ind}_{M \cap w(L)}^M w(\sigma_{L \cap w^{-1}M})$$

where $W^{L,M}$ is the set of $w \in S_{2n+3}$ such that w is increasing on $I_1 = \{1, \dots, n_1\}$, $I_2 = \{n_1 + 1, \dots, n_1 + n_2\}$, etc. and w^{-1} is increasing on $J_{-s} = \{1, \dots, N_s\}$, $J_{-s+1} = \{N_s + 1, \dots, N_s + N_{s-1}\}$, etc. Fix the irreducible representation of $\mathrm{GL}_{N_s} \times \dots \times \mathrm{GL}_{N_1}$

$$\sigma_1 = \bigotimes_{j=1}^s \mathrm{Ind}_{T_j}^{\mathrm{GL}_{N_j}} \bigotimes_{i \mid n_i \geq 2j} \chi_i |\cdot|^{j-\nu_i}$$

where T_j is the standard maximal torus of GL_{N_j} , $\nu_i = \begin{cases} 0 & n_i \text{ odd} \\ 1/2 & n_i \text{ even} \end{cases}$.

There is a unique w such that $\mathrm{Ind}_{M \cap w(L)}^M w(\sigma_{L \cap w^{-1}M})$ admits a subquotient of the form $\sigma_1 \otimes \sigma_0 \otimes \theta(\sigma_1)$ as above, moreover $\mathrm{Ind}_{M \cap w(L)}^M w(\sigma_{L \cap w^{-1}M})$ is irreducible, and

$$\sigma_1 = \mathrm{Ind}_{T_0}^{\mathrm{GL}_{N_0}} \bigotimes_{i \mid n_i \text{ odd}} \chi_i$$

Specifically, w maps the first element of I_i in $J_{-\lfloor (n_i+1)/2 \rfloor}$, the second in $J_{-\lfloor (n_i+1)/2 \rfloor} + 1$, \dots , the central element (if n_i is odd) in J_0 , etc.

Let M' be the parabolic subgroup of Sp_{2n+2}/K corresponding to M , i.e.

$$M' = \mathrm{GL}_{N_s} \times \dots \times \mathrm{GL}_{N_1} \times \mathrm{Sp}_{N_0-1}$$

By [Art11, 2.2.6], $\sum_i \mathrm{Tr} \Pi_i$ is a stable transfer of $\mathrm{Tr}_{\mathrm{GL}_{2n+3}^+} \tau$. By [MW06, Lemme 4.2.1] (more accurately, the proof of the lemma),

$$\sum_i \mathrm{Tr} ((\Pi_i)_{M'}^{\mathrm{ss}}[\sigma_1])$$

is a stable transfer of $\mathrm{Tr} \left(\tau_{M'}^{\mathrm{ss}}[\sigma_1] \right)$ (where $\cdot[\cdot]$ denotes the isotypical component on the factor $\mathrm{GL}_{N_s} \times \dots \times \mathrm{GL}_{N_1}$).

Since $\tau_{M'}^{\mathrm{ss}}[\sigma_1] = \sigma_1 \otimes \sigma_0 \otimes \theta(\sigma_1)$, the stable transfer of $\mathrm{Tr} \left(\tau_{M'}^{\mathrm{ss}}[\sigma_1] \right)$ is equal to the product of $\mathrm{Tr}(\sigma_1)$ and $\sum_l \mathrm{Tr} \Pi'_l$ where the Π'_l are the elements of the A-packet associated with the parameter

$$\bigoplus_{i \mid n_i \text{ odd}} \chi_i$$

At least one representation Π'_l is unramified for some hyperspecial compact subgroup of $\mathrm{Sp}_{N_0-1}(K)$, and so a Jacquet module of a Π_i contains a nonzero vector fixed by an Iwahori subgroup. This proves that at least one of the Π_i has Iwahori-invariants. \square

Assumption 6.1.2. *Let F_0 be a totally real field, and let π be a regular, L -algebraic, self-dual, cuspidal (RLASDC) representation of $\mathrm{GL}_{2n}(\mathbb{A}_{F_0})$. Assume that for any place $v|p$ of F_0 , π_v has vectors fixed under an Iwahori subgroup of $\mathrm{GL}_{2n}(\mathbb{A}_{F_0,v})$. Then there exists a RLASDC representation π_0 of $\mathrm{GL}_3(\mathbb{A}_{F_0})$, a totally real extension F/F_0 which is trivial, quadratic or quartic, and an automorphic representation Π of $\mathbf{G}(\mathbb{A}_F)$ such that*

1. For any place $v|p$ of F_0 , $\pi_{0,v}$ is unramified.
2. $\mathrm{BC}_{F/F_0}(\pi)$ and $\mathrm{BC}_{F/F_0}(\pi_0)$ remain cuspidal.

3. For any place v of F above p , Π_v has invariants under the action of the Iwahori subgroup G_0 of $\mathbf{G}(F_v)$.
4. For any finite place v of F such that $\mathrm{BC}_{F/F_0}(\pi)_v$ and $\mathrm{BC}_{F/F_0}(\pi_0)_v$ are unramified, Π_v is unramified, and via the inclusion $\mathrm{SO}_{2n+3}(\mathbb{C}) \hookrightarrow \mathrm{GL}_{2n+3}(\mathbb{C})$, the Satake parameter of Π_v is equal to the direct sum of those of $\mathrm{BC}_{F/F_0}(\pi)_v$ and $\mathrm{BC}_{F/F_0}(\pi_0)_v$.

Let us comment briefly on the proof to come. First we construct π_0 . Let δ be a cuspidal automorphic representation of PGL_2/F_0 which is unramified at the p -adic places, Steinberg at the ℓ -adic places for some arbitrary prime $\ell \neq p$, and whose local Langlands parameters at the real places are of the form $\mathrm{Ind}_{W_{\mathbb{C}}}^{W_{\mathbb{R}}}(z \mapsto (z/\bar{z})^a)$ where a is a half-integer big enough with respects to their analogues appearing in the local Langlands parameters of π . Such a representation exists thanks to [Clo86, Theorem 1B]. Let π_0 be the automorphic representation of GL_3/F_0 obtained by functoriality from δ through the adjoint representation of $\widehat{\mathrm{PGL}}_2(\mathbb{C}) = \mathrm{SL}_2(\mathbb{C})$ on its Lie algebra. The representation π_0 exists and is cuspidal by [GJ78, Theorem 9.3]. The condition at the ℓ -adic places ensures that no nontrivial twist of δ (seen as a representation of GL_2/F_0) is isomorphic to δ , and the cuspidality of π_0 follows. We can twist π_0 by the central character of π , to ensure that $\pi \oplus \pi_0$ has trivial central character. Clearly π_0 is a RLASDC representation of GL_3/F_0 .

Note that for $\mathrm{BC}_{F/F_0}(\pi)$ and $\mathrm{BC}_{F/F_0}(\pi_0)$ to remain cuspidal, it is enough for F/F_0 to be totally ramified above a finite place of F_0 at which π and π_0 are unramified. To begin with one can choose such a quadratic extension of F_0 , in order to define \mathbf{G} . The automorphic representation $\Psi := \mathrm{BC}_{F/F_0}(\pi) \oplus \mathrm{BC}_{F/F_0}(\pi_0)$ can be seen as a global, orthogonal parameter. This determines a global packet P_{Ψ} of representations of $\mathbf{G}(\mathbb{A}_F)$, and Arthur's results shall attach to each $\Pi \in P_{\Psi}$ a character of $S_{\Psi} \simeq \mathbb{Z}/2\mathbb{Z}$, and characterize the automorphic Π 's as the ones whose character is trivial. We can choose the components Π_v at the finite places of F not lying above p to be associated with a trivial character of S_{Ψ_v} , and taking a quadratic extension split above the p -adic and real places of F (at which Π_v is imposed) allows to “double” the contribution of the characters, thus yielding a trivial global character.

Proposition 6.1.3. *Let F be a totally real field, and let π be a regular, L -algebraic, self-dual, cuspidal representation of $\mathrm{GL}_{2n}(\mathbb{A}_F)$. Suppose that for any place v of F above p , π_v has invariants under an Iwahori subgroup. Then for any complex conjugation $c \in G_F$, $\mathrm{Tr}(\rho_{\iota_p, \iota_{\infty}}(\pi)(c)) = 0$.*

Proof. By the previous assumption, up to a (solvable) base change to a totally real extension (which only restricts the Galois representation to this totally real field, so that we get even more complex conjugations), we can take a RLASDC representation π_0 of $\mathrm{GL}_3(\mathbb{A}_F)$ and transfer $\pi \oplus \pi_0$ to an automorphic representation Π of $\mathbf{G}(\mathbb{A}_F)$. The representation Π defines (at least) one point x of the eigenvariety \mathcal{X} defined by \mathbf{G} (and by an open subgroup U of $\mathbf{G}(\mathbb{A}_{F,f}^{(p)})$). Of course, by the Čebotarev density theorem and the compatibility of the transfer at the unramified places, the representation associated with Π is equal to $\rho_{\iota_p, \iota_{\infty}}(\pi) \oplus \rho_{\iota_p, \iota_{\infty}}(\pi)$. Since the Hodge-Tate weights of $\rho_{\iota_p, \iota_{\infty}}(\pi)|_{G_{F_v}}$ are non-zero

for any place $v|p$, $\rho_{\ell_p, \ell_\infty}(\pi)$ does not contain an Artin character. By [BR92], $\rho_{\ell_p, \ell_\infty}(\pi_0)$ is irreducible and thus does not contain any character. There are only finitely many Artin characters taking values in $\{\pm 1\}$ and unramified at all the finite places at which Π is unramified. For any such character η , the pseudocharacter T on the eigenvariety is such that $T_x - \eta$ is not a pseudocharacter, hence we can find $g_{\eta, 1}, \dots, g_{\eta, 2n+3}$ such that

$$t_\eta := \sum_{\sigma \in S_{2n+3}} (T_x - \eta)_\sigma(g_{\eta, 1}, \dots, g_{\eta, 2n+3}) \neq 0$$

Let us choose N greater than all the $v_p(t_\eta)$ and such that $p^N > 2n + 4$. Let Π' be an automorphic representation of $\mathbf{G}(\mathbb{A}_F)$ satisfying the requirements of Theorem 4.2.2 for this choice of N . Then the $\text{Tr}(\rho_{\ell_p, \ell_\infty}(\Pi')) - \eta$ are not pseudocharacters, thus $\rho_{\ell_p, \ell_\infty}(\Pi')$ does not contain an Artin character and by Theorem 4.2.2 it is irreducible. This Galois representation is (by construction in the proof of Corollary 4.1.3) the direct sum of representations associated with cuspidal representations. Since it is irreducible, there is only one of them, and it has the property that its associated Galois representations is irreducible, so that the theorem of [Tay] can be applied: for any complex conjugation $c \in G_F$, $\text{Tr}(\rho_{\ell_p, \ell_\infty}(\Pi')(c)) = \pm 1$. Since $\det \rho_{\ell_p, \ell_\infty}(\Pi') = 1$, $\text{Tr}(\rho_{\ell_p, \ell_\infty}(\Pi')(c)) = (-1)^{n+1}$.

As $p^N > 2n+4$ and $|\text{Tr}(\rho_{\ell_p, \ell_\infty}(\Pi)(c)) - \text{Tr}(\rho_{\ell_p, \ell_\infty}(\Pi')(c))| \leq 2n+4$, we can conclude that $\text{Tr}(\rho_{\ell_p, \ell_\infty}(\Pi)(c)) = (-1)^{n+1}$, and hence that $\text{Tr}(\rho_{\ell_p, \ell_\infty}(\pi)(c)) + \text{Tr}(\rho_{\ell_p, \ell_\infty}(\pi_0)(c)) = (-1)^{n+1}$. We also know that $\det \rho_{\ell_p, \ell_\infty}(\pi_0) = \det \rho_{\ell_p, \ell_\infty}(\pi)(c) = (-1)^n$, and that $\text{Tr}(\rho_{\ell_p, \ell_\infty}(\pi_0)(c)) = \pm 1$ by Taylor's theorem, from which we can conclude that $\text{Tr}(\rho_{\ell_p, \ell_\infty}(\pi_0)(c)) = (-1)^{n+1}$. Thus $\text{Tr}(\rho_{\ell_p, \ell_\infty}(\pi)(c)) = 0$. \square

6.2 Regular, L-algebraic, self-dual, cuspidal representations of $\text{GL}_{2n+1}(\mathbb{A}_F)$ having Iwahori-invariants

In this subsection, \mathbf{G} is the orthogonal reductive group defined in section 5, of dimension $2n + 2$ if n is odd, $2n + 4$ if n is even.

Lemma 6.2.1. *Let K be a finite extension of \mathbb{Q}_p . Let $\phi : W_K \times \text{SU}(2) \rightarrow \text{SO}_{2m}(\mathbb{C})$ be a Langlands parameter. Assume that the subgroup $I \times \{1\}$ (I being the inertia subgroup of W_K) is contained in the kernel of ϕ .*

Then the packet of representations of the split group $\text{SO}_{2m}(K)$ associated with ϕ by Arthur contains a representation having a non-zero vector fixed under the Iwahori subgroup.

Proof. Of course this result is very similar to 6.1.1. However Mœglin and Waldspurger have not put their lemma in writing in this case, and the transfer factors are no longer trivial, so that one needs to modify the definition of “stable transfer”. For this one needs to use the transfer factors $\Delta_{\widetilde{\text{GL}}_{2m}, \text{SO}_{2m}}(\cdot, \cdot)$ defined in [KS99]. They depend in general on the choice of an inner class of inner twistings [KS99, 1.2] (in our case an inner class of isomorphisms between GL_{2m}/K and its quasi-split inner form defined over \bar{K} , which we just take to be the identity), and a Whittaker datum of the quasi-split inner form. Arthur chooses the standard splitting of GL_{2m} and an arbitrary character $K \rightarrow \mathbb{C}^\times$, but this will not matter to us since both GL_{2m} and SO_{2m} are *split*, so that the factor $\langle z_J, s_J \rangle$ of

[KS99, 4.2] (by which the transfer factors are multiplied when another splitting is chosen) is trivial. Indeed to compute this factor we can choose the split torus T_H of SO_{2m}/K , which is a norm group (see [KS99, Lemma 3.3B]) for the split torus T of GL_{2m}/K , and thus, using the notations of [KS99, 4.2], T^x is split and $H^1(K, T^x)$ is trivial, so that $z' = 1$ (z_J is the image of z' in $H^1(K, J)$, so that it is trivial). Since both groups are split the ϵ -factor of [KS99, 5.3] is also trivial, so the transfer factors are canonical.

Let $H = \mathrm{SO}_{2m}(K)$, $\tilde{\tau}$ the representation of $\widetilde{\mathrm{GL}}_{2m}^+$ associated with ϕ , and τ^H the sum of the elements of the packet associated by Arthur with ϕ . This packet is in general bigger than the conjectural A-packet, because it corresponds to the $O_{2m}(\mathbb{C})$ -conjugacy class of the parameter ϕ , and thus it can be the disjoint union of two A-packets. Arthur shows ([Art11, 8.3]) that the following character identity holds:

$$\Theta_{\tau^H}(h)\Delta(h, g) = \Theta_{\tilde{\tau}}(g) \quad (6.2.1)$$

whenever the stable $O_{2m}(K)$ -conjugacy class h in $\mathrm{SO}_{2m}(K)$ is the (in this case, there is at most one) norm of the conjugacy class g in $\widetilde{\mathrm{GL}}_{2m}(K)$, both assumed to be strongly $\widetilde{\mathrm{GL}}_{2m}^+$ -regular. Here the terms Δ_{IV} are left out of the product defining the transfer factors Δ , as in [MW06]. This is the natural generalization of the notion of “stable transfer” of [MW06].

Let

$$M = \mathrm{GL}_{N_s} \times \dots \times \mathrm{GL}_{N_1} \times \mathrm{GL}_{N_0} \times \mathrm{GL}_{N_1} \times \dots \times \mathrm{GL}_{N_s}$$

be a θ -stable Levi subgroup of GL_{2m} , and $M' = \mathrm{GL}_{N_s} \times \dots \times \mathrm{GL}_{N_1} \times \mathrm{SO}_{N_0}$ the corresponding parabolic subgroup of SO_{2m} . To mimic the proof of 6.1.1, we only need to show that $\mathrm{Tr}(\tau_{M'}^H)$ is a stable transfer of $\mathrm{Tr}_{\widetilde{M}}(\tau_{\widetilde{M}})$, where “stable transfer” has the above meaning, that is the character identity 6.2.1 involving transfer factors. Note that \widetilde{M}^+ has a factor $\mathrm{GL}_{m-N_0/2} \times \mathrm{GL}_{m-N_0/2}$ together with the automorphism $\theta(a, b) = (\theta(b), \theta(a))$, for which the theory of endoscopy is trivial: θ -conjugacy classes are in bijection with conjugacy classes in $\mathrm{GL}_{m-N_0/2}$ (over K or \bar{K}) via $(a, b) \mapsto a\theta(b)$ and the θ -invariant irreducible representations are the ones of the form $\sigma \otimes \theta(\sigma)$.

So we need to check that if $g = (g_1, g_0)$ is a strongly regular $\mathrm{GL}_{2m}(K)$ -conjugacy class in $\widetilde{\mathrm{GL}}_{2m}(K)$ determined by a conjugacy class g_1 in $\mathrm{GL}_{m-N_0/2}(K)$ and a $\mathrm{GL}_{N_0}(K)$ -conjugacy class g_0 in $\widetilde{\mathrm{GL}}_{N_0}(K)$, and if h_0 is the $O_{2m}(\bar{K})$ -conjugacy class in $\mathrm{SO}_{2m}(K)$ corresponding to g_0 , then

$$\Delta_{\widetilde{\mathrm{GL}}_{N_0}, \mathrm{SO}_{N_0}}(h_0, g_0) = \Delta_{\widetilde{\mathrm{GL}}_{2m}, \mathrm{SO}_{2m}}((g_1, h_0), (g_1, g_0))$$

Fortunately the transfer factors have been computed by Waldspurger in [Wal10]. We recall his notations and formulas. The conjugacy class g_1 , being regular enough, is parametrized by a finite set I_1 , a collection of finite extensions $K_{\pm i}$ of K for $i \in I_1$, and (regular enough, i.e. generating $K_{\pm i}$ over K) elements $x_{i,1} \in K_{\pm i}$. As in [Wal10], g_0 is parametrized by a finite set I_0 , finite extensions $K_{\pm i}$ of K , $K_{\pm i}$ -algebras K_i , and $x_i \in K_i$. Each K_i is either a quadratic field extension of $K_{\pm i}$ or $K_{\pm i} \times K_{\pm i}$, and x_i is determined only modulo $N_{K_i/K_{\pm i}} K_i^\times$. Then g is parametrized by $I = I_1 \sqcup I_0$, with $K_i = K_{\pm i} \times K_{\pm i}$ and $x_i = (x_{i,1}, 1)$ for $i \in I_1$, and the same data for I_0 . Let τ_i be the non-trivial $K_{\pm i}$ -automorphism of K_i ,

and $y_i = -x_i/\tau_i(x_i)$. Let I^* be the set of $i \in I$ such that K_i is a field (so $I^* \subset I_0$). For any $i \in I$, let Φ_i be the set of K -morphisms $K_i \rightarrow \bar{K}$, and let $P_I(T) = \prod_{i \in I} \prod_{\phi \in \Phi_i} (T - \phi(y_i))$. Define P_{I_0} similarly. For $i \in I^*$ (resp. I_0^*), let $C_i = x_i^{-1} P'_I(y_i) P_I(-1) y_i^{1-m} (1 + y_i)$ (resp. $C_{i,0} = x_i^{-1} P'_{I_0}(y_i) P_{I_0}(-1) y_i^{1-m} (1 + y_i)$). We have dropped the factor η of [Wal10, 1.10], because as remarked above, the transfer factors do not depend on the chosen splitting. Observe also that the factors computed by Waldspurger are really the factors Δ_0/Δ_{IV} of [KS99, 5.3], but the ϵ factor is trivial so they are complete.

Waldspurger shows that

$$\Delta_{\widetilde{\mathrm{GL}}_{2m}, \mathrm{SO}_{2m}}((g_1, h_0), (g_1, g_0)) = \prod_{i \in I^*} \mathrm{sign}_{K_i/K_{\pm i}}(C_i)$$

where $\mathrm{sign}_{K_i/K_{\pm i}}$ is the nontrivial character of $K_{\pm i}^\times/N_{K_i/K_{\pm i}}K_i^\times$. We are left to show that $\prod_{i \in I^*} \mathrm{sign}_{K_i/K_{\pm i}}(C_i/C_{i,0}) = 1$.

$$\begin{aligned} C_i/C_{i,0} &= y_i^{N_0/2-m} \prod_{j \in I_1} \prod_{\phi \in \Phi_j} (y_i - \phi(y_j)) (-1 - \phi(y_j)) \\ &= \prod_{j \in I_1} \prod_{\phi \in \Phi_{\pm j}} y_i^{-1} (y_i + \phi(x_{j,1})) (y_i + \phi(x_{j,1})^{-1}) (\phi(x_{j,1}) - 1) (\phi(x_{j,1})^{-1} - 1) \\ &= (-1)^{m-N_0/2} N_{K_i/K_{\pm i}} \left(\prod_{j \in I_1} \prod_{\phi \in \Phi_{\pm j}} (y_i + \phi(x_{j,1})) (\phi(x_{j,1})^{-1} - 1) \right) \end{aligned}$$

where $\Phi_{\pm j}$ is the set of K -morphisms $K_{\pm j} \rightarrow \bar{K}$. Thus

$$\begin{aligned} \prod_{i \in I^*} \mathrm{sign}_{K_i/K_{\pm i}}(C_i/C_{i,0}) &= \prod_{i \in I^*} \mathrm{sign}_{K_i/K_{\pm i}}|_{K^\times} \left((-1)^{m-N_0/2} \right) \\ &= 1 \end{aligned}$$

since $\prod_{i \in I^*} \mathrm{sign}_{K_i/K_{\pm i}}|_{K^\times}$ is easily checked to be equal to the Hilbert symbol with the discriminant of our special orthogonal group, which is 1 (this is the condition for g_0 to have a norm in the special orthogonal group). \square

Assumption 6.2.2. *Let F_0 be a totally real field, and let π be a regular, L -algebraic, self-dual, cuspidal representation of $\mathrm{GL}_{2n+1}(\mathbb{A}_{F_0})$. Assume that for any place $v|p$ of F_0 , π_v has vectors fixed under the Iwahori. Then there exists a RLASDC representation π_0 of $\mathrm{GL}_1(\mathbb{A}_{F_0})$ if n is odd (resp. $\mathrm{GL}_3(\mathbb{A}_{F_0})$ if n is even), a totally real extension F/F_0 which is trivial or quadratic, and an automorphic representation Π of $\mathbf{G}(\mathbb{A}_F)$ such that*

1. *For any place $v|p$ of F_0 , $\pi_{0,v}$ is unramified.*
2. *$\mathrm{BC}_{F/F_0}(\pi)$ and $\mathrm{BC}_{F/F_0}(\pi_0)$ remain cuspidal.*
3. *For any place v of F above p , Π_v has invariants under the action of the Iwahori subgroup of $\mathbf{G}(F_v)$.*
4. *For any finite place v of F such that $\mathrm{BC}_{F/F_0}(\pi)_v$ and $\mathrm{BC}_{F/F_0}(\pi_0)_v$ are unramified, Π_v is unramified, and via the inclusion $\mathrm{SO}_{2n+2}(\mathbb{C}) \hookrightarrow \mathrm{GL}_{2n+2}(\mathbb{C})$ (resp. $\mathrm{SO}_{2n+4}(\mathbb{C}) \hookrightarrow \mathrm{GL}_{2n+2}(\mathbb{C})$), the Satake parameter of Π_v is equal to the direct sum of those of $\mathrm{BC}_{F/F_0}(\pi)_v$ and $\mathrm{BC}_{F/F_0}(\pi_0)_v$.*

This is very similar to Assumption 6.1.2. In fact in this case the group S_Ψ is trivial, which explains why it is enough to take a quadratic extension of F_0 . This is necessary to be able to define the group \mathbf{G} . The crucial observation is that the local Langlands parameters of $\mathrm{BC}_{F/F_0}(\pi) \oplus \mathrm{BC}_{F/F_0}(\pi_0)$ at the infinite places correspond to parameters for the compact groups $\mathrm{SO}_{2n+2}/\mathbb{R}$ (resp. SO_{2n+4}). These parameters are of the form

$$\epsilon^n \oplus \bigoplus_{i=1}^n \mathrm{Ind}_{W_{\mathbb{R}}}^{W_{\mathbb{C}}} (z \mapsto (z/\bar{z})^{r_i})$$

($r_1 > \dots > r_n > 0$) for $\mathrm{BC}_{F/F_0}(\pi)$, and

$$\begin{cases} 1 & \text{if } n \text{ is odd} \\ \epsilon \oplus \mathrm{Ind}_{W_{\mathbb{R}}}^{W_{\mathbb{C}}} (z \mapsto (z/\bar{z})^r) & \text{if } n \text{ is even} \end{cases}$$

so that the direct sum of the two is always of the form

$$1 \oplus \epsilon \oplus \bigoplus_{i=1}^{k-1} \mathrm{Ind}_{W_{\mathbb{R}}}^{W_{\mathbb{C}}} (z \mapsto (z/\bar{z})^{r_i})$$

for distinct, positive r_i . This is the Langlands parameter corresponding to the representation of $\mathrm{SO}_{2k}(\mathbb{R})$ having highest weight $\sum_{i=1}^k (r_i - (k-i))e_i$ with $r_k = 0$, where the root system consists of the $\pm e_i \pm e_j$ ($i \neq j$) and the simple roots are $e_1 - e_2, \dots, e_{k-1} - e_k, e_{k-1} + e_k$.

Note that, contrary to the symplectic case, there is one outer automorphism of the even orthogonal group, and so there may be two choices for the Satake parameters of Π_v , mapping to the same conjugacy class in the general linear group. Fortunately we only need the existence.

Proposition 6.2.3. *Let F be a totally real field, and let π be an L -algebraic, self-dual, cuspidal representation of $\mathrm{GL}_{2n+1}(\mathbb{A}_F)$. Suppose that for any place v of F above p , π_v has invariants under an Iwahori. Then for any complex conjugation $c \in G_F$, $\mathrm{Tr}(\rho_{\iota_p, \iota_\infty}(\pi)(c)) = \pm 1$.*

Proof. The proof is similar to that of Proposition 6.1.3. We use the previous assumption to be able to assume (after base change) that there is a representation π_0 (of $\mathrm{GL}_1(\mathbb{A}_F)$ if n is odd, $\mathrm{GL}_3(\mathbb{A}_F)$ if n is even) such that $\pi \oplus \pi_0$ transfers to an automorphic representation Π of $\mathbf{G}(\mathbb{A}_F)$, with compatibility at the unramified places. The representation Π has Iwahori-invariants at the p -adic places of F , and thus it defines a point of the eigenvariety \mathcal{X} associated with \mathbf{G} (and an idempotent defined by an open subgroup of $\mathbf{G}(\mathbb{A}_{F,f}^{(p)})$). By Theorem 5.0.3, Π is congruent (at all the complex conjugations, and modulo arbitrarily big powers of p) to another automorphic representation Π' of \mathbf{G} , and $\rho_{\iota_p, \iota_\infty}(\Pi')$ is irreducible. Hence $\rho_{\iota_p, \iota_\infty}(\Pi') = \rho_{\iota_p, \iota_\infty}(\pi')$ for some RLASDC π' of $\mathrm{GL}_{2k}(\mathbb{A}_F)$, which is unramified at all the p -adic places of F , and we can apply Proposition 6.1.3 to π' . This proves that $\mathrm{Tr}(\rho_{\iota_p, \iota_\infty}(\pi)(c)) = -\mathrm{Tr}(\rho_{\iota_p, \iota_\infty}(\pi_0)(c)) = \pm 1$. \square

solvable extension F''/F' such that $\mathrm{BC}_{F''/F'}(\pi)$ has Iwahori invariants at all the places of F'' above p . In general $\mathrm{BC}_{F''/F'}(\pi)$ is not cuspidal, but only induced by cuspids: $\mathrm{BC}_{F''/F'}(\pi) = \pi_1 \boxplus \dots \boxplus \pi_k$. However it is self-dual, and the particular form of the Langlands parameters at the infinite places imposes that all π_i be self-dual. We can then apply Propositions 6.1.3 and 6.2.3 to the π_i , and conclude by induction that for any complex conjugation $c \in G_F$, the conjugacy class of $\rho_{\nu_p, \nu_\infty}(\pi)(c)$ is given by the recipe found in [BG10, Lemma 2.3.2], that is to say $|\mathrm{Tr} \rho_{\nu_p, \nu_\infty}(\pi)(c)| \leq 1$. \square

Remark 6.3.3. *The case n even, $\eta_\infty(-1) = (-1)^{q+1}$ is trivial. The case n even, q odd and $\eta_\infty(-1) = -1$ remains open.*

For the sake of clarity, we state the theorem using the more common normalization of \mathbb{C} -algebraic representations.

Theorem 6.3.4. *Let $n \geq 2$, F a totally real number field, π a regular, algebraic, essentially self-dual, cuspidal representation of $\mathrm{GL}_n(\mathbb{A}_F)$, such that $\pi^\vee \simeq \eta |\det|^q \pi$, where η is an Artin character. Suppose that one of the following conditions holds*

1. n is odd.
2. n is even, q is odd, and $\eta_\infty(-1) = 1$.

Then for any complex conjugation $c \in G_F$, $|\mathrm{Tr}(r_{\nu_p, \nu_\infty}(\pi)(c))| \leq 1$.

Proof. Apply the previous theorem to $\pi |\det|^{(n-1)/2}$. \square

6.4 A supplementary, non-regular case

In this subsection \mathbf{G} is the orthogonal group of section 5.

Assumption 6.4.1. *Let F_0 be a totally real field, and let π be an L -algebraic, self-dual, cuspidal representation of $\mathrm{GL}_{2n}(\mathbb{A}_{F_0})$. Assume that for any place $v|p$ of F_0 , π_v has vectors fixed under the Iwahori, and that for any real place v of F_0 ,*

$$\mathcal{L}\mathcal{L}(\pi_v) \simeq \bigoplus_{i=1}^n \mathrm{Ind}_{W_{\mathbb{R}}}^{W_{\mathbb{C}}} (z \mapsto (z/\bar{z})^{r_i})$$

where $r_n > \dots > r_1 \geq 0$ are integers (note that π is not regular if $r_1 = 0$). Then there exists a totally real extension F/F_0 which is trivial or quadratic, and an automorphic representation Π of $\mathbf{G}(\mathbb{A}_F)$ such that

1. $\mathrm{BC}_{F/F_0}(\pi)$ remains cuspidal.
2. For any place v of F above p , Π_v has invariants under the action of the Iwahori subgroup of $\mathbf{G}(F_v)$.
3. For any finite place v of F such that $\mathrm{BC}_{F/F_0}(\pi)_v$ is unramified, Π_v is unramified, and via the inclusion $\mathrm{SO}_{2n+2}(\mathbb{C}) \hookrightarrow \mathrm{GL}_{2n}(\mathbb{C})$, the Satake parameter of Π_v is equal to the one of $\mathrm{BC}_{F/F_0}(\pi)_v$.

Of course this is very similar to Assumptions 6.1.2 and 6.2.2, and as in the latter case the group S_Ψ is trivial.

For L-algebraic, self-dual, cuspidal automorphic representations of GL_{2n} having “almost regular” Langlands parameter at the archimedean places as above, the corresponding p -adic Galois representation is known to exist by [Gol]. Exactly as in the previous subsection, we have the following:

Theorem 6.4.2. *Let $n \geq 2$, F a totally real number field, π an L-algebraic, essentially self-dual, cuspidal representation of $\mathrm{GL}_{2n}(\mathbb{A}_F)$, such that $\pi^\vee \simeq \eta\pi$, where η is an Artin character. Assume that at any real place v of F , $\eta_v(-1) = 1$ and*

$$\mathcal{LL}(\pi_v) \simeq \bigoplus_{i=1}^n \mathrm{Ind}_{W_{\mathbb{R}}}^{W_{\mathbb{C}}} (z \mapsto (z/\bar{z})^{r_i})$$

where $r_n > \dots > r_1 \geq 0$ are integers. Then for any complex conjugation $c \in G_F$, $\mathrm{Tr}(\rho_{\iota_p, \iota_\infty}(\pi)(c)) = 0$.

Proof. Identical to that of Theorem 6.3.2. □

References

- [Art06] James Arthur, *A note on l -packets*, Pure and Applied Mathematics Quarterly **2** (2006), no. 1, 199–217, Special Issue: In honor of John H. Coates.
- [Art11] ———, *The Endoscopic Classification of Representations: Orthogonal and Symplectic groups*, TODO, 2011.
- [BC11] Joël Bellaïche and Gaëtan Chenevier, *The sign of galois representations attached to automorphic forms for unitary groups*, Compositio Math. **147** (2011), 1337–1352.
- [BG10] Kevin Buzzard and Toby Gee, *The conjectural connections between automorphic representations and Galois representations*, <http://arxiv.org/abs/1009.0785v1>, 2010.
- [BLGGT] Thomas Barnet-Lamb, Toby Gee, David Geraghty, and Richard Taylor, *Potential automorphy and change of weight*, <http://arxiv.org/abs/1010.2561>.
- [BM02] Christophe Breuil and Ariane Mézard, *Multiplicités modulaires et représentations de $GL_2(\mathbb{Z}_p)$ et de $\text{Gal}(\overline{\mathbb{Q}_p}/\mathbb{Q}_p)$ en $\ell = p$* , Duke Math. J. **115** (2002), no. 2, 205–310, With an appendix by Guy Henniart.
- [Bos09] Siegfried Bosch, *Half a century of rigid analytic spaces*, Pure Appl. Math. Q. **5** (2009), no. 4, Special Issue: In honor of John Tate. Part 1, 1435–1467.
- [BR92] Don Blasius and Jonathan D. Rogawski, *Tate classes and arithmetic quotients of the two-ball.*, The zeta functions of Picard modular surfaces. CRM Workshop, Montreal / Can. 1988, 421-444 (1992)., 1992.
- [Buz07] Kevin Buzzard, *Eigenvarieties, L -functions and Galois representations*, London Math. Soc. Lecture Notes, vol. 320, Cambridge Univ. Press, 2007, Durham, 2004, pp. 59–120.
- [BZ77] Joseph Bernstein and Andrei Zelevinsky, *Induced representations of reductive p -adic groups. I*, Ann. Sci. École Norm. Sup. **10** (1977), no. 4, 441–472.
- [Cas] William Casselman, *Introduction to the theory of admissible representations of p -adic reductive groups*, available at <http://www.math.ubc.ca/~cass/research/pdf/p-adic-book.pdf>.
- [Cas80] ———, *The unramified principal series of p -adic groups. I. the spherical function*, Compositio Mathematica **40** (1980), no. 3, 387–406.
- [CG] Frank Calegari and Toby Gee, *Irreducibility of automorphic Galois representations of $GL(n)$, n at most 5*, <http://arxiv.org/abs/1104.4827>.

- [CH] Gaëtan Chenevier and Michael Harris, *Stabilisation de la formule des traces, variétés de shimura, et applications arithmétiques*, vol. 2, ch. Construction of automorphic Galois representations, II, ?, ?
- [Che51] Claude Chevalley, *Deux Théorèmes d'Arithmétique*, J. Math. Soc. Japan **3** (1951), no. 1, 36–44.
- [Che04] Gaëtan Chenevier, *Familles p -adiques de formes automorphes pour GL_n* , J. Reine Angew. Math. **570** (2004), 143–217.
- [Che09] ———, *Une application des variétés de Hecke des groupes unitaires*, Part of the Book Project under the direction of Michael Harris at Paris 7: <http://fa.institut.math.jussieu.fr/node/29>, 2009.
- [Clo86] Laurent Clozel, *On limit multiplicities of discrete series representations in spaces of automorphic forms*, Invent. Math. **83** (1986), no. 2, 265–284.
- [Clo88] Laurent Clozel, *Motifs et Formes Automorphes: Applications du Principe de Fonctorialité*, Automorphic Forms, Shimura Varieties, and L -functions (Laurent Clozel and James S. Milne, eds.), vol. 1, 1988, Proceedings of a Conference held at the University of Michigan, Ann Arbor.
- [Con99] Brian Conrad, *Irreducible components of rigid spaces*, Ann. Inst. Fourier **49** (1999), no. 2, 473–541.
- [GJ78] Stephen Gelbart and Hervé Jacquet, *A relation between automorphic representations of $GL(2)$ and $GL(3)$* , Annales scientifiques de l'ENS **11** (1978), no. 4, 471–542.
- [Gol] Wushi Goldring, *Galois representations associated to holomorphic limits of discrete series*, preprint.
- [IM65] Nagayoshi Iwahori and Hideya Matsumoto, *On some Bruhat decomposition and the structure of the Hecke rings of p -adic Chevalley groups*, Publications Mathématiques de l'IHÉS **25** (1965), 5–48.
- [Jon] Owen Jones, *An analogue of the BGG resolution for locally analytic principal series*, <http://arxiv.org/abs/0912.4756>.
- [Kot86] Robert Kottwitz, *Stable trace formula: elliptic singular terms*, Mathematische Annalen **275** (1986), no. 3, 365–399.
- [KS99] Robert E. Kottwitz and Diana Shelstad, *Foundations of twisted endoscopy*, Astérisque (1999), no. 255, vi+190.
- [Loe11] David Loeffler, *Overconvergent algebraic automorphic forms*, Proc. London Math. Soc. **102** (2011), no. 2, 193–228.

- [MW89] Colette Moeglin and Jean-Loup Waldspurger, *Le spectre résiduel de $GL(n)$* , Annales scientifiques de l'ENS **22** (1989), no. 4, 605–674.
- [MW06] ———, *Sur le transfert des traces d'un groupe classique p -adique à un groupe linéaire tordu*, Selecta Math. **12** (2006), no. 3-4, 433–515.
- [Tad94] Marko Tadić, *Representations of p -adic symplectic groups*, Compositio Mathematica **90** (1994), no. 2, 123–181.
- [Tay] Richard Taylor, *The image of complex conjugation in ℓ -adic representations associated to automorphic forms*, to appear A.N.T.
- [Tay91] ———, *Galois representations associated to Siegel modular forms of low weight*, Duke Math. J. **63** (1991), no. 2, 281–332.
- [Wal10] Jean-Loup Waldspurger, *Les facteurs de transfert pour les groupes classiques: un formulaire*, Manuscripta Math. **133** (2010), no. 1-2, 41–82.