



HAL
open science

Paleoproterozoic evolution of Senegal in the Birimian Framework

Hervé Théveniaut, Claude Delor

► **To cite this version:**

Hervé Théveniaut, Claude Delor. Paleoproterozoic evolution of Senegal in the Birimian Framework. 34th International Geological Congress, Aug 2012, Brisbane, Australia. hal-00711189

HAL Id: hal-00711189

<https://brgm.hal.science/hal-00711189>

Submitted on 22 Jun 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Paleoproterozoic evolution of Senegal in the Birimian Framework

Hervé THEVENIAUT¹, Claude DELOR¹, P M NDIAYE², D P DIALLO², E DIOH³, S SERGEEV⁵

¹ BRGM, GEO/GSO, 3 avenue Claude-Guillemin, BP 36009, 45060 Orléans Cedex 2, France. Email h.theveniaut@brgm.fr

² UCAD, Département de géologie, Faculté des Sciences, Dakar-Fann, Sénégal

³ IFAN – 1, place Soweto – BP 206 – DAKAR, Sénégal

⁴ VSEGEI, 74 Sredny prospect, 199106 St.-Petersburg, Russia

The whole of East-Senegal Birimian terrains have been investigated with a pluri-thematic approach combining airborne geophysics, field mapping, geochemical and isotopic data. Birimian Easternmost Paleoproterozoic terrains begins with the Mako Group of volcanics and metavolcanics, all assumed to have been emplaced at 2.20-2.18 Ga as a juvenile crust.

Reworking of this Eo-Rhyacian crust is recorded with evidence of melting processes and associated products. At the scale of the Mako Group, the Mesorhyacian Sandikounda-Soukouta granodioritic suite and coeval intermediate to acid volcanism (2.18-2.14 Ga) reflect the consumption of the Eorhyacian juvenile crust during early collision stages.

The breakup of Eo-Rhyacian rocks, between 2.14 and 2.10 Ga has resulted in the formation of a main detrital basin that is now referred as the Dialé-Daléma Group. Main lithologies include silicoclastic deposits among which carbonate formations have been found. The upper part of this Group is characterized by the presence of acid to intermediate volcanics dated at ca. 2.1 Ga

The Saraya granitic suite dated at 2.10-2.08 Ga, is mainly recorded in the DD Group with emplacement of the main Saraya batholith and peripheric plutons. It is characterized as a syntectonic magmatic pulse generated in a tectonic context dominated by strong senestral shearing and melting of silicoclastic sources. The specific emplacement of the more Mg-K Boboti magmatic suite is recorded between 2.08 and 2.06 on the easternmost Birimian areas, with evidence of high temperature magma.