



HAL
open science

Hyperbolicity, automorphic forms and Siegel modular varieties

Erwan Rousseau

► **To cite this version:**

Erwan Rousseau. Hyperbolicity, automorphic forms and Siegel modular varieties. 2013. hal-00790243v1

HAL Id: hal-00790243

<https://hal.science/hal-00790243v1>

Preprint submitted on 19 Feb 2013 (v1), last revised 27 Feb 2015 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

HYPERBOLICITY, AUTOMORPHIC FORMS AND SIEGEL MODULAR VARIETIES

ERWAN ROUSSEAU

ABSTRACT. We study the hyperbolicity of compactifications of quotients of bounded symmetric domains by arithmetic groups. We prove that, up to an étale cover, they are Kobayashi hyperbolic modulo the boundary. Applying our techniques to Siegel modular varieties, we improve some former results of Nadel on the non-existence of certain level structures on abelian varieties over complex function fields.

1. INTRODUCTION

Any complex space Y can be equipped with an intrinsic pseudo-distance d_Y , the Kobayashi pseudo-distance [7]. It is the largest pseudo-distance such that every holomorphic map $f : \Delta \rightarrow Y$ from the unit disc, equipped with the Poincaré metric, is distance decreasing.

Recall [7] that a complex space Y is said to be (Kobayashi) hyperbolic modulo $W \subset Y$ if for every pair of distinct points p, q of Y we have $d_Y(p, q) > 0$ unless both are contained in W . In the case where $W = \emptyset$, Brody's lemma gives a criterion for hyperbolicity: Y is hyperbolic if and only if there is no non-constant entire curves $f : \mathbb{C} \rightarrow Y$.

In the case where $W \neq \emptyset$, we do not know any Brody's criterion for hyperbolicity modulo W . It is still an open question if Brody hyperbolicity modulo W implies Kobayashi hyperbolicity modulo W . The reason is that the classical Brody's lemma does not give any precision about where the entire curve is.

In this article, we investigate hyperbolic properties of compactifications of quotients of bounded symmetric domains by arithmetic groups. It is well known that such quotients may be far from being hyperbolic, as rational Hilbert modular surfaces show.

We obtain the following result

Theorem 1.1. *Let $X := \mathbb{B}/\Gamma$ be a smooth quotient of a bounded symmetric domain by an arithmetic subgroup $\Gamma \subset \text{Aut}(\mathbb{B})$. Then there exists a finite étale cover $X_1 = \mathbb{B}/\Gamma_1$ determined by a subgroup $\Gamma_1 \subset \Gamma$ such that $\overline{X_1}$, a compactification of X_1 , is Kobayashi hyperbolic modulo $D_1 := \overline{X_1} \setminus X_1$.*

As a corollary, we obtain a new simple proof of the main result of [9], where it is proved that $\overline{X_1}$ is Brody hyperbolic modulo D_1 : the image $f(\mathbb{C})$ of any nonconstant holomorphic map $f : \mathbb{C} \rightarrow \overline{X_1}$ is contained in D_1 .

2010 *Mathematics Subject Classification.* Primary: 32Q45, 32M15; Secondary: 11G99, 14K15.
Partially supported by the ANR project "POSITIVE", ANR-2010-BLAN-0119-01.

When Nadel's proof [9] was based on Nevanlinna theory, our proof consists in using the fundamental distance decreasing property of the Kobayashi pseudo metric mentioned above. To prove Theorem 1.1, we will construct pseudo distances on $\overline{X_1}$ for which holomorphic maps from the unit disc are distance decreasing.

As an application, we study Siegel modular varieties from this point of view. Their geometry has attracted a lot of attention (see [5] for a survey). In particular, the birational geometry of \mathcal{A}_g and $\mathcal{A}_g(n)$, the moduli spaces of principally polarized abelian varieties with level structures, has been extensively investigated. The Kodaira dimension of compactifications $\overline{\mathcal{A}_g(n)}$ has been studied by Tai, Freitag, Mumford, Hulek and others proving the following result

Theorem 1.2 (Tai, Freitag, Mumford, Hulek). *$\overline{\mathcal{A}_g(n)}$ is of general type for the following values of g and $n \geq n_0$:*

g	1	2	3	4	5	6	≥ 7
n_0	7	4	3	2	2	2	1

Therefore it is very natural to study hyperbolicity of these spaces in light of the Green-Griffiths-Lang conjecture

Conjecture 1.3 (Green-Griffiths, Lang). *Let X be a projective variety of general type. Then there exists a proper algebraic subvariety $Y \subset X$ such that X is hyperbolic modulo Y .*

In this setting, using the theory of automorphic forms, we can apply the same strategy as above and obtain

Theorem 1.4. *$\overline{\mathcal{A}_g(n)}$ is hyperbolic modulo $D := \overline{\mathcal{A}_g(n)} \setminus \mathcal{A}_g(n)$ if $n > 6g$.*

As a corollary we have

Theorem 1.5. *If A is a principally polarized abelian variety of dimension $g \geq 1$ which is defined and nonconstant over k , a complex function field of genus ≤ 1 , then A does not admit a level- n structure for $n > 6g$.*

This improves the previous bound, quadratic in g , obtained by Nadel in [9].

Acknowledgments. The author friendly thanks Carlo Gasbarri and Xavier Roulleau for several interesting and fruitful discussions and remarks. He also thanks the CNRS for the opportunity to spend a semester in Montreal at the UMI CNRS-CRM and the hospitality of UQAM-Cirget where part of this work was done.

2. DISTANCE DECREASING PSEUDO DISTANCES

Let $X := \mathbb{B}/\Gamma$ be a quotient of a bounded symmetric domain by an arithmetic subgroup $\Gamma \subset \text{Aut}(\mathbb{B})$. We first give a criterion to ensure the existence of a pseudo distance \tilde{g} on \overline{X} non degenerate at a given $x \in X$ and satisfying the distance decreasing property: every holomorphic map $f : (\Delta, g_P) \rightarrow (\overline{X}, \tilde{g})$ from the unit disc, equipped with the Poincaré metric g_P , is distance decreasing i.e.

$$f^*\tilde{g} \leq g_P.$$

By passing to a subgroup of finite index if necessary, we can always suppose that Γ is neat [3]. In particular, X admits a smooth toroidal compactification \overline{X} where $D = \overline{X} \setminus X$ is a normal crossings divisor [1].

Let g denote the Bergmann metric on \mathbb{B} such that $\text{Ricci}(g) = -g$. Let $\gamma > 0$ be a rational number such that g has holomorphic sectional curvature $\leq -\gamma$. We want to construct a continuous function $\psi : \overline{X} \rightarrow \mathbb{R}^+$ such that $\psi.g$ defines a pseudo metric on \overline{X} .

Theorem 2.1. *Let $x \in X$ and suppose there is a section $s \in H^0(\overline{X}, l(K_{\overline{X}} + D))$ such that*

- $s(x) \neq 0$
- s is vanishing on D with multiplicity $m > \frac{l}{\gamma}$.

Then there exists a pseudo metric \tilde{g} non-degenerate at x satisfying the distance decreasing property. In particular, the Kobayashi pseudo metric is non degenerate at x .

We will prove this theorem in 3 steps.

From [8], we know that the Bergmann metric induces a *good* singular metric $h := (\det g)^{-1}$ on $K_{\overline{X}} + D$. Let $x \in X$ and s given as above. Let $0 < \epsilon$ such that

$$\gamma - \epsilon > \frac{l}{m}.$$

We define

$$\psi := \|s\|_h^{\frac{2(\gamma-\epsilon)}{l}}.$$

Despite the singularities of h , we have

Proposition 2.2. *ψ is a continuous function on \overline{X} vanishing on D .*

Proof. We know [8] that $K_{\overline{X}} + D$ is a *good extension* of $K_{\overline{X}}$ with the metric h . This implies that near D , which locally is given by $\{z_1 \dots z_k = 0\}$, we have

$$(2.1) \quad \|s\|_h = O(|z_1 \dots z_k|^m \log^{2N} |z_1 \dots z_k|),$$

which suffices to obtain the conclusion. □

The second step is

Proposition 2.3. *$\psi.g$ is a continuous pseudo metric on \overline{X} vanishing on D .*

Proof. We know that g has Poincaré growth near D (Lemma 2.1 [9]) i.e.

$$g = O(g_P),$$

where

$$g_P := \sum_{i=1}^k \frac{dz_i \otimes d\bar{z}_i}{|z_i|^2 \log^2 |z_i|^2} + \sum_{i=k+1}^n dz_i \otimes d\bar{z}_i$$

is the metric obtained by taking the Poincaré metric on the punctured disc. Therefore if we combine this with (2.1), we have

$$\|s\|_h^{\frac{2(\gamma-\epsilon)}{l}} . g_{ij} = O(|z_1 \dots z_k|^{\frac{2m(\gamma-\epsilon)}{l} - 2} \log^\alpha |z_1 \dots z_k|).$$

This estimate clearly implies the proposition. □

Finally as a third step, we are in position to construct a distance decreasing pseudo distance which finishes the proof of Theorem 2.1.

Proposition 2.4. *There exists a constant $\beta > 0$ such that $\tilde{g} := \beta \cdot \psi \cdot g$ is a distance decreasing pseudo-distance: for any holomorphic map $f : \Delta \rightarrow \bar{X}$ from the unit disc equipped with the Poincaré metric g_P , we have*

$$f^* \tilde{g} \leq g_P.$$

Proof. Let

$$u(z) := (1 - |z|^2)^2 \cdot f^* \tilde{g}.$$

Restricting to a smaller disk, we may assume that u vanishes on the boundary of Δ . Take a point z_0 at which u is maximum. $z_0 \in \Delta \setminus f^{-1}(\{\psi = 0\})$ otherwise u would vanish identically. Therefore at z_0 we have

$$i\partial\bar{\partial} \log u(z)_{z=z_0} \leq 0.$$

We have

$$i\partial\bar{\partial} \log u(z) = i\partial\bar{\partial} \log f^* \psi + i\partial\bar{\partial} \log f^* g + i\partial\bar{\partial} \log(1 - |z|^2)^2.$$

On X , we have

$$i\partial\bar{\partial} \log \psi = (\gamma - \epsilon) \text{Ricci}(g) = -(\gamma - \epsilon) \cdot g$$

by the Kähler-Einstein property.

g has holomorphic sectional curvature $\leq -\gamma$ therefore

$$i\partial\bar{\partial} \log f^* g \geq \gamma \cdot f^* g.$$

Moreover

$$i\partial\bar{\partial} \log(1 - |z|^2)^2 = \frac{-1}{(1 - |z|^2)^2}.$$

Combining the above inequalities, we obtain

$$-(\gamma - \epsilon) \cdot f^* g(z_0) + \gamma \cdot f^* g - \frac{1}{(1 - |z_0|^2)^2} \leq 0.$$

So

$$\epsilon \cdot f^* g(z_0) \cdot (1 - |z_0|^2)^2 \leq 1.$$

Finally, we have

$$u(z) \leq u(z_0) = (1 - |z_0|^2)^2 \cdot \beta \cdot \psi(f(z_0)) \cdot f^* g(z_0) \leq \frac{\beta \cdot \psi(f(z_0))}{\epsilon}.$$

Therefore if we take

$$\beta = \frac{\epsilon}{\sup \psi},$$

we obtain the distance decreasing property. □

3. PROOF OF THEOREM 1.1

To prove Theorem 1.1 it is now sufficient to show the existence of an étale cover $X_1 = \mathbb{B}/\Gamma_1$ satisfying the hypothesis of Theorem 2.1 for every $x \in X_1$.

First we recall the following result

Proposition 3.1 ([6], 4.2). *For any point $x \in X$, there exists a section $s \in H^0(\overline{X}, ((\binom{n+1}{2} + 2)(K_{\overline{X}} + D)))$ such that $s(x) \neq 0$ and $s|_D \equiv 0$, where $D := \overline{X} \setminus X$.*

Let $l := \binom{n+1}{2} + 2$ and take a positive integer m such that

$$m > \frac{l}{\gamma}.$$

Now, we have

Proposition 3.2. *There exists a covering X_1 of X such that for any $x \in X_1$, there is a section $s \in H^0(\overline{X}_1, l(K_{\overline{X}_1} + D_1))$ vanishing on $D_1 := \overline{X}_1 \setminus X_1$ with multiplicity at least m such that $s(x) \neq 0$.*

Proof. By [8], one can find a covering X_1 such that $\pi : \overline{X}_1 \rightarrow \overline{X}$ is ramified to order at least m along D_1 . Let $x \in X_1$. From Proposition 3.1, there is a section $s_0 \in H^0(\overline{X}, l(K_{\overline{X}} + D))$ such that $s_0(\pi(x)) \neq 0$ and $s_0|_D \equiv 0$. We just have to take

$$s := \pi^*(s_0) \in H^0(\overline{X}_1, l(K_{\overline{X}_1} + D_1) - mD_1).$$

□

As a consequence, we obtain a proof of Theorem 1.1

Proof. The previous Proposition and Theorem 2.1 imply that for any $x \in X_1$, there is a pseudo-distance ρ non zero at x such that

$$d_{\overline{X}_1} \geq \rho.$$

This implies that \overline{X}_1 is Kobayashi hyperbolic modulo D_1 .

□

4. SIEGEL MODULAR VARIETIES

We will apply Theorem 2.1 to the particular case of quotients of the Siegel upper half space by principal congruence subgroups.

Here $\mathbb{B} = \mathbb{H}_g$, the Siegel upper half space of rank g . The arithmetic group Γ is $Sp(2g, \mathbb{Z})$ and $\Gamma(n)$ is the kernel of $Sp(2g, \mathbb{Z}) \rightarrow Sp(2g, \mathbb{Z}/n)$.

The space $\mathcal{A}_g(n) := \mathbb{H}_g/\Gamma(n)$ is the moduli space of principally polarized abelian varieties with a level- n structure. Recall that if A is an abelian variety of dimension g over a field k of characteristic 0, a level- n structure is a $2g$ -tuple (x_1, \dots, x_{2g}) of points in $A(k)$ which generate the subgroup of all n -torsion points in $A(\overline{k})$.

It is known that the Bergmann metric on \mathbb{H}_g has holomorphic sectional curvature $\leq \frac{-2}{g(g+1)}$, we can therefore take

$$\gamma = \frac{2}{g(g+1)}.$$

We are going to see that in this situation, we can use the theory of automorphic forms to obtain the section needed to apply Theorem 2.1.

First, recall (see for example [5]) that the automorphy factor of modular forms on \mathbb{H}_g defines a line bundle L on the quotient $X := \mathbb{H}_g/\Gamma(n)$, the (\mathbb{Q} -) line bundle of modular forms of weight 1. If the spaces $\overline{X} = \overline{\mathcal{A}_g(n)}$ are smooth (for example $\Gamma(n)$ is neat when $n \geq 3$), we have

$$K_{\overline{X}} = (g+1)L - D,$$

where D is the boundary divisor.

Let F be a modular form with respect to the full modular group $Sp(2g, \mathbb{Z})$. Then following [4], we define the order $o(F)$ of F as the quotient of the vanishing order of F at the boundary divided by the weight of F .

Particularly useful for us, is the following result of Weissauer on the existence of cusp forms of small slope which do not vanish at a given point of the Siegel space.

Theorem 4.1 ([10]). *For every point $\tau \in \mathbb{H}_g$ and every $\epsilon > 0$ there exists a modular form F of order $o(F) \geq \frac{1}{12+\epsilon}$ which does not vanish at τ .*

We obtain as a corollary

Proposition 4.2. *Let $X := \mathcal{A}_g(n)$, $x \in X$ and suppose $n > 6g$. Then there is a section $s \in H^0(\overline{X}, l(K_{\overline{X}} + D))$ such that*

- $s(x) \neq 0$
- s is vanishing on D with multiplicity $m > \frac{l}{\gamma}$.

Proof. We consider the natural map

$$\pi : \overline{\mathcal{A}_g(n)} \rightarrow \overline{\mathcal{A}_g}.$$

This map is ramified to order at least n over the divisor at infinity [9]. We choose some $\epsilon > 0$ such that $\epsilon < \frac{2}{g}$. Now, we take a modular form given by Weissauer's theorem non vanishing at $\pi(x)$. In terms of line bundles, it gives a section $\tilde{s} \in H^0(\overline{\mathcal{A}_g}, kL - mD)$ such that $\frac{m}{k} \geq \frac{1}{12+\epsilon}$ non vanishing at $\pi(x)$. We take $s := \pi^*(\tilde{s})$. It gives a section in $H^0(\overline{X}, kL - mnD)$ non vanishing at x , where we make the slight abuse of denoting the line bundles corresponding to the modular forms and the boundary again by L and D . Therefore, using the equality $K_{\overline{X}} = (g+1)L - D$, we have obtained a section in $H^0(\overline{X}, l(K_{\overline{X}} + D))$ with $l = \frac{k}{g+1}$ vanishing on D with multiplicity mn . Finally, we observe that

$$mn \geq \frac{(6g+1)k}{12+\epsilon} > \frac{kg}{2} = \frac{l}{\gamma}.$$

□

Theorem 1.4 is an immediate corollary of Proposition 4.2 and Theorem 2.1.

Theorem 1.5 is a consequence of Theorem 1.4, since a nonconstant principally polarized abelian variety over k , a complex function field of genus ≤ 1 , which admits a level- n structure induces a nonconstant holomorphic map $\mathbb{C} \rightarrow \overline{\mathcal{A}_g(n)}$ whose image does not lie in D .

REFERENCES

- [1] Ash A., Mumford, D., Rapoport M., Tai Y., *Smooth compactification of locally symmetric varieties*, Lie Groups: History, Frontiers and Applications, Vol. IV. Math. Sci. Press, Brookline, Mass., 1975. iv+335 pp.
- [2] Baily W. L., Jr.; Borel A., *Compactification of arithmetic quotients of bounded symmetric domains*, Ann. of Math. (2) 84 1966 442–528.
- [3] Borel A., *Introduction aux groupes arithmétiques*, Hermann, Paris, 1969.
- [4] Hulek K., *Nef Divisors on Moduli Spaces of Abelian Varieties.*, In: Th. Peternell, F.-O. Schreyer (Eds.), *Complex Analysis and Algebraic Geometry*, de Gruyter, (2000) 255–274.
- [5] Hulek K., Sankaran G.K., *The geometry of Siegel modular varieties*, In: S. Mori, Y. Miyaoka (Eds.), *Higher Dimensional Birational Geometry*, Advanced Studies in Pure Mathematics 35 (2002), 89–156.
- [6] Hwang J.-M., To W.-K., *Uniform boundedness of level structures on abelian varieties over complex function fields*, Math. Ann. 335 (2006), no. 2, 363–377.
- [7] Kobayashi S., *Hyperbolic complex spaces*, Grundlehren der Mathematischen Wissenschaften 318, Springer Verlag, Berlin, 1998, xiv+471 pp.
- [8] Mumford D., *Hirzebruch's proportionality theorem in the noncompact case*, Invent. Math. 42 (1977), 239–272.
- [9] Nadel A. M., *The nonexistence of certain level structures on abelian varieties over complex function fields*, Ann. of Math. (2) 129 (1989), no. 1, 161–178.
- [10] Weissauer R., *Untervarietäten der Siegelschen Modulmannigfaltigkeiten von allgemeinem Typ.*, Math. Ann. 275 (1986), 207–220.

Erwan Rousseau,
erwan.rousseau@cmi.univ-mrs.fr
Laboratoire d'Analyse, Topologie, Probabilités
Université d'Aix-Marseille et CNRS
39, rue Frédéric Joliot-Curie
13453 Marseille Cedex 13
France