



HAL
open science

Caractérisation des mouvements oscillants dans l'atmosphère stable d'une vallée encaissée

Chantal Staquet, Yann Largeron, Charles Chemel

► **To cite this version:**

Chantal Staquet, Yann Largeron, Charles Chemel. Caractérisation des mouvements oscillants dans l'atmosphère stable d'une vallée encaissée. Symposium OGOA, May 2013, Lyon, France. hal-00854444

HAL Id: hal-00854444

<https://hal.science/hal-00854444>

Submitted on 27 Aug 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Chantal Staquet (LEGI)

Caractérisation des mouvements oscillants dans l'atmosphère stable d'une vallée encaissée

C. Staquet, Y. Largeron (University of Grenoble & LEGI, Grenoble, France)

C. Chemel (NCAS-Weather, University of Hefordshire, UK)

In a valley sheltered from strong synoptic effects, the dynamics of the valley atmosphere at night are dominated by katabatic winds. In a stably stratified atmosphere, these winds undergo temporal oscillations, whose frequency is given by $N\sin\theta$ for an infinitely long slope of constant slope angle θ , N being the buoyancy frequency. Such an unsteady flow in a stably stratified atmosphere may also generate internal gravity waves (IGWs) (Figure 1). The numerical study by Chemel et al. (Meteorol Atmos Phys 203:187-194, 2009) showed that, in the stable atmosphere of a deep valley, the oscillatory motions associated with the IGWs generated by katabatic winds are distinct from those of the katabatic winds. The IGW frequency was found to be independent of α and about $0.8N$.




Figure 1: Vertical cross-section of a valley of idealized shape in a stably-stratified atmosphere. Contours of the vertical component of the velocity field are displayed, which attest of the generation of an internal gravity wave field from the katabatic flow (flowing along the slopes but hardly visible on the figure). The wave amplitude is about ten times smaller than the katabatic flow amplitude.

Their study did not consider the effects of the background stratification and valley geometry on these results. These effects are investigated in the present study for a wide range of stratification and slope angles, through numerical simulations for a deep valley. The two oscillatory systems are reproduced in the simulations. The frequency of the oscillations of the katabatic winds is found to be equal to N times the sine of the maximum slope angle. Remarkably, the IGW frequency is found to also vary as $C_w N$, with C_w in the range $0.7 - 0.95$. These values for C_w are similar to those reported for IGWs radiated by any turbulent field with no dominant frequency component. Results suggest that the IGW wavelength is controlled by the valley depth.

1. Largeron Y., Staquet C. & Chemel C. 2013 Characterization of oscillating motions in the stable atmosphere of a deep valley. *Boundary Layer Meteorology*, DOI 10.1007/s10546-013-9825-y.