



HAL
open science

Complexity of Existential Positive First-Order Logic

Manuel Bodirsky, Miki Hermann, Florian Richoux

► **To cite this version:**

Manuel Bodirsky, Miki Hermann, Florian Richoux. Complexity of Existential Positive First-Order Logic. *Journal of Logic and Computation*, 2013, 23 (4), pp.753-760. 10.1093/logcom/exr043. hal-00870985

HAL Id: hal-00870985

<https://hal.science/hal-00870985>

Submitted on 21 Oct 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Complexity of Existential Positive First-Order Logic

Manuel Bodirsky, Miki Hermann

LIX (UMR CNRS 7161), École Polytechnique, 91128 Palaiseau, France
{bodirsky, hermann}@lix.polytechnique.fr

Florian Richoux*

JFLI, CNRS - University of Tokyo, Japan
richoux@jfli.itc.u-tokyo.ac.jp

Abstract

Let Γ be a (not necessarily finite) structure with a finite relational signature. We prove that deciding whether a given existential positive sentence holds in Γ is in LOGSPACE or complete for the class $\text{CSP}(\Gamma)_{\text{NP}}$ under deterministic polynomial-time many-one reductions. Here, $\text{CSP}(\Gamma)_{\text{NP}}$ is the class of problems that can be reduced to the *constraint satisfaction problem* of Γ under *non-deterministic* polynomial-time many-one reductions.

Key words: Computational Complexity, Existential Positive First-Order Logic, Constraint Satisfaction Problems

1 Introduction

We study the computational complexity of the following class of computational problems. Let Γ be a structure with finite or infinite domain and with a finite relational signature. The model-checking problem for existential positive first-order logic, parametrized by Γ , is the following problem.

Problem: EXPOS(Γ)

Input: An existential positive first-order sentence Φ .

Question: Does Γ satisfy Φ ?

Existential positive first-order formula over Γ are first-order formulas without universal quantifiers, equalities, and negation symbols, and formally defined as follows:

- if R is a relation symbol of a relation from Γ with arity k and x_1, \dots, x_k are (not necessarily distinct) variables, then $R(x_1, \dots, x_k)$ is an existential positive first-order formula (such formulas are called *atomic*);
- if φ and ψ are existential positive first-order formulas, then $\varphi \wedge \psi$ and $\varphi \vee \psi$ are existential positive first-order formulas;
- if φ is an existential positive first-order formula with a free variable x then $\exists x.\varphi$ is an existential positive first-order formula.

*This work was done during the PhD studies of the third author at École Polytechnique.

An *existential positive first-order sentence* is an existential positive first-order formula without free variables.

Note that we do not allow the equality symbol in the existential positive sentences; this only makes our results stronger, since one might always add a relation symbol $=$ for the equality relation to the signature of Γ to obtain the result for the case where the equality symbol is allowed. Also note that adding a symbol for equality to Γ might change the complexity of $\text{EXPOS}(\Gamma)$. Consider for example $\Gamma := (\mathbb{N}; \neq)$; here, $\text{EXPOS}(\Gamma)$ can be reduced to the Boolean formula evaluation problem (which is known to be in LOGSPACE) as follows: atomic formulas in Φ of the form $x \neq y$ are replaced by *true*, and atomic formulas of the form $x \neq x$ are replaced by *false*. The resulting Boolean formula is equivalent to true if and only if Φ is true in Γ . However, the problem $\text{EXPOS}(\Gamma')$ for $\Gamma' := (\mathbb{N}; \neq, =)$ is NP-complete. Similar examples exist over finite domains.

The *constraint satisfaction problem* $\text{CSP}(\Gamma)$ for Γ is defined similarly, but its input consists of a *primitive positive* sentence, that is, a existential positive sentence without disjunctions. Constraint satisfaction problems frequently appear in many areas of computer science, and have attracted a lot of attention, in particular in combinatorics, artificial intelligence, finite model theory and universal algebra; we refer to the recent collection of survey articles on this subject [1]. The class of constraint satisfaction problems for infinite structures Γ is a rich class of problems; it can be shown that for every computational problem there exists a relational structure Γ such that $\text{CSP}(\Gamma)$ is equivalent to that problem under polynomial-time Turing reductions [2].

In this paper, we show that the complexity classification for existential positive first-order sentences over infinite structures can be reduced to the complexity classification for constraint satisfaction problems. For finite structures Γ , our result implies that $\text{EXPOS}(\Gamma)$ is in LOGSPACE or NP-complete. The LOGSPACE -solvable cases of $\text{EXPOS}(\Gamma)$ are in this case precisely those relational structures Γ with an element a such that all non-empty relations in Γ contain the tuple (a, \dots, a) ; in this case, $\text{EXPOS}(\Gamma)$ is called *a-valid*. Interestingly, this is no longer true for infinite structures Γ . To see this, consider again the structure $\Gamma := (\mathbb{N}; \neq)$, which is clearly not *a-valid*, but in LOGSPACE as we have noticed above.

A universal-algebraic study of the model-checking problem for finite structures Γ and various other syntactic restrictions of first-order logic (for instance positive first-order logic) can be found in [9].

A preliminary version of this article appeared in [3]. The present version differs in that the main proof has been simplified and now also works without the relation symbol for equality; moreover, Proposition 3 and Section 4 have been added.

2 Main Result

We write $L \leq_m L'$ if there exists a deterministic polynomial-time many-one reduction from L to L' .

Definition 1 (from [6]) A problem A is *non-deterministic polynomial-time many-one reducible* to a problem B ($A \leq_{\text{NP}} B$) if there is a nondeterministic polynomial-time Turing machine M such that $x \in A$ if and only if there exists a computation of M that outputs y on input x , and $y \in B$. We denote by A_{NP} the smallest class that contains A and is downward closed under \leq_{NP} .

Observe that \leq_{NP} is transitive [6]. To state the complexity classification for existential positive first-order logic, we need the following concept. The Γ -*localizer* $F(\psi)$ of a formula ψ is defined as follows:

- $F(\exists x.\psi) = F(\psi)$
- $F(\varphi \wedge \psi) = F(\varphi) \wedge F(\psi)$
- $F(\varphi \vee \psi) = F(\varphi) \vee F(\psi)$

- When ψ is atomic, then $F(\psi) = \begin{cases} true & \text{if } \psi \text{ is satisfiable in } \Gamma \\ false & \text{otherwise} \end{cases}$

Definition 2 We call a structure Γ *locally refutable* if every existential positive sentence Φ is true in Γ if and only if the Γ -localizer $F(\Phi)$ is logically equivalent to *true*.

Proposition 3 A structure Γ is locally refutable if and only if every unsatisfiable conjunction of atomic formulas contains an unsatisfiable conjunct.

Proof: First suppose that Γ is locally refutable, and let φ be a conjunction of atomic formulas with variables x_1, \dots, x_n . Then every conjunct of φ is satisfiable in Γ if and only if $F(\varphi)$ is true. By local refutability of Γ this is the case if and only if $\exists x_1, \dots, x_n. \varphi$ is true in Γ , which shows the claim.

Now suppose that Γ is not locally refutable, that is, there is an existential positive sentence Φ that is false in Γ such that $F(\Phi)$ is true. Define recursively for each subformula ψ of Φ where $F(\psi)$ is true the formula $T(\psi)$ as follows. If ψ is of the form $\psi_1 \vee \psi_2$, then for some $i \in \{1, 2\}$ the formula $F(\psi_i)$ must be true, and we set $T(\psi)$ to be $T(\psi_i)$. If ψ is of the form $\psi_1 \wedge \psi_2$, then for both $i \in \{1, 2\}$ the formula $F(\psi_i)$ must be true, and we set $T(\psi)$ to be $T(\psi_1) \wedge T(\psi_2)$.

Each conjunct φ in $T(\Phi)$ is satisfiable in Γ since $F(\Phi)$ is true. But since Φ is false in Γ , $T(\Phi)$ must be unsatisfiable. \square

In Section 3, we will show the following result.

Theorem 4 Let Γ be a structure with a finite relational signature τ . If Γ is locally refutable then the problem $\text{EXPOS}(\Gamma)$ to decide whether an existential positive sentence is true in Γ is in LOGSPACE . If Γ is not locally refutable, then $\text{EXPOS}(\Gamma)$ is complete for the class $\text{CSP}(\Gamma)_{\text{NP}}$ under polynomial-time many-one reductions.

In particular, $\text{EXPOS}(\Gamma)$ is in LOGSPACE or is NP-hard (under deterministic polynomial-time many-one reductions). If Γ is finite, then $\text{EXPOS}(\Gamma)$ is in LOGSPACE or NP-complete, because finite domain constraint satisfaction problems are clearly in NP. The observation that $\text{EXPOS}(\Gamma)$ is in LOGSPACE or NP-complete has previously been made in [5] and independently in [8]. However, our proof remains the same for finite domains and is simpler than the previous proofs.

3 Proof

Before we prove Theorem 4, we start with the following simpler result.

Theorem 5 Let Γ be a structure with a finite relational signature τ . If Γ is locally refutable, then the problem $\text{EXPOS}(\Gamma)$ to decide whether an existential positive sentence is true in Γ is in LOGSPACE . If Γ is not locally refutable, then $\text{EXPOS}(\Gamma)$ is NP-hard (under polynomial-time many-one reductions).

To prove Theorem 5, we need first to prove the following lemma.

Lemma 6 A structure Γ is not locally refutable if and only if there are existential positive formulas ψ_0 and ψ_1 with the property that

- ψ_0 and ψ_1 define non-empty relations over Γ ;
- $\psi_0 \wedge \psi_1$ defines the empty relation over Γ .

Proof: The “if”-part of the statement is immediate. To show the “only if”-part, suppose that Γ is not locally refutable. Then by Proposition 3 there is an unsatisfiable conjunction ψ of satisfiable atomic formulas. Among all such formulas ψ , let ψ be one of minimal length. Let ψ_0 be one of the atomic formulas in ψ , and let ψ_1 be the conjunction over the remaining conjuncts in ψ . Since ψ was chosen to be minimal, the formula ψ_1 must be satisfiable. By construction ψ_0 is also satisfiable and ψ is unsatisfiable, which is what we had to show. \square

Proof of Theorem 5: If Γ is locally refutable, then $\text{EXPOS}(\Gamma)$ can be reduced to the positive Boolean formula evaluation problem, which is known to be LOGSPACE -complete. We only have to construct from an existential positive sentence Φ a Boolean formula $F := F_\Gamma(\Phi)$ as described before Definition 2. Clearly, this construction can be performed with logarithmic work-space. We evaluate F , and reject if F is false, and accept otherwise.

If Γ is not locally refutable, we show NP-hardness of $\text{EXPOS}(\Gamma)$ by reduction from 3-SAT. Let I be a 3-SAT instance. We construct an instance Φ of $\text{EXPOS}(\Gamma)$ as follows. Let ψ_0 and ψ_1 be the formulas from Lemma 6 (suppose they are d -ary). Let v_1, \dots, v_n be the Boolean variables in I . For each v_i we introduce d new variables $\bar{x}_i = x_i^1, \dots, x_i^d$. Let Φ be the instance of $\text{EXPOS}(\Gamma)$ that contains the following conjuncts:

- For each $1 \leq i \leq n$, the formula $\psi_0(\bar{x}_i) \vee \psi_1(\bar{x}_i)$
- For each clause $l_1 \vee l_2 \vee l_3$ in I , the formula $\psi_{i_1}(\bar{x}_{j_1}) \vee \psi_{i_2}(\bar{x}_{j_2}) \vee \psi_{i_3}(\bar{x}_{j_3})$ where $i_p = 0$ if l_p equals $\neg x_{j_p}$ and $i_p = 1$ if l_p equals x_{j_p} , for all $p \in \{1, 2, 3\}$.

It is clear that Φ can be computed in deterministic polynomial time from I , and that Φ is true in Γ if and only if I is satisfiable. \square

Applied to finite relational structures Γ , we obtain the result from [5] and [8], that is, $\text{EXPOS}(\Gamma)$ is in LOGSPACE if Γ is a -valid and NP-complete otherwise. We prove in the following proposition that, over a finite domain D , Γ is locally refutable if and only if it is a -valid for an element $a \in D$.

Proposition 7 *Let Γ be a relational structure with a finite domain D . Then Γ is locally refutable if and only if it is a -valid for an element $a \in D$.*

Proof: Suppose that Γ is a -valid, and let Φ be an existential positive sentence over the signature of Γ . To show that Γ is locally refutable, we only have to show that Φ is true in Γ when $F(\Phi)$ is equivalent to true (since the other direction holds trivially). But this follows from the fact that if an atomic formula $R(x_1, \dots, x_n)$ is satisfiable in Γ then in fact this formula can be satisfied by setting all variables to a .

For the opposite direction of the statement, let $D = \{a_1, \dots, a_n\}$, and suppose that for all $a \in D$ the structure Γ is not a -valid. That is, for each $a_i \in D$ there exists a non-empty relation R_i of arity r_i in Γ such that $(a_i, \dots, a_i) \notin R_i$. Let r be $\sum_{i=1}^n r_i$, and let x_1, \dots, x_{rn} be distinct variables. Consider the formula

$$\psi = \bigwedge_{\bar{y} \in \{x_1, \dots, x_{rn}\}^r} R_1(y_1, \dots, y_{r_1}) \wedge \dots \wedge R_n(y_{r-r_n+1}, \dots, y_r) \quad (1)$$

By the pigeonhole principle, for every mapping $f: \{x_1, \dots, x_{rn}\} \rightarrow D$ at least r variables are mapped to the same value, say to a_i . For a vector \bar{y} that contains exactly these r variables, for some l there is a conjunct $R_l(y_{l+1}, \dots, y_{l+r_l})$ in ψ ; but by assumption, R_l does not contain the tuple (a_i, \dots, a_i) . This shows that $\exists x_1, \dots, x_{rn}. \psi$ is not true in Γ . On the other hand, since each relation R_i is non-empty, it is clear that the Boolean formula $F(\exists x_1, \dots, x_{rn}. \psi)$ is true. Therefore, Γ is not locally refutable. \square

Remark 8 In the proof of Theorem 4 it will be convenient to assume that Γ has a single relation R . When we study the problem $\text{CSP}(\Gamma)$, this is without loss of generality, since we can always find a CSP which is deterministic polynomial-time equivalent and where the template is of this form: if $\Gamma = (D; R_1, \dots, R_n)$ where R_i has arity r_i and is not empty, then $\text{CSP}(\Gamma)$ is equivalent to $\text{CSP}(D; R_1 \times \dots \times R_n)$ where $R_1 \times \dots \times R_n$ is the $\sum_{i=1}^n r_i$ -ary relation defined as the Cartesian product of the relations R_1, \dots, R_n . Similarly, $\text{EXPOS}(\Gamma)$ is equivalent to $\text{EXPOS}(D; R_1 \times \dots \times R_n)$.

Proof of Theorem 4: If Γ is locally refutable then the statement has been shown in Theorem 5. Suppose that Γ is not locally refutable. To show that $\text{EXPOS}(\Gamma)$ is contained in $\text{CSP}(\Gamma)_{\text{NP}}$, we construct a non-deterministic Turing machine T which takes as input an instance Φ of $\text{EXPOS}(\Gamma)$, and which outputs an instance $T(\Phi)$ of $\text{CSP}(\Gamma)$ as follows.

On input Φ the machine T proceeds recursively as follows:

- if Φ is of the form $\exists x.\varphi$ then return $\exists x.T(\varphi)$;
- if Φ is of the form $\varphi_1 \wedge \varphi_2$ then return $T(\varphi_1) \wedge T(\varphi_2)$;
- if Φ is of the form $\varphi_1 \vee \varphi_2$ then non-deterministically return either $T(\varphi_1)$ or $T(\varphi_2)$;
- if Φ is of the form $R(x_1, \dots, x_k)$ then return $R(x_1, \dots, x_k)$.

The output of T can be viewed as an instance of $\text{CSP}(\Gamma)$, since it can be transformed to a primitive positive sentence (by moving all existential quantifiers to the front). It is clear that T has polynomial running time, and that Φ is true in Γ if and only if there exists a computation of T on Φ that computes a sentence that is true in Γ .

We now show that $\text{EXPOS}(\Gamma)$ is hard for $\text{CSP}(\Gamma)_{\text{NP}}$ under \leq_m -reductions. Let L be a problem with a non-deterministic polynomial-time many-one reduction to $\text{CSP}(\Gamma)$, and let M be the non-deterministic Turing machine that computes the reduction. We have to construct a deterministic Turing machine M' that computes for any input string s in polynomial time in $|s|$ an instance Φ of $\text{EXPOS}(\Gamma)$ such that Φ is true in Γ if and only if there exists a computation of M on s that computes a satisfiable instance of $\text{CSP}(\Gamma)$.

Say that the running time of M on s is in $O(|s|^e)$ for a constant e . Hence, there are constants s_0 and c such that for $|s| > s_0$ the running time of M and hence also the number of constraints in the input instance of $\text{CSP}(\Gamma)$ produced by the reduction is bounded by $t := c|s|^e$. The non-deterministic computation of M can be viewed as a deterministic computation with access to non-deterministic advice bits as shown in [4]. We also know that for $|s| > s_0$, the machine M can access at most t non-deterministic bits. If w is a sufficiently long bit-string, we write M_w for the deterministic Turing machine obtained from M by using the bits in w as the non-deterministic bits, and $M_w(s)$ for the instance of $\text{CSP}(\Gamma)$ computed by M_w on input s .

If $|s| \leq s_0$, then M' returns $\exists \bar{x}.\psi_1(\bar{x})$ if there is an $w \in \{0, 1\}^*$ such that $M_w(s)$ is a satisfiable instance of $\text{CSP}(\Gamma)$, and M' returns $\exists \bar{x}(\psi_0(\bar{x}) \wedge \psi_1(\bar{x}))$ otherwise (i.e., it returns a false instance of $\text{EXPOS}(\Gamma)$; ψ_0 and ψ_1 are defined in Lemma 6). Since s_0 is a fixed finite value, M' can perform these computations in constant time.

By Remark 8 made above, we can assume without loss of generality that Γ has just a single relation R . Let l be the arity of R . Then instances of $\text{CSP}(\Gamma)$ with variables x_1, \dots, x_n can be encoded as sequences of numbers that are represented by binary strings of length $\lceil \log t \rceil$ as follows: the i -th number m in this sequence indicates that the $((i-1) \bmod l) + 1$ -st variable in the $((i-1) \text{div } l) + 1$ -st constraint is x_m .

For $|s| > s_0$, we use a construction from the proof of Cook's theorem given in [4]. In this proof, a computation of a non-deterministic Turing machine T accepting a language L is encoded by Boolean variables that represent the state and the position of the read-write head of T at time r , and the content of

the tape at position j at time r . The tape content at time 0 consists of the input x , written at positions 1 through n , and the non-deterministic advice bit string w , written at positions -1 through $-|w|$. The proof in [4] specifies a deterministic polynomial-time computable transformation f_L that computes for a given string s a SAT instance $f_L(s)$ such that there is an accepting computation of T on s if and only if there is a satisfying truth assignment for $f_L(s)$.

In our case, the machine M computes a reduction and thus computes an output string. Recall our binary representation of instances of the CSP M writes on the output tape a sequence of numbers represented by binary strings of length $\lceil \log t \rceil$. It is straightforward to modify the transformation f_L given in the proof of Theorem 2.1 in [4] to obtain for all positive integers a, b, c where $a \leq t$, $b \leq l$, $c \leq \lceil \log t \rceil$, and $d \in \{0, 1\}$, a deterministic polynomial-time transformation $g_{a,b,c}^d$ that computes for a given string s a SAT instance $g_{a,b,c}^d(s)$ with distinguished variables z_1, \dots, z_p , $p \leq t$ for the non-deterministic bits in the computation of M such that the following are equivalent:

- $g_{a,b,c}^d(s)$ has a satisfying assignment where z_i is set to $w_i \in \{0, 1\}$ for $1 \leq i \leq p$;
- the c -th bit in the b -th variable of the a -th constraint in $M_w(s)$ equals d .

We use the transformations $g_{a,b,c}^d$ to define M' as follows. The machine M' first computes the formulas $g_{a,b,c}^d(s)$. For every Boolean variable v in these formulas we introduce a new conjunct $\psi_0(\bar{x}_v) \vee \psi_1(\bar{x}_v)$ where \bar{x}_v is a d -tuple of fresh variables and ψ_0 and ψ_1 are the two formulas defined in Lemma 6. Then, every positive literal v in the original conjuncts of the formula is replaced by $\psi_1(\bar{x}_v)$, and every negative literal $l = \neg v$ by $\psi_0(\bar{x}_v)$. We then existentially quantify over all variables except for $\bar{x}_{z_1}, \dots, \bar{x}_{z_p}$. Let $\psi_{a,b,c}^d(s)$ denote the resulting existential positive formula. For positive integers k and i , we denote as $k[i]$ the i -th bit in the binary representation of k . Let n be the total number of variables in the CSP instance $M_w(s)$ (in particular, $n \leq t$). It is clear that the formula

$$\exists y_1, \dots, y_n, \bar{x}_{z_1}, \dots, \bar{x}_{z_p}. \bigwedge_{1 \leq a, k_1, \dots, k_l \leq t} \left(\left(\bigwedge_{b \leq l, c} \psi_{a,b,c}^{k_b[c]}(s) \right) \rightarrow R(y_{k_1}, \dots, y_{k_l}) \right)$$

can be re-written in existential positive form Φ without blow-up: we can replace implications $\alpha \rightarrow \beta$ by $\neg \alpha \vee \beta$, and then move the negation to the atomic level, where we can remove negation by exchanging the role of φ_0 and φ_1 . Hence, Φ can be computed by M' in polynomial time.

We claim that the formula Φ is true in Γ if and only if there exists a computation of M on s that computes a satisfiable instance of $\text{CSP}(\Gamma)$. To see this, let w be a sufficiently long bit-string such that $M_w(s)$ is a satisfiable instance of $\text{CSP}(\Gamma)$. Suppose for the sake of notation that the n variables in $M_w(s)$ are the variables y_1, \dots, y_n . Let a_1, \dots, a_n be a satisfying assignment to those n variables. Then, if for $1 \leq i \leq n$ the variable y_i in the formula Φ is set to a_i , and for $1 \leq i \leq p$ the variables \bar{x}_{z_i} are set to a tuple that satisfies ψ_d where d is the i -th bit in w , we claim that the inner part of Φ is true in Γ . The reason is that, due to the way how we set the variables of the form \bar{x}_{z_i} , the precondition $\left(\bigwedge_{b \leq l, c} \psi_{a,b,c}^{k_b[c]}(s) \right)$ is true if and only if $R(y_{k_1}, \dots, y_{k_l})$ is a constraint in $M_w(s)$. Therefore, all the atomic formulas of the form $R(y_{k_1}, \dots, y_{k_l})$ are satisfied due to the way how we set the variables y_i , and hence Φ is true in Γ . It is straightforward to verify that the opposite implication holds as well, and this shows the claimed equivalence. \square

4 Structures With Function Symbols

In this section, we briefly discuss the complexity of $\text{EXPOS}(\Gamma)$ when Γ might also contain functions. That is, we assume that the signature of Γ consists of a finite set of relation and function symbols, and that the input formulas for the problem $\text{EXPOS}(\Gamma)$ are existential positive first-order formulas over this signature. It is easy to see from the proofs in the previous section that when Γ is not locally refutable, then $\text{EXPOS}(\Gamma)$ is still NP-hard (with the same definition of local refutability as before).

The case when Γ is locally refutable becomes more intricate when Γ has functions. We present an example of a locally refutable structure Γ where $\text{EXPOS}(\Gamma)$ is NP-hard. Let the signature of Γ be the structure $(2^{\mathbb{N}}; \neq, \cap, \cup, c, \mathbf{0}, \mathbf{1})$ where \neq is the binary disequality relation, \cap and \cup are binary functions for intersection and union, respectively, c is a unary function for complementation, and $\mathbf{0}, \mathbf{1}$ are constants (i.e., 0-ary functions) for the empty set and the full set \mathbb{N} , respectively.

Proposition 9 *The structure $(2^{\mathbb{N}}; \neq, \cap, \cup, c, \mathbf{0}, \mathbf{1})$ is locally refutable.*

Proof: By Lemma 6 it suffices to show that if Ψ is a conjunction of atomic formulas that are satisfiable in Γ , then Ψ is satisfiable over Γ . Since the only relation symbol in the structure is \neq , every conjunct in Ψ is of the form $t_1 \neq t_2$, where t_1 and t_2 are terms formed by variables and the function symbols $\cap, \cup, c, \mathbf{1}$ and $\mathbf{0}$. By Boole's fundamental theorem of Boolean algebras, $t = t'$ can be re-written as $t'' = \mathbf{0}$. Therefore, Ψ can be written as $t_1 \neq \mathbf{0} \wedge \dots \wedge t_n \neq \mathbf{0}$. Since Γ is an infinite Boolean algebra, Theorem 5.1 in [7] shows that if $t_i \neq \mathbf{0}$ is satisfiable in Γ for all $i \leq n$, then Ψ is satisfiable in Γ as well. \square

Proposition 10 *The problem $\text{EXPOS}(2^{\mathbb{N}}; \neq, \cap, \cup, c, \mathbf{0}, \mathbf{1})$ is NP-hard.*

Proof: The proof is by reduction from SAT. Given a Boolean formula Ψ in CNF with variables x_1, \dots, x_n , we replace each conjunction in Ψ by \cap , each disjunction by \cup , and each negation by c . Let t be the resulting term over the signature $\{\cap, \cup, c\}$ and variables x_1, \dots, x_n . It is easy to verify that $\exists x_1, \dots, x_n. t \neq \mathbf{0}$ is true in Γ if and only if Ψ is a satisfiable Boolean formula. \square

5 Conclusion

In this paper, we proved that for an arbitrary (finite or infinite) relational structure the problem $\text{EXPOS}(\Gamma)$ is in LOGSPACE if Γ is locally refutable, or otherwise complete for the class $\text{CSP}(\Gamma)_{\text{NP}}$ under deterministic polynomial-time many-one reductions. In particular, if Γ is not locally refutable then the problem $\text{EXPOS}(\Gamma)$ is NP-hard. Structures with a finite domain are locally refutable if and only if they are a -valid for some value a of the domain D . Finally, we present an example of a structure that shows that our result cannot be straightforwardly extended to structures Γ with function symbols, since local refutability of Γ no longer implies that $\text{EXPOS}(\Gamma)$ is in LOGSPACE when Γ contains function symbols.

Acknowledgment

We would like to thank V́ctor Dalmau for helpful suggestions, and Moritz Mller for the encouragement to study the case where the structure Γ contains function symbols.

References

- [1] N. Creignou, Ph. G. Kolaitis, and H. Vollmer, editors. *Complexity of Constraints — An Overview of Current Research Themes*, volume 5250 of Lecture Notes in Computer Science, Springer Verlag, 2008.
- [2] M. Bodirsky and M. Grohe. Non-Dichotomies in Constraint Satisfaction Complexity. *Proceedings 35th International Colloquium on Automata, Languages and Programming (ICALP 2008), Part II, Reykjavik (Iceland)*, volume 5126 of Lecture Notes in Computer Science, 184–196, 2008.
- [3] M. Bodirsky, M. Hermann and F. Richoux. Complexity of Existential Positive First-Order Logic. *Proceedings 5th Conference on Computability in Europe (CiE 2009), Heidelberg (Germany)*, 31–36, 2009.
- [4] M. R. Garey and D. S. Johnson. *Computers and Intractability: A Guide to the Theory of NP-Completeness*. W.H. Freeman and Co, 1979.
- [5] M. Hermann and F. Richoux. On the Computational Complexity of Monotone Constraint Satisfaction Problems. *Proceedings 3rd Annual Workshop on Algorithms and Computation (WALCOM 2009), Kolkata (India)*, 286–297, 2009.
- [6] R. E. Ladner, N. A. Lynch and A. L. Selman. A Comparison of Polynomial-Time Reducibilities. *Theoretical Computer Science*, 1(2), 103–124, 1975.
- [7] K. Marriott and M. Odersky. Negative boolean constraints. *Theoretical Computer Science*, 160(1&2), 365–380, 1996.
- [8] B. Martin. Dichotomies and Duality in First-order Model Checking Problems. *CoRR abs/cs/0609022*, 2006.
- [9] B. Martin. First-Order Model Checking Problems Parameterized by the Model. *Proceedings 4th Conference on Computability in Europe (CiE 2008), Athens (Greece)*, 417–427, 2008.