



HAL
open science

Yet again on two examples by Iyama and Yoshino

Daniele Faenzi

► **To cite this version:**

| Daniele Faenzi. Yet again on two examples by Iyama and Yoshino. 2014. hal-01062912v1

HAL Id: hal-01062912

<https://hal.science/hal-01062912v1>

Preprint submitted on 10 Sep 2014 (v1), last revised 20 Jul 2019 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

YET AGAIN ON TWO EXAMPLES BY IYAMA AND YOSHINO

DANIELE FAENZI

ABSTRACT. We give an elementary proof of Iyama-Yoshino's classification of rigid MCM modules on Veronese embeddings in \mathbb{P}^9 .

INTRODUCTION

The beautiful theory of cluster tilting in triangulated categories has been developed by Iyama and Yoshino; as an important outcome of this the authors gave in [IY08, Theorem 1.2 and Theorem 1.3] the classification of rigid indecomposable MCM modules over two Veronese embeddings in \mathbb{P}^9 given, respectively, by plane cubics and space quadrics. Another proof, that makes use of Orlov's singularity category, appears in [KMVdB11], where the link between power series Veronese rings and the graded rings of the corresponding varieties is also explained. Also, [KR08] contains yet another argument.

The goal of this note is to present a simple proof of Iyama-Yoshino's classification of rigid MCM modules over the aforementioned Veronese rings, making use of vector bundles and Beilinson's theorem. This proof works over a field k which is algebraically closed or finite.

Consider the embedding of the projective space \mathbb{P}^n given by homogeneous forms of degree d , i.e. the d -fold Veronese variety. A coherent sheaf E on \mathbb{P}^n is arithmetically Cohen-Macaulay (ACM) with respect to this embedding if and only if E is locally free and has *no intermediate cohomology*:

$$(1) \quad H^i(\mathbb{P}^n, E(dt)) = 0, \quad \text{for all } t \in \mathbb{Z} \text{ and all } 0 < i < n.$$

This is equivalent to ask that the module of global sections associated with E is MCM over the corresponding Veronese ring. For d -fold Veronese embeddings of \mathbb{P}^n in \mathbb{P}^9 (i.e. $\{n, d\} = \{2, 3\}$), we are going to classify ACM bundles E which are *rigid*, i.e. $\text{Ext}_{\mathbb{P}^n}^1(E, E) = 0$. We set $\ell = \binom{n+1}{2}$.

To state the classification, we define the *Fibonacci numbers* $a_{\ell, k}$ by the relations: $a_{\ell, 0} = 0$, $a_{\ell, 1} = 1$ and $a_{\ell, k+1} = \ell a_{\ell, k} - a_{\ell, k-1}$. For instance $(a_{3, k})$ is given by the odd values of the usual Fibonacci sequence:

$$a_{3, k} = 0, 1, 3, 8, 21, 55, 144, \dots \quad \text{for } k = 0, 1, 2, 3, 4, 5, 6, \dots$$

Theorem 1 ($\{n, d\} = \{2, 3\}$). *Let E be an indecomposable bundle on \mathbb{P}^n satisfying (1).*

i) *If E has no endomorphism factoring through $\mathcal{O}_{\mathbb{P}^n}(t)$, then there are $a, b \geq 0$ such that, up to a twist by $\mathcal{O}_{\mathbb{P}^n}(s)$, E or E^* is the cokernel of an injective map:*

$$\Omega_{\mathbb{P}^n}^2(1)^b \rightarrow \mathcal{O}_{\mathbb{P}^n}(-1)^a$$

ii) *If E is indecomposable and rigid, then there is $k \geq 1$ such that, up to tensoring with $\mathcal{O}_{\mathbb{P}^n}(s)$, E or E^* is the cokernel of an injective map:*

$$\Omega_{\mathbb{P}^n}^2(1)^{a_{\ell, k-1}} \rightarrow \mathcal{O}_{\mathbb{P}^n}(-1)^{a_{\ell, k}}$$

2010 *Mathematics Subject Classification.* 14F05; 13C14; 14J60.

Key words and phrases. Rigid Maximal Cohen-Macaulay modules, Veronese varieties.

Author partially supported by ANR GEOLMI ANR-11-BS03-0011.

iii) Conversely for any $k \geq 1$, there is a unique indecomposable bundle E_k having a resolution of the form:

$$0 \rightarrow \Omega_{\mathbb{P}^n}^2(1)^{a_{\ell,k-1}} \rightarrow \mathcal{O}_{\mathbb{P}^n}(-1)^{a_{\ell,k}} \rightarrow E_k \rightarrow 0,$$

and both E_k and E_k^* are ACM and exceptional.

In the previous statement, it is understood that a bundle E is *exceptional* if it is rigid, *simple* (i.e. $\mathrm{Hom}_{\mathbb{P}^n}(E, E) \simeq \mathbf{k}$) and $\mathrm{Ext}_{\mathbb{P}^n}^i(E, E) = 0$ for $i \geq 2$. Next, we write $\Omega_{\mathbb{P}^n}^p = \wedge^p \Omega_{\mathbb{P}^n}^1$ for the bundle of differential p -forms on \mathbb{P}^n .


Remark 2. Part (i) of Theorem 1 is a version of Iyama-Yoshino's general results on Veronese rings [IY08, Theorem 9.1 and 9.3], to the effect that for $\{n, d\} = \{2, 3\}$ the stable category of MCM modules is equivalent to the category of representations of a certain Kronecker quiver. However our result is algorithmic, for it provides the representation associated with an MCM module via Beilinson spectral sequence applied to the corresponding ACM bundle.

Remark 3. The rank of the bundle E_k is given by the Fibonacci number between $a_{3,k-1}$ and $a_{3,k}$ in case $(n, d) = (2, 3)$. In this case E_{2k} (respectively, E_{2k+1}) is the k -th sheafified syzygy occurring in the resolution of $\mathcal{O}_{\mathbb{P}^2}(1)$ (respectively, of $\mathcal{O}_{\mathbb{P}^2}(2)$) over the Veronese ring, twisted by $\mathcal{O}_{\mathbb{P}^2}(3(k-1))$. A similar result holds for $(n, d) = (3, 2)$.

As for notation, we write small letters for the dimension of a space in capital letter, for instance $h^i(\mathbb{P}^n, E) = \dim_{\mathbf{k}} H^i(\mathbb{P}^n, E)$. We also write $\chi(E, F) = \sum (-1)^i \mathrm{ext}_{\mathbb{P}^n}^i(E, F)$ and $\chi(E) = \chi(\mathcal{O}_{\mathbb{P}^n}, E)$. $\delta_{i,j}$ is Kronecker's delta.

1. FIBONACCI BUNDLES

1.1. Let us write Υ_{ℓ} for the ℓ -th Kronecker quiver, namely the oriented graph with two vertices \mathbf{e}_0 and \mathbf{e}_1 , and ℓ arrows from \mathbf{e}_0 to \mathbf{e}_1 . A representation R of Υ_{ℓ} , with dimension vector (a, b) is the choice of ℓ matrices of size $a \times b$.



We identify a basis of $H^0(\mathbb{P}^n, \Omega_{\mathbb{P}^n}(2))$ with the set of $\ell = \binom{n+1}{2}$ arrows of Υ_{ℓ} . Then the derived category of finite-dimensional representations of Υ_{ℓ} embeds into the derived category of $\mathcal{O}_{\mathbb{P}^n}$ -modules by sending R to the cone $\Phi(R)$ of the morphism e_R associated with R according to this identification:

$$\Phi(R)[-1] \rightarrow \mathcal{O}_{\mathbb{P}^n}(-1)^a \xrightarrow{e_R} \Omega_{\mathbb{P}^n}(1)^b,$$

where we denote by $[-1]$ the shift to the right of complexes. It is clear that:

$$\mathrm{Ext}_{\mathbb{P}^n}^i(\Phi(R), \Phi(R)) \simeq \mathrm{Ext}_{\Upsilon_{\ell}}^i(R, R), \quad \text{for all } i.$$

1.2. We will use Kac's classification of rigid Υ_{ℓ} -modules as Schur roots (hence the restriction on \mathbf{k}), which is also one of the main ingredients in Iyama-Yoshino's proof. By [Kac80, Theorem 4], any non-zero rigid Υ_{ℓ} -module is a direct sum of rigid simple representations of the form R_k , for some $k \in \mathbb{Z}$, where R_k is defined as the unique indecomposable representation of Υ_{ℓ} with dimension vector $(a_{\ell,k-1}, a_{\ell,k})$ for $k \geq 1$, or $(a_{\ell,-k}, a_{\ell,1-k})$ for $k \leq 0$.

Set $F_k = \Phi(R_k)$ for $k \geq 1$, and $F_k = \Phi(R_k)[-1]$ for $k \leq 0$. It turns out that F_k is an exceptional locally free sheaf, called a *Fibonacci bundle*, cf. [Bra08]. We rewrite the defining exact sequences of F_k :

$$(2) \quad \begin{aligned} 0 \rightarrow \mathcal{O}_{\mathbb{P}^n}(-1)^{a_{\ell,k-1}} \rightarrow \Omega_{\mathbb{P}^n}(1)^{a_{\ell,k}} \rightarrow F_k \rightarrow 0, & \quad \text{for } k \geq 1, \\ 0 \rightarrow F_k \rightarrow \mathcal{O}_{\mathbb{P}^n}(-1)^{a_{\ell,-k}} \rightarrow \Omega_{\mathbb{P}^n}(1)^{a_{\ell,1-k}} \rightarrow 0, & \quad \text{for } k \leq 0. \end{aligned}$$

1.3. Here is a lemma on the cohomology of Fibonacci bundles.

Lemma 4. *For $k \geq 1$, the only non-vanishing intermediate cohomology of F_k is:*

$$h^1(\mathbb{P}^n, F_k(-1)) = a_{\ell, k-1} \quad h^{n-1}(\mathbb{P}^n, F_k(-n)) = a_{\ell, k}.$$

Proof. We consider the left and right mutation endofunctors of the derived category of coherent sheaves on \mathbb{P}^n , that associate with a pair (E, F) of complexes, two complexes denoted respectively by $R_F E$ and $L_E F$. These are the cones of the natural evaluation maps $f_{E,F}$ and $g_{E,F}$:

$$E \xrightarrow{f_{E,F}} \mathrm{RHom}_{\mathbb{P}^n}(E, F)^* \otimes F \rightarrow R_F E, \quad L_E F \rightarrow \mathrm{RHom}_{\mathbb{P}^n}(E, F) \otimes E \xrightarrow{g_{E,F}} F.$$

It is well-known (cf. [Bra08]) that the Fibonacci bundles F_k can be defined recursively from $F_0 = \mathcal{O}_{\mathbb{P}^n}(-1)$ and $F_1 = \Omega_{\mathbb{P}^n}(1)$ by setting:

$$(3) \quad F_{k+1} = R_{F_k} F_{k-1}, \quad \text{for } k \geq 1,$$

$$(4) \quad F_{k-1} = L_{F_k} F_{k+1}, \quad \text{for } k \leq 0.$$

This way, for any $k \in \mathbb{Z}$ we get a natural exact sequence:

$$(5) \quad 0 \rightarrow F_{k-1} \rightarrow (F_k)^\ell \rightarrow F_{k+1} \rightarrow 0.$$

Over \mathbb{P}^n , we consider the full exceptional sequence:

$$(\mathcal{O}_{\mathbb{P}^n}(-1), \Omega_{\mathbb{P}^n}(1), \mathcal{O}_{\mathbb{P}^n}, \mathcal{O}_{\mathbb{P}^n}(2), \mathcal{O}_{\mathbb{P}^n}(3), \dots, \mathcal{O}_{\mathbb{P}^n}(n-1)),$$

obtained from the standard collection $(\mathcal{O}_{\mathbb{P}^n}(-1), \dots, \mathcal{O}_{\mathbb{P}^n}(n-1))$, by the mutation $\Omega_{\mathbb{P}^n}(-1) \simeq L_{\mathcal{O}_{\mathbb{P}^n}} \mathcal{O}_{\mathbb{P}^n}(1)$ (all the terminology and results we need on exceptional collections are contained in [Bon89]). By (3), we can replace the previous exceptional sequence with:

$$(F_{k-1}, F_k, \mathcal{O}_{\mathbb{P}^n}, \mathcal{O}_{\mathbb{P}^n}(2), \dots, \mathcal{O}_{\mathbb{P}^n}(n-1)),$$

Right-mutating F_{k-1} through the full collection, we must get back $F_{k-1} \otimes \omega_{\mathbb{P}^n}^* \simeq F_{k-1}(n+1)$. So, using (5), we get a long exact sequence:

$$(6) \quad 0 \rightarrow F_{k+1} \rightarrow \mathcal{O}_{\mathbb{P}^n}^{u_1} \rightarrow \mathcal{O}_{\mathbb{P}^n}(2)^{u_2} \rightarrow \dots \rightarrow \mathcal{O}_{\mathbb{P}^n}(n-1)^{u_{n-1}} \rightarrow F_{k-1}(n+1) \rightarrow 0,$$

for some integers u_i . Now by (2) we get:

$$H^i(\mathbb{P}^n, F_k(t)) = 0 \quad \text{for:} \quad \begin{cases} 2 \leq i \leq n-2, & \forall t, \\ i = 0, & t \leq 0, \\ i = 1, & t \neq -1, \\ i = n-1, & t \geq 1-n. \end{cases}$$

The required non-vanishing cohomology of F_k appears again from (2). So it only remains to check that $H^{n-1}(\mathbb{P}^n, F_k(t)) = 0$ for $t \leq -n-1$. But this is clear by induction once we twist (6) by $\mathcal{O}_{\mathbb{P}^n}(t)$, and take cohomology. \square

1.4. We compute the Ext groups between pairs of Fibonacci bundles.

Lemma 5. *For any pair of integers $j \geq k+1$ we have:*

$$\mathrm{ext}_{\mathbb{P}^n}^i(F_j, F_k) = \delta_{1,i} a_{\ell, j-k-1}, \quad \mathrm{ext}_{\mathbb{P}^n}^i(F_k, F_j) = \delta_{0,i} a_{\ell, j-k+1}.$$

Proof. The formulas hold for $k = j$ since F_k is exceptional, and we easily compute $\chi(F_j, F_k) = -a_{\ell, j-k-1}$ and $\chi(F_k, F_j) = a_{\ell, j-k+1}$ (for instance by computing χ of Υ_ℓ -modules via the Cartan form and using faithfulness of Φ).

The second formula is proved once we show $\mathrm{Ext}_{\mathbb{P}^n}^i(F_k, F_j) = 0$ for $i \geq 1$. In fact, since the category of Υ_ℓ -representations is hereditary, the second formula holds if $k \leq 0$ for in this case $F_k \simeq \Phi(R_k)[-1]$. By the same reason, we only have to check it for $i = 1$. Using (5), this vanishing holds for j if it does for $j-1$ and $j-2$. Since the statement is clear when extended to $j = k$, it suffices to check $\mathrm{Ext}_{\mathbb{P}^n}^1(F_k, F_{k-1}) = 0$.

Since $\chi(F_k, F_{k-1}) = 0$, $\text{Hom}_{\mathbb{P}^n}(F_k, F_{k-1}) = 0$ will do the job. However, any nonzero map $F_k \rightarrow F_{k-1}$ would give, again by (5), a non-scalar endomorphism of F_k , which cannot exist since F_k is simple. The second formula is now proved.

As for the first formula, again we see that it holds if $k \leq 0$ and $j \geq 1$ once we check it for $i = 0$. However using repeatedly (5) we see that a non-zero map $F_j \rightarrow F_k$ leads to an endomorphism of F_j which factors through F_k this is absurd for F_j is simple. When j, k have the same sign, the first formula has to be checked for $i = 2$ only. Moreover, we have just proved the statement for $j = k + 1$, and using (5) and exceptionality of F_k we get it for $j = k + 2$. By iterating this argument we get the statement for any $j \geq k + 1$. \square

Remark 6. This lemma holds more generally (with the same proof) for any exceptional pair (F_0, F_1) of objects on a projective k -variety X , with $\text{hom}_X(F_0, F_1) = \ell$, by defining recursively F_k for all $k \in \mathbb{Z}$ by (3) and (4).

2. RIGID ACM BUNDLES ON THE THIRD VERONESE SURFACE

We prove here Theorem 1 in case $(n, d) = (2, 3)$.

2.1. Let us first prove (i). So let E be an indecomposable vector bundle on \mathbb{P}^2 satisfying (1). Without loss of generality, we may replace E by $E(s)$, where s is the smallest integer such that $h^0(\mathbb{P}^2, E(s)) \neq 0$. Set $\alpha_{i,j} = h^i(\mathbb{P}^2, E(-j))$. Of course, $\alpha_{0,j} = 0$ if and only if $j \geq 0$. The Beilinson complex F associated with E (see for instance [Huy06, Chapter 8]) reads:

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^2}(-1)^{\alpha_{1,2}} \xrightarrow{d_0} \begin{array}{c} \mathcal{O}_{\mathbb{P}^2}(-1)^{\alpha_{2,2}} \\ \oplus \\ \Omega_{\mathbb{P}^2}(1)^{\alpha_{1,1}} \\ \oplus \\ \mathcal{O}_{\mathbb{P}^2}^{\alpha_{0,0}} \end{array} \xrightarrow{d_1} \begin{array}{c} \Omega_{\mathbb{P}^2}(1)^{\alpha_{2,1}} \\ \oplus \\ \mathcal{O}_{\mathbb{P}^2}^{\alpha_{1,0}} \end{array} \xrightarrow{d_2} \mathcal{O}_{\mathbb{P}^2}^{\alpha_{2,0}} \rightarrow 0.$$

The term consisting of three summands in the above complex sits in degree 0 (we call it *middle term*), and the cohomology of this complex is E . By condition (1), at least one of the $\alpha_{1,j}$ is zero, for $j = 0, 1, 2$.

If $\alpha_{1,2} = 0$, then $d_0 = 0$. By minimality of the Beilinson complex the restriction of d_1 to the summand $\mathcal{O}_{\mathbb{P}^2}^{\alpha_{0,0}}$ of the middle term is also zero. Therefore $\mathcal{O}_{\mathbb{P}^2}^{\alpha_{0,0}}$ is a direct summand of E , so $E \simeq \mathcal{O}_{\mathbb{P}^2}$ by indecomposability of E .

If $\alpha_{1,1} = 0$, then the non-zero component of d_0 is just a map $\mathcal{O}_{\mathbb{P}^2}(-1)^{\alpha_{1,2}} \rightarrow \mathcal{O}_{\mathbb{P}^2}^{\alpha_{0,0}}$, and a direct summand of E is the cokernel of this map. By indecomposability of E , in this case $E(-1)$ has a resolution of the desired form with $a = \alpha_{0,0}$ and $b = \alpha_{1,2}$.

2.2. It remains to look at the case $\alpha_{1,0} = 0$. Note that the restriction of d_1 to $\Omega_{\mathbb{P}^2}(1)^{\alpha_{1,1}} \oplus \mathcal{O}_{\mathbb{P}^2}^{\alpha_{0,0}}$ is zero, which implies that a direct summand of E (hence all of E by indecomposability) has the resolution:

$$(7) \quad 0 \rightarrow \mathcal{O}_{\mathbb{P}^2}(-1)^{\alpha_{1,2}} \xrightarrow{d_0} \Omega_{\mathbb{P}^2}(1)^{\alpha_{1,1}} \oplus \mathcal{O}_{\mathbb{P}^2}^{\alpha_{0,0}} \rightarrow E \rightarrow 0$$

and $\alpha_{2,j} = 0$ for $j = 0, 1, 2$. We compute $\chi(E(-3)) = 3\alpha_{1,2} - 3\alpha_{1,1} + \alpha_{0,0}$, so:

$$h^0(\mathbb{P}^2, E^*) = h^2(\mathbb{P}^2, E(-3)) = 3\alpha_{1,2} - 3\alpha_{1,1} + \alpha_{0,0}.$$

If this value is positive, then there is a non-trivial morphism $g : E \rightarrow \mathcal{O}_{\mathbb{P}^2}$, and since $\alpha_{0,0} \neq 0$ there also exists $0 \neq f : \mathcal{O}_{\mathbb{P}^2} \rightarrow E$. So E has an endomorphism factoring through $\mathcal{O}_{\mathbb{P}^2}$, a contradiction.

Hence we may assume $3\alpha_{1,2} - 3\alpha_{1,1} + \alpha_{0,0}$, in other words $\alpha_{0,3} = 0$. Therefore, the Beilinson complex associated with $E(-1)$ gives a resolution:

$$0 \rightarrow E(-1) \rightarrow \Omega_{\mathbb{P}^2}(1)^{\alpha_{1,2}} \rightarrow \mathcal{O}_{\mathbb{P}^2}^{\alpha_{1,1}} \rightarrow 0.$$

It is easy to convert this resolution into the form we want by the diagram:

$$\begin{array}{ccccccc} & & 0 & & 0 & & \\ & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & E(-1) & \longrightarrow & \Omega_{\mathbb{P}^2}(1)^{\alpha_{1,2}} & \longrightarrow & \mathcal{O}_{\mathbb{P}^2}^{\alpha_{1,1}} \rightarrow 0 \\ & & \downarrow & & \downarrow & & \parallel \\ 0 & \longrightarrow & \mathcal{O}_{\mathbb{P}^2}^{3\alpha_{1,2}-\alpha_{1,1}} & \longrightarrow & \mathcal{O}_{\mathbb{P}^2}^{3\alpha_{1,2}} & \longrightarrow & \mathcal{O}_{\mathbb{P}^2}^{\alpha_{1,1}} \rightarrow 0 \\ & & \downarrow & & \downarrow & & \\ & & \mathcal{O}_{\mathbb{P}^2}(1)^{\alpha_{1,2}} & = & \mathcal{O}_{\mathbb{P}^2}(1)^{\alpha_{1,2}} & & \\ & & \downarrow & & \downarrow & & \\ & & 0 & & 0 & & \end{array}$$

From the leftmost column, it follows that E^* has a resolution of the desired form, with $a = 3\alpha_{1,2} - \alpha_{1,1}$ and $b = \alpha_{1,2}$. Claim (i) is thus proved.

2.3. Let us now prove (ii). We may assume that $\text{rk}(E) > 1$. We check that E being indecomposable and rigid forces E to be simple, so that it has no endomorphism factoring through $\mathcal{O}_{\mathbb{P}^2}(t)$ and thus (i) applies. In the previous proof, we used this condition only for $\alpha_{1,0} = 0$, and (i) will apply if $\alpha_{0,3} = 0$.

2.4. We work with $\alpha_{1,0} = 0$. Let e be the restricted map $e : \mathcal{O}_{\mathbb{P}^2}(-1)^{\alpha_{1,2}} \rightarrow \Omega_{\mathbb{P}^2}(1)^{\alpha_{1,1}}$ extracted from d_0 and let F be its cone, shifted by 1:

$$(8) \quad F \rightarrow \mathcal{O}_{\mathbb{P}^2}(-1)^{\alpha_{1,2}} \xrightarrow{e} \Omega_{\mathbb{P}^2}(1)^{\alpha_{1,1}}.$$

This is a complex with two terms, and its cohomology is concentrated in degrees zero and one, namely $\mathcal{H}^0 F \simeq \ker(e)$ and $\mathcal{H}^1 F \simeq \text{coker}(e)$. From (7) we easily see that F fits into a distinguished triangle:

$$(9) \quad F \rightarrow \mathcal{O}_{\mathbb{P}^2}^{\alpha_{0,0}} \rightarrow E.$$

Applying $\text{Hom}_{\mathbb{P}^2}(\mathcal{O}_{\mathbb{P}^2}, -)$ to (8), we get $\text{Ext}_{\mathbb{P}^2}^i(\mathcal{O}_{\mathbb{P}^2}, F) = 0$ for all i , so:

$$\text{Ext}_{\mathbb{P}^2}^i(F, F) \simeq \text{Ext}_{\mathbb{P}^2}^i(E, F[1]), \quad \text{for all } i.$$

Also, we know that $H^2(\mathbb{P}^2, E^*) = 0$, so applying $\text{Hom}_{\mathbb{P}^2}(E, -)$ to (9) we get:

$$\text{Ext}_{\mathbb{P}^2}^1(E, E) \rightarrow \text{Ext}_{\mathbb{P}^2}^1(E, F[1]) \rightarrow 0.$$

Putting this together, we obtain a surjection:

$$\text{Ext}_{\mathbb{P}^2}^1(E, E) \twoheadrightarrow \text{Ext}_{\mathbb{P}^2}^1(F, F) \simeq \text{Ext}_{\mathbb{P}^2}^1(R, R),$$

with $F \simeq \Phi(R)$. We understand now that, if E is rigid, then also R is.

2.5. If R is rigid, then by §1.2, R is a direct sum of rigid simple representations of the form R_k . Therefore, cohomology of (9) gives an exact sequence:

$$0 \rightarrow \bigoplus_{i \leq 0} F_i^{r_i} \rightarrow \mathcal{O}_{\mathbb{P}^2}^{\alpha_{0,0}} \rightarrow E \rightarrow \bigoplus_{i \geq 1} F_i^{r_i} \rightarrow 0,$$

for some integers r_i . If only R_i with $i \leq 0$ appear, then we are done by §2.3. Indeed, in that case E is globally generated, so $H^0(\mathbb{P}^2, E) \neq 0$ implies $H^0(\mathbb{P}^2, E^*) = 0$ for otherwise $\mathcal{O}_{\mathbb{P}^2}$ would be a direct summand of E .

If some R_i appears with $i \geq 1$, we call I the (non-zero) image of the middle map in the previous exact sequence, and we check $\text{Ext}_{\mathbb{P}^2}^1(F_j, I) = 0$ for all $j \geq 1$, which

contradicts E being indecomposable. To check this, note that $\mathrm{Hom}_{\mathbb{P}^2}(F_j, -)$ gives an exact sequence:

$$\mathrm{Ext}_{\mathbb{P}^2}^1(F_j, \mathcal{O}_{\mathbb{P}^2})^{\alpha_{0,0}} \rightarrow \mathrm{Ext}_{\mathbb{P}^2}^1(F_j, I) \rightarrow \bigoplus_{i \leq 0} \mathrm{Ext}_{\mathbb{P}^2}^2(F_j, F_i)^{r_i}.$$

The leftmost term vanishes by Serre duality and Lemma 4. The rightmost term is zero by Lemma 5. Part (ii) is now proved.

2.6. The statement (iii) is clear by Lemma 4 and by exceptionality of Fibonacci bundles. The fact that E^* is also ACM is obvious by Serre duality.

Remark 7. If \mathbf{k} is algebraically closed of characteristic zero, we may apply [Dre86, Corollaire 7], to the effect that a rigid bundle is a direct sum of exceptional bundles. So, at the price of relying on this result, from (i) we may deduce directly (ii) via Kac's theorem.

3. ACM BUNDLES ON THE SECOND VERONESE THREEFOLD

The techniques we have just seen apply to the embedding of \mathbb{P}^3 in \mathbb{P}^9 by quadratic forms. Again we replace E with the $E(s)$, where s is the smallest integer such that E has non-zero global sections, and set $\alpha_{i,j} = h^i(\mathbb{P}^3, E(-j))$. If (1) gives $\alpha_{1,1} = \alpha_{2,1} = 0$, then $E(-1)$ has the desired resolution. On the other hand, if (1) tells $\alpha_{1,0} = \alpha_{2,0} = \alpha_{1,2} = \alpha_{2,2} = 0$, then we are left with a resolution of the form:

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^3}(-1)^{\alpha_{1,3}} \xrightarrow{d_0} \Omega_{\mathbb{P}^3}(1)^{\alpha_{1,1}} \oplus \mathcal{O}_{\mathbb{P}^3}^{\alpha_{0,0}} \rightarrow E \rightarrow 0.$$

This time we also have $\alpha_{0,4} = 0$, and $\alpha_{1,4} = \alpha_{2,4} = 0$ again by (1), and simplicity of E gives $\alpha_{3,4} = h^0(\mathbb{P}^3, E^*) = 0$. So $E(-1)$ has a resolution like:

$$0 \rightarrow E(-1) \rightarrow \Omega_{\mathbb{P}^3}^2(2)^{\alpha_{1,3}} \rightarrow \mathcal{O}_{\mathbb{P}^3}^{\alpha_{1,1}} \rightarrow 0.$$

Then, using the same trick as in the proof of the previous theorem, we see that E^* has the desired resolution, with $a = 6\alpha_{1,3} - \alpha_{1,1}$ and $b = \alpha_{1,3}$.

This proves the first statement. The rest follows by the same path. Drezet's theorem as shortcut for (i) \Rightarrow (ii) may be replaced by [HZ13].

Remark 8. It should be noted that, in [IY08, Theorem 1.2 and Theorem 1.3], the ACM bundle E on the given Veronese variety is assumed to have a rigid module of global sections. This implies, respectively, $\mathrm{Ext}_{\mathbb{P}^2}^1(E, E(3t)) = 0$, or $\mathrm{Ext}_{\mathbb{P}^3}^1(E, E(2t)) = 0$, for all $t \in \mathbb{Z}$. A priori, this is a stronger requirement than just $\mathrm{Ext}_{\mathbb{P}^n}^1(E, E) = 0$. However, our proof shows that the two conditions are equivalent for ACM bundles.

4. RIGID ACM BUNDLES HIGHER VERONESE SURFACES

Assume \mathbf{k} algebraically closed. The next result shows that, for $d \geq 4$, the class of rigid ACM bundles on d -fold Veronese surfaces contains the class of exceptional bundles on \mathbb{P}^2 , which is indeed quite complicated, cf. [DLP85]. At least if $\mathrm{char}(\mathbf{k}) = 0$, the two classes coincide by [Dre86, Corollaire 7].

Theorem 9. *Let F be an exceptional bundle on \mathbb{P}^2 and fix $d \geq 4$. Then there is an integer t such that $E = F(t)$ satisfies (1).*

Proof. It is known that F is actually stable by [DLP85]. This implies that F has natural cohomology by [HL93], i.e. for all $t \in \mathbb{Z}$ there is at most one i such that $H^i(\mathbb{P}^2, F(t)) \neq 0$. Then, $H^1(\mathbb{P}^2, F(t)) \neq 0$ if and only if $\chi(F(t)) < 0$.

Let now that r , c_1 and c_2 be the rank and the Chern classes of F . Computing χ by additivity, we see that $\chi(F(t))$ is a polynomial of degree 2 in t , of dominant term $r/2$, whose discriminant is:

$$\Delta = c_1^2(1-r) + r(2c_2 + r/4) = -\chi(F, F) + 5r^2/4.$$

So, using $\chi(F, F) = 1$, we get $\Delta = -1 + 5r^2/4$. Therefore, the roots of $\chi(F(t))$ differ at most by:

$$\lceil \frac{2\sqrt{\Delta}}{r} \rceil = \lceil \frac{\sqrt{5r^2-4}}{r} \rceil \leq 3.$$

Then, there are at most three consecutive integers t_0 , $t_0 + 1$, $t_0 + 2$ such that $H^1(\mathbb{P}^2, F(t_0 + j)) \neq 0$ for $j = 0, 1, 2$. This means that $E = F(t_0 - 1)$ satisfies (1) for any choice of $d \geq 4$. \square

Acknowledgements. I would like to thank F.-O. Schreyer, J. Pons Llopis and M. C. Brambilla for useful comments and discussions.

REFERENCES

- [Bon89] ALEXEI I. BONDAL. *Representations of associative algebras and coherent sheaves*. Izv. Akad. Nauk SSSR Ser. Mat., 53(1):25–44, 1989.
- [Bra08] MARIA CHIARA BRAMBILLA. *Cokernel bundles and Fibonacci bundles*. Math. Nachr., 281(4):499–516, 2008.
- [DLP85] JEAN-MARC DREZET AND JOSEPH LE POTIER. *Fibrés stables et fibrés exceptionnels sur \mathbb{P}_2* . Ann. Sci. École Norm. Sup. (4), 18(2):193–243, 1985.
- [Dre86] JEAN-MARC DREZET. *Fibrés exceptionnels et suite spectrale de Beilinson généralisée sur $\mathbb{P}_2(\mathbb{C})$* . Math. Ann., 275(1):25–48, 1986.
- [HL93] ANDRÉ HIRSCHOWITZ AND YVES LASZLO. *Fibrés génériques sur le plan projectif*. Math. Ann., 297(1):85–102, 1993.
- [Huy06] DANIEL HUYBRECHTS. *Fourier-Mukai transforms in algebraic geometry*. Oxford Mathematical Monographs. The Clarendon Press Oxford University Press, Oxford, 2006.
- [HZ13] DIETER HAPPEL AND DAN ZACHARIA. *A note on sheaves without self-extensions on the projective n -space*. Proc. Amer. Math. Soc., 141(10):3383–3390, 2013.
- [IY08] OSAMU IYAMA AND YUJI YOSHINO. *Mutation in triangulated categories and rigid Cohen-Macaulay modules*. Invent. Math., 172(1):117–168, 2008.
- [Kac80] VICTOR G. KAC. *Infinite root systems, representations of graphs and invariant theory*. Invent. Math., 56(1):57–92, 1980.
- [KMVdB11] BERNHARD KELLER, DANIEL MURFET, AND MICHEL VAN DEN BERGH. *On two examples by Iyama and Yoshino*. Compos. Math., 147(2):591–612, 2011.
- [KR08] BERNHARD KELLER AND IDUN REITEN. *Acyclic Calabi-Yau categories*. Compos. Math., 144(5):1332–1348, 2008. With an appendix by Michel Van den Bergh.

E-mail address: daniele.faenzi@univ-pau.fr

UNIVERSITÉ DE PAU ET DES PAYS DE L'ADOUR
 AVENUE DE L'UNIVERSITÉ - BP 576
 64012 PAU CEDEX - FRANCE
 URL: <http://univ-pau.fr/~faenzi/>