



HAL
open science

Construction of Hadamard states by characteristic Cauchy problem

Christian Gérard, Michal Wrochna

► **To cite this version:**

Christian Gérard, Michal Wrochna. Construction of Hadamard states by characteristic Cauchy problem. 2014. hal-01067150v1

HAL Id: hal-01067150

<https://hal.science/hal-01067150v1>

Preprint submitted on 23 Sep 2014 (v1), last revised 3 Jan 2017 (v4)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

CONSTRUCTION OF HADAMARD STATES BY CHARACTERISTIC CAUCHY PROBLEM

C. GÉRARD AND M. WROCHNA

ABSTRACT. We construct Hadamard states for Klein-Gordon fields in a spacetime M_0 equal to the interior of the future lightcone C from a base point p in a globally hyperbolic spacetime (M, g) .

Under some regularity conditions at future infinity of C , we identify a boundary symplectic space of functions on C , which allows to construct states for Klein-Gordon quantum fields in M_0 from states on the CCR algebra associated to the boundary symplectic space. We formulate the natural microlocal condition on the boundary state on C ensuring that the bulk state it induces in M_0 satisfies the Hadamard condition.

Using pseudodifferential calculus on the cone C we construct a large class of Hadamard boundary states on the boundary with pseudodifferential covariances, and characterize the pure states among them. We then show that these pure boundary states induce pure Hadamard states in M_0 .

1. INTRODUCTION

Hadamard states are widely accepted as physically admissible states for non-interacting quantum fields on a curved spacetime, one of the main reasons being their link with the renormalization of the stress-energy tensor, a basic step in the formulation of semi-classical Einstein equations.

For Klein-Gordon fields, the construction of Hadamard states amounts to finding bi-solutions of the Klein-Gordon equation with a specified wave front set (that is, verifying the *microlocal spectrum condition*) and satisfying additionally a positivity property.

There exist several methods to construct Hadamard states for Klein-Gordon fields: the first method relies on the Fulling-Narowich-Wald deformation argument [FNW], which reduces the construction of Hadamard states on an arbitrary spacetime to the case of ultrastatic spacetimes, where vacuum or thermal states are easily shown to be Hadamard states.

The second approach, used in [GW], uses pseudodifferential calculus on a fixed Cauchy surface Σ in (M, g) and relies on the construction of a parametrix for the Cauchy problem on Σ . To use pseudodifferential calculus, some restrictions on Σ and on the behavior of the metric g at spatial infinity are necessary. On the other hand, the method in [GW] produces a large classes of rather explicit Hadamard states, whose covariances, expressed in terms of Cauchy data are pseudodifferential operators.

Another method, initiated by Moretti [Mo1, Mo2] applies to *conformal field equations*, like the conformal wave equation, on an *asymptotically flat vacuum spacetime* (M_0, g_0) . By asymptotic flatness, there exists a metric \tilde{g}_0 , conformal to g_0 , and a spacetime (M, \tilde{g}) such that (M_0, \tilde{g}_0) can be causally embedded as an open set in (M, \tilde{g}) , with the boundary $C = \partial M_0$ of M_0 being *null* in (M, \tilde{g}) . States on the

2010 *Mathematics Subject Classification.* 81T13, 81T20, 35S05, 35S35.

Key words and phrases. Hadamard states, microlocal spectrum condition, pseudo-differential calculus, characteristic Cauchy problem, curved spacetimes.

boundary symplectic space, containing the traces on C of solutions of the wave equation in M_0 , naturally induce states inside M_0 .

This method has been successfully applied in [Mo1, Mo2] to construct a distinguished Hadamard state for asymptotically flat vacuum spacetimes with past time infinity and then extended to several other geometrical situations in [DMP1, DMP2]. Further results also include generalization to Maxwell fields [DS] and linearized gravity [BDM].

In the present paper we rework systematically the above strategy in order to construct a large class of Hadamard states (instead of a preferred single one) and to characterize the pure ones. For the sake of clarity, we do not impose geometrical assumptions on M_0 that allow to correctly embed it in a larger spacetime M .

Instead we go the other way around and work in an a priori arbitrary globally hyperbolic spacetime M , fix a base point p and consider the interior of the future lightcone

$$C := \partial J^+(p) \setminus \{p\}$$

as the spacetime M_0 of main interest, i.e. $M_0 := I^+(p)$.

We make the following assumption on the geometry of C .

Hypothesis 1.1. *We assume that there exists $f \in C^\infty(M)$ such that:*

- (1) $C \subset f^{-1}(\{0\})$, $\nabla_a f \neq 0$ on C , $\nabla_a f(p) = 0$, $\nabla_a \nabla_b f(p) = -2g_{ab}(p)$,
- (2) *the vector field $\nabla^a f$ is complete on C .*

Using Hypothesis 1.1 one can construct coordinates (f, s, θ) near C , such that $C \subset \{f = 0\}$ and

$$g|_C = -2df ds + h(s, \theta)d\theta^2,$$

where $h(s, \theta)d\theta^2$ is a Riemannian metric on \mathbb{S}^{d-1} .

Such choice of coordinates allows one to identify C with $\tilde{C} := \mathbb{R} \times \mathbb{S}^{d-1}$. A natural space of smooth functions on \tilde{C} is then provided by $\mathcal{H}(\tilde{C})$ — the intersection of Sobolev spaces of all orders, defined using the standard metric $m(\theta)d\theta^2$ on \mathbb{S}^{d-1} .

The *bulk-to-boundary correspondence* can be expressed in this setup as follows. For an appropriate choice of $\beta(s, \theta) \in C^\infty(M_0)$, the restriction map

$$\rho\phi := (\beta^{-1}\phi)|_C, \quad \phi \in C_{\text{sc}}^\infty(M_0)$$

is a monomorphism¹ between the symplectic space of smooth, space-compact solutions of $P_0 := P|_{M_0}$ (endowed with the usual symplectic form induced by the causal propagator) and $\mathcal{H}(\tilde{C})$, equipped with the symplectic form

$$(1.1) \quad \bar{g}_1 \sigma_C g_2 := \int_{\mathbb{R} \times \mathbb{S}^{d-1}} (\partial_s \bar{g}_1 g_2 - \bar{g}_1 \partial_s g_2) |m|^{\frac{1}{2}}(\theta) ds d\theta, \quad g_1, g_2 \in \mathcal{H}(\tilde{C}).$$

Thus, a quasi-free state on $(\mathcal{H}(\tilde{C}), \sigma_C)$ with two-point functions λ^\pm induces a unique quasi-free state on the usual symplectic space associated to P_0 .

Product-type pseudodifferential operators. In [GW] we have constructed Hadamard states whose two-point functions on a Cauchy surface Σ are pseudodifferential operators. In the present case, the obvious difference is that on the cone C , the coordinate s is distinguished both from the point of view of the microlocal spectrum condition (from now on abbreviated (μsc)) and in the expression (1.1) for the symplectic form. This suggests that one should rather consider *product-type* pseudodifferential operators $\Psi^{p_1, p_2}(\tilde{C})$ with symbols satisfying estimates:

$$|\partial_s^{\alpha_1} \partial_\sigma^{\beta_1} \partial_\theta^{\alpha_2} \partial_\eta^{\beta_2} a(s, \theta, \sigma, \eta)| \in O(\langle \sigma \rangle^{p_1 - |\beta_1|} \langle \eta \rangle^{p_2 - |\beta_2|})$$

¹By monomorphism of symplectic spaces we mean an injective linear map that intertwines the symplectic forms.

in the covariables $\xi = (\sigma, \eta)$ relative to the decomposition $\tilde{C} = \mathbb{R} \times \mathbb{S}^{d-1}$. Actually, to cope with the issue that σ_C is defined using an operator $D_s := i^{-1}\partial_s$ whose spectrum is not separated from $\{0\}$ (analogously to the infrared problem in massless theories), we need to introduce a larger class $\tilde{\Psi}^{p_1, p_2}(\tilde{C})$ that includes some operators whose symbol is discontinuous at $\eta = 0$. Namely, we set

$$\tilde{\Psi}^{p_1, p_2}(\tilde{C}) := \Psi^{p_1, p_2}(\tilde{C}) + B^{-\infty}\Psi^{p_2}(\tilde{C}),$$

where $B^{-\infty}\Psi^{p_2}(\tilde{C})$ is the class of pseudodifferential operators of order p_2 (in the θ variables) with values in operators on \mathbb{R} that infinitely increase Sobolev regularity. Then for instance $|D_s| \otimes \mathbb{1}_\theta \in \tilde{\Psi}^{1,0}(\tilde{C})$ although it is not in the pseudodifferential class $\Psi^{1,0}(\tilde{C})$.

Summary of results. Our main results can be summarized as follows. We always assume Hypothesis 1.1. If E, F are topological vector spaces, we write $T : E \rightarrow F$ to mean $T : E \rightarrow F$ is linear and continuous.

- 1) For pairs² of two-point functions λ^\pm on C satisfying $\lambda^\pm : \mathcal{H}(C) \rightarrow \mathcal{H}(C)$, we give in Thm. 5.3 conditions on $\text{WF}(\lambda^\pm)$ that guarantee that the corresponding two-point functions on M_0 satisfy (μsc) . This is essentially an adaptation of the results of [Mo2] to our framework.
- 2) In Thm. 7.4 we construct a large class of Hadamard states by specifying their two-point functions $\lambda^\pm \in \tilde{\Psi}^{0,0}(\tilde{C})$ on the cone.
- 3) In Thm. 8.2 we characterize the subclass of Hadamard states constructed in **2)**, which additionally are pure on the symplectic space $(\mathcal{H}(\tilde{C}), \sigma_C)$ on the cone. It turns out that they can be parametrized by a single operator in $\tilde{\Psi}^{-\infty,0}(\tilde{C})$.
- 4) In Thm. 8.4 we prove that if $\dim M \geq 4$, then the pure states considered in **3)** induce pure states in the interior M_0 of the cone.

In Subsect. 2.3 we argue that Hypothesis 1.1 covers the case when M_0 is an asymptotically flat vacuum spacetime with future time infinity, after a conformal transformation. Thus, our result **4)** solves an open question by Moretti [Mo2] for $\dim M \geq 4$.

Characteristic Cauchy problem. The proof of our main result **4)** relies on the existence of a unique solution for the *characteristic Cauchy problem* (also called *Goursat problem* in the literature) in appropriate Sobolev spaces. Since it is of independent interest, we present below this auxiliary result.

Let Σ be a Cauchy surface in the future of $\{p\}$ and $\Sigma_0 := \Sigma \cap M_0$. Then Σ_0 individuates a compact region of the interior of the cone, namely

$$C_0 := (J^-(\Sigma_0; M) \cap C) \cup \{p\}.$$

Both Σ_0 and C_0 are compact sets in M_0 with smooth boundary $\partial\Sigma_0 = \partial C_0$. We denote by $H_0^1(\Sigma_0)$, $H_0^1(C_0)$ the respective restricted Sobolev spaces of order 1, i.e. the space of distributions in $H^1(\Sigma_0)$, $H^1(C_0)$ that vanish on the boundary. Let us denote by

$$U_{\Sigma_0} : H^1(\Sigma_0) \oplus L^2(\Sigma_0) \rightarrow C^0(\mathbb{R}, H^1(C_0)) \cap C^1(\mathbb{R}, L^2(C_0))$$

the map which assigns to Cauchy data on Σ_0 the corresponding finite energy solution for P_0 . In Subsect. 8.3 we prove the following result.

²We work with charged fields, in which case it is natural to associate a *pair* of two-point functions to a quasi-free state, cf. 3.2.1. The charged and neutral approaches are equivalent.

Theorem 1.1. *The map*

$$T : H_0^1(\Sigma_0) \oplus L^2(\Sigma_0) \rightarrow H_0^1(C_0) \\ \varphi \mapsto (U_{\Sigma_0} \varphi)|_{C_0}$$

is a homeomorphism. Moreover, if $\dim M \geq 4$ then $T(C_0^\infty(\Sigma_0) \oplus C_0^\infty(\Sigma_0))$ is dense in $|D_s|^{\frac{1}{2}}L^2(\tilde{C})$.

The first part of Thm. 1.1 provides in fact the solution to the characteristic Cauchy problem

$$\begin{cases} P_0 u = 0, \\ \rho u = \varphi, \quad \varphi \in H_0^1(C_0). \end{cases}$$

Our proof proceeds by reduction to a case already considered by Hörmander in [Hö2], namely when the characteristic surface is the graph of a Lipschitz function defined on a *compact* domain.

The second part of Thm. 1.1 asserts that there is no loss of information on the level of purity of states when going from the cone C to its interior M_0 . The precise form of the statement comes from the fact that the one-particle Hilbert space associated to our Hadamard states, i.e. the completion of $\mathcal{H}(\tilde{C})$ for the inner product $(\cdot | (\lambda^+ + \lambda^-) \cdot)$, equals $|D_s|^{\frac{1}{2}}L^2(\tilde{C})$. The validity of such result appears to be very delicate, it would be for instance problematic for $|D_s|^\alpha L^2(\tilde{C})$ with $\alpha < \frac{1}{2}$ instead of $\alpha = \frac{1}{2}$ and we do not know whether it holds for $d < 3$. The generalization of Thm. 1.1 to other geometrical situations is thus an interesting open problem, particularly relevant for the quantum field theoretical bulk-to-boundary correspondence.

At this point it is worth mentioning that beside Hörmander's work [Hö2] there is a considerable literature on the characteristic Cauchy problem for the Klein-Gordon equation, to mention only [BW, Ca, Do]. However, known results require either more regularity or conditions on the support of the solution (usually both) and as such cannot be directly applied in our problem. It is possible, though, that our method presented in Subsect. 8.4 can be used to bypass the often made space-compactness assumption.

Plan of the paper. In Sect. 2 we fix the geometric setup and outline the construction of null coordinates near the cone C . In Sect. 3 we briefly review the Klein-Gordon field in M_0 and the definition of Hadamard states. Sect. 4 is devoted to the so-called bulk-to-boundary correspondence, i.e. to the definition of a convenient symplectic space $(\mathcal{H}(\tilde{C}), \sigma_C)$ of functions on C , containing the traces on C of space-compact solutions in M_0 .

In Sect. 5, we formulate the Hadamard condition on C , i.e. the natural microlocal condition on the two-point functions of a quasi-free state on $(\mathcal{H}(\tilde{C}), \sigma_C)$ which ensures that the induced state in M_0 is a Hadamard state.

Sect. 6 is devoted to the pseudodifferential calculus on $\mathbb{R} \times \mathbb{S}^{d-1}$, more precisely to the ‘product-type’ classes, associated to bi-homogeneous symbols. We also describe more general operator classes which are pseudodifferential only in the variables in \mathbb{S}^{d-1} .

In Sect. 7 we construct large classes of Hadamard states on the cone, whose covariances belong to the operator classes introduced in Sect. 6. In Sect. 8 we characterize *pure* Hadamard states, and show that they induce pure states in M_0 . Finally in Sect. 9 we discuss the invariance of our classes of Hadamard states under change of null coordinates on C . Various technical results are collected in Appendix A.

2. GEOMETRIC SETUP

In this section we describe our geometrical setup and construct null coordinates near the cone C .

2.1. Future lightcone. We consider a globally hyperbolic spacetime (M, g) of dimension $\dim M = d + 1$.

As outlined in the introduction, we fix a base point $p \in M$, and consider

$$C = \partial J^+(p) \setminus \{p\}, \quad M_0 = I^+(p),$$

so that C is the future lightcone from p , with tip removed, and M_0 is the interior of C . From [Wa, Sect. 8.1] we know that M_0 is open, with

$$J^+(p) = \overline{M_0}, \quad \partial M_0 = \partial J^-(p) = C \cup \{p\}.$$

We assume Hypothesis 1.1, i.e. that there exists $f \in C^\infty(M)$ such that:

- (1) $C \subset f^{-1}(\{0\})$, $\nabla_a f \neq 0$ on C , $\nabla_a f(p) = 0$, $\nabla_a \nabla_b f(p) = -2g_{ab}(p)$,
- (2) the vector field $\nabla^a f$ is complete on C .

It follows that C is a smooth hypersurface, although \overline{C} is not smooth. Moreover since C is a null hypersurface, $\nabla^a f$ is tangent to C .

2.2. Causal structure. We now collect some useful results on the causal structure of M_0 and M .

Lemma 2.1. *Let $K \subset M_0$ be compact. Then:*

$$(2.1) \quad J^-(K) \cap J^+(p) \text{ is compact,}$$

$$(2.2) \quad J^+(K) \cap C = \emptyset.$$

Proof. (2.1) follows from [BGP, Lemma A.5.7]. Moreover if $V \subset M_0$ is open with $K \subset V$, we have $J^+(K) \subset I^+(V) \subset M_0$. Since $\partial J^-(p) = \partial M_0$ and M_0 is open, this implies (2.2). \square

The following lemma is due to Moretti [Mo1, Thm. 4.1 (a)]. If $K \subset M_0$, the notation $J^\pm(K; M_0)$ or $J^\pm(K; M)$ is used in the place of $J^\pm(K)$ to specify which causal structure one refers to.

Lemma 2.2. *The Lorentzian manifold (M_0, g) is globally hyperbolic. Moreover*

$$(2.3) \quad J^+(K; M_0) = J^+(K; M), \quad J^-(K; M_0) = J^-(K; M) \cap M_0, \quad \forall K \subset M_0.$$

The next proposition is also due to Moretti [Mo2, Lemma 4.3].

Proposition 2.3. *Let $K \subset M_0$ be compact. Then there exists a neighborhood U_1 of p in M such that no null geodesic starting from K intersects $\overline{C} \cap U_1$.*

2.3. Asymptotically flat spacetimes. In what follows we explain the relation between Hypothesis 1.1 and the geometrical assumptions met in the literature on Hadamard states [Mo1, Mo2, DS, BDM].

Let us consider two globally hyperbolic spacetimes (M_0, g_0) and (M, g) , where M_0 is an embedded submanifold of M . One introduces the following set of assumptions.

Hypothesis 2.1. *Suppose the spacetime (M, g) is such that:*

- (1) there exists $\Omega \in C^\infty(M)$ with $\Omega > 0$ on M_0 and $g|_{M_0} = \Omega^2 \upharpoonright_{M_0} g_0$,
- (2) there exists $i^- \in M$ such that $J^+(i^-; M)$ is closed and

$$M_0 = J^+(i^-; M) \setminus \partial J^+(i^-; M),$$

(3) g_0 solves the vacuum Einstein equations at least in a neighborhood of \mathcal{I}^-

$$\mathcal{I}^- := \partial J^+(i^-; M) \setminus \{i^-\},$$

(4) $\Omega = 0$ and $d\Omega \neq 0$ on \mathcal{I}^- , $d\Omega(i^-) = 0$, $\nabla_a \nabla_b \Omega(i^-) = -2g_{ab}(i^-)$,

(5) if $n^a := g^{ab} \nabla_b \Omega$, then there exists $\omega \in C^\infty(M)$, with $\omega > 0$ on $M_0 \cup \mathcal{I}^-$ and

(a) $\nabla_a(\omega^4 n^a) = 0$ on \mathcal{I}^- ,

(b) the vector field $\omega^{-1}n$ is complete on \mathcal{I}^- ,

Above, the symbols ∇_a refer to the metric g .

One says that (M_0, g_0) is an *asymptotically flat vacuum spacetime with past time infinity* i^- if there exists a spacetime (M, g) such that M_0 is an embedded submanifold of M and Hypothesis 2.1 is satisfied³.

Lemma 2.4. *Suppose (M_0, g_0) is an asymptotically flat vacuum spacetime with past time infinity i^- and let (M, g) satisfy Hypothesis 2.1. Then Hypothesis 1.1 is satisfied for $p := i^-$ and $f = \omega\Omega$.*

Note that actually only conditions (1), (2), (4) and (5b) in Hypothesis 2.1 are needed in Lemma 2.4.

In the present paper we construct Hadamard states in $(M_0, g|_{M_0})$. This yields however also Hadamard states on (M_0, g_0) since the two metrics are conformally related.

2.4. Null coordinates near C . For later use it is convenient to introduce null coordinates near C . The construction seems to be well-known, we sketch it for the reader's convenience. Note however the estimates in Lemma 2.5, which will be useful later on.

We first choose normal coordinates (y^0, \bar{y}) at p such that on a neighborhood U_1 of p , $C = \{(y^0)^2 - |\bar{y}|^2 = 0, y^0 > 0\}$.

Set

$$(2.4) \quad v := y^0 + |\bar{y}|, \quad w := y^0 - |\bar{y}|, \quad \psi := \frac{\bar{y}}{|\bar{y}|} \in \mathbb{S}^{d-1},$$

so that on a neighborhood of p one has $C = \{w = 0, v > 0\}$. Abusing notation slightly, we denote by $\psi^1, \dots, \psi^{d-1}$ coordinates on \mathbb{S}^{d-1} , and use the same letter for their pullback to local coordinates on M near p . We set

$$(2.5) \quad S := \{w = 0, v = \epsilon_0\}$$

where $\epsilon_0 > 0$ will be chosen small enough. Note that $S \subset C$ is diffeomorphic to \mathbb{S}^{d-1} .

Lemma 2.5. (1) *There exists a unique solution $s \in C^\infty(C)$ of:*

$$\begin{cases} \nabla^a f \nabla_a s = -1, \\ s|_S = 0. \end{cases}$$

(2) *There exists unique solutions $\theta^j \in C^\infty(C)$, $1 \leq j \leq d-1$ of:*

$$\begin{cases} \nabla^a f \nabla_a \theta^j = 0, \\ \theta^j|_S = \psi^j. \end{cases}$$

(3) *Moreover there exists $0 < \epsilon_0 < \epsilon_1$ and $k, \tilde{\theta}^j \in C^\infty(]-\epsilon_1, \epsilon_1[\times \mathbb{S}^{d-1})$ such that*

$$s(v, \psi) = \ln(v) + k(v, \psi), \quad \theta^j(v, \psi) = \tilde{\theta}^j(v, \psi), \quad \text{on }]-\epsilon_1, 0[\times \mathbb{S}^{d-1}.$$

³Note that we consider here only globally hyperbolic spacetimes, cf. [Mo2, App. A] for a more general definition.

Proof. The proof is given in Appendix A.4. \square

It remains to extend s, θ^j to smooth functions on a neighborhood of C .

We argue as in [Wa, Sect. 11.1]: for $s_0 \in \mathbb{R}$, the submanifold $S_{s_0} = \{s = s_0\} \subset C$ is spacelike, of codimension 2 in M . At a given point of S_{s_0} the orthogonal to its tangent space is two dimensional, timelike, and hence contain two null lines. One of them is generated by $\nabla^a f$, the other is transverse to C . We extend (s, θ) to a neighborhood of C by imposing that (s, θ) are constant along the above family of null geodesics, transverse to C .

Lemma 2.6. *The functions (f, s, θ) constructed above are a system of local coordinates near C with $C \subset \{f = 0\}$ and*

$$(2.6) \quad g|_C = -2dfds + h_{ij}(s, \theta)d\theta^i d\theta^j,$$

where $h_{ij}(s, \theta)d\theta^i d\theta^j$ is a smooth, s -dependent Riemannian metric on \mathbb{S}^{d-1} .

Proof. The proof will be given in Appendix A.3. \square

2.5. Estimates on traces. In this subsection we derive estimates, in the coordinates (s, θ) on C constructed above, for the restriction to C of a smooth, space compact function in M . These estimates will be applied later to traces on C of solutions of the Klein-Gordon equation in M_0 .

Clearly the only task is to control what happens near p , i.e. when $s \rightarrow -\infty$. We first derive estimates in the coordinates (v, ψ) introduced in (2.4), in a neighborhood of $v = 0$.

For $m \in \mathbb{N}$ we denote by S^m the space of functions g such that for some $\epsilon, R > 0$:

$$(2.7) \quad |\partial_v^\beta \partial_\psi^\alpha g(v, \psi)| \leq C_{\alpha, \beta} |v|^{(m-\alpha)_+}, \forall |\alpha| + |\beta| \geq 1, \text{ uniformly on } [-\epsilon, R] \times \mathbb{S}^{d-1}.$$

where $n_+ = \max(n, 0)$. Clearly $g(v, \psi) = v^m \in S^m$ and $v^m g \in S^{m+p}$ if $g \in S^p$.

Lemma 2.7. (1) *if $\phi \in C_{sc}^\infty(M)$ then $\hat{\phi}(v, \psi) := \phi|_C(v, \psi)$ belongs to S^1 .*

(2) *Let $|h| = \det[h_{ij}]$. Then $|h|(v, \psi) = v^{2(d-1)} r_0(v, \psi)$ for $r_0, r_0^{-1} \in S^0$.*

Proof. The function ϕ is smooth in the normal coordinates (y^0, \bar{y}) hence $\hat{\phi}(v, \psi) = \phi(\frac{1}{2}v, \frac{1}{2}v\psi)$. Considering the map $\chi : \mathbb{S}^{d-1} \ni \psi \mapsto \psi \in \mathbb{R}^d$ and denoting still by ψ some coordinates on \mathbb{S}^{d-1} we have:

$$\partial_v \tilde{\phi} = \partial_{y^0} \phi - \psi \cdot \partial_{\bar{y}} \phi, \quad \partial_\psi \tilde{\phi} = -v \partial_\psi \chi^i \partial_{\bar{y}^i} \phi.$$

From this we obtain (1). To prove (2) we need to express $h_{ij} = \langle \partial_{\theta^i} |g \partial_{\theta^j} \rangle$ on C . An easy computation using the estimates in Lemma 2.5 shows that on C we have:

$$\partial_{\theta^i} = \alpha_i^j(v, \psi) \partial_{\psi^j} + v r_0(v, \psi) \partial_v,$$

where $\alpha_i^j, r_0 \in S^0$ and $[\alpha^{ij}](v, \psi)$ invertible. Plugging this into (A.9), we obtain

$$[h_{ij}](v, \psi) = v^2 \left({}^t [a_i^j](v, \psi) [m_{ij}](\psi) [a_i^j](v, \psi) + v [b_{ij}](v, \psi) \right),$$

where $b_{ij} \in S^0$. This implies (2). \square

We will also need later the following lemma. We denote by $m_{ij}(\theta)d\theta^i d\theta^j$ the standard Riemannian metric on \mathbb{S}^{d-1} and set:

$$(2.8) \quad \beta(s, \theta) := |m|^{\frac{1}{4}}(\theta) |h|^{-\frac{1}{4}}(s, \theta),$$

Lemma 2.8. *Let*

$$\tilde{\phi}(s, \theta) := \beta^{-1}(s, \theta) \phi|_C(s, \theta), \quad \phi \in C_0^\infty(M),$$

Then for all $s_1 \in \mathbb{R}$ one has:

$$\tilde{\phi} \in O(e^{s(d-1)/2}), \quad \partial_s^\alpha \partial_\theta^\beta \tilde{\phi} \in O(e^{s(d+1)/2}), \quad s \in]-\infty, s_1], \quad \forall |\alpha| + |\beta| \geq 1.$$

Proof. We note that $\beta^{-1} = v^{(d-1)/2}r_0(v, \psi)$, for $r_0, r_0^{-1} \in S^0$. From this and Lemma 2.7 it follows that if $\phi \in C_0^\infty(M)$, then $\tilde{\phi}(v, \psi) \in v^{(d-1)/2}S^0$. It remains to estimate the derivatives of $\tilde{\phi}$ w.r.t. s and θ . By a standard computation we obtain for $g \in C_0^\infty(C)$:

$$\begin{aligned}\partial_{\theta^i} g &= a_i^j(v, \psi) \partial_{\psi^j} g + v r_i(v, \psi) \partial_v g, \\ \partial_s g &= v(1 + v r_0(v, \psi)) \partial_v g + v b^j(v, \psi) \partial_{\psi^j} g,\end{aligned}$$

for $r_0, r_i, b^j, a_i^j \in S^0$, and $[a_i^j]$ invertible. From this point on the lemma is a routine computation. \square

3. KLEIN-GORDON FIELDS INSIDE THE FUTURE LIGHTCONE

3.1. Klein-Gordon equation in M_0 . We fix a smooth real function $r \in C^\infty(M)$ and consider the *Klein-Gordon operator* on (M, g) :

$$P(x, D_x) = -\nabla^a \nabla_a + r(x), \text{ acting on } C^\infty(M).$$

We denote by $E_\pm \in \mathcal{D}'(M \times M)$ the retarded/advanced Green's functions for P , by $E = E_+ - E_- \in \mathcal{D}'(M \times M)$ the Pauli-Jordan commutator function, and by $\text{Sol}_{\text{sc}}(P)$ the space of smooth, complex valued, space-compact solutions of

$$P(x, D_x)\phi = 0 \text{ in } M.$$

Recall that we have set in Subsect. 2.1:

$$M_0 := I^+(p),$$

and by Lemma 2.2 we know that (M_0, g) is globally hyperbolic.

We denote by $P_0 = -\nabla^a \nabla_a + r(x)$ the restriction of P to M_0 , by $E_0 \in \mathcal{D}'(M_0 \times M_0)$ the Pauli-Jordan function for P_0 , and by $\text{Sol}_{\text{sc}}(P_0)$ the space of smooth, complex valued, space-compact solutions of

$$P_0(x, D_x)\phi_0 = 0 \text{ in } M_0.$$

By the global hyperbolicity of (M_0, g) we know that $\text{Sol}_{\text{sc}}(P_0) = E_0 \mathcal{D}(M_0)$. From (2.3) and the uniqueness of $E_{0\pm}$ we obtain that $E_{0\pm} = E_\pm \upharpoonright_{M_0 \times M_0}$, hence

$$E_0 = E \upharpoonright_{M_0 \times M_0} .$$

It follows that any $\phi_0 \in \text{Sol}_{\text{sc}}(P_0)$ uniquely extends to $\phi \in \text{Sol}_{\text{sc}}(P)$, in fact

$$(3.1) \quad \phi_0 = E_0 f_0, \quad f_0 \in \mathcal{D}(M_0) \Rightarrow \phi = E f_0 \upharpoonright_{M_0} .$$

As usual we equip $\text{Sol}_{\text{sc}}(P_0)$ with the symplectic form

$$(3.2) \quad \bar{\phi}_1 \sigma_0 \phi_2 := \int_{\Sigma_0} \overline{\nabla_a \phi_1} \phi_2 - \bar{\phi}_1 \nabla_a \phi_2 n^a d\sigma_h,$$

where $\Sigma_0 \subset M_0$ is a Cauchy hypersurface for (M_0, g) (see Subsect. A.1 for notation). It is well known that

$$E_0 : (C_0^\infty(M_0)/P_0 C_0^\infty(M_0), E_0) \rightarrow (\text{Sol}_{\text{sc}}(P_0), \sigma_0)$$

is a symplectomorphism.

3.2. Hadamard states in M_0 . We first briefly recall some standard facts, and refer for example to [GW, Sect. 2] for details and notation.

3.2.1. *Covariances of a quasi-free state.* If (\mathcal{Y}, σ) is a complex symplectic space, the *complex covariances* $\Lambda^\pm \in L_{\mathbb{H}}(\mathcal{Y}, \mathcal{Y}^*)$ of a (gauge invariant) quasi-free state ω on $\text{CCR}(\mathcal{Y}, \sigma)$ (the polynomial CCR *-algebra of (\mathcal{Y}, σ)) are defined by:

$$\omega(\psi(y_1)\psi^*(y_2)) =: (y_1|\Lambda^+y_2), \quad \omega(\psi^*(y_2)\psi(y_1)) =: (y_1|\Lambda^-y_2), \quad y_1, y_2 \in \mathcal{Y}.$$

From the CCR we obtain that $\Lambda^+ - \Lambda^- = i\sigma =: q$, and the necessary and sufficient condition for Λ^\pm to be the complex covariances of a (gauge invariant) quasi-free state is that $\Lambda^\pm \geq 0$.

If $(\mathcal{Y}, \sigma) = (C_0^\infty(M_0)/PC_0^\infty(M_0), E_0)$, the complex covariances of a state ω are induced from *two-point functions*, still denoted by Λ^\pm such that

$$\Lambda^\pm \in \mathcal{D}'(M_0 \times M_0), \quad P\Lambda^\pm = \Lambda^\pm P = 0,$$

where we identify operators on $C_0^\infty(M_0)$ with sesquilinear forms using the scalar product

$$(u|v) := \int_{M_0} \bar{u}v \, d\mu_g, \quad u, v \in C_0^\infty(M_0).$$

3.2.2. *Hadamard condition.* We now recall the *Hadamard condition* for quasi-free states. We denote by T^*M the cotangent bundle of M and $Z = \{(x, 0)\} \subset T^*M$ the zero section. The *principal symbol* of P is $p(x, \xi) = \xi_a g^{ab}(x) \xi_b$, the set

$$\mathcal{N} := \{(x, \xi) \in T^*M \setminus Z : p(x, \xi) = 0\}$$

is called the *characteristic manifold* of p .

The Hamilton vector field of p will be denoted by H_p , whose integral curves inside \mathcal{N} are called *bicharacteristics*.

We will use the notation $X = (x, \xi)$ for points in $T^*M \setminus Z$ and write $X_1 \sim X_2$ if $X_1 = (x_1, \xi_1)$ and $X_2 = (x_2, \xi_2)$ are in \mathcal{N} and X_1 and X_2 lie on the same bicharacteristic of p .

Let us fix a time orientation and denote by $V_{x\pm} \subset T_x M$ for $x \in M$, the open future/past light cones and $V_{x\pm}^*$ the dual cones

$$V_x^{*\pm} := \{\xi \in T_x^* M : \xi \cdot v > 0, \forall v \in V_{x\pm}, v \neq 0\}.$$

The set \mathcal{N} has two connected components invariant under the Hamiltonian flow of p , namely:

$$\mathcal{N}^\pm := \{X \in \mathcal{N} : \xi \in V_x^{*\pm}\}.$$

Definition 3.1. A quasi-free state ω on $\text{CCR}(C_0^\infty(M_0)/PC_0^\infty(M_0), E_0)$ with two-point functions Λ^\pm satisfies the microlocal spectrum condition if:

$$(\mu\text{sc}) \quad \text{WF}(\Lambda^\pm)' \subset \mathcal{N}^\pm \times \mathcal{N}^\pm.$$

Quasi-free states satisfying (μsc) are called Hadamard states.

We refer the reader to [Wr] and references therein for a discussion on equivalent formulations of the microlocal spectrum condition.

4. BULK-TO-BOUNDARY CORRESPONDENCE

4.1. **Boundary symplectic space.** We equip C with the coordinates (s, θ) constructed in Subsect. 2.4 and hence identify C with

$$(4.3) \quad \tilde{C} := \mathbb{R} \times \mathbb{S}^{d-1}.$$

We denote by $H^k(\tilde{C})$, $k \in \mathbb{N}$ the Sobolev space

$$H^k(\tilde{C}) := \left\{ g \in \mathcal{D}'(\mathbb{R} \times \mathbb{S}^{d-1}) : \int |\partial_s^\alpha \partial_\theta^\beta g|^2 |m|^{\frac{1}{2}} ds d\theta < \infty, \alpha + |\beta| \leq k \right\},$$

and extend the definition of $H^k(\tilde{C})$ to $k \in \mathbb{R}$ in the usual way. The space $H^0(\tilde{C})$ will be denoted simply by $L^2(\tilde{C})$. We set also:

$$\mathcal{H}(\tilde{C}) := \bigcap_{k \in \mathbb{R}} H^k(\tilde{C}), \quad \mathcal{H}'(\tilde{C}) := \bigcup_{k \in \mathbb{R}} H^k(\tilde{C}),$$

equipped with their canonical topologies.

We set

$$(4.4) \quad \bar{g}_1 \sigma_C g_2 := \int_{\mathbb{R} \times \mathbb{S}^{d-1}} (\partial_s \bar{g}_1 g_2 - \bar{g}_1 \partial_s g_2) |m|^{\frac{1}{2}}(\theta) ds d\theta, \quad g_1, g_2 \in \mathcal{H}(\tilde{C}).$$

Introducing the *charge* $q := i\sigma_C$ we have:

$$\bar{g}_1 q g_2 = 2(g_1 |D_s g_2|_{L^2(\tilde{C})}), \quad g_1, g_2 \in \mathcal{H}(\tilde{C}),$$

where $D_s = i^{-1} \partial_s$ is selfadjoint on $L^2(\tilde{C})$ on its natural domain. Clearly $(\mathcal{H}(\tilde{C}), \sigma_C)$ is a complex symplectic space.

4.2. Bulk-to-boundary correspondence.

Definition 4.1. Let $\beta \in C^\infty(\tilde{C})$ be defined in (2.8). We set

$$\begin{aligned} \rho : \text{Sol}_{\text{sc}}(P_0) &\rightarrow C^\infty(\mathbb{R} \times \mathbb{S}^{d-1}) \\ \phi &\mapsto \beta^{-1}(s, \theta) \phi|_C(s, \theta). \end{aligned}$$

Proposition 4.2. (1) ρ maps $\text{Sol}_{\text{sc}}(P_0)$ into $\mathcal{H}(\tilde{C})$;

(2) $\rho : (\text{Sol}_{\text{sc}}(P_0), \sigma) \rightarrow (\mathcal{H}(\tilde{C}), \sigma_C)$ is a monomorphism, i.e.:

$$\overline{\rho\phi_1} \sigma_C \rho\phi_2 = \overline{\phi_1} \sigma \phi_2, \quad \forall \phi_1, \phi_2 \in \text{Sol}_{\text{sc}}(P_0).$$

Proof. Let ϕ_0, ϕ as in (3.1). By Lemma 2.1 and the support properties of E , we see that $\text{supp}\phi \cap \tilde{C}$ is compact in M . Therefore the restriction of ϕ to C equals the restriction of a smooth compactly supported function to C . By Lemma 2.8 and the fact that $\rho\phi_0$ is supported in $] -\infty, s_1] \times \mathbb{S}^{d-1}$ for some s_1 , we obtain that $\rho\phi_0 \in \mathcal{H}(\tilde{C})$, which proves (1).

We now prove (2). Let $\phi_{i,0} \in \text{Sol}_{\text{sc}}(P_0)$, $i = 1, 2$ which are restrictions to M_0 of $\phi_i \in \text{Sol}_{\text{sc}}(P)$. We fix a Cauchy surface Σ_0 for (M_0, g) such that $\text{supp}\phi_{i,0} \cap \Sigma_0 \subset K \Subset M_0$. We can find a Cauchy surface Σ for (M, g) such that $\Sigma \cap K = \Sigma_0 \cap K$. Denoting by

$$J_a(\phi_1, \phi_2) := \overline{\phi_1} \nabla_a \phi_2 - \overline{\nabla_a \phi_1} \phi_2,$$

the conserved current, we have:

$$\overline{\phi_{1,0}} \sigma_0 \phi_{2,0} = \overline{\phi_1} \sigma \phi_2,$$

where

$$\overline{\phi_1} \sigma \phi_2 = - \int_{\Sigma} J_a(\phi_1, \phi_2) n^a d\sigma_h,$$

is the symplectic form on $\text{Sol}_{\text{sc}}(P)$. We now apply Stokes formula in the form (A.6) to the domain $U \subset M$ bounded by $\Sigma \cap K$, \tilde{C} and $\partial J^+(\Sigma \cap K)$, using that $\nabla_a J^a(\phi_1, \phi_2) = 0$. The boundary term on $\Sigma \cap K$ yields $-\overline{\phi_1} \sigma \phi_2$, the boundary term on $\partial J^+(\Sigma \cap K)$ vanishes. To express the boundary term on \tilde{C} , we use the coordinates (f, s, θ) constructed in Subsect. 2.4. We formally obtain the quantity:

$$\bar{g}_1 \hat{\sigma} g_2 = \int_{\mathbb{R} \times \mathbb{S}^{d-1}} (\partial_s \bar{g}_1 g_2 - \bar{g}_1 \partial_s g_2) |h|^{\frac{1}{2}}(s, \theta) ds d\theta$$

for $g_i = (\phi_i)|_C$. This equals $\overline{\rho\phi_1} \sigma_C \rho\phi_2$ by an easy computation.

To justify the use of Stokes formula, we need to take care of the fact that \tilde{C} is not smooth at p . This can be done as follows: for $0 < \epsilon \ll 1$ we denote by U_ϵ some ϵ -neighborhood of p . We replace \tilde{C} by a smooth hypersurface C_ϵ , obtained

by smoothly gluing $C \setminus U_\epsilon$ to a piece of a Cauchy surface Σ'_ϵ passing through U_ϵ . The contribution of the integral on Σ_ϵ is written using (A.4), and converges to 0 when $\epsilon \rightarrow 0$, using that ϕ_i are smooth functions. The contribution of the integral on $C \setminus U_\epsilon$ converges to $\overline{\rho\phi_1}\sigma_C\rho\phi_2$, using that $\rho\phi_i \in \mathcal{H}(\tilde{C})$. This completes the proof of the proposition. \square

4.3. Pullback of states from the boundary. Since

$$\rho : (\text{Sol}_{\text{sc}}(P_0), \sigma_0) \rightarrow (\mathcal{H}(\tilde{C}), \sigma_C)$$

is a monomorphism, we can pullback a quasi-free state ω_C on $\text{CCR}(\mathcal{H}(\tilde{C}), \sigma_C)$ to a quasi-free state ω_0 on $\text{CCR}(C_0^\infty(M_0)/P_0C_0^\infty(M_0), E_0)$ by setting:

$$(4.5) \quad \omega_0(\psi(u_1)\psi^*(u_2)) := \omega_C(\psi(\rho \circ E_0 u_1)\psi^*(\rho \circ E_0 u_2)), \quad u_1, u_2 \in C_0^\infty(M_0).$$

If $\lambda^\pm \in L_h(\mathcal{H}(\tilde{C}), \mathcal{H}(\tilde{C})^*)$ are the complex covariances of ω_C , then the complex covariances of ω_0 are (formally) given by:

$$(4.6) \quad \Lambda^\pm := (\rho \circ E_0)^* \circ \lambda^\pm \circ (\rho \circ E_0).$$

5. HADAMARD CONDITION ON THE CONE

In this section we formulate the natural boundary version of the bulk Hadamard condition (μsc).

5.1. Preparations. We recall that $p(x, \xi)$ denotes the principal symbol of the Klein-Gordon operator P (or P_0).

Let $C \subset M$ be the backward lightcone introduced in Subsect. 2.1. We denote by $N^*C \subset T^*M \setminus Z$ the *conormal bundle* to C , i.e.

$$N^*C := \{(x, \xi) \in T^*M \setminus Z : x \in C, \xi = 0 \text{ on } T_x C\}.$$

The fact that C is characteristic is equivalent to

$$(5.1) \quad N^*C \subset \mathcal{N},$$

where \mathcal{N} is the characteristic manifold of p . Since N^*C is Lagrangian, it is well known that (5.1) implies that N^*C is invariant under the flow of H_p . The projections on M of bicharacteristics starting from N^*C are (modulo reparametrization) *characteristic curves*, i.e. integral curves of the vector field $v^a = \nabla^a f$, if $f \in C^\infty(M)$ is some defining function of C , i.e. $f = 0$, $df \neq 0$ on C .

We will use the coordinates (f, s, θ) introduced in Subsect. 2.4, which, for ease of notation, will be denoted by $x = (r, s, y) \in \mathbb{R} \times \mathbb{R} \times \mathbb{S}^{d-1}$. The dual coordinates are denoted $\xi = (\varrho, \sigma, \eta)$, elements of T^*M will sometimes be denoted by $X = (x, \xi)$ and elements of T^*C will be denoted by $Y = ((s, y), (\sigma, \eta))$.

In the above coordinates, we have

$$C = \{r = 0\}, \quad N^*C = \{r = 0, \sigma = \eta = 0\},$$

and from (2.6) we obtain that:

$$(5.2) \quad p(x, \xi)|_C = -2\varrho\sigma + h(s, y, \eta),$$

where we set $h(s, y, \eta) = h^{ij}(0, s, y)\eta_i\eta_j$. Note that $h(s, y, \eta)$ is *elliptic*, i.e. $h(s, y, \eta) \geq c_0|\eta|^2$, for $c_0 > 0$, locally in (s, y) , since $h_{ij}dy^i dy^j$ is Riemannian.

For later use let us extend the notation $X_1 \sim X_2$ introduced in 3.2.2. For $Y = (s, y, \sigma, \eta) \in T^*C$, $X = (x, \xi) \in T^*M$, we will write $Y \sim X$ if

$$(5.3) \quad \sigma \neq 0, \quad ((0, s, y), ((2\sigma)^{-1}h(s, y, \eta), \sigma, \eta)) \sim X.$$

Recall also that the positive/negative energy components \mathcal{N}^\pm of \mathcal{N} were defined in 3.2.2.

Lemma 5.1. *Let $Y_1 = (s_1, y_1, \sigma_1, \eta_1) \in T^*C$, $X_2 = (x_2, \xi_2) \in T^*M$ with $x_2 \notin C$. Then:*

(1) *there exists $\varrho_1 \in \mathbb{R}$ such that*

$$X_1 := ((0, s_1, y_1), (\varrho_1, \sigma_1, \eta_1)) \sim (x_2, \xi_2) =: X_2$$

iff $\sigma_1 \neq 0$ and then $\varrho_1 = (2\sigma_1)^{-1}h(s_1, y_1, \eta_1)$ and $Y_1 \sim X_2$.

(2) *if $Y_1 \sim X_2$ then $X_2 \in \mathcal{N}^\pm$ iff $\pm\sigma_1 > 0$.*

Proof. Let $X_1 = ((0, s_1, y_1), (\varrho_1, \sigma_1, \eta_1)) \in \mathcal{N}$. By (5.2) we have

$$-2\varrho_1\sigma_1 + h(s_1, y_1, \eta_1) = 0.$$

If $\sigma_1 = 0$ then $h(s_1, y_1, \eta_1) = 0$ hence $\eta_1 = 0$ by ellipticity of h . Therefore $\sigma_1 = 0$ implies $X_1 \in N^*C$. Since $X_2 \sim X_1$ and N^*C is invariant under the flow of H_p , we have also $X_2 \in N^*C$ which contradicts the hypothesis that $x_2 \notin C$. Therefore necessarily $\sigma_1 \neq 0$ and hence $\varrho_1 = (2\sigma_1)^{-1}h(s_1, y_1, \eta_1)$ and $Y_1 \sim X_2$. This proves (1).

To prove (2) we have to show that

$$(5.4) \quad \pm\sigma_1 > 0 \Leftrightarrow ((0, s_1, y_1), ((2\sigma_1)^{-1}h(s_1, y_1, \eta_1), \sigma_1, \eta_1)) \in \mathcal{N}^\pm.$$

Let us fix $(y_1, \eta_1) \in T^*\mathbb{S}^{d-1}$, $\sigma_1 \in \mathbb{R}$. Since \mathcal{N}^\pm are the two connected components of \mathcal{N} , it suffices by connexity to prove (5.4) for s_1 in a neighborhood of $-\infty$, i.e. in a neighborhood of p in M . Recall that we introduced Gaussian normal coordinates (y^0, \bar{y}) near p with ∂_{y^0} future oriented. Let α be the one form $(2\sigma_1)^{-1}h(s_1, y_1, \eta_1)dr + \sigma_1 ds + \eta_1 dy$. Then

$$((0, s_1, y_1), ((2\sigma_1)^{-1}h(s_1, y_1, \eta_1), \sigma_1, \eta_1)) \in \mathcal{N}^\pm \Leftrightarrow \mp \langle \alpha | g^{-1} dy^0 \rangle > 0.$$

Since it suffices to check the sign of $\langle \alpha | g^{-1} dy^0 \rangle$ near p , we can, by a simple approximation argument (see e.g. (A.9)) replace g by the flat metric at p . We have then (see Lemma 2.5 and recall that $s = u, r = f$):

$$y^0 = v + w, \quad v = e^s, \quad w = e^{-s}r,$$

hence

$$\mp \langle \alpha | g^{-1} dy^0 \rangle = \pm 2(e^{-s_1}\sigma_1 + e^{s_1}(2\sigma_1)^{-1}h(s_1, y_1, \eta_1)),$$

has the same sign as $\pm\sigma_1$, which proves (5.4). \square

Recall that $E \in \mathcal{D}'(M \times M)$ is the Pauli-Jordan commutator function for P and $\rho : \mathcal{D}(M) \ni u \mapsto u|_C \in C^\infty(\tilde{C})$ is (modulo a smooth, non-zero multiplicative factor) the operator of restriction to C , defined in Def. 4.1.

Proposition 5.2. *Let $\chi \in C_0^\infty(M)$ with $\text{supp } \chi \subset M \setminus C$ and $\psi \in C_0^\infty(\tilde{C})$. Then:*

(1) $\text{WF}(\psi\rho \circ E\chi)' \subset \{(Y_1, X_2) : y_1 \in \text{supp } \psi, x_2 \in \text{supp } \chi, Y_1 \sim X_2\}$,

where the notation $Y \sim X$ is defined in (5.3).

(2) $\psi\rho \circ E\chi : \mathcal{D}(M) \rightarrow \mathcal{D}(\tilde{C})$ extends continuously as $\psi\rho \circ E\chi : \mathcal{D}'(M) \rightarrow \mathcal{D}'(\tilde{C})$.

Proof. It is well-known that:

$$(5.5) \quad \begin{aligned} \text{supp } E &\subset \{(x_1, x_2) : x_1 \in J(x_2)\}, \\ \text{WF}(E)' &= \{(X_1, X_2) \in \mathcal{N} \times \mathcal{N} : X_1 \sim X_2\}. \end{aligned}$$

On the other hand the distributional kernel of ρ equals

$$\delta(r_2) \otimes \delta(s_1, y_1, s_2, y_2) \beta^{-1}(s_1, y_1) \in \mathcal{D}'(\tilde{C} \times M).$$

It follows that:

$$(5.6) \quad \begin{aligned} \text{WF}(\rho)' &= \{(Y_1, X_2) : r_2 = 0, (s_1, y_1) = (s_2, y_2), \\ &\quad (\sigma_1, \eta_1) = (\sigma_2, \eta_2), (\sigma_2, \eta_2) \neq (0, 0)\}. \end{aligned}$$

Since $E : \mathcal{D}(M) \rightarrow \mathcal{E}(M)$ we see that $\psi\rho \circ E\chi : \mathcal{D}(M) \rightarrow \mathcal{D}(\tilde{C})$. Moreover there exists $\chi_1 \in C_0^\infty(M)$ such that $\psi\rho \circ E\chi = \psi\rho \circ \chi_1 E\chi$. The notations ${}_{M_1}\Gamma, \Gamma_{M_2}$ for $\Gamma \subset T^*M_1 \times T^*M_2$ are defined in [GW, Subsect. 3.2]. We have then

$${}_{\tilde{C}}\text{WF}(\rho)' = \text{WF}(E)'_M = \emptyset,$$

and it follows from [Hö1, Chap. 8] and (5.5), (5.6) that:

$$\begin{aligned} \text{WF}(\psi\rho \circ E\chi)' &\subset \text{WF}(\psi\rho)' \circ \text{WF}(E\chi)' \\ &\subset \{(Y_1, X_2) : \exists \varrho_1 \text{ s.t. } ((0, s_1, y_1), (\varrho_1, \sigma_1, \eta_1)) \sim X_2, x_2 \in \text{supp}\chi\}. \end{aligned}$$

Using that $\text{supp}\chi \cap C = \emptyset$ and Lemma 5.1 (1), this implies (1). Moreover (1) implies that

$$(5.7) \quad \text{WF}(\psi\rho \circ E\chi)'_M = \emptyset.$$

Again by [Hö1] this implies that $\psi\rho \circ E\chi : \mathcal{D}(M) \rightarrow \mathcal{D}(\tilde{C})$ extends continuously as $\psi\rho \circ E\chi : \mathcal{D}'(M) \rightarrow \mathcal{D}'(\tilde{C})$. \square

5.2. Hadamard condition on the cone. Recall from Subsect. 4.2 that to a quasi-free state ω_C on $\text{CCR}(\mathcal{H}(\tilde{C}), \sigma_C)$ we can associate a quasi-free state ω_0 on $\text{CCR}(C_0^\infty(M_0)/PC_0^\infty(M_0), E_0)$. In this subsection we give natural conditions on the covariances λ^\pm of ω_C which ensure that the induced state ω_0 satisfies the microlocal spectrum condition (μsc).

Recall that we denote by $Y = ((s, y), (\sigma, \eta))$ the points in $T^*\tilde{C}$. We also denote by Δ the diagonal in $T^*\tilde{C} \times T^*\tilde{C}$.

Theorem 5.3. *Let $\lambda^\pm : \mathcal{H}(\tilde{C}) \rightarrow \mathcal{H}(\tilde{C})$ and*

$$\Lambda^\pm := (\rho \circ E_0)^* \circ \lambda^\pm \circ (\rho \circ E_0).$$

Then:

- (1) $\Lambda^\pm \in \mathcal{D}'(M_0 \times M_0)$.
- (2) *If*

- i)* $\text{WF}(\lambda^\pm)' \cap \{(Y_1, Y_2) : \pm\sigma_1 < 0 \text{ or } \pm\sigma_2 < 0\} = \emptyset$,
- ii)* $\text{WF}(\lambda^+ - \lambda^-)' \cap \{(Y_1, Y_2) : \sigma_1 \text{ and } \sigma_2 \neq 0\} \subset \Delta$,

then:

- iii)* $\text{WF}(\lambda^\pm)' \cap \{(Y_1, Y_2) : \pm\sigma_1 > 0 \text{ and } \pm\sigma_2 > 0\} \subset \Delta$.

- (3) *Assume moreover that $\lambda^\pm : \mathcal{H}(\tilde{C}) \rightarrow \mathcal{H}(\tilde{C})$, ${}_{\tilde{C}}\text{WF}(\lambda^\pm)' = \text{WF}(\lambda^\pm)'_{\tilde{C}} = \emptyset$.*

Then if i) and iii) in (2) hold, Λ^\pm satisfy (μsc).

Proof. To prove (1) it suffices to check that $\rho \circ E_0 : \mathcal{D}(M_0) \rightarrow \mathcal{H}(\tilde{C})$. If $\chi \in C_0^\infty(M_0)$, then by Lemma 2.1 $\rho \circ E_0\chi = \rho \circ \chi_1 E\chi$ for some $\chi_1 \in C_0^\infty(M)$. Since $E : \mathcal{D}(M) \rightarrow \mathcal{E}(M)$ and $\rho : \mathcal{D}(M) \rightarrow \mathcal{H}(\tilde{C})$ are continuous, this proves (1).

To prove (2) we write:

$$\begin{aligned} &\text{WF}(\lambda^\pm)' \cap \{\pm\sigma_1 > 0, \pm\sigma_2 > 0\} \\ &\subset (\text{WF}(\lambda^\mp)' \cap \{\pm\sigma_1 > 0, \pm\sigma_2 > 0\}) \cup (\text{WF}(\lambda^+ - \lambda^-)' \cap \{\pm\sigma_1 > 0, \pm\sigma_2 > 0\}) \\ &\subset (\text{WF}(\lambda^\mp)' \cap \{\pm\sigma_1 > 0, \pm\sigma_2 > 0\}) \cup (\text{WF}(\lambda^+ - \lambda^-)' \cap \{\sigma_1, \sigma_2 \neq 0\}). \end{aligned}$$

The first set in the last line is empty by *i)*, and the second is contained in Δ by *ii)*.

To prove (3) we follow an argument due to Moretti [Mo2]. We treat only the case of λ^+ , the case of λ^- being similar, and omit the + superscript. Let $\chi_i \in C_0^\infty(M_0)$,

$i = 1, 2$. By Prop. 2.3 there exists $\psi_i \in C_0^\infty(C)$ (and hence $\psi_i \equiv 0$ near p) such that any null geodesic starting from $\text{supp}\chi_i$ intersects C in $\{\psi_i = 1\}$. We have:

$$\begin{aligned} \chi_1 \Lambda \chi_2 &= \chi_1 (\rho \circ E)^* \psi_1 \circ \lambda \circ \psi_2 (\rho \circ E) \chi_2 \\ &\quad + \chi_1 (\rho \circ E)^* \psi_1 \circ \lambda \circ (1 - \psi_2) (\rho \circ E) \chi_2 \\ &\quad + \chi_1 (\rho \circ E)^* (1 - \psi_1) \circ \lambda \circ \psi_2 (\rho \circ E) \chi_2 \\ &\quad + \chi_1 (\rho \circ E)^* (1 - \psi_1) \circ \lambda \circ (1 - \psi_2) (\rho \circ E) \chi_2 \\ &=: \Lambda_1 + \Lambda_2 + \Lambda_3 + \Lambda_4. \end{aligned}$$

By the properties of χ_i, ψ_i , we can find $\tilde{\chi}_i \in C_0^\infty(M)$ supported near p such that:

- (a) $(1 - \psi_i) (\rho \circ E) \chi_i = (1 - \psi_i) \rho \circ \tilde{\chi}_i E \chi_i$,
- (b) no null geodesic from $\text{supp}\chi_i$ intersects $\text{supp}\tilde{\chi}_i$.

It follows from (b) and (5.5) that $\tilde{\chi}_i E \chi_i$ has a smooth compactly supported kernel, hence $\tilde{\chi}_i E \chi_i : \mathcal{D}'(M) \rightarrow \mathcal{D}(M)$. Since $(1 - \psi_i) \rho : \mathcal{D}(M) \rightarrow \mathcal{H}(\tilde{C})$ we see that

$$(5.8) \quad (1 - \psi_i) \rho \circ E \chi_i : \mathcal{D}'(M) \rightarrow \mathcal{H}(\tilde{C}),$$

hence

$$(5.9) \quad \chi_i (\rho \circ E)^* (1 - \psi_i) : \mathcal{H}'(\tilde{C}) \rightarrow \mathcal{D}(M).$$

It remains to examine the properties of $\psi_i (\rho \circ E) \chi_i$. By Prop. 5.2 we know that $\psi_i (\rho \circ E) \chi_i : \mathcal{D}'(M) \rightarrow \mathcal{E}'(\tilde{C})$. Since $\mathcal{E}'(\tilde{C}) \subset \mathcal{H}'(\tilde{C})$ continuously, we have

$$(5.10) \quad \psi_i (\rho \circ E) \chi_i : \mathcal{D}'(M) \rightarrow \mathcal{H}'(\tilde{C}),$$

hence:

$$(5.11) \quad \chi_i (\rho \circ E)^* \psi_i : \mathcal{H}(\tilde{C}) \rightarrow \mathcal{D}(M).$$

From (5.8), ..., (5.11) and the assumption that $\lambda : \mathcal{H}(\tilde{C}) \rightarrow \mathcal{H}(\tilde{C})$ it follows that $\Lambda_i : \mathcal{D}'(M_0) \rightarrow \mathcal{D}(M_0)$ hence has a smooth kernel for $i = 2, 3, 4$, and $\text{WF}(\chi_1 \Lambda \chi_2)' = \text{WF}(\Lambda_1)'$.

To bound $\text{WF}(\Lambda_1)'$ we choose $\tilde{\psi}_i \in C_0^\infty(\tilde{C})$ such that $\tilde{\psi}_i \psi_i = \psi_i$ and write

$$\Lambda_1 = (\chi_1 (\rho \circ E) \psi_1) \circ (\tilde{\psi}_1 \lambda \tilde{\psi}_2) \circ (\psi_2 (\rho \circ E) \chi_2) =: K_1^* \circ d \circ K_2,$$

where $K_i = \psi_i (\rho \circ E) \chi_i \in \mathcal{E}'(M \times \tilde{C})$, $d = \tilde{\psi}_1 \lambda \tilde{\psi}_2 \in \mathcal{E}'(\tilde{C} \times \tilde{C})$. The distributions K_1, K_2 and d have compact support. Moreover we have

$$\text{WF}(d)'_{\tilde{C}} =_{\tilde{C}} \text{WF}(d)' = \text{WF}(K_1)'_M =_M \text{WF}(K_2)' = \emptyset.$$

In fact the first two equalities follow from the corresponding hypothesis on $\text{WF}(c)'$, the last two from (5.7). We can then apply the results in [Hö1, Chap. 8] on the composition of kernels, and obtain that $K_2^* \circ d \circ K_1$ is well defined and

$$\text{WF}(K_2^* \circ d \circ K_1) \subset \text{WF}(K_2^*)' \circ \text{WF}(d)' \circ \text{WF}(K_1)'$$

Now we apply Prop. 5.2 (1), the fact that $\text{WF}(d)' \subset \text{WF}(\lambda)'$ and Lemma 5.1 (1). We obtain that if $(X_1, X_2) \in \text{WF}(\Lambda)'$, necessarily $X_1, X_2 \in \mathcal{N}_+$ and $X_1 \sim X_2$, which is exactly condition (μsc). \square

6. PSEUDODIFFERENTIAL CALCULUS

In this section we collect rather standard results on the pseudodifferential calculus on $\tilde{C} = \mathbb{R} \times \mathbb{S}^{d-1}$. We will however need to consider *bi-homogeneous* symbols on $\mathbb{R} \times \mathbb{S}^{d-1}$, i.e. symbols having different homogeneities in the covariables σ and η , dual to s and θ .

The reason for this is that the charge $q = -2D_s$ is not an elliptic differential operator in the usual sense (considered on \tilde{C}), hence operators like $(q - z)^{-1}$ for $z \in \mathbb{C} \setminus \mathbb{R}$ are not in the usual pseudodifferential classes.

For $k, k' \in \mathbb{R}$ we denote by $H^k(\mathbb{R})$, $H^{k'}(\mathbb{S}^{d-1})$ the Sobolev space on \mathbb{R} , \mathbb{S}^{d-1} or order k , k' , and by $\|\cdot\|_k$, $\|\cdot\|_{k'}$ their respective norms. Furthermore, we denote by $H^{k,k'}(\mathbb{R} \times \mathbb{S}^{d-1})$ the Sobolev space on $\mathbb{R} \times \mathbb{S}^{d-1}$ of bi-order (k, k') , i.e. the completion of $C_0^\infty(\mathbb{R} \times \mathbb{S}^{d-1})$ for the norm

$$\|\psi\|_{k,k'} := \|\langle D_s \rangle^k \langle D_\theta \rangle^{k'} \psi\|_2.$$

We set also for $p \in \mathbb{R}$:

$$B^p(\mathbb{R}) = \bigcap_{k \in \mathbb{R}} B(H^k(\mathbb{R}), H^{k-p}(\mathbb{R})),$$

equipped with its natural topology.

 6.1. Pseudodifferential operators on $\mathbb{R} \times \mathbb{R}^{d-1}$.

Definition 6.1. Let $p_1, p_2 \in \mathbb{R}$.

- (1) we denote by $S^{p_1, p_2}(\mathbb{R} \times \mathbb{R}^{d-1})$ the space of symbols $a \in C^\infty(T^*\mathbb{R} \times T^*\mathbb{R}^{d-1})$ such that

$$|\partial_s^{\alpha_1} \partial_\sigma^{\beta_1} \partial_y^{\alpha_2} \partial_\eta^{\beta_2} a| \in O(\langle \sigma \rangle^{p_1 - |\beta_1|} \langle \eta \rangle^{p_2 - |\beta_2|}), \quad \alpha_1, \beta_1 \in \mathbb{N}, \quad \alpha_2, \beta_2 \in \mathbb{N}^{d-1}.$$

- (2) we denote by $B^{p_1} S^{p_2}(\mathbb{R} \times \mathbb{S}^{d-1})$ the space of $a \in C^\infty(T^*\mathbb{R}^{d-1}, B^{p_1}(\mathbb{R}))$ such that:

$$\|\partial_y^{\alpha_2} \partial_\eta^{\beta_2} a\|_{p_1, k_1} \in O(\langle \eta \rangle^{p_2 - |\beta_2|}), \quad \alpha_2, \beta_2 \in \mathbb{N}^{d-1},$$

where $\|\cdot\|_{p_1, k_1}$ is any seminorm of a in $B^{p_1}(\mathbb{R})$.

Using the Weyl quantization on $\mathbb{R} \times \mathbb{R}^{d-1}$, we obtain a map

$$S^{p_1, p_2}(\mathbb{R} \times \mathbb{R}^{d-1}) \ni a \mapsto \text{Op}(a) \in B(C_0^\infty(\mathbb{R} \times \mathbb{R}^{d-1}), C^\infty(\mathbb{R} \times \mathbb{R}^{d-1})),$$

whose range, denoted by $\Psi^{p_1, p_2}(\mathbb{R} \times \mathbb{R}^{d-1})$ is the space of pseudodifferential operators on $\mathbb{R} \times \mathbb{R}^{d-1}$ of bi-order (p_1, p_2) . Similarly using the Weyl quantization on \mathbb{R}^{d-1} we obtain a map

$$B^{p_1} S^{p_2}(\mathbb{R} \times \mathbb{R}^{d-1}) \ni a \mapsto \text{Op}(a) \in B(C_0^\infty(\mathbb{R} \times \mathbb{R}^{d-1}), C^\infty(\mathbb{R} \times \mathbb{R}^{d-1})),$$

whose range will be denoted by $B^{p_1} \Psi^{p_2}(\mathbb{R} \times \mathbb{R}^{d-1})$.

6.2. Pseudodifferential operators on \tilde{C} . Let $A : C_0^\infty(\tilde{C}) \rightarrow C^\infty(\tilde{C})$. If $\chi_i \in C^\infty(\mathbb{S}^{d-1})$, $i = 1, 2$ are cutoff functions supported in chart open sets $\Omega_i \subset \mathbb{S}^{d-1}$ and $\phi_i : \Omega_i \rightarrow \mathbb{R}^{d-1}$ are coordinate charts, then $\phi_1^* \circ \chi_1 A \chi_2 \circ \phi_2^{-1*} : C_0^\infty(\mathbb{R} \times \mathbb{R}^{d-1}) \rightarrow C^\infty(\mathbb{R} \times \mathbb{R}^{d-1})$.

- Definition 6.2.** (1) We denote by $\Psi^{p_1, p_2}(\tilde{C})$ the space of operators $A : C_0^\infty(\tilde{C}) \rightarrow C^\infty(\tilde{C})$ such that for any χ_i, ϕ_i as above $\phi_1^* \circ \chi_1 A \chi_2 \circ \phi_2^{-1*} \in \Psi^{p_1, p_2}(\mathbb{R} \times \mathbb{R}^{d-1})$.
 (2) We denote by $B^{p_1} \Psi^{p_2}(\tilde{C})$ the space of operators $A : C_0^\infty(\tilde{C}) \rightarrow C^\infty(\tilde{C})$ such that for any χ_i, ϕ_i as above $\phi_1^* \circ \chi_1 A \chi_2 \circ \phi_2^{-1*} \in B^{p_1} \Psi^{p_2}(\mathbb{R} \times \mathbb{R}^{d-1})$.
 (3) We set

$$\Psi^{-\infty, p_2}(\tilde{C}) = \bigcap_{p_1 \in \mathbb{R}} \Psi^{p_1, p_2}(\tilde{C}), \quad B^{-\infty} \Psi^{p_2}(\tilde{C}) = \bigcap_{p_1 \in \mathbb{R}} B^{p_1} \Psi^{p_2}(\tilde{C}).$$

(4) We set

$$\tilde{\Psi}^{p_1, p_2}(\tilde{C}) = \Psi^{p_1, p_2}(\tilde{C}) + B^{-\infty}\Psi^{p_2}(\tilde{C}).$$

Note that if one defines analogously $\tilde{\Psi}^{-\infty, p_2}(\tilde{C}) := \bigcap_{p_1 \in \mathbb{R}} \tilde{\Psi}^{p_1, p_2}(\tilde{C})$, then actually $\tilde{\Psi}^{-\infty, p_2}(\tilde{C}) = B^{-\infty}\Psi^{p_2}(\tilde{C})$. Moreover it is easy to check that

$$\tilde{\Psi}^{p_1, p_2}(\tilde{C}) \circ \tilde{\Psi}^{q_1, q_2}(\tilde{C}) \subset \tilde{\Psi}^{p_1+p_2, q_1+q_2}(\tilde{C}).$$

We refer the reader to [Ro, BS, RT] and references therein for more details on the pseudo-differential calculus on products of manifolds⁴.

6.3. Beals criterion. Let us denote by $\Psi^p(\mathbb{S}^{d-1})$ the classes of standard pseudo-differential operators on \mathbb{S}^{d-1} . It is well-known that $\Psi^p(\mathbb{S}^{d-1})$ can be characterized by the Beals criterion, namely an operator $A : C^\infty(\mathbb{S}^{d-1}) \rightarrow C^\infty(\mathbb{S}^{d-1})$ belongs to $\Psi^p(\mathbb{S}^{d-1})$ iff

$$(6.1) \quad \text{ad}_{f_1} \cdots \text{ad}_{f_n} \text{ad}_{X_1} \cdots \text{ad}_{X_m} A : H^k(\mathbb{S}^{d-1}) \rightarrow H^{k-p+n}(\mathbb{S}^{d-1}), \quad n, m \in \mathbb{N}, \quad k \in \mathbb{Z},$$

for any $f_i \in C^\infty(\mathbb{S}^{d-1})$ and X_j smooth vector fields on \mathbb{S}^{d-1} [RT]. Moreover one can find a finite set of such f_i and X_j such that the topology on $\Psi^p(\mathbb{S}^{d-1})$ given by the collection of the norms of the multi-commutators is equivalent to the standard topology on $\Psi^p(\mathbb{S}^{d-1})$, given by the symbol space topologies of the pullbacks $\phi_i^* \circ \chi_i A \chi_j \circ \phi_j$ in Def. 6.2, for a fixed covering of \mathbb{S}^{d-1} by chart neighborhoods U_i .

These characterizations immediately carry over to the classes $B^{p_1} \Psi^{p_2}(\tilde{C})$. In fact it is easy to see that $A \in B^{p_1} \Psi^{p_2}(\tilde{C})$ iff

$$(6.2)$$

$$\text{ad}_{f_1} \cdots \text{ad}_{f_n} \text{ad}_{X_1} \cdots \text{ad}_{X_m} A : H^{k, k'}(\tilde{C}) \rightarrow H^{k-p_1, k'-p_2+n}(\mathbb{S}^{d-1}), \quad n, m \in \mathbb{N}, \quad k, k' \in \mathbb{Z}.$$

This result can be deduced from the previous one by considering the operators

$$(|u_1| \otimes \mathbb{1}_{\mathbb{S}^{d-1}}) \circ A \circ (|u_2| \otimes \mathbb{1}_{\mathbb{S}^{d-1}}) : C^\infty(\mathbb{S}^{d-1}) \rightarrow C^\infty(\mathbb{S}^{d-1})$$

for $u_1 \in H^{-k+p_1}(\mathbb{R})$, $u_2 \in H^k(\mathbb{R})$, which belongs to $\Psi^{p_2}(\mathbb{S}^{d-1})$ if (6.2) holds. Applying the result recalled above about the equivalence of the standard topology and the topology given by the multicommutator norms, one obtains that $A \in B^{p_1} \Psi^{p_2}(\tilde{C})$ if (6.2) holds.

In the usual case one can deduce from the Beals criterion standard results on the functional calculus for pseudo-differential operators, for example on complex powers of elliptic Ψ DOs [Bo]. These results are easy to extend to the classes $B^{p_1} \Psi^{p_2}(\tilde{C})$. We will need only a very simple one, which we now state. Recall that $\tilde{\Psi}^{-\infty, 0}(\tilde{C}) = B^{-\infty} \Psi^0(\tilde{C}) \subset B(L^2(\tilde{C}))$. The spectrum of $b \in B(L^2(\tilde{C}))$ is denoted $\text{spec}(b)$.

Proposition 6.3. *Let $b \in \tilde{\Psi}^{-\infty, 0}(\tilde{C})$ and F holomorphic near $\text{spec}(b)$ with $F(0) = 0$. Then $F(b) \in \tilde{\Psi}^{-\infty, 0}(\tilde{C})$.*

Proof. The proof consists of expressing $F(b)$ as a contour integral and applying the Beals criterion to the resolvent $(b - z)^{-1}$. \square

6.4. Essential support. We denote by $\Psi_{\text{ph}}^p(\mathbb{R})$, $p \in \mathbb{R}$ the class of global pseudo-differential operators on \mathbb{R} with poly-homogenous symbols.

Definition 6.4. *The essential support of $a \in \Psi^{p_1, p_2}(\tilde{C})$, denoted by $\text{ess supp}(a) \subset T^*\mathbb{R} \setminus Z$ is defined by:*

$(s_0, \sigma_0) \notin \text{ess supp}(a)$ if there exists $b \in \Psi_{\text{ph}}^0(\mathbb{R})$, elliptic at (s_0, σ_0) such that $b \circ a \in \Psi^{-\infty, p_2}(\tilde{C})$.

Clearly $\text{ess supp}(a)$ is a closed conic subset of $T^*\mathbb{R} \setminus Z$. Moreover one can equivalently require that $a \circ b \in \Psi^{-\infty, p_2}(\tilde{C})$ for some $b \in \Psi_{\text{ph}}^0(\mathbb{R})$, elliptic at (s_0, σ_0) .

⁴Note however that the literature discusses mostly the case when both manifolds are compact.

6.5. Wavefront set of kernels. For $N = \mathbb{R}, \mathbb{S}^{d-1}, \mathbb{R} \times \mathbb{S}^{d-1}$, we denote by Δ_N the diagonal in $T^*N \times T^*N$, and by Z_N the zero section in T^*N .

For an operator $a \in \Psi^{p_1, p_2}(\mathbb{R} \times \mathbb{S}^{d-1})$ it is in general not true that $\text{WF}(a)'$ is contained in the full diagonal $\Delta_{\mathbb{R} \times \mathbb{S}^{d-1}}$ (as would be the case for an operator in $\Psi^p(\mathbb{R} \times \mathbb{S}^{d-1})$). Instead one has the following estimate, which can be thought as a natural generalization of the usual estimate for the wave front set of tensor products of distributions (in this case Schwartz kernels) [BS].

Lemma 6.5. *Let $a \in \Psi^{p_1, p_2}(\mathbb{R} \times \mathbb{S}^{d-1})$. Then:*

$$\text{WF}(a)' \subset \Delta_{\mathbb{R}} \times \Delta_{\mathbb{S}^{d-1}} \cup \Delta_{\mathbb{R}} \times (Z_{\mathbb{S}^{d-1}} \times Z_{\mathbb{S}^{d-1}}) \cup (Z_{\mathbb{R}} \times Z_{\mathbb{R}}) \times \Delta_{\mathbb{S}^{d-1}}.$$

Less precise estimates are valid for the $\tilde{\Psi}^{p_1, p_2}(\mathbb{R} \times \mathbb{S}^{d-1})$ classes:

Lemma 6.6. (1) *Let $a \in B^{-\infty} \Psi^{p_2}(\tilde{C})$. Then*

$$\text{WF}(a)' \cap \{(Y_1, Y_2) : \sigma_1 \neq 0 \text{ or } \sigma_2 \neq 0\} = \emptyset.$$

(2) *Let $a \in \tilde{\Psi}^{p_1, p_2}(\tilde{C})$. Then*

$$\tilde{C} \text{WF}(a)' = \text{WF}(a)'_{\tilde{C}} = \emptyset.$$

The proof is given in Subsect. A.5.

6.6. Toeplitz pseudo-differential operators on \tilde{C} . We recall that $\mathcal{H}(\tilde{C}) = \bigcap_{m \in \mathbb{R}} H^m(\tilde{C}) = \bigcap_{k \in \mathbb{R}} H^{k, k}(\tilde{C})$. Let us set

$$L_{\pm}^2(\tilde{C}) := \mathbb{1}_{\mathbb{R}^{\pm}}(D_s) L^2(\tilde{C})$$

and denote by $i_{\pm} : L_{\pm}^2(\tilde{C}) \rightarrow L^2(\tilde{C})$ the corresponding isometric injection, so that $\pi_{\pm} := i_{\pm} i_{\pm}^* = \mathbb{1}_{\mathbb{R}^{\pm}}(D_s)$ is the orthogonal projection on $L_{\pm}^2(\tilde{C})$ in $L^2(\tilde{C})$. We set also

$$(6.3) \quad \mathcal{H}_{\pm}(\tilde{C}) := i_{\pm}^* \mathcal{H}(\tilde{C}) \subset \mathcal{H}(\tilde{C}).$$

We will see in Sect. 7 that this provides a useful setup for the discussion of the positivity condition $\lambda^{\pm} \geq 0$ for the two-point functions of a Hadamard state.

Writing $\mathbb{1}_{\mathbb{R}^{\pm}} = \chi \mathbb{1}_{\mathbb{R}^{\pm}} + (1 - \chi) \mathbb{1}_{\mathbb{R}^{\pm}}$ for a cutoff function $\chi \in C_0^{\infty}(\mathbb{R})$ equal to 1 near 0, we see that

$$(6.4) \quad \pi_{\pm} \in \tilde{\Psi}^{0,0}(\tilde{C}).$$

For $\alpha, \beta \in \{+, -\}$ and $p_1, p_2 \in \mathbb{R}$ we set:

$$\tilde{\Psi}_{\alpha\beta}^{p_1, p_2}(\tilde{C}) := i_{\alpha} \circ \tilde{\Psi}^{p_1, p_2}(\tilde{C}) \circ i_{\beta}^*.$$

By (6.3) we see that $\tilde{\Psi}_{\alpha\beta}^{p_1, p_2}(\tilde{C}) : \mathcal{H}_{\beta}(\tilde{C}) \rightarrow \mathcal{H}_{\alpha}(\tilde{C})$. Moreover if we set:

$$R_{\alpha\beta} : \tilde{\Psi}^{p_1, p_2}(\tilde{C}) \ni a \mapsto i_{\alpha}^* \circ a \circ i_{\beta} \in \tilde{\Psi}_{\alpha\beta}^{p_1, p_2}(\tilde{C}),$$

then using (6.4), we see that $R_{\alpha\beta}$ has right inverse

$$T_{\alpha\beta} : \tilde{\Psi}_{\alpha\beta}^{p_1, p_2}(\tilde{C}) \ni a \mapsto i_{\alpha} \circ a \circ i_{\beta}^* \in \tilde{\Psi}^{p_1, p_2}(\tilde{C}),$$

which allows to identify $\tilde{\Psi}_{\alpha\beta}^{p_1, p_2}(\tilde{C})$ with $\text{Ran} T_{\alpha\beta} \subset \tilde{\Psi}^{p_1, p_2}(\tilde{C})$. From (6.4) we also have:

$$(6.5) \quad \tilde{\Psi}_{\alpha\beta}^{p_1, p_2}(\tilde{C}) \circ \tilde{\Psi}_{\beta\gamma}^{q_1, q_2}(\tilde{C}) \subset \tilde{\Psi}_{\alpha\gamma}^{p_1+q_1, p_2+q_2}(\tilde{C}).$$

7. CONSTRUCTION OF HADAMARD STATES ON THE CONE

From the discussion in Subsect. 5.2, in particular Thm. 5.3, we are led to the following definition.

Definition 7.1. *A pair of maps $\lambda^\pm : \mathcal{H}(\tilde{C}) \rightarrow \mathcal{H}(\tilde{C})$ is called a pair of Hadamard two-point functions on the cone C if:*

$$\begin{aligned}
 & i) \quad \tilde{C}\text{-WF}(\lambda^\pm)' = \text{WF}(\lambda^\pm)'_{\tilde{C}} = \emptyset, \\
 & ii) \quad \text{WF}(\lambda^\pm)' \cap \{(Y_1, Y_2) : \pm\sigma_1 < 0 \text{ or } \pm\sigma_2 < 0\} = \emptyset, \\
 & iii) \quad \lambda^+ - \lambda^- = 2D_s, \\
 & iv) \quad \lambda^\pm \geq 0 \text{ on } \mathcal{H}(\tilde{C}).
 \end{aligned}
 \tag{Had}$$

As the name suggests, if λ^\pm are Hadamard two-point functions on C in the sense of the above definition, then Λ^\pm defined in (4.6) are Hadamard two-point functions on M_0 (as follows from Thm. 5.3).

We now discuss in more detail the various conditions in (Had). It is natural to consider pseudodifferential two-point functions, i.e. to assume that $\lambda^\pm \in \tilde{\Psi}^{p_1, p_2}(\tilde{C})$. Moreover to analyze conditions (Had) *iii*), *iv*) it is convenient to reduce oneself to λ^\pm of the form:

$$(7.6) \quad \lambda^\pm = (2|D_s|)^{\frac{1}{2}} c^\pm (2|D_s|)^{\frac{1}{2}}, \text{ where } c^\pm \in \tilde{\Psi}^{p_1, p_2}(\tilde{C}),$$

for $p_1, p_2 \in \mathbb{R}$. Note that writing $(2|D_s|)^{\frac{1}{2}}$ as $\chi(D_s)(2|D_s|)^{\frac{1}{2}} + (1 - \chi(D_s))(2|D_s|)^{\frac{1}{2}}$ for $\chi \in C_0^\infty(\mathbb{R})$ equal to 1 near 0, we see that (7.6) implies that $\lambda^\pm \in \tilde{\Psi}^{p_1+1, p_2}(\tilde{C})$.

7.1. Wavefront set. We first analyze conditions (Had) *i*), *ii*).

Proposition 7.2. *Assume that*

$$\begin{aligned}
 & (7.7) \quad \lambda^\pm = a^\pm + r^\pm, \quad a^\pm \in \Psi^{p_1, p_2}(\tilde{C}), \quad r^\pm \in \tilde{\Psi}^{-\infty, p_2}(\tilde{C}), \\
 & \quad (\mathbb{R} \times \mathbb{R}^\mp) \cap \text{ess sup}(a^\pm) = \emptyset.
 \end{aligned}$$

*Then λ^\pm satisfies conditions (Had) *i*), *ii*).*

Proof. The fact that λ^\pm satisfy *i*) follows from Lemma 6.6 (2). Also since by Lemma 6.6 (1) r^\pm satisfy *ii*) we can assume that $\lambda^\pm = a^\pm$. We treat only the case of λ^+ and use the notation in the proof of Lemma 6.6. Let $\tilde{Y}_1, \tilde{Y}_2 \in T^*\tilde{C} \setminus Z$ with $\tilde{\sigma}_1 \neq 0$ or $\tilde{\sigma}_2 \neq 0$. Let us assume that $\tilde{\sigma}_1 \neq 0$, the case $\tilde{\sigma}_2 \neq 0$ being similar, using the remark after Def. 6.4.

Since $(\mathbb{R} \times \mathbb{R}^+) \cap \text{ess sup}(a^+) = \emptyset$, we can find a cutoff function χ_1 with $\chi_1(\tilde{s}_1) \neq 0$, a neighborhood V_1 of $\tilde{\sigma}_1$ and some $m_1 \in \Psi_{\text{ph}}^0(\mathbb{R})$ elliptic at $(\tilde{s}_1, \tilde{\sigma}_1)$ such that $(1 - m_1)(s, D_s)v_{\sigma, \lambda} \in O(\langle \lambda \rangle^{-\infty})$ in all $H^k(\mathbb{R})$ and $m_1(s, D_s) \circ a \in \tilde{\Psi}^{-\infty, p_2}(\tilde{C})$. The fact that $(\tilde{Y}_1, \tilde{Y}_2) \notin \text{WF}(a)'$ follows then from the same arguments as in the proof of Lemma 6.6. \square

In terms of c^\pm appearing in (7.6), a natural condition implying (7.7) is

$$(\mu_{SC}) \quad \mathbb{1}_{\mathbb{R}^\mp}(D_s)c^\pm \in \tilde{\Psi}^{-\infty, p_2}(\tilde{C}),$$

which clearly implies that λ^\pm satisfy (7.7).

Lemma 7.3. *Let λ^\pm be given by (7.6) such that (μ_{SC}) holds. Then*

$$c^\pm = \mathbb{1}_{\mathbb{R}^\pm}(D_s) + \tilde{\Psi}^{-\infty, p_2}(\tilde{C}).$$

Proof. In terms of c^\pm (Had) *iii*) becomes $c^+ - c^- = \text{sgn}(D_s)$. Let $\chi^\pm \in C^\infty(\mathbb{R})$ be cutoff functions equal to 1 near $\pm\infty$ and to 0 near $\mp\infty$. From condition $(\mu\text{sc})_C$ and pseudodifferential calculus we obtain that

$$(7.8) \quad c^\pm = \chi^\pm(D_s)c^\pm\chi^\mp(D_s) + \tilde{\Psi}^{-\infty,p_2}(\tilde{C}).$$

Using successively (7.8) and $c^+ - c^- = \text{sgn}(D_s)$ we obtain

$$\begin{aligned} c^\pm &= \chi^\pm c^\pm \chi^\pm + \tilde{\Psi}^{-\infty,p_2}(\tilde{C}) \\ &= \chi^\pm(c^\mp \pm \text{sgn}(D_s))\chi^\pm + \tilde{\Psi}^{-\infty,p_2}(\tilde{C}) \\ &= \chi^\pm c^\mp \chi^\pm + \chi^\pm \chi^\pm + \tilde{\Psi}^{-\infty,p_2}(\tilde{C}) \\ &= \chi^\pm \chi^\mp c^\mp \chi^\mp \chi^\pm + \chi^\pm + \tilde{\Psi}^{-\infty,p_2}(\tilde{C}) \\ &= \chi^\pm + \tilde{\Psi}^{-\infty,p_2}(\tilde{C}) \\ &= \mathbb{1}_{\mathbb{R}^\pm}(D_s) + \tilde{\Psi}^{-\infty,p_2}(\tilde{C}). \quad \square \end{aligned}$$

7.2. Positivity. We now discuss conditions (Had) *iii*), *iv*). In terms of c^\pm they become:

$$(7.9) \quad \begin{aligned} \textit{iii)} \quad & c^+ - c^- = \text{sgn}(D_s), \\ \textit{iv)} \quad & c^\pm \geq 0 \text{ on } \mathcal{H}(\tilde{C}). \end{aligned}$$

To analyze (7.9) we use the framework of Subsect. 6.6. We denote c^+ simply by c and set

$$c_{\alpha\beta} = i_\alpha^* \circ c \circ i_\beta, \quad \alpha, \beta \in \{+, -\},$$

so that:

$$(7.10) \quad c = \sum_{\alpha, \beta \in \{+, -\}} i_\alpha c_{\alpha\beta} i_\beta^*.$$

Then (7.9) is equivalent to:

$$(7.11) \quad \begin{pmatrix} c_{++} & c_{+-} \\ c_{-+} & c_{--} \end{pmatrix} \geq 0, \quad \begin{pmatrix} c_{++} - \mathbb{1} & c_{+-} \\ c_{-+} & c_{--} + \mathbb{1} \end{pmatrix} \geq 0 \text{ on } \mathcal{H}_+(\tilde{C}) \oplus \mathcal{H}_-(\tilde{C}),$$

which is equivalent to:

$$\begin{aligned} \textit{i)} \quad & c_{++} \geq 0, \quad c_{--} \geq \mathbb{1}, \quad c_{-+} = c_{+-}^*, \\ \textit{ii)} \quad & |(u_+|c_{+-}u_-)| \leq (u_+|c_{++}u_+)^{\frac{1}{2}}(u_-|c_{--}u_-)^{\frac{1}{2}}, \\ & |(u_+|c_{+-}u_-)| \leq (u_+|(c_{++} - \mathbb{1})u_+)^{\frac{1}{2}}(u_-|(c_{--} + \mathbb{1})u_-)^{\frac{1}{2}}, \quad u_\pm \in \mathcal{H}_\pm(\tilde{C}). \end{aligned}$$

Condition *ii*) above is implied by

$$|(u_+|c_{+-}u_-)| \leq (u_+|(c_{++} - \mathbb{1})u_+)^{\frac{1}{2}}(u_-|c_{--}u_-)^{\frac{1}{2}}, \quad u_\pm \in \mathcal{H}_\pm(\tilde{C}).$$

We are now in position to prove the following theorem, which is the analog of [GW, Thm. 7.5] in the present situation. It provides a rather large class of Hadamard two-point functions on C , hence by Thm. 5.3, of Hadamard states on M_0 .

Theorem 7.4. *Assume that*

$$\begin{aligned} c_{++} &= \mathbb{1} + a_+^* a_+, \quad c_{--} = a_-^* a_-, \\ c_{+-} &= c_{-+}^* = a_+^* d a_-, \end{aligned}$$

for $a_+ \in \tilde{\Psi}_{++}^{-\infty,0}(\tilde{C})$, $a_- \in \tilde{\Psi}_{--}^{-\infty,0}(\tilde{C})$, $d \in \tilde{\Psi}_{+-}^{0,0}(\tilde{C})$ with $\|d\|_{B(L^2_-(\tilde{C}), L^2_+(\tilde{C}))} \leq \mathbb{1}$.

Let c be given by (7.10) and $\lambda^+ = (2|D_s|)^{\frac{1}{2}}c(2|D_s|)^{\frac{1}{2}}$, $\lambda^- = \lambda^+ - 2D_s$. Then λ^\pm is a pair of Hadamard two-point functions on the cone.

Proof. We set as before $\lambda^\pm = (2|D_s|)^{\frac{1}{2}}c^\pm(2|D_s|)^{\frac{1}{2}} \in \tilde{\Psi}^{1,0}(\tilde{C})$, so that $c^+ = c$, $c^- = c - \text{sgn}(D_s)$. Condition (7.9) follows from the above discussion. It remains to check condition (μ_{sc_C}) . We embed the spaces $\tilde{\Psi}_{\alpha\beta}^{p_1,p_2}(\tilde{C})$ into $\tilde{\Psi}^{p_1,p_2}(\tilde{C})$ as explained at the end of Subsect. 6.6, and we have:

$$c^+ = a_+^*a_+ + a_+^*da_- + a_-^*d^*a_+ + a_-^*a_- + \mathbb{1}_{\mathbb{R}^+}(D_s),$$

$$c^- = a_+^*a_+ + a_+^*da_- + a_-^*d^*a_+ + a_-^*a_- + \mathbb{1}_{\mathbb{R}^-}(D_s),$$

hence

$$\mathbb{1}_{\mathbb{R}^-}(D_s)c^+ = a_+^*a_+ + a_+^*da_- \in \tilde{\Psi}^{-\infty,0}(\tilde{C}),$$

$$\mathbb{1}_{\mathbb{R}^+}(D_s)c^- = a_-^*d^*a_+ + a_-^*a_- \in \tilde{\Psi}^{-\infty,0}(\tilde{C}),$$

and condition (μ_{sc_C}) is satisfied. \square

Remark 7.5. *The special choice of vanishing a_+ , a_- and d in Thm. 7.4 gives two-point functions*

$$\lambda^\pm = \pm 2\mathbb{1}_{\mathbb{R}^\pm}(D_s)D_s.$$

In the setting of asymptotically flat spacetimes with past time infinity i^- these correspond to the Hadamard state found and further studied in [Mo1, Mo2].

8. PURE HADAMARD STATES

In this section we first characterize *pure* Hadamard states on the cone C . We then prove that any pure Hadamard state ω_C on C induces a pure Hadamard state ω_0 in M_0 .

8.1. An abstract criterion for purity. Let (\mathcal{Y}, σ) a complex symplectic space and ω a gauge invariant quasi-free state on $\text{CCR}(\mathcal{Y}, \sigma)$, with complex covariances λ^\pm .

Let \mathcal{Y}^{cpl} the completion of \mathcal{Y} for the norm

$$(8.1) \quad \|y\|_\omega := (\bar{y} \cdot \lambda^+ y + \bar{y} \cdot \lambda^- y)^{\frac{1}{2}}.$$

Let us introduce the hermitian form $q = i\sigma \in L_h(\mathcal{Y}, \mathcal{Y}^*)$. Clearly q, λ^\pm extend uniquely to \mathcal{Y}^{cpl} . Then ω is pure iff [AS]:

- (1) q is non-degenerate on \mathcal{Y}^{cpl} ,
- (2) there exists an involution $\kappa : \mathcal{Y}^{\text{cpl}} \rightarrow \mathcal{Y}^{\text{cpl}}$ such that $\kappa^*q\kappa = q$, $q\kappa \geq 0$ and $\lambda^\pm = \frac{1}{2}q(\kappa \pm \mathbb{1})$.

From this discussion we obtain immediately the following lemma.

Lemma 8.1. *Let $(\mathcal{Y}_i, \sigma_i)$, $i = 1, 2$ be two complex symplectic spaces and $\rho : \mathcal{Y}_1 \rightarrow \mathcal{Y}_2$ an injective map such that $\rho^*\sigma_2\rho = \sigma_1$. Let ω_2 be a pure, gauge-invariant quasi-free state on $\text{CCR}(\mathcal{Y}_2, \sigma_2)$. Let ω_1 the gauge invariant quasi-free state on $\text{CCR}(\mathcal{Y}_1, \sigma_1)$ defined by the complex covariances*

$$\lambda_1^\pm = \rho^*\lambda_2^\pm\rho.$$

Then if $\rho\mathcal{Y}_1$ is dense in \mathcal{Y}_2 for the norm $\|\cdot\|_{\omega_2}$ defined in (8.1), the state ω_1 is pure on $\text{CCR}(\mathcal{Y}_1, \sigma_1)$.

8.2. Pure Hadamard states on the cone. The following theorem is the exact analog of [GW, Thm. 7.10]. In what follows we will use the notations introduced in Subsect. 6.6.

Theorem 8.2. *Let λ^\pm be the two-point functions of a state ω_C on $(\mathcal{H}(\tilde{C}), \sigma_C)$ of the form (7.6) and satisfying (μ_{sc_C}) . Then ω_C is pure iff there exists $a \in \tilde{\Psi}_{+-}^{-\infty,0}(\tilde{C})$ such that*

$$c^+ =$$

Proof. We consider the pair c^\pm obtained from λ^\pm , denote as before c^+ by c and identify c with the matrix $\begin{pmatrix} c_{++} & c_{+-} \\ c_{-+} & c_{--} \end{pmatrix}$. Arguing as in the proof of [GW, Thm. 7.10], we obtain that the state ω_C on $(\mathcal{H}(\tilde{C}), \sigma_C)$ with covariances λ^\pm is pure iff

$$(8.2) \quad c = \begin{pmatrix} \mathbb{1} + a^*a & a^*(\mathbb{1} + aa^*)^{\frac{1}{2}} \\ (\mathbb{1} + aa^*)^{\frac{1}{2}}a & aa^* \end{pmatrix},$$

for some $a : L_+^2(\tilde{C}) \rightarrow L_-^2(\tilde{C})$. This proves \Leftarrow .

Let us now prove \Rightarrow . Since we assumed that $c^\pm \in \tilde{\Psi}^{0,0}(\tilde{C})$ satisfy (μ_{sc_C}) , we obtain that

$$(8.3) \quad a^*a \in \tilde{\Psi}_{++}^{-\infty,0}(\tilde{C}), \quad (\mathbb{1} + aa^*)^{\frac{1}{2}}a \in \tilde{\Psi}_{+-}^{-\infty,0}(\tilde{C}).$$

We claim that

$$(8.4) \quad (\mathbb{1} + aa^*)^{-\frac{1}{2}} \in \mathbb{1} + \tilde{\Psi}_{++}^{-\infty,0}(\tilde{C}).$$

Let us prove (8.4). We use the operators $R_{\alpha\beta}, T_{\alpha\beta}$ defined at the end of Subsect. 6.6. We first embed aa^* into $\tilde{\Psi}^{-\infty,0}(\tilde{C})$, i.e. consider $b = T_{--}(a^*a)$. Then $b \geq 0$ on $L^2(\tilde{C})$ and applying Prop. 6.3 to $F(z) = (1+z)^{\frac{1}{2}} - 1$ we obtain that $(\mathbb{1} + b)^{-\frac{1}{2}} - \mathbb{1} \in \tilde{\Psi}^{-\infty,0}(\tilde{C})$. Writing b as a 2×2 matrix acting on $L_+^2(\tilde{C}) \oplus L_-^2(\tilde{C})$ we see that $R_{++} \left((\mathbb{1} + b)^{\frac{1}{2}} \right) = (\mathbb{1} + aa^*)^{\frac{1}{2}}$, which proves (8.4). From (8.4) and (8.3) we obtain that $a \in \tilde{\Psi}_{+-}^{-\infty,0}(\tilde{C})$. \square

In the next lemma we identify the completion of $\mathcal{H}(\tilde{C})$ for the norm (8.1) associated to any Hadamard state considered in Thm. 8.2.

Let us first fix some notation. For $a : L_+^2(\tilde{C}) \rightarrow L_-^2(\tilde{C})$ we denote by $c^+(a)$ the operator defined in (8.2) set $c^-(a) = c^+(a) - \text{sgn}(D_s)$, and

$$(8.5) \quad \lambda^\pm(a) = (2|D_s|)^{\frac{1}{2}} c^\pm(a) (2|D_s|)^{\frac{1}{2}}.$$

If \mathcal{H} is a Hilbert space and $h \geq 0$ is a selfadjoint operator on \mathcal{H} with $\text{Ker}h = \{0\}$, we denote by $h\mathcal{H}$ the completion of $\text{Dom}h^{-1}$ (i.e. the range of h) for the norm $\|h^{-1}u\|_{\mathcal{H}}$.

Lemma 8.3. *Let $a : L_+^2(\tilde{C}) \rightarrow L_-^2(\tilde{C})$. Then the completion of $\mathcal{H}(\tilde{C})$ for the norm $(\cdot | (\lambda^+(a) + \lambda^-(a)) \cdot)^{\frac{1}{2}}$ equals $|D_s|^{\frac{1}{2}} L^2(\tilde{C})$.*

Proof. By (8.5) and the definition of $|D_s|^{\frac{1}{2}} L^2(\tilde{C})$ it suffices to prove that the completion of $\mathcal{H}(\tilde{C})$ for the norm $(u | (c^+(a) + c^-(a)) u)^{\frac{1}{2}}$ equals $L^2(\tilde{C})$. Let

$$u(a) = \begin{pmatrix} (\mathbb{1} + aa^*)^{\frac{1}{2}} & a \\ a^* & (\mathbb{1} + a^*a)^{\frac{1}{2}} \end{pmatrix},$$

and note that

$$(8.6) \quad u(a)^* c^\pm(0) u(a) = c^\pm(a).$$

Moreover using the identity $af(a^*a) = f(aa^*)a$, valid for any Borel function f , we obtain that $u(a)^{-1} = u(-a)$, hence $u(a) : L^2(\tilde{C}) \rightarrow L^2(\tilde{C})$ is boundedly invertible. By (8.6) it suffices to treat the case $a = 0$ which is obvious since $c^+(0) + c^-(0) = \mathbb{1}$. \square

8.3. Pure Hadamard states in M_0 . Our main result concerns the purity of the states induced in the bulk. We postpone the introduction of the key technical ingredients of the proof to Subsect. 8.4 for the sake of self-consistency of our results on the characteristic Cauchy problem.

Theorem 8.4. *Assume that $\dim M \geq 4$. Let ω_C be a pure Hadamard state on $\text{CCR}(\mathcal{H}(\tilde{C}), \sigma_C)$ as in Thm. 8.2. Then the state ω induced by ω_C on $\text{CCR}(C_0^\infty(M_0)/PC_0^\infty(M_0), E_0)$ is a pure state.*

Proof. The proof relies on Lemma 8.1 and on some results on the characteristic Cauchy problem in M_0 , proved below in Subsect. 8.4. Recall that the map $\rho : \text{Sol}_{\text{sc}}(P_0) \rightarrow \mathcal{H}(\tilde{C})$ was introduced in Def. 4.1. By Lemmas 8.1 and 8.3 it suffices to check that $\rho(\text{Sol}_{\text{sc}}(P_0))$ is dense in $|D_s|^{\frac{1}{2}}L^2(\tilde{C})$. Since $C_0^\infty(\mathbb{R} \times \mathbb{S}^{d-1})$ is dense in $|D_s|^{\frac{1}{2}}L^2(\tilde{C})$ it suffices for $w \in C_0^\infty(\mathbb{R} \times \mathbb{S}^{d-1})$ to find a sequence $\phi_n \in \text{Sol}_{\text{sc}}(P_0)$ such that $\rho\phi_n \rightarrow w$ in $|D_s|^{\frac{1}{2}}L^2(\tilde{C})$.

We will use freely the notation introduced below in Subsect. 8.4. We first fix a Cauchy surface Σ in (M, g) as in 8.4.2 to the future of $\text{supp} w$. Note that since w vanishes near $s = -\infty$, we know that w belongs to the space $\tilde{H}_0^1(\tilde{C}_0)$ introduced in Prop. 8.8. By Thm. 8.7 and Prop. 8.8, there exists f in the energy space $\mathcal{E}_0(\Sigma_0)$ such that $w = R \circ T f$. Since $C_0^\infty(\Sigma_0) \oplus C_0^\infty(\Sigma_0)$ is dense in $\mathcal{E}_0(\Sigma_0)$, there exists a sequence $f_n \in C_0^\infty(\Sigma_0) \oplus C_0^\infty(\Sigma_0)$ such that $f_n \rightarrow f$ in $\mathcal{E}_0(\Sigma_0)$. By Thm. 8.7 and Prop. 8.8 we have $R \circ T f_n \rightarrow w$ in $\tilde{H}_0^1(\tilde{C}_0)$, hence also $R \circ T f_n \rightarrow w$ in $|D_s|^{\frac{1}{2}}L^2(\tilde{C})$, by Remark 8.9.

Let $\phi_n \in \text{Sol}_{\text{sc}}(P_0)$ the solution with Cauchy data f_n on Σ_0 . Then $\rho\phi_n = R \circ T f_n \rightarrow w$ in $|D_s|^{\frac{1}{2}}L^2(\tilde{C})$, which completes the proof of the theorem. \square

8.4. A characteristic Cauchy problem in M_0 . From Lemma 8.1, we see that to deduce purity of the bulk state from the purity of the boundary state, the range of ρ in $\mathcal{H}(\tilde{C})$ should be sufficiently large. One way to ensure this is to solve a *characteristic Cauchy problem* in M_0 , i.e. to construct an inverse for ρ . If M has a *compact* Cauchy surface, the characteristic problem was shown to be well posed in *energy spaces* by Hörmander [Hö2]. With some care the results of [Hö2] can be used in our situation.

8.4.1. Characteristic Cauchy problem for compact Cauchy surfaces. We recall an important result of Hörmander [Hö2] on the characteristic Cauchy problem in energy spaces. The framework of [Hö2] is as follows:

One considers a spacetime (\tilde{M}, \tilde{g}) for $\tilde{M} = \mathbb{R} \times \tilde{\Sigma}$, $\tilde{\Sigma}$ a smooth *compact* manifold and $\tilde{g} = -\tilde{\beta}(t, x)dt^2 + \tilde{h}_{ij}(t, x)dx^i dx^j$. One also fixes a real function $\tilde{r} \in C^\infty(\tilde{M})$.

If $\tilde{\Sigma}_1$ is a Cauchy hypersurface in (\tilde{M}, \tilde{g}) , we will denote by

$$\tilde{U}_{\tilde{\Sigma}_1} : C^\infty(\tilde{\Sigma}_1) \oplus C^\infty(\tilde{\Sigma}_1) \rightarrow C^\infty(\tilde{M})$$

the Cauchy evolution operator for $-\square_{\tilde{g}} + \tilde{r}$, so that $\phi = \tilde{U}_{\tilde{\Sigma}_1} f$ solves

$$\begin{cases} -\square_{\tilde{g}}\phi + \tilde{r}\phi = 0, \\ \phi|_{\tilde{\Sigma}_1} = f^0, \quad n^\mu \nabla_\mu \phi|_{\tilde{\Sigma}_1} = f^1. \end{cases}$$

A hypersurface \tilde{C} of the form

$$(8.7) \quad \tilde{C} = \{(F(x), x) : x \in \tilde{\Sigma}\}, \quad F \text{ Lipschitz,}$$

is called *space-like* (resp. *weakly space-like*) if

$$\sup_{x \in \tilde{\Sigma}} (-\beta^{-1}(F(x), x) + \partial_i F(x) h^{ij}(F(x), x) \partial_j F(x)) < 0, \quad (\text{resp. } \leq 0).$$

If F is smooth then of course \tilde{C} is space-like (resp. weakly space-like) iff all tangent vectors at each point of \tilde{C} are space-like (resp. space-like or null).

Since $\tilde{\Sigma}$ is compact and F Lipschitz, the Sobolev space $H^1(\tilde{C})$ and of course $L^2(\tilde{C})$ are well defined, for example by identifying \tilde{C} with $\tilde{\Sigma}$ and using the Riemannian metric $\tilde{h}_{ij}(0, \mathbf{x})dx^i dx^j$ on $\tilde{\Sigma}$ to equip \tilde{C} with a density $d\nu_{\tilde{C}}$.

One also needs the measure

$$d\nu_{\tilde{C}}^0 = (\beta^{-1} - h^{ij}\partial_i \tilde{F}\partial_j \tilde{F})d\nu_{\tilde{C}},$$

which vanishes if \tilde{C} is a null hypersurface.

We set now

$$(8.8) \quad \mathcal{E}(\tilde{C}) := H^1(\tilde{C}) \oplus L^2(\tilde{C}, d\nu_{\tilde{C}}^0).$$

Note that if \tilde{C} is space-like (i.e. a Cauchy hypersurface), then $\mathcal{E}(\tilde{C}) = H^1(\tilde{C}) \oplus L^2(\tilde{C})$.

The result of [Hö2] is the following theorem:

Theorem 8.5 ([Hö2]). *Let $\tilde{\Sigma}_1$ be any Cauchy hypersurface in \tilde{M} and \tilde{C} be weakly space-like of the form (8.7). Then the map*

$$\begin{aligned} \tilde{T} : \mathcal{E}(\tilde{\Sigma}_1) &\rightarrow \mathcal{E}(\tilde{C}) \\ f &\mapsto ((\tilde{U}_{\tilde{\Sigma}_1} f)|_{\tilde{C}}, (\beta^{-1}\partial_t \tilde{U}_{\tilde{\Sigma}_1} f)|_{\tilde{C}}) \end{aligned}$$

is a homeomorphism.

Note that if \tilde{C} is characteristic, then $L^2(\tilde{C}, d\nu_{\tilde{C}}^0) = \{0\}$ and $\mathcal{E}(\tilde{C}) = H^1(\tilde{C})$, so one obtains as a particular case the solvability of the characteristic Cauchy problem in energy spaces.

8.4.2. *Embedding M_0 into \tilde{M} .* We will use Hörmander's result recalled above to solve a characteristic Cauchy problem in M_0 , in an arbitrary neighborhood of p . The first task is to locally embed M into a spacetime \tilde{M} as above.

We fix a Cauchy hypersurface Σ to the future of p and identify M with $\mathbb{R} \times \Sigma$ with $g = -\beta(t, \mathbf{x})dt^2 + h_{ij}(t, \mathbf{x})dx^i dx^j$. We set $\Sigma_0 = \Sigma \cap M_0$ and fix an open, precompact set U such that $J^-(\Sigma_0) \cap J^+(p) \subset U$.

The following lemma shows that over U , C can be parametrized by Σ .

Lemma 8.6. *There exists a bounded, Lipschitz function F defined on Σ such that*

$$\overline{C} \cap U = \{(t, \mathbf{x}) : t = F(\mathbf{x})\} \cap U.$$

Proof. The proof is given in Appendix A.6. \square

We next embed Σ_0 into a smooth compact manifold $\tilde{\Sigma}$. We consider the spacetime $\tilde{M} = \mathbb{R} \times \tilde{\Sigma}$ and extend F to a Lipschitz function \tilde{F} on $\tilde{\Sigma}$, g to a metric \tilde{g} as in 8.4.1. We set

$$\tilde{C} = \{t = \tilde{F}(\mathbf{x})\} \subset \tilde{M},$$

and define:

$$(8.9) \quad C_0 := (J^-(\Sigma_0; M) \cap C) \cup \{p\}.$$

C_0 is an open subset of \overline{C} , with \overline{C}_0 compact in M and

$$(8.10) \quad \partial\Sigma_0 = \partial C_0.$$

We claim that we can choose the embedding $\Sigma_0 \subset \tilde{\Sigma}$ and the extensions \tilde{F} and \tilde{g} so that:

$$(8.11) \quad J^-(\tilde{\Sigma} \setminus \overline{\Sigma}_0; \tilde{M}) \cap \overline{C}_0 = \emptyset,$$

$$(8.12) \quad \tilde{C} \text{ is weakly space-like in } \tilde{M}.$$

This is clearly possible by modifying Σ , F and g only outside a large open set U , and using that the embedding of (M_0, g) into (M, g) is causally compatible, see (2.3).

The situation is summarized in Fig. 1 below. Identification symbols (a single and double bar) are used to stress that $\tilde{\Sigma}$ is compact.




Fig. 1: The modified cone \tilde{C} .

8.4.3. Sobolev spaces. We now recall some well-known facts about Sobolev spaces. If Ω is a relatively compact open set in a compact manifold X with smooth boundary $\partial\Omega$, then $H_0^1(\Omega)$, defined as the closure of $C_0^\infty(\Omega)$ in $H^1(\Omega)$ can also be characterized as $H_0^1(\Omega) = \{u \in H^1(\Omega) : u|_{\partial\Omega} = 0\}$. The restriction operator $r_\Omega : H^1(X) \rightarrow H^1(\Omega)$ is surjective from $H_{\partial\Omega}^1(X) = \{u \in H^1(X) : u|_{\partial\Omega} = 0\}$ to $H_0^1(\Omega)$, with right inverse $e_\Omega : H_0^1(\Omega) \rightarrow H_{\partial\Omega}^1(X)$ equal to the extension by 0 in $X \setminus \Omega$.

We set $\mathcal{E}_0(\Omega) := H_0^1(\Omega) \oplus L^2(\Omega)$, and $\mathcal{E}_{\partial\Omega}(X) = H_{\partial\Omega}^1(X) \oplus L^2(X)$. The operator $r_\Omega \oplus r_\Omega : \mathcal{E}_{\partial\Omega}(X) \rightarrow \mathcal{E}_0(\Omega)$ will still be denoted by r_Ω , and $e_\Omega \oplus e_\Omega : \mathcal{E}_0(\Omega) \rightarrow \mathcal{E}_{\partial\Omega}(X)$ by e_Ω .

We will use these facts for $\Omega = \Sigma_0, C_0$ and $X = \tilde{\Sigma}, \tilde{C}$. If $\Omega = C_0$, then we use the notation in (8.8), i.e. $\mathcal{E}_0(C_0) = H_0^1(C_0) \oplus \{0\} \sim H_0^1(C_0)$, since C_0 is characteristic.

8.4.4. Characteristic Cauchy problem.

Theorem 8.7. *The map*

$$\begin{aligned} T : \mathcal{E}_0(\Sigma_0) &\rightarrow \mathcal{E}_0(C_0) \\ f &\mapsto (U_{\Sigma_0} f)|_{C_0} \end{aligned}$$

is a homeomorphism.

Proof. We will prove the theorem by reducing ourselves to Thm. 8.5. We first claim that

$$(8.13) \quad T = r_{C_0} \circ \tilde{T} \circ e_{\Sigma_0}.$$

In fact this follows from the fact that $e_{\Sigma_0} : \mathcal{E}_0(\Sigma_0) \rightarrow \mathcal{E}(\tilde{\Sigma})$ is the extension by 0.

By Thm. 8.5, this implies that $T : \mathcal{E}_0(\Sigma_0) \rightarrow \mathcal{E}(C_0)$. Moreover by Huyghens principle, if $f \in C_0^\infty(\Sigma_0) \oplus C_0^\infty(\Sigma_0)$, then Tf vanishes near ∂C_0 , hence T maps continuously $\mathcal{E}_0(\Sigma_0)$ into $\mathcal{E}_0(C_0)$.

We next claim that $S = r_{\Sigma_0} \circ \tilde{T}^{-1} \circ e_{C_0}$ is a right inverse to T . In fact let $g \in \mathcal{E}_0(C_0)$ and $\tilde{f} = \tilde{T}^{-1} \circ e_{C_0} g = (\tilde{f}^0, \tilde{f}^1) \in \mathcal{E}(\tilde{\Sigma})$. Since $\partial\Sigma_0 = \partial C_0$ we have $\tilde{f}^0|_{\partial\Sigma_0} = g|_{\partial C_0} = 0$ hence $e_{\Sigma_0} \circ r_{\Sigma_0} \tilde{f} \in \mathcal{E}(\tilde{\Sigma})$. Since $\tilde{f} - e_{\Sigma_0} \circ r_{\Sigma_0} \tilde{f}$ vanishes on $\tilde{\Sigma}_0$, we obtain by (8.11) and Huyghens principle that

$$r_{C_0} \circ \tilde{T}(\tilde{f} - e_{\Sigma_0} \circ r_{\Sigma_0} \tilde{f}) = 0,$$

hence $T \circ Sg = r_{C_0} \circ \tilde{T} \tilde{f} = r_{C_0} \circ e_{C_0} g = g$. This completes the proof of the theorem. \square

8.5. Sobolev space on the cone in null coordinates. Let us set

$$R : C^\infty(C) \ni g \mapsto \beta^{-1}g(s, \theta) \in C^\infty(\mathbb{R} \times \mathbb{S}^{d-1}).$$

The goal in this subsection is to describe more precisely the image of $H_0^1(C_0)$ under R .

We will denote by $\tilde{C}_0 \subset \mathbb{R} \times \mathbb{S}^{d-1}$ the image of C_0 under the map $C \ni q \mapsto (s(q), \theta(q))$ where the coordinates (s, θ) are constructed in Lemma 2.5. Using that $\partial C_0 = \partial \Sigma_0$ is space-like and included in C , we easily obtain from Lemma 2.6 that \tilde{C}_0 is of the form:

$$\tilde{C}_0 = \{(s, \theta) \in \mathbb{R} \times \mathbb{S}^{d-1} : s < s_0(\theta)\},$$

for some smooth function s_0 . To simplify notation the measure $|m|^{\frac{1}{2}}(\theta)d\theta$ on \mathbb{S}^{d-1} will be simply denoted by $d\theta$. We also set $r = e^s$.

Proposition 8.8. *The image of $H_0^1(C_0)$ under R equals to the completion of $C_0^\infty(\tilde{C}_0)$ under the norm:*

$$\|\psi\|_1 := \left(\int_{\tilde{C}_0} (r^{-1}|\partial_s \psi|^2 + r^{-1}|\partial_\theta \psi|^2 + r^{-1}|\psi|^2) ds d\theta \right)^{\frac{1}{2}}.$$

We will denote this space by $\tilde{H}_0^1(\tilde{C}_0)$.

Remark 8.9. *Since $r \leq r_0$ on C_0 , we see that $\tilde{H}_0^1(\tilde{C}_0)$ injects continuously into $|D_s|^{\frac{1}{2}}L^2(\mathbb{R} \times \mathbb{S}^{d-1})$.*

Proof. We recall that (v, ψ) (see (2.4)) are coordinates on C such that the topology in $H_0^1(C_0)$ is given by the norm

$$\left(\int_{C_0} (|v|^{d-1}|\partial_v g|^2 + |v|^{d-3}|\partial_\psi g|^2 + |v|^{d-1}|g|^2) dv d\psi \right)^{\frac{1}{2}}.$$

Recall that we have set $r = e^s$. A function $g \in H_0^1(C_0)$ expressed in the coordinates (s, θ) or (r, θ) will be still denoted by g . Similarly the image of \tilde{C}_0 under the map $(s, \theta) \mapsto (e^s, \theta)$ will still be denoted by \tilde{C}_0 .

From Lemma 2.5 (3) and a routine computation, we see that an equivalent norm on $H_0^1(C_0)$ is:

$$(8.14) \quad \left(\int_{\tilde{C}_0} (r^{d-1}|\partial_r g|^2 + r^{d-3}|\partial_\theta g|^2 + r^{d-1}|g|^2) dr d\theta \right)^{\frac{1}{2}}.$$

Since $d = \dim M - 1 \geq 3$, the Hardy's inequality $-\Delta \geq C|x|^{-2}$ holds on $L^2(\mathbb{R}^d)$. Considering (r, θ) as polar coordinates on \mathbb{R}^d we obtain that:

$$\int_{\tilde{C}_0} r^{d-1}|\partial_r g|^2 + r^{d-3}|\partial_\theta g|^2 dr d\theta \geq C \int_{\tilde{C}_0} r^{d-3}|g|^2 dr d\theta, \quad g \in H_0^1(C_0).$$

Therefore adding a term $r^{d-3}|g|^2$ under the integral sign in (8.14) yields an equivalent norm on $H_0^1(C_0)$. Since r is bounded on \tilde{C}_0 this term dominates the term $r^{d-1}|g|^2$ and we finally obtain that the topology of $H^1(C_0)$ is given by the norm

$$\left(\int_{\tilde{C}_0} (r^{d-1}|\partial_r g|^2 + r^{d-3}|\partial_\theta g|^2 + \alpha r^{d-3}|g|^2) dr d\theta \right)^{\frac{1}{2}},$$

where the constant $\alpha > 0$ can be chosen arbitrarily large. Going back to coordinates (s, θ) we obtain the norm

$$(8.15) \quad \left(\int_{\tilde{C}_0} (r^{d-2}|\partial_s g|^2 + r^{d-2}|\partial_\theta g|^2 + \alpha r^{d-2}|g|^2) ds d\theta \right)^{\frac{1}{2}}.$$

For two functions $m, n \in C^\infty(\mathcal{C}_0)$ we write $m \sim n$ if $m = r_0 n$, for $r_0, r_0^{-1} \in S^0$, where the class S^0 is defined in Subsect. 2.5. We have $\beta \sim r^{-(d-1)/2}$, hence

$$(8.16) \quad \partial_s \beta, \partial_\theta \beta \sim r^{-(d-1)/2}.$$

Setting $\psi = Rg = \beta^{-1}g$ we have:

$$\partial_s g = \beta \partial_s \psi + (\partial_s \beta) \psi, \quad \partial_\theta g = \beta \partial_\theta \psi + (\partial_\theta \beta) \psi.$$

Using then (8.16), and choosing $\alpha \gg 1$ in (8.15), we obtain that (8.15) is equivalent to

$$(8.17) \quad \left(\int_{\tilde{\mathcal{C}}_0} (r^{-1} |\partial_s \psi|^2 + r^{-1} |\partial_\theta \psi|^2 + r^{-1} |\psi|^2) ds d\theta \right)^{\frac{1}{2}}.$$

This completes the proof of the proposition. \square

9. CHANGE OF NULL COORDINATES

The map $\rho : \text{Sol}_{\text{sc}}(P_0) \rightarrow \mathcal{H}(\tilde{C})$ introduced in Def. 4.1 depends on the choice of the null coordinates (s, θ) on C , i.e. on the choice of the initial hypersurface S , used in Lemma 2.5 to construct (s, θ) . In this section we discuss how our class of Hadamard states depends on the above choice.

9.1. New null coordinates. We fix a reference hypersurface S in C , yielding null coordinates (s, θ) near C such that $g|_C$ is given by (2.6) and $S = \{f = s = 0\}$.

We choose another hypersurface \tilde{S} transverse to $\nabla^a f$ in C , hence:

$$(9.1) \quad \tilde{S} = \{f = 0, s = b(\theta)\}, \text{ for some } b \in C^\infty(\mathbb{S}^{d-1}).$$

Since $\nabla^a f|_C = \partial_s$, we obtain that the new coordinates $(\tilde{s}, \tilde{\theta})$ obtained from Lemma 2.5 with S replaced by \tilde{S} are given by:

$$(9.2) \quad \tilde{\theta} = \theta, \quad \tilde{s}(s, \theta) = s - b(\theta).$$

We have then

$$g|_C = -2df d\tilde{s} + \tilde{h}_{ij}(\tilde{s}, \theta) d\theta^i d\theta^j,$$

and a standard computation shows that $|h|(\tilde{s}, \theta) = |h|(s, \theta)$, hence $\tilde{\beta}(\tilde{s}, \theta) = \beta(s, \theta)$. Denoting by $\tilde{\rho}$ the analog of ρ in Def. 4.1 for the new coordinates (\tilde{s}, θ) we have then

$$(9.3) \quad \tilde{\rho} \phi = U \rho \phi, \quad \phi \in \text{Sol}_{\text{sc}}(P_0),$$

where:

$$U : \begin{array}{l} \mathcal{H}(\tilde{C}) \rightarrow \mathcal{H}(\tilde{C}) \\ g \mapsto Ug(s, \theta) = g(s + b(\theta), \theta). \end{array}$$

The map U is symplectic on $(\mathcal{H}(\tilde{C}), \sigma_C)$ and unitary on $L^2(\tilde{C})$ with $U^* D_s U = D_s$.

Proposition 9.1. *If $A \in \tilde{\Psi}^{-\infty, p}(\tilde{C})$ then $UAU^{-1} \in \tilde{\Psi}^{-\infty, p}(\tilde{C})$.*

Remark 9.2. *Note that the above invariance property does not hold for the classes $\Psi^{m, p}(\tilde{C})$, since for example the classes $\Psi^{m, p}(\mathbb{R} \times \mathbb{R}^{d-1})$ are not even preserved by linear changes of variables $(s, y) \mapsto (s + Ay, y)$.*

Proof. We will use the Beals criterion explained in Subsect. 6.3, which implies that $B \in \tilde{\Psi}^{-\infty, p}(\tilde{C})$ iff for any functions $g_1, \dots, g_n \in C^\infty(\mathbb{S}^{d-1})$ and smooth vector fields X_1, \dots, X_m on \mathbb{S}^{d-1} and for any $N \in \mathbb{N}, k, k' \in \mathbb{R}$ one has:

$$(9.4) \quad \text{ad}_{X_1} \cdots \text{ad}_{X_m} \text{ad}_{g_1} \cdots \text{ad}_{g_n} B : H^{k, k'}(\tilde{C}) \rightarrow H^{k+N, k'-p+n}(\tilde{C}).$$

To simplify notation, we rewrite (9.4) as

$$(9.5) \quad \text{ad}_X^\alpha \text{ad}_g^\beta B : H^{k, k'}(\tilde{C}) \rightarrow H^{k+N, k'+p+|\beta|}(\tilde{C}),$$

denoting by \bar{X} , resp. \bar{g} an arbitrary n -uple of vector fields, resp. m -uple of functions.

If g is a function on \mathbb{S}^{d-1} , considered as a multiplication operator, and if X is a vector field on \mathbb{S}^{d-1} we have:

$$(9.6) \quad U^{-1}gU = g, \quad U^{-1}XU = X + (X \cdot db)\partial_s, \quad U^{-1}\partial_s U = \partial_s.$$

Let now $A \in \tilde{\Psi}^{-\infty, p}(\tilde{C})$. For $\psi \in C^\infty(\mathbb{S}^{d-1} \times \mathbb{S}^{d-1})$, let us denote by A_ψ the operator with distributional kernel $A(s_1, s_2, \theta_1, \theta_2)\psi(\theta_1, \theta_2)$. By the well-known properties of the pseudodifferential calculus on \mathbb{S}^{d-1} we know that if $\psi = 1$ in some neighborhood of the diagonal, then $A - A_\psi \in \tilde{\Psi}^{-\infty, -\infty}(\tilde{C})$, or equivalently maps $H^{k, k'}(\tilde{C})$ into $H^{k+N, k'+N}(\tilde{C})$ for any k, k', N . Using (9.6) this implies that $U(A - A_\psi)U^{-1}$ has the same property, hence belongs to $\tilde{\Psi}^{-\infty, -\infty}(\tilde{C})$.

Therefore we can replace A by A_ψ , and assume that the kernel of A is supported in $\mathbb{R} \times \mathbb{R} \times \Omega$, where Ω is an arbitrarily small neighborhood of the diagonal in $\mathbb{S}^{d-1} \times \mathbb{S}^{d-1}$. Introducing a smooth partition of unity $1 = \sum_1^M \chi_i$ on \mathbb{S}^{d-1} , we see that we can replace A by $\chi A \chi$, where $\chi \in C^\infty(\mathbb{S}^{d-1})$ is supported in a small neighborhood of a point $\theta_0 \in \mathbb{S}^{d-1}$. We pick local coordinates $\theta_1, \dots, \theta_{d-1}$ near θ_0 and rewrite (9.5) as:

$$(9.7) \quad \langle \partial_s \rangle^{k+N} \langle \partial_\theta \rangle^{k'-p+|\beta|} \text{ad}_{\bar{X}}^\alpha \text{ad}_{\bar{g}}^\beta A \langle \partial_s \rangle^{-k} \langle \partial_\theta \rangle^{-k'} \in B(L^2(\tilde{C})).$$

We set now $A' = UAU^{-1}$. Note first that if the kernel of A is supported in $\mathbb{R} \times \mathbb{R} \times \Omega$, then so is the kernel of A' , hence by the above discussion it suffices to check that A' satisfies (9.7). Let us set $U^{-1}XU = X'$ if X is a vector field on \mathbb{S}^{d-1} , and in particular $\partial'_\theta = U^{-1}\partial_\theta U = \partial_\theta + \partial_\theta b \partial_s$. Then an easy computation yields:

$$(9.8) \quad \begin{aligned} & \langle \partial_s \rangle^{k+N} \langle \partial_\theta \rangle^{k'-p+|\beta|} \text{ad}_{\bar{X}}^\alpha \text{ad}_{\bar{g}}^\beta UAU^{-1} \langle \partial_s \rangle^{-k} \langle \partial_\theta \rangle^{-k'} \\ &= U \langle \partial_s \rangle^{k+N} \langle \partial'_\theta \rangle^{k'-p+|\beta|} \text{ad}_{\bar{X}'}^\alpha \text{ad}_{\bar{g}'}^\beta A \langle \partial_s \rangle^{-k} \langle \partial'_\theta \rangle^{-k'} U^{-1}. \end{aligned}$$

Using (9.6) and the fact that $A \in \tilde{\Psi}^{-\infty, p}(\tilde{C})$, we obtain that

$$\text{ad}_{\bar{X}}^\alpha \text{ad}_{\bar{g}}^\beta A \in \tilde{\Psi}^{-\infty, p-|\beta|}(\tilde{C}),$$

$$\langle \partial_s \rangle^N \langle \partial_\theta \rangle^{k'-p+|\beta|} \text{ad}_{\bar{X}}^\alpha \text{ad}_{\bar{g}}^\beta A \langle \partial_s \rangle^N \langle \partial_\theta \rangle^{-k'} \in B(L^2(\tilde{C})),$$

for any $N \in \mathbb{N}$. It follows that the l.h.s. of (9.8) belongs to $B(L^2(\tilde{C}))$ if for any $s \in \mathbb{R}$ there exists $N \in \mathbb{N}$ such that:

$$(9.9) \quad \langle \partial_s \rangle^{-N} \langle \partial'_\theta \rangle^s \langle \partial_\theta \rangle^{-s}, \quad \langle \partial_s \rangle^{-N} \langle \partial_\theta \rangle^s \langle \partial'_\theta \rangle^{-s} \in B(L^2(\tilde{C})).$$

Let us now prove (9.9). The first statement of (9.9) is easy to check for $s \in \mathbb{N}$, using that $\partial'_\theta = \partial_\theta + \partial_\theta b \partial_s$. Conjugation by U gives the second statement for $s \in \mathbb{N}$. By duality and interpolation we obtain then (9.9) for arbitrary s , which completes the proof of the proposition. \square

From Prop. 9.1 and the fact that $U^*D_s U = D_s$, we obtain immediately the following result.

Proposition 9.3. *The classes of Hadamard states obtained in Thms. 7.4, 8.2 are independent on the choice of the null coordinates (s, θ) .*

APPENDIX A

A.1. Stokes formula. Let (M, g) an orientable, oriented pseudo-Riemannian manifold of dimension n . We denote by $d\text{Vol}_g \in \Lambda^n(M)$ the associated volume form and by $d\mu_g = |d\text{Vol}_g|$ the associated density.

Let $\Sigma \subset M$ a smooth submanifold of codimension 1 and $\iota : \Sigma \rightarrow M$ the natural injection, which induces $\iota^* : \Lambda(M) \rightarrow \Lambda(\Sigma)$. From the orientation of M and a continuous transverse vector field $v \in T_\Sigma M$ we obtain an induced orientation of Σ . If $\Sigma \subset \partial U$ for an open set $U \subset M$, we choose v pointing outwards.

If $\omega \in \Lambda^n(M)$ and $X \in TM$, then $X \lrcorner \omega \in \Lambda^{n-1}(M)$ and one sets:

$$\iota_X^* \omega := \iota^*(X \lrcorner \omega) \in \Lambda^{n-1}(\Sigma).$$

Similarly if $\mu = |\omega|$ is a density on M , we set $\iota_X^* \mu := |\iota_X \omega|$, which is a density on Σ .

If ∇_a is the Levi-Civita connection associated to g then:

$$\nabla_a X^a d\text{Vol}_g = d(X \lrcorner d\text{Vol}_g),$$

which applying Stokes formula:

$$(A.1) \quad \int_U d\omega = \int_{\partial U} \iota^* \omega, \quad \omega \in \Lambda^{n-1}(M),$$

to $\omega = \iota_X^* d\text{Vol}_g$, yields:

$$(A.2) \quad \int_U \nabla_a X^a d\text{Vol}_g = \int_{\partial U} \iota_X^* d\text{Vol}_g.$$

A.1.1. Non-characteristic boundaries. Assume first $\Sigma \subset \partial U$ is *non characteristic*, that is the one-dimensional space:

$$T_x(\Sigma)^{\text{ann}} \subset T_x M^*$$

is not *null* (the superscript ^{ann} denotes the annihilator). It follows that the metric $h := \iota^* g$ on Σ is non-degenerate (in the Lorentzian case, one typically assume that Σ is space-like, then $h = \iota^* g$ is Riemannian). Let $n \in T_\Sigma M$ be the unit, outward pointing normal vector field to σ . Then :

$$(A.3) \quad d\text{Vol}_h = \iota_n^* d\text{Vol}_g, \quad \iota_X^* d\text{Vol}_g = X^a n_a d\text{Vol}_h,$$

hence

$$\int_\Sigma \iota_X^* d\text{Vol}_g := \int_\Sigma X^a n_a d\sigma_h.$$

If all of ∂U is non-characteristic, then from (A.2) we obtain Gauss' formula:

$$(A.4) \quad \int_U \nabla_a X^a d\mu_g = \int_\Sigma X^a n_a d\sigma_h,$$

where $d\sigma_h = |d\text{Vol}_h|$.

A.1.2. Characteristic boundaries. Assume now that Σ is characteristic. Then there is no normal vector field anymore. To express the r.h.s. of (A.2), one chooses a defining function f for Σ , i.e. such that $f = 0, df \neq 0$ on Σ , and complete f with coordinates y^1, \dots, y^{n-1} such that $df \wedge dy^1 \wedge \dots \wedge dy^{n-1}$ is positively oriented. Then computing in the coordinates f, y^1, \dots, y^{n-1} one sees that

$$\iota_X^* d\text{Vol}_g = X^a \nabla_a f |g|^{\frac{1}{2}} dy^1 \wedge \dots \wedge dy^{n-1},$$

hence:

$$(A.5) \quad \int_\Sigma \iota_X^* d\text{Vol}_g = \int_\Sigma X_a \nabla^a f |g|^{\frac{1}{2}} dy^1 \wedge \dots \wedge dy^{n-1}$$

In the general case we can for example split ∂U as $\Sigma_1 \cup \Sigma_2$, where Σ_1 is non-characteristic, Σ_2 is characteristic and obtain:

$$(A.6) \quad \int_U \nabla_a X^a d\mu_g = \int_{\Sigma_1} X^a n_a d\sigma_h + \int_{\Sigma_2} X_a \nabla^a f |g|^{\frac{1}{2}} dy^1 \wedge \dots \wedge dy^{n-1}.$$

A.2. Conformal transformations. In this section we briefly discuss conformal transformations of a globally hyperbolic spacetime (M, g) . Let $\omega \in C^\infty(M)$ be strictly positive and consider the conformally related metric

$$g' = \omega^2 g.$$

Set $P = -\nabla^a \nabla_a + \frac{n-2}{4(n-1)} R$. For this special choice of the lower order terms, the conformal transformation $g \rightarrow g'$ amounts to

$$P' = \omega^{-n/2-1} P \omega^{n/2-1}.$$

This entails that the causal propagators are related by $E' = \omega^{-n/2+1} E \omega^{n/2+1}$. One concludes that multiplication by $\omega^{-n/2+1}$ induces a symplectic map

$$(A.7) \quad (\text{Sol}_{\text{sc}}(P), \sigma) \xrightarrow{\omega^{-n/2+1}} (\text{Sol}_{\text{sc}}(P'), \sigma'),$$

where σ, σ' are defined as in (3.2) using the respective volume densities.

We apply this discussion to (M_0, g) and the conformally related spacetime with metric $g' = \omega^2 g$. In the setting of Sect. 4.1, there is a monomorphism of symplectic spaces

$$(\text{Sol}_{\text{sc}}(P_0), \sigma_0) \xrightarrow{\rho} (\mathcal{H}(\tilde{C}), \sigma_C).$$

By (A.7) we also have a monomorphism

$$(\text{Sol}_{\text{sc}}(P'_0), \sigma'_0) \xrightarrow{\rho \circ \omega^{n/2-1}} (\mathcal{H}(\tilde{C}), \sigma_C).$$

Therefore, one can construct states for the conformally related spacetime using the bulk-to-boundary correspondence with a modified trace map $\rho' = \rho \circ \omega^{n/2-1}$.

A.3. Proof of Lemma 2.6. We fix a point $q \in C$ and complete the coordinate $x^0 = f$ by local coordinates $\bar{x} = (x^1, \dots, x^d)$ near q . The functions s, θ_k defined on C are denoted by $s(\bar{x}), \theta_k(\bar{x})$, since \bar{x} are local coordinates on C . We denote by $h(\bar{x})$ the restriction of g^{-1} to T^*C . Note that the fact that C is null implies that $g^{00}(0, \bar{x}) \equiv 0$ and that from Lemma 2.5 we have:

$$(A.8) \quad g^{i0}(\bar{x}) \partial_i s(\bar{x}) = -1, \quad g^{i0}(\bar{x}) \partial_i \theta_k(\bar{x}) = 0.$$

If X is a null vector, orthogonal to $C \cap \{u(\bar{x}) = u(q)\}$ and transverse to C , we obtain that

$$g^{-1} X = \lambda \left(\frac{1}{2} \nabla_i \nabla^i s, \nabla_i s \right), \quad \lambda \in \mathbb{R}.$$

Let us denote for the moment by $\tilde{s}, \tilde{\theta}_k$ the extensions of s, θ_k outside C , which are constant along the flow of X . We obtain that on C :

$$d\tilde{s} = \left(\frac{1}{2} ds \cdot h ds, ds \right), \quad d\tilde{\theta}_k = (ds \cdot h d\theta_k, d\theta_k).$$

Using also $df = (1, 0, \dots, 0)$ and (A.8), a routine computation leads to the following identities on C :

$$\begin{aligned} df \cdot g^{-1} df &= d\tilde{s} \cdot g^{-1} d\tilde{s} = df \cdot g^{-1} d\tilde{\theta}_k = d\tilde{s} \cdot g^{-1} d\tilde{\theta}_k = 0, \\ df \cdot g^{-1} d\tilde{s} &= d\tilde{s} \cdot g^{-1} df = -1, \\ d\tilde{\theta}_k \cdot g^{-1} d\tilde{\theta}_l &= \partial_i \theta_k h^{ij} \partial_j \theta_l. \end{aligned}$$

This implies that g is of the form (2.6) on C . \square

A.4. **Proof of Lemma 2.5.** Since (y^0, \bar{y}) are normal coordinates, we have:

$$(A.9) \quad g|_C = -dvdw + v^2 m_{ij}(\psi) d\psi^i d\psi^j + v^2 g_1,$$

where $m_{ij}(\psi) d\psi^i d\psi^j$ is the standard Riemannian metric on \mathbb{S}^{d-1} and g_1 is a smooth pseudo-Riemannian metric in the arguments dv, dw and $vd\psi^i$.

We start by expressing f in the normal coordinates (y^0, \bar{y}) . By Malgrange's preparation theorem [Hö1, Thm. 7.5.6] one can write

$$f(y^0, \bar{y}) = m(y^0, \bar{y})((y^0)^2 - |\bar{y}|^2) + a(\bar{y})y^0 + b(\bar{y}),$$

for m , resp. $a, b \in C^\infty$ near $(0, 0)$, resp. near 0 . Since $C \subset f^{-1}(\{0\})$, we obtain that $b(\bar{y}) = a(\bar{y})|\bar{y}|$, and since $b \in C^\infty(\mathbb{R}^d)$, necessarily $a \in O(|\bar{y}|^\infty)$. Moreover from the Hessian of f at p we obtain that $m(0, 0) = 1$.

Going to coordinates (v, w, ψ) , we obtain:

$$f(v, w, \psi) = m(v, w, \psi)vw + wa(v, w, \psi),$$

for $a \in O(|w - v|^\infty)$. Using also that $m(0, 0, \psi) = 1$, it follows that:

$$\partial_v f(v, 0, \psi) = \partial_{\psi^i} f(v, 0, \psi) = 0, \quad \partial_w f(v, 0, \psi) = v + r(v, \psi),$$

for $r \in O(|v|^2)$. Using (A.9) to express $(g^{-1})|_C$ we obtain after an easy computation that:

$$(A.10) \quad \nabla^a f = (v + v^2 a^0(v, \psi))\partial_v + v^2 a^i(v, \psi)\partial_{\psi^i},$$

where a^0, a^i are smooth, bounded functions near $v = 0$.

Let us now prove (1). Using (A.10) we obtain the equation near p :

$$(v + v^2 a^0(v, \psi))\partial_v s + v^2 a^i(v, \psi)\partial_{\psi^i} s = -1,$$

for smooth functions a^0, a^i . We set $s = \ln(vh(v, \psi))$ and obtain after an elementary computation

$$(1 + va^0)\partial_v h + a^0 h + va^i(v, \psi)\partial_{\psi^i} h = 0,$$

which we can uniquely solve on $[-\epsilon_1, \epsilon_1] \times \mathbb{S}^{d-1}$ by fixing $h(0, \psi)$. We may fix $h(0, \psi) > 0$ to ensure that $s(\epsilon_0, \psi) = 0$. We obtain $s = \ln(v) + \ln h(v, \psi)$ for $h \in C^\infty([-\epsilon_1, \epsilon_1] \times \mathbb{S}^{d-1})$, $h > 0$.

It remains to extend s globally to C . To do this it suffices to check that for any $q \in C$, the integral curve of $\nabla^a f$ through q crosses S at one and only one point. By [Wa, Corollary to Thm. 8.1.2] we know that q can be joined to p by a null geodesic γ . Locally a null geodesic on C is, modulo reparametrization, an integral curve of $\nabla^a f$. Since $\nabla^a f$ is complete, the whole $\gamma \setminus \{p\}$ is an integral curve of $\nabla^a f$. Hence the integral curve of $\nabla^a f$ through q crosses S . Choosing ϵ_0 in (2.5) small enough, we can ensure that $\nabla^a f \nabla_a v > 0$ on S , hence the integral curve through q crosses S at only one point. We can hence extend s globally to C , as a C^∞ function.

The proof of (2) is similar. We obtain the equation near p :

$$(v + v^2 a^0(v, \psi))\partial_v \theta^j + v^2 a^i(v, \psi)\partial_{\psi^i} \theta^j = 0,$$

or equivalently:

$$(1 + va^0(v, \psi))\partial_v \theta^j + va^i(v, \psi)\partial_{\psi^i} \theta^j = 0,$$

which we can solve in $]-\epsilon_1, \epsilon_1[\times \mathbb{S}^{d-1}$ by imposing $\theta^j(\epsilon_0, \psi) = \psi^j$. The estimate (3) on θ^j is immediate. We extend θ^j to all of C by the same argument as before.

□

A.5. Proof of Lemma 6.6. We use the characterization of the wavefront set of kernels using oscillatory test functions, which we now recall:

let $(\tilde{s}, \tilde{y}) \in C$ and $\lambda \geq 1$. We set for $(\sigma, \eta) \in \mathbb{R} \times \mathbb{R}^{d-1}$:

$$(A.11) \quad v_{\sigma, \lambda}(\cdot) = \chi(\cdot) e^{i\lambda \langle \cdot, \sigma \rangle} \in C_0^\infty(\mathbb{R}), \quad w_{\eta, \lambda}(\cdot) = \psi(\cdot) e^{i\lambda \langle \cdot, \eta \rangle} \in C^\infty(\mathbb{S}^{d-1}),$$

where $\chi \in C_0^\infty(\mathbb{R})$, resp. $\psi \in C^\infty(\mathbb{S}^{d-1})$ are supported near \tilde{s} , resp. \tilde{y} . We set $u_{(\sigma, \eta), \lambda} = v_{\sigma, \lambda} \otimes w_{\eta, \lambda}$. Note that if V resp. W are small neighborhoods of $\tilde{\sigma} \in \mathbb{R}$, resp. $\tilde{\eta} \in \mathbb{R}^{d-1}$ then for $n_+ = \max(n, 0)$, we have uniformly on $U = V \times W$:

$$(A.12) \quad \|u_{(\sigma, \eta), \lambda}\|_{k, k'} \in \begin{cases} O(\langle \lambda \rangle^{k_+ + k'_+}) \\ O(\langle \lambda \rangle^{k_+ + k'_+}) \text{ if } \sigma_0 \neq 0, \\ O(\langle \lambda \rangle^{k_+ + k'_+}) \text{ if } \eta_0 \neq 0. \end{cases}$$

Let now $\tilde{Y}_1, \tilde{Y}_2 \in T^*C$. Then $(\tilde{Y}_1, \tilde{Y}_2) \notin \text{WF}(a)'$ if there exists cutoff functions χ_i , resp. ψ_i with $\chi_i(\tilde{s}_i), \psi_i(\tilde{y}_i) \neq 0$ and neighborhoods $U_i = V_i \times W_i$ of $(\tilde{\sigma}_i, \tilde{\eta}_i)$ such that:

$$(A.13) \quad (u_{(\sigma_1, \eta_1), \lambda} | a u_{(\sigma_2, \eta_2), \lambda})_{L^2(C)} \in O(\langle \lambda \rangle^{-\infty}), \text{ uniformly for } (\sigma_i, \eta_i) \in U_i.$$

We first prove (1). Let $a \in B^{-\infty} \Psi^{p_2}(C)$ and $\tilde{Y}_1, \tilde{Y}_2 \in T^*C$ such that $\tilde{\sigma}_1 \neq 0$ or $\tilde{\sigma}_2 \neq 0$. Then (A.13) follows from (A.12) and the fact that $a : H^{k_1, k_2} \rightarrow H^{k_1+m, k_2+p_2}$ for any $m \geq 0$.

We now prove (2). If $a \in \Psi^{p_1, p_2}(C)$ the statement follows from Lemma 6.5. It remains to consider the case $a \in B^{-\infty} \Psi^{p_2}(C)$, and to prove that (A.13) holds if $(\tilde{\sigma}_1, \tilde{\eta}_1) = (0, 0)$ and $(\tilde{\sigma}_2, \tilde{\eta}_2) \neq 0$ or vice versa. If $\tilde{\sigma}_1 \neq 0$ or $\tilde{\sigma}_2 \neq 0$ we have already proved (A.13).

Assume now that $\tilde{\eta}_1 = 0$ and $\tilde{\eta}_2 \neq 0$, the other case being similar. Then we can find cutoff functions $g_i \in C_0^\infty(\mathbb{R}^{d-1})$ supported near $\tilde{\eta}_i$, with disjoint supports such that $(1 - g_i(\lambda^{-1} D_y)) u_{(\sigma_i, \eta_i), \lambda} \in O(\lambda^{-\infty})$ in all $H^{k, k'}$, uniformly for $(\sigma_i, \eta_i) \in U$. It follows that

$$\begin{aligned} & (u_{(\sigma_1, \eta_1), \lambda} | a u_{(\sigma_2, \eta_2), \lambda})_{L^2(C)} \\ &= (u_{(\sigma_1, \eta_1), \lambda} | g_1(\lambda^{-1} D_y) a g_2(\lambda^{-1} D_y) u_{(\sigma_2, \eta_2), \lambda})_{L^2(C)} + O(\langle \lambda \rangle^{-\infty}), \end{aligned}$$

uniformly for $(\sigma_i, \eta_i) \in U_i$. By pseudodifferential calculus on \mathbb{S}^{d-1} , we know that $g_1(\lambda^{-1} D_y) a g_2(\lambda^{-1} D_y) \in O(\langle \lambda \rangle^{-\infty})$ in $B(H^{k, k'})$ for any $k, k' \in \mathbb{R}$. Combined with (A.12), we obtain (A.13) also if $\tilde{\eta}_1 = 0$, $\tilde{\eta}_2 \neq 0$. This completes the proof of the lemma. \square

A.6. Proof of Lemma 8.6. Set $\gamma_x = \{(s, x) : s \leq 0\}$, $x \in \Sigma$. To prove that \overline{C} is the graph of a function F over Σ we have to show that for each $x \in \Sigma$, γ_x intersects \overline{C} at one and only one point. Then we have:

$$F(x) = \inf\{s \leq 0 : (s, x) \in I^+(p)\}.$$

If $F(x) = -\infty$ then $\gamma_x \subset I^+(p) \cap J^-(0, x) \subset J^+(p) \cap J^-(0, x)$. This last set is compact by global hyperbolicity, which is a contradiction. Hence γ_x intersects \overline{C} . Moreover if $(t_1, x) \in \overline{C}$, then $(s, x) \in J^-(p)$ for all $t_1 \leq s \leq 0$. This shows that γ_x intersects \overline{C} at only one point, hence the function F is well defined, and bounded.

Let (T^0, x^0) the coordinates of p . For $x \neq x^0$, C is smooth near $(F(x), x)$ and ∂_t is transverse to C . By the implicit function theorem this implies that F is smooth near x . Moreover if $K_1 \subset \Sigma$ is a compact set then dF is uniformly bounded on $K_1 \setminus \{x^0\}$. To prove this it suffices to introduce normal coordinates at p such that near p , C becomes a neighborhood of the tip of the flat lightcone. \square

Acknowledgments. The work of M.W. was partially supported by the FMJH (French Government Program: ANR-10-CAMP-0151-02).

REFERENCES

- [AS] H. Araki, M. Shiraishi, *On quasi-free states of canonical commutation relations I*, Publ. RIMS Kyoto Univ. 7 (1971/72), 105-120.
- [Bo] J.M. Bony, *Caractérisation des opérateurs pseudo-différentiels*, Séminaire EDP, Centre de Mathématiques Laurent Schwartz, (1996–1997).
- [BDM] M. Benini, C. Dappiaggi, S. Murro, *Radiative observables for linearized gravity on asymptotically flat spacetimes and their boundary induced states*, [arXiv:1404.4551](#) (2014).
- [BGP] C. Bär, N. Ginoux, F. Pfäffle, *Wave equation on Lorentzian manifolds and quantization*, ESI Lectures in Mathematics and Physics, EMS (2007).
- [BS] M. Borsero, R. Schulz, *Microlocal properties of bisingular operators*, Journal of Pseudo-Differential Operators and Applications (2013), 1-25.
- [BW] C. Bär, R.T. Wafo, *Initial value problems for wave equations on manifolds*, [arXiv:1408.4995](#) (2014).
- [Ca] F. Cagnac, *Problème de Cauchy sur un conoïde caractéristique pour des équations quasi-linéaires*, Ann. Mat. Pura Appl. 129 (1982), 13-41.
- [Do] M. Dossa, *Solutions C^∞ d'une classe de problèmes de Cauchy quasi-linéaires hyperboliques du second ordre sur un conoïde caractéristique*, Ann. Fac. Sci. Toulouse Math. 11 (2002), 351376.
- [DMP1] C. Dappiaggi, V. Moretti and N. Pinamonti, *Distinguished quantum states in a class of cosmological spacetimes and their Hadamard property*, J. Math. Phys. 50 (2009) 062304.
- [DMP2] C. Dappiaggi, V. Moretti and N. Pinamonti, *Rigorous construction and Hadamard property of the Unruh state in Schwarzschild spacetime*, Adv. Theor. Math. Phys. 15 (2011) 355.
- [DS] C. Dappiaggi, D. Siemssen, *Hadamard States for the Vector Potential on Asymptotically Flat Spacetimes*, Rev. Math. Phys. 25, 1350002 (2013).
- [FNW] S.A. Fulling, F.J. Narcowich, R.M. Wald, *Singularity structure of the two-point function in quantum field theory in curved space-time, II*, Annals of Physics, **136** (1981), 243-272.
- [GW] C. Gérard, M. Wrochna, *Construction of Hadamard states by pseudo-differential calculus*, Comm. Math. Phys. **325**, (2014), 713-755.
- [Mo1] V. Moretti, *Uniqueness theorem for BMS-invariant states of scalar QFT on the null boundary of asymptotically flat spacetimes and bulk-boundary observable algebra correspondence*, Comm. Math. Phys. **268** (2006), 727-756.
- [Mo2] V. Moretti, *Quantum out-states holographically induced by asymptotic flatness: invariance under space-time symmetries, energy positivity and Hadamard property*, Comm. Math. Phys. **279** (2008), 31-75.
- [Hö1] L. Hörmander, *The Analysis of Linear Partial Differential Operators I. Distribution Theory and Fourier Analysis*, Springer, Berlin Heidelberg New York (1985).
- [Hö2] L. Hörmander, *A remark on the characteristic Cauchy problem*, J. Funct. Anal. **93**, (1990), 270-277.
- [Ro] L. Rodino, *A class of pseudo differential operators on the product of two manifolds and applications*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (1975), no. 4, 287–302.
- [RT] M. Ruzhansky, V. Turunen, *Pseudo-differential Operators and Symmetries: Background Analysis and Advanced Topics*, Springer (2009).
- [Wa] R.M. Wald, *General Relativity*, University of Chicago Press (1984).
- [Wr] M. Wrochna, *Singularities of two-point functions in Quantum Field Theory*, PhD thesis, University of Göttingen (2013).

DÉPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ PARIS-SUD XI, 91405 ORSAY CEDEX, FRANCE
E-mail address: christian.gerard@math.u-psud.fr

UNIVERSITÉ JOSEPH FOURIER (GRENOBLE 1), INSTITUT FOURIER, UMR 5582 CNRS, BP 74
 38402 SAINT-MARTIN D'HÈRES CEDEX, FRANCE
E-mail address: michal.wrochna@ujf-grenoble.fr