



# On stochastic recursive equations and infinite server queues

Eitan Altman

## ► To cite this version:

Eitan Altman. On stochastic recursive equations and infinite server queues. RR-5211, INRIA. 2004, pp.13. inria-00070782

**HAL Id: inria-00070782**

**<https://inria.hal.science/inria-00070782>**

Submitted on 19 May 2006

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# *On stochastic recursive equations and infinite server queues*

Eitan Altman

**N° 5211**

May 2004

\_\_\_\_\_ Thème COM \_\_\_\_\_



*rapport  
de recherche*





## On stochastic recursive equations and infinite server queues

Eitan Altman\*

Thème COM — Systèmes communicants  
Projet Maestro

Rapport de recherche n° 5211 — May 2004 — 13 pages

**Abstract:** The purpose of this paper is to investigate some performance measures of the discrete time  $G/G/\infty$  queue under a general arrival process. We assume more precisely that at each time unit a batch with a random size may arrive, where the sequence of batch sizes need not be i.i.d. All we request is that it would be stationary ergodic and that the service duration has a phase type distribution. Our goal is to obtain explicit expressions for the first two moments of number of customers in steady state. We obtain this by computing the first two moments of some generic stochastic recursive equations that our system satisfies. We then show that these class of recursive equations allow to solve not only the  $G/PH/\infty$  queue but also a network of such queues. We finally investigate the process of residual activity time in a  $G/G/\infty$  queue under general stationary ergodic assumptions, obtain the unique stationary solution and establish coupling convergence to it from any initial state.

**Key-words:** Stochastic Recursive Equations, Branching Processes with Migration, Infinite Server Queues

\* This work was supported by the EURO-NGI network of excellence

# Équations stochastiques récursives et des files d'attente à serveur infini

**Résumé :** L'objectif de cet article est d'étudier des mesures de performances de la file d'attente  $G/G/\infty$  en temps discret sous des processus d'arrivées généraux. Nous supposons plus précisément qu'à chaque unité de temps une rafale de taille aléatoire arrive. Le processus des tailles des rafales est supposé stationnaire ergodique et la distribution des temps de services est de type de Cox. Notre but est d'obtenir explicitement les deux premiers moments du nombre de clients dans la file dans l'état stationnaire. Nous l'obtenons en calculant les deux premiers moments de certaines équations récursives stochastiques satisfaites par notre système. Puis nous montrons que cette classe d'équations récursives permet d'analyser non seulement une seule file mais aussi un réseau entier. Nous étudions enfin le processus de temps d'activité résiduel de la file  $G/G/\infty$  sous des hypothèses de temps d'arrivées et de services stationnaires et ergodiques. Nous obtenons le régime stationnaire unique et montrons le couplage à ce régime à partir de tout état initial.

**Mots-clés :** Équations Stochastiques Récursives, Processus de Branchement avec Migration, Files d'Attente à Infinité de Serveurs

# On stochastic recursive equations and infinite server queues

Eitan Altman

INRIA, BP93, 06902 Sophia Antipolis, France

The purpose of this paper is to investigate some performance measures of the discrete time  $G/G/\infty$  queue under a general arrival process. We assume more precisely that at each time unit a batch with a random size may arrive, where the sequence of batch sizes need not be i.i.d. All we request is that it would be stationary ergodic and that the service duration has a phase type distribution. Our goal is to obtain explicit expressions for the first two moments of number of customers in steady state. We obtain this by computing the first two moments of some generic stochastic recursive equations that our system satisfies. We then show that these class of recursive equations allow to solve not only the  $G/PH/\infty$  queue but also a network of such queues. We finally investigate the process of residual activity time in a  $G/G/\infty$  queue under general stationary ergodic assumptions, obtain the unique stationary solution and establish coupling convergence to it from any initial state.

## 1 Introduction

Most explicit expressions for performance measures in queueing networks are known under independence assumptions on the driving processes (service and interarrival times). An interesting challenge is to obtain explicit expressions for the case in which the independence is relaxed and only stationarity and ergodicity of some components of the driving sequences are assumed. One line of research that allows to handle stationary ergodic sequences is based on identifying measures that are insensitive to correlations. For example, the probability of finding a  $G/G/1$  queue empty is just the ratio between the expected service time and the expected interarrival time (which follows directly from Little's Law). The expected cycle duration in a polling system (under fairly general condition) too, depends on the interarrival, service and vacation times only through their expectations under general stationary ergodic assumptions (see e.g. [5]). An example of performance measures that depend on the whole distribution of service times but is insensitive to correlations is the growth rate of number of customers or of sojourn time in a (discriminatory) processor sharing queue in overload [4, 13].

---

\*This work was supported by the EURO-NGI network of excellence

In this paper we study a queueing problem under a stationary ergodic arrival process, in which the correlations indeed influence the performance but in which despite the dependence between arrival times, explicit expressions are obtained for the two first moments of the stationary number of customers. More precisely, we study the discrete time  $G/G/\infty$  queue in which at each time unit a random batch with a random size may arrive, where the sequence of batch sizes is stationary ergodic and service durations have a phase type distribution. We first compute the two moments of some generic stochastic recursive equations that our system satisfies. These are simplified versions of stochastic recursions introduced in [2] which already enabled us to study polling systems [2, 12] and queues with vacations [2] in which vacation times are correlated, and are related to branching process with migration [1]. We then show that these class of recursive equations allow to solve not only the  $G/PH/\infty$  queue but also a network of such queues. We finally investigate the process of residual activity time in a  $G/G/\infty$  queue under general stationary ergodic assumptions, obtain the unique stationary solution and establish coupling convergence to it from any initial state.

The infinite server queue which is the topic of our paper has had various applications in Teletraffic and in networking modeling. The output process of an  $M/GI/\infty$  queue has been used to model long range dependent traffic, c.f. in video applicationskruntz. In [17] the connectivity of ad-hoc networks on a line has been considered. The distribution of distance covered by a connected set of mobiles has been shown to correspond to a busy period in the  $GI/GI/\infty$  queue and its distribution was computed for various channel conditions. Furthermore the distribution of the number of connected mobiles has been computed using its correspondence to the number of customers served in a busy period of a  $GI/GI/\infty$  queue. Finally, the infinite server queue has also been used in the context of communication networks and distributed computer systems, see e.g. [14].

The structure of the paper is as follows. We introduce in Section 2 generic stochastic recursive equation corresponding to a branching type process in non Markov random environment with migration. The first and second moments of the corresponding state variables are introduced in Section 3. This allows us to derive in Section 4 explicit performance measures for the  $G/PH/\infty$  discrete time queue. Further stability results for the  $G/G/\infty$  queue are presented in Section 5 followed by a concluding section.

## 2 The model

Consider a column vector  $Y_n$  whose entries are  $Y_n^i$ ,  $i = 1, \dots, N$  where  $Y_n^i$  take values on the nonnegative integers. Consider the following stochastic recursive equation:

$$Y_{n+1} = A_n(Y_n) + B_n \quad (1)$$

where the  $i$  the element of the column vector  $A_n(Y_n)$  is given by

$$[A_n(Y_n)]_i = \sum_{j=1}^N \sum_{k=1}^{Y_n^j} \xi_{ji}^{(k)}(n)$$

where  $\xi^{(k)}(n)$ ,  $k = 1, 2, 3, \dots$ ,  $n = 1, 2, 3, \dots$  are i.i.d. random matrices of size  $N \times N$ . Each of its element is a nonnegative integer. Denote  $E[\xi_{ij}^{(k)}(n)] = A_{ji}$ . The  $N$ -dimensional vector  $B_n$  is a stationary ergodic stochastic whose entries  $B_n^i$ ,  $i = 1, \dots, N$  are nonnegative integers.

$A_n(y)$  has a divisibility property: if for some  $k$ ,  $y = y^0 + y^1 + \dots + y^k$  where  $y^m$  are integers, then  $A_n(y)$  can be represented as

$$A_n(y) = \sum_{i=0}^k A_n^{(i)}(y^i)$$

where  $\{A_n^{(i)}\}_{i=0,1,2,\dots,k}$  are i.i.d. with the same distribution as  $A_n(\cdot)$ . Note also that  $A_n(0) = 0$ . The divisibility property allows us to use the framework of [2] to characterize the distribution of  $Y_n$  and its limiting behavior.

We shall understand below  $\prod_{i=n}^k A_i(x) = x$  whenever  $k < n$ , and  $\prod_{i=n}^k A_i(x) = A_k A_{k-1} \dots A_n$  whenever  $k > n$ .

We note that although (1) is not linear in  $Y_n$ , it is linear in expectation; if we let  $y$  be a column vector then

$$E[A_n(y)] = Ay. \quad (2)$$

Moreover, we have for  $j > 1$  by Wald's equation

$$E\left[\left(\prod_{i=1}^j A_j\right)(y)\right] = A^j y \quad (3)$$

We recall the following property of our system:

**Theorem 1** (i)  $Y_n$  can be written in the form

$$Y_n = \sum_{j=0}^{n-1} \left( \prod_{i=n-j}^{n-1} A_i^{(n-j)} \right) (B_{n-j-1}) + \left( \prod_{i=0}^{n-1} A_i^{(0)} \right) (Y_0), \quad n > 0 \quad (4)$$

(ii) there is a unique stationary solution  $Y_n^*$  of (1), distributed like

$$Y_n^* =_d \sum_{j=0}^{\infty} \left( \prod_{i=n-j}^{n-1} A_i^{(n-j)} \right) (B_{n-j-1}), \quad n \in \mathbb{Z}, \quad (5)$$

The sum on the right side of (5) converges absolutely  $P$ -almost surely. Furthermore, for all initial conditions  $Y_0$ ,  $\|Y_n - Y_n^*\| \rightarrow 0$ ,  $P$ -almost surely on the same probability space. In particular, the distribution of  $Y_n$  converges to that of  $Y_0^*$  as  $n \rightarrow \infty$ .

**Proof.** (4) is obtained by iterating (1). Theorem 2 and Lemma 1 in [2] imply (ii). ■



### 3 First and second moments

Denote by  $y_i$  and  $y_i^{(2)}$  the first and second moment of the  $i$ th element of  $Y_n^*$ . Denote  $\text{cov}(Y)_{ij} = E[(Y_0^*)_i(Y_0^*)_j] - y_i y_j$ . Let  $b_i$  and  $b_i^{(2)}$  denote the two first moments of  $B_n^i$ . Denote  $\text{cov}(\xi)_{jk}^i = E(\xi_{ij}^{(0)} \xi_{ik}^{(0)}) - A_{ji} A_{ki}$  and define the following  $N \times N$  matrices:  $\mathcal{B}(k)$  is the matrix whose  $ij$ th entry equals  $E[B_0^i B_k^j]$ , where  $k$  is an integer.  $\hat{B}$  is the matrix whose  $ij$ th entry equals  $b_i b_j$ ,  $\text{cov}(B)$  is the matrix whose  $ij$ th entry equals  $E[B_0^i B_0^j]$ . Define  $\hat{\mathcal{B}} := \mathcal{B} - \hat{B}$ .

**Theorem 2** (i) The first moment of  $Y_n^*$  is given by

$$E[Y_0^*] = (I - A)^{-1} b, \quad (6)$$

(ii) Define  $Q$  to be the matrix whose  $(ij)$ th entry is

$$Q_{ij} = \sum_{k=1}^N y_k (\text{cov}(\xi)_{ij}^k).$$

Then the matrix  $\text{cov}(Y)$  is the unique solution of the set of linear equations:

$$\text{cov}(Y) = \text{cov}(B) + \sum_{r=1}^{\infty} \left( A^r \hat{\mathcal{B}}(r) + \left[ A^r \hat{\mathcal{B}}(r) \right]^T \right) + A \text{cov}(Y) A^T + Q. \quad (7)$$

**Proof.** (i) Taking the first moment at stationary regime of (1) we obtain (6).

(ii) To obtain the covariance, we first compute

$$\begin{aligned} E[(A_0(Y_0))_i (A_0(Y_0))_j] &= E(E[(A_0(Y_0))_i (A_0(Y_0))_j | Y_0]) \\ &= E\left(\sum_{k=1}^N Y_0^k A_{ki} \sum_{m \neq k} Y_0^m A_{jm}\right) + E\left(\sum_{k=1}^N E\left[\sum_{r=1}^{Y_0^k} \sum_{s=1}^{Y_0^k} \xi_{ki}^{(r)} \xi_{kj}^{(s)} \middle| Y_0^k\right]\right) \\ &= \sum_{k=1}^N \sum_{m \neq k} A_{ik} A_{jm} E[Y_0^k Y_0^m] + E\left(\sum_{k=1}^N E\left[\sum_{r=1}^{Y_0^k} \sum_{s=1, s \neq r}^{Y_0^k} \xi_{ki}^{(r)} \xi_{kj}^{(s)} \middle| Y_0^k\right]\right) \\ &\quad + E\left(\sum_{k=1}^N E\left[\sum_{r=1}^{Y_0^k} \xi_{ki}^{(r)} \xi_{kj}^{(r)} \middle| Y_0^k\right]\right) \\ &= \sum_{k=1}^N \sum_{m \neq k} A_{ik} A_{jm} E[Y_0^k Y_0^m] + \sum_{k=1}^N [(y_k^{(2)} - y_k)] A_{ik} A_{jk} + \sum_{k=1}^N y_k E[\xi_{ki}^{(0)} \xi_{kj}^{(0)}] \\ &= \sum_{k=1}^N \sum_{m=1}^N A_{ik} A_{jm} E[Y_0^k Y_0^m] + \sum_{k=1}^N y_k \text{cov}(\xi)_{ij}^k \end{aligned}$$

$$\begin{aligned}
E[(Y_0)_i B_0^r] &= \sum_{j=0}^{\infty} E \left\{ \left[ \left( \prod_{i=-j}^{-1} A_i^{(-j)} \right) (B_{-j-1}) \right]_i B_0^r \right\} \\
&= \sum_{j=0}^{\infty} E \left( E \left\{ \left[ \left( \prod_{i=-j}^{-1} A_i^{(-j)} \right) (B_{-j-1}) \right]_i B_0^r \right\} \middle| B_0, B_{-1}, B_{-2}, \dots \right) \\
&= \sum_{j=0}^{\infty} E \left( (A^j B_{-j-1})_i B_0^r \right) = \sum_{j=0}^{\infty} \sum_{s=1}^N (A^j)_{is} \mathcal{B}(j+1)_{s,r}
\end{aligned}$$

(where the last equality follows from (3)),

$$\begin{aligned}
E[(A_0(Y_0))_i B_0^r] &= E \left[ ((A_0(Y_0))_i B_0^r \middle| Y_0, B_0) \right] = \sum_{k=1}^N A_{ik} E[(Y_0)_k B_0^r] \\
&= \sum_{j=1}^{\infty} (A^j \mathcal{B}(j))_{i,r}
\end{aligned}$$

We thus obtain

$$\begin{aligned}
E[Y_0^i Y_0^j] &= E[B_0^i B_0^j] + E[(A_0(Y_0))_i B_0^j] + E[(A_0(Y_0))_j B_0^i] \\
&\quad + \sum_{k=1}^N \sum_{m=1}^N E[Y_0^k Y_0^m] A_{ik} A_{jm} + Q_{ij} \\
&= E[B_0^i B_0^j] + \sum_{r=1}^{\infty} \left( A^r \mathcal{B}(r) + [A^r \mathcal{B}(r)]^T \right)_{i,j} + \sum_{k=1}^N \sum_{m=1}^N E[Y_0^k Y_0^m] A_{ik} A_{jm} + Q_{ij}
\end{aligned}$$

so that

$$\text{cov}(Y)_{ij} = \text{cov}(B)_{ij} + \sum_{r=1}^{\infty} \left( A^r \hat{\mathcal{B}}(r) + [A^r \hat{\mathcal{B}}(r)]^T \right)_{i,j} + \sum_{k=1}^N \sum_{m=1}^N \text{cov}(Y)_{km} A_{ik} A_{jm} + Q_{ij}$$

We conclude that  $\text{cov}(Y)$  is a solution of (7). Next we show uniqueness. Let  $Z_1$  and  $Z_2$  be two solutions of (7) and define  $Z = Z_1 - Z_2$ . Then  $Z$  satisfies  $Z = A^T Z A$ . Iterating that we obtain that

$$Z = \lim_{n \rightarrow \infty} A^n Z (A^T)^n = 0$$

where the last equality follows since  $\|A\| < 1$ . This implies the uniqueness. ■

## 4 The G/PH/ $\infty$ queue

We now consider a discrete time G/PH/ $\infty$  queue. We shall first apply the general theory of previous sections in order to compute the steady state moments of some performance measures. We shall then strengthen the stability results (corresponding to Theorem 1) while relaxing further the statistical assumptions.

Let  $B_n = (B_n^1, \dots, B_n^N)^T$  be a column vector for each integer  $n$ , where  $B_n^i$  is the number of arrivals at the  $n$ th time slot that start their service at phase  $i$ .  $B_n$  is assumed to be a stationary ergodic sequence.

Service times are considered to be i.i.d. and independent of the arrival process. We represent the service time as the discrete time analogous of a phase type distribution: there are  $N$  possible service phases. The initial phase  $k$  is chosen at random according to some probability  $p(k)$ . If at the beginning of slot  $n$  a customer is in a service phase  $i$  then it will move at the end of the slot to a service phase  $j$  with probability  $P_{ij}$ . With probability  $1 - \sum_{j=1}^N P_{ij}$  it ends service and leaves the system at the end of the time slot.  $P$  is a substochastic matrix (it has nonnegative elements and its largest eigenvalue is strictly smaller than 1), which means that services ends in finite time w.p.1. and that  $(I - P)$  is invertible.

Let  $Y_n^i$  denote the number of customers in phase  $i$  at time  $n$ . Let  $\xi^{(k)}(n)$ ,  $k = 1, 2, 3, \dots$ ,  $n = 1, 2, 3, \dots$  be i.i.d. random matrices of size  $N \times N$ . Each of its element can take values of 0 or 1, and the elements are all independent. We assume  $E[\xi_{ij}^{(k)}(n)] = P_{ij}$ .

With these notation the queue dynamics can be written as (1) and we can apply the results of the previous sections to get the first two moments as well as the general distribution at stationary regime.

**Corollary 1** (i) *Theorems 1 and 2 hold for the G/PH/ $\infty$  queue.*

(ii) *The first and second moments of the number of customers at the system in stationary regime are given respectively by  $\mathbf{1}^T(I - A)^{-1}b$  and  $\mathbf{1}^T \text{cov}(Y)\mathbf{1}$ , respectively, where  $\mathbf{1}$  is a column vector with all entries 1's.*

**Remark 1** We present a simple interpretation of the first moment of the number of customers at the system. Denote by  $\lambda$  the expected number of arrivals per slot. Clearly  $\lambda = |b|$  where  $|b|$  is the sum of entries of the vector  $b$ . Define  $\zeta$  to be the expected service time of an arbitrary customer and let  $\rho = \lambda\zeta$ . We shall first compute  $\zeta$ . The  $ij$ th element of the matrix  $(I - A)^{-1}$  has the interpretation of the total expected number of slots that a customer that had arrived at service phase  $j$  spent at state  $i$ . Thus the  $j$ th entry of the vector  $\mathbf{1}^T(I - A)^{-1}$  has the interpretation of the total expected number of slots that a customer that had arrived at service phase  $j$  spent in the system. and let the vector  $\beta$  be the vector whose  $i$ th entry is  $b_i/|b|$ . Then

$$\zeta = \mathbf{1}^T(I - A)^{-1}\beta$$

and

$$\rho = (\mathbf{1}^T(I - A)^{-1}\beta)|b| = \mathbf{1}^T(I - A)^{-1}b,$$

which is our expression for the first moment of the number of customers at the system. This relation is known to hold in fact for general G/G/ $\infty$  queues, see e.g. [6, p. 134].

#### 4.1 Departure process

One can use the same methodology to describe the departure process. To do that, we can augment the system with a new "phase" which we call "d" (for departure), and update the phase transitions as follows:

$$\begin{aligned}\bar{P}_{ij} &= P_{ij}, \quad i, j \in \{1, \dots, N\}, \\ \bar{P}_{id} &= 1 - \sum_{j=1}^N P_{ij}, \quad i \in \{1, \dots, N\} \\ \bar{P}_{di} &= 0, \quad i \in \{1, \dots, N, d\}\end{aligned}$$

Quantities corresponding to the new system are denoted by adding a bar. We set  $\bar{B}_n^i = B_n^i$  for  $i = 1, \dots, N$  and  $\bar{B}_n^d = 0$  for all integers  $n$ . Since  $P$  is assumed to be sub-stochastic, so is  $\bar{P}$ . Note that in our new system, only customers in phases  $1, \dots, N$  correspond to those really present in the original system, whereas customers at phase  $d$  are already out of the system.

#### 4.2 Extension to a network

Consider now  $M$  stations, each with infinite number of servers. The service time at station  $i$  has a set  $\mathcal{N}_i$  of  $N_i$  phases. Let  $N = N_1 + \dots + N_M$ . For any  $j = 1, \dots, N$  let  $s(j)$  denote the station to which  $j$  corresponds, i.e. if  $j \in \mathcal{N}_i$  then  $s(j) = i$ .

If at time  $n$  a customer was at phase  $j$  in station  $s(j)$  then it either moves to another phase at the same station or moves to another phase in another station; the next phase  $k$  (either at the same station or at another one) is chosen with probability  $P_{jk}$ ; with probability  $1 - \sum_{k=1}^N P_{jk}$  the customer leaves the system. Again we assume that the choice of next phase are independent.

Let  $B_n = (B_n^1, \dots, B_n^N)^T$  be a column vector for each integer  $n$ , where  $B_n^i$  is the number of arrivals at the  $n$ th time slot that start their service at phase  $i$  in station  $s(i)$ .  $B_n$  is assumed to be a stationary ergodic sequence.

With this description we see that we can identify the whole network as a single server station problem with infinite number of servers and with  $N$  phases. Thus we can apply all previous results.

### 5 Residual activity time in the $G/G/\infty$ queue

Define the residual activity time at a given instant as the total time till the system empties from that instant onwards if new arrivals do not occur.

We shall analyze in this section the residual activity of a  $G/G/\infty$  queue under weaker statistical assumptions than those used so far. We shall obtain the existence of a stationary regime as well as convergence to it in the coupling-convergence sense (see e.g. Borovkov [7, 8]).

**The model** The  $n$ th arrival event occurs at time  $T_n$ : a batch of  $B_n$  customers arrive. Denote  $\tau_n = T_{n+1} - T_n$ ; they replace the fixed slots we had before. Let  $\sigma_n$  be the largest service time required among the  $B_n$  customers that arrive at time  $T_n$ . We shall assume that the joint sequence  $(\tau_n, \sigma_n)$  is stationary ergodic and that  $E[\tau_0]$  and  $E[\sigma_0]$  are finite and strictly positive.  $\sigma_n$  in particular, need not have a "phase type distribution" as before. Let  $V_n$  be the residual activity time just before  $T_n$ . Then  $V_n$  can be written recursively as:

$$V_{n+1} = \left( \max(V_n, \sigma_n) - \tau_n \right)^+$$

where  $(x)^+ := \max(x, 0)$ .

Iterating this relation gives:

$$\begin{aligned} V_{n+2} &= \left( \max \left\{ \left[ \max(V_n, \sigma_n) - \tau_n \right]^+, \sigma_{n+1} \right\} - \tau_{n+1} \right)^+ \\ &= \max \left( \max(V_n, \sigma_n) - \tau_n - \tau_{n+1}, \sigma_{n+1} - \tau_{n+1}, 0 \right). \end{aligned}$$

Further iterating directly yields:

$$V_{n+k} = \max(Z_n, Z_{n+1}, \dots, Z_{n+k-1}, 0)$$

where

$$Z_n = \max(V_n, \sigma_n) - \sum_{i=0}^{k-1} \tau_{n+i}, \quad Z_{n+j} = \sigma_{n+j} - \sum_{i=j}^{k-1} \tau_{n+i}, \quad j = 1, \dots, k-1.$$

**Stationary solution** We use the Loynes' type scheme [16] to obtain the stationary regime and the convergence to it.

**Theorem 3**  $V_n$  converges a.s. to a unique stationary regime that is given by

$$V_n^* := \left( \max_{j < n} \left[ \sigma_j - \sum_{i=j}^{n-1} \tau_i \right] \right)^+. \quad (8)$$

from any initial  $V_0$ . Moreover  $V_n^*$  is  $P$ -a.s. finite.

**Proof:** Define on the same probability space as the process  $V_n$  the shifted processes  $V_n^{[m]}$ , where  $m$  are integers:

$$V_{-m}^{[m]} = 0, \quad V_{n+1}^{[m]} = \left( \max(V_n^{[m]}, \sigma_n) - \tau_n \right)^+, \quad n \geq -m.$$

Then as before, we can write for  $n > -m$ :

$$V_n^{[m]} = \left( \max_{-m \leq j < n} \left[ \sigma_j - \sum_{i=j}^{n-1} \tau_i \right] \right)^+$$

which monotonely increases to the sequence  $V_n^*$  given in (8). Clearly  $V_n^*$  is a stationary ergodic process. We shall show that it is  $P - a.s.$  finite. Indeed, since  $(\tau_n, \sigma_n)$  is stationary ergodic, the Cesaro sums converge to the expectation  $P - a.s.$  and hence there is some R.V.  $J_0$  which is finite  $P - a.s.$  such that for all  $j > J_0$ ,

$$\sigma_{-j} < jE[\tau_0]/3 \text{ and } \sum_{i=-j}^{-1} \tau_i > j2E[\tau_0]/3$$

Hence the term in brackets in (8) is negative for all  $-j > J_0$  so that  $V_0^*$  is finite  $P - a.s.$  Due to stationarity this is true for  $V_n^*$  for all  $n$ .

**Coupling** We show that for any initial value  $V_0$  there is a time  $N_0$  which is finite  $P - a.s.$  such  $V_n$  coincides with  $V_n^*$  for all  $n > N_0$ . Indeed, fix  $V_0$  and define

$$N_0 := \inf \left\{ l : \max(V_0, V_0^*) < \sum_{i=0}^{l-1} \tau_i \right\}.$$

$N_0$  is clearly finite  $P - a.s.$  due to the ergodicity of  $\tau_i$ . Moreover, it is clear from the explicit expressions we have for  $V_0$  and for  $V_0^*$  that they coincide for  $n > N_0$ . Uniqueness of the stationary regime follows from the fact that coupling has been established for arbitrary initial state. ■

**Remark 2** Our construction establishes in fact that we have strong coupling convergence in the sense of [7, 8].

**Remark 3** A stability result is already given in [6, p. 133] for a general G/G/ $\infty$  queue. Namely, it is shown that  $V_0$  is finite almost surely but the form of the stationary regime and the convergence results are not given.

## 6 Concluding comments

In this paper we have studied and used stochastic recursive equations to investigate the discrete infinite server queue with batch arrivals where the size of the batches follow a general stationary ergodic process. We obtained explicit expressions for the first and second moments of the state variables appearing in the stochastic recursive equations and applied them to solve the infinite server queue problem. Other stochastic recursive equations have been used to study the stability of the queue under even more general probabilistic assumptions and convergence to a unique stationary regime has been established. The simple explicit expressions obtained makes our results appealing to various applications of the infinite server queue. For example, they can be used to represent the first and second moments of the number of connected mobiles at an arbitrary location in the one dimensional ad-hoc network of [17], using the equivalence between the ad-hoc network and an infinite server queue given in [17].

## References

- [1] S. R. Adke and V. G. Gadag, "A new class of branching processes", *Branching Processes: Proceedings of the First World Congress*, C.C.Heyde (Editor), 1-13, Springer Lecture Notes 99, 1995.
- [2] E. Altman, "Stochastic recursive equations with applications to queues with dependent vacations", *Annals of Operations Research*, 112(1): 43-61; Apr, 2002.
- [3] E. Altman and A. Hordijk, "Applications of Borovkov's Renovation Theory to Non-Stationary Stochastic Recursive Sequences and their Control", *Advances of Applied Probability* **29**, pp. 388-413, 1997.
- [4] E. Altman, T. Jimenez, D. Kofman, "DPS queues with stationary ergodic service times and the performance of TCP in overload", *Proceedings of IEEE Infocom*, Hong-Kong, March 2004.
- [5] E. Altman, P. Konstantopoulos, Z. Liu, "Stability, Monotonicity and Invariant Quantities in General Polling Systems", *Queueing Systems* **11**, pp. 35-57, 1992
- [6] F. Baccelli and P. Brémaud, *Elements of Queueing Theory*, Springer, second edition, 2003.
- [7] A. A. Borovkov, *Asymptotic Methods in Queueing Theory*, John Wiley & Sons, 1984 (translated from Russian).
- [8] A. A. Borovkov and S. G. Foss, "Stochastically recursive sequences and their generalizations", *Siberian Advances in Mathematics*, **2**, No. 1, pp. 16-81, 1992.
- [9] A. Brandt, "The stochastic equation  $Y_{n+1} = A_n Y_n + B_n$  with stationary coefficients", *Advances in Applied Probability*, Vol. 18, 1986.
- [10] A. Brandt, P. Franken and B. Lisek, *Stationary Stochastic Models*, Akademie-Verlag, Berlin, 1992.
- [11] P. Glasserman and D. D. Yao, "Stochastic vector difference equations with stationary coefficients", *J. Appl. Prob.*, Vol 32, pp 851-866, 1995.
- [12] R. Groenevelt and E. Altman, "An alternating priority server with correlated and cross correlated vacations", manuscript.
- [13] A. Jean-Marie and P. Robert, *On the transient behavior of the processor sharing queue*, *QUESTA* **17**, No. 1-2, pp. 129-136, 1994.
- [14] H. Kameda and Y. Zhang, "Uniqueness of the solution for optimal static routing in open BCMP queueing networks", *Math. Comput. Modelling* **22**, No. 10-12, 119-130, 1995.

- 
- [15] M. Kruntz and A. Makowski, "Modeling video traffic using M/G/infinity input processes: A compromise between Markovian and LRD models," *IEEE Journal on Selected Areas in Communications (JSAC)*, pp. 733-748, Vol. 16, No. 5, June 1998.
  - [16] R. Loynes, "The stability of a queue with non-independent inter-arrival and service times", *Proc. Cambr. Phil. Soc.* **58**, No. 3, pp. 497-520, 1962.
  - [17] D. Miorandi and E. Altman, "Connectivity in Ad-Hoc Networks: a Queueing Theoretical Approach", *Proceedings of WiOpt workshop*, Cambridge, UK, March 2004.





---

Unité de recherche INRIA Sophia Antipolis  
2004, route des Lucioles - BP 93 - 06902 Sophia Antipolis Cedex (France)

Unité de recherche INRIA Futurs : Parc Club Orsay Université - ZAC des Vignes  
4, rue Jacques Monod - 91893 ORSAY Cedex (France)

Unité de recherche INRIA Lorraine : LORIA, Technopôle de Nancy-Brabois - Campus scientifique  
615, rue du Jardin Botanique - BP 101 - 54602 Villers-lès-Nancy Cedex (France)

Unité de recherche INRIA Rennes : IRISA, Campus universitaire de Beaulieu - 35042 Rennes Cedex (France)

Unité de recherche INRIA Rhône-Alpes : 655, avenue de l'Europe - 38334 Montbonnot Saint-Ismier (France)

Unité de recherche INRIA Rocquencourt : Domaine de Voluceau - Rocquencourt - BP 105 - 78153 Le Chesnay Cedex (France)

---

Éditeur  
INRIA - Domaine de Voluceau - Rocquencourt, BP 105 - 78153 Le Chesnay Cedex (France)  
<http://www.inria.fr>  
ISSN 0249-6399