

A Cancellation Operator Suitable for Identification of Nonlinear Volterra Models

Céline Casenave, Gérard Montseny

► **To cite this version:**

Céline Casenave, Gérard Montseny. A Cancellation Operator Suitable for Identification of Nonlinear Volterra Models. IFAC Workshop on Control of Distributed Parameter Systems, CDPS 2009, Jul 2009, Toulouse, France. 2009. <hal-01061526>

HAL Id: hal-01061526

<https://hal.inria.fr/hal-01061526>

Submitted on 7 Sep 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A Cancellation Operator Suitable for Identification of Nonlinear Volterra Models

C. Casenave* and G. Montseny*

LAAS - CNRS, Université de Toulouse.

Abstract

We present a new identification method for nonlinear possibly singular Volterra models based on a suitable operatorial transformation of the problem with the property that some nonlinear terms are cancelled, allowing to get an equivalent formulation well adapted to least square identification.

Keywords

Nonlinear Volterra Equation, Diffusive Representation, Pseudo-Inverse.

1 Introduction

Consider a nonlinear Volterra model of the form:

$$H(\partial_t)X = F(X) + G(X)u, \quad (1)$$

where $u(t)$, $X(t) \in \mathbb{R}$, $t > 0$, F and G are possibly singular functions defined on \mathbb{R} , and $H(\partial_t)$ is an invertible causal convolution operator of diffusive type [3] such that $|\frac{H(i\omega)}{\omega}| \rightarrow 0$ when $|\omega| \rightarrow \infty$. We suppose that problem (1) is well-posed.

Let ψ_X the γ -diffusive representation of X , that is the unique solution of the following problem on $(t, \xi) \in \mathbb{R}^{+*} \times \mathbb{R}$ [3]:

$$\partial_t \psi_X = \gamma \psi_X + X, \quad \psi_X(0, \cdot) = 0, \quad (2)$$

with $\gamma : \mathbb{R} \rightarrow \mathbb{C}^-$. With μ the γ -symbol associated with $H(\partial_t) \circ \partial_t^{-1}$, the γ -diffusive formulation of equation (1) is [3]:

$$\langle \mu, \gamma \psi_X + X \rangle_{\Delta'_\gamma, \Delta_\gamma} = F(X) + G(X)u. \quad (3)$$

The problem under consideration in the sequel is to build estimations of $H(\partial_t)$ via the identification of the γ -symbol μ of $H(\partial_t) \circ \partial_t^{-1}$ from (noised) data $(\tilde{X}, \tilde{u}) = (X^k + \eta^k, u^k)_{k=1:n}$, with (X^k, u^k) solution of (1) and η^k continuous measurement noises.

2 The \mathfrak{D}_ε -transformed problem

Given \mathbf{X}, \mathbf{Y} two Banach spaces, let $(x, y) \in (C^0([t_0, T]; \mathbf{X}))^n \times (C^0([t_0, T]; \mathbf{Y}))^n$ and $\varepsilon \in \mathbb{R}^+$; we denote:

$$\Omega_{x,\varepsilon} := \bigcup_{i,j} \{(i, j)\} \times \Omega_{x,\varepsilon}^{i,j}, \quad \text{where } \Omega_{x,\varepsilon}^{i,j} := \{(t, \tau) \in [t_0, T]^2; \|x^i(t) - x^j(\tau)\| \leq \varepsilon\}.$$

The ε -cancellation operator \mathfrak{D}_ε is defined by:

$$(x, y) \mapsto \mathfrak{D}_\varepsilon(x, y) : \quad \begin{array}{ccc} \Omega_{x,\varepsilon} & \rightarrow & \mathbf{Y} \\ (i, j, t, \tau) & \mapsto & y^i(t) - y^j(\tau). \end{array} \quad (4)$$

*7 avenue du Colonel Roche, 31077 Toulouse Cedex 4, France. e-mail: casenave@laas.fr, montseny@laas.fr

Proposition 2.1. (1) \mathfrak{D}_ε is continuous. (2) For any continuous $f : \mathbf{X}_0 \subset \mathbf{X} \rightarrow \mathbf{Y}$, we have the *cancellation property*: $\mathfrak{D}_0(x, f \circ x) = 0$.

Thanks to proposition 2.1 and by application of operator $\mathfrak{D}_0(X, \cdot)$ to both members of the equation, model (3) is rewritten ($n = 1$):

$$\begin{cases} \langle \mu, \mathfrak{D}_0(X, \gamma \psi_X + X) \rangle = \mathfrak{D}_0(X, G(X)u) \\ \langle \mu, \gamma \psi_X(t_0, \cdot) + X(t_0) \rangle - F(X(t_0)) = G(X(t_0))u(t_0). \end{cases} \quad (5)$$

The interest of this new formulation is that up to the quantity $F_0 := F(X(t_0))$, the γ -symbol μ can be identified *independently* of F by (for simplicity, G is here supposed to be known):

$$(\mu^*, F_0^*) = \mathcal{Y}_\varepsilon^\dagger \left(\mathfrak{D}_\varepsilon(\tilde{X}, G(\tilde{X})\tilde{u}), G(\tilde{X}(t_0))\tilde{u}(t_0) \right),$$

where $\mathcal{Y}_\varepsilon^\dagger$ is the pseudo-inverse, in a suitable sense, of the operator \mathcal{Y}_ε defined by:


$$(\mu, F_0) \mapsto \left(\left\langle \mu, \mathfrak{D}_\varepsilon(\tilde{X}, \gamma \psi_{\tilde{X}} + \tilde{X}) \right\rangle, -F_0 + \left\langle \mu, \gamma \psi_{\tilde{X}}(t_0, \cdot) + \tilde{X}(t_0) \right\rangle \right). \quad (6)$$

By means of standard regression methods, the function F can then be easily estimated from the "pseudo graph" of F :

$$\mathcal{G}_F = \bigcup_{k,j} \left\{ \left(\tilde{X}^k(t_j), \left\langle \mu^*, \gamma \psi_{\tilde{X}^k}(t_j, \cdot) + \tilde{X}^k(t_j) \right\rangle - G(\tilde{X}^k(t_j))\tilde{u}^k(t_j) \right) \right\}. \quad (7)$$

3 A concrete example

This method has been tested on data elaborated from numerical simulation of the following model of spherical flames [2]: $X \partial_t^{1/2} X = X \ln(X) + u$, $u \geq 0$, $X \geq 0$. The identified symbol $H^*(i\omega) = \int \frac{\mu^*(\xi)}{i\omega - \gamma(\xi)} d\xi$ [3] and function F^* are given here-after. More details and extensions will be found in [1] and in a further publication.



(a) Frequency response of $H^*(i\omega)$.

(b) Identified F^* .

References

- [1] C. Casenave, G. Montseny, *Diffusive Identification of Volterra Models by Cancellation of the Nonlinear Term*, IFAC Symposium on System Identification, (SYSID 2009), Saint-Malo, July 6-8, 2009.
- [2] G. Joulin, *Point-source initiation of lean spherical flames of light reactants: an asymptotic theory*, Comb. Sci. and Tech., pp 99-113, 1985.
- [3] G. Montseny, *Représentation Diffusive*, Hermes-Science, Paris, 2005.