

Bindweeds or random walks in random environments on multiplexed trees and their asymptotics

Mikhail Menshikov, Dimitri Petritis, Serguei Popov

► **To cite this version:**

Mikhail Menshikov, Dimitri Petritis, Serguei Popov. Bindweeds or random walks in random environments on multiplexed trees and their asymptotics. Cyril Banderier and Christian Krattenthaler. Discrete Random Walks, DRW'03, 2003, Paris, France. Discrete Mathematics and Theoretical Computer Science, DMTCS Proceedings vol. AC, Discrete Random Walks (DRW'03), pp.205-216, 2003, DMTCS Proceedings. <hal-00000952v2>

HAL Id: hal-00000952

<https://hal.inria.fr/hal-00000952v2>

Submitted on 12 Aug 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Bindweeds or random walks in random environments on multiplexed trees and their asymptotics

Mikhail Menshikov¹ and Dimitri Petritis² and Serguei Popov^{3†}

¹ *Department of Mathematical Sciences, University of Durham, South Road, Durham DH1 3LE, United Kingdom*
Mikhail.Menshikov@durham.ac.uk

² *Institut de Recherche Mathématique, Université de Rennes I and CNRS UMR 6625, 35042 Rennes Cedex, France*
Dimitri.Petritis@univ-rennes1.fr

³ *Instituto de Matemática e Estatística, Universidade de São Paulo, Rua do Matão 1010, CEP 05508-090, São Paulo SP, Brasil*
popov@ime.usp.br

We report on the asymptotic behaviour of a new model of random walk, we term the bindweed model, evolving in a random environment on an infinite multiplexed tree. The term *multiplexed* means that the model can be viewed as a nearest neighbours random walk on a tree whose vertices carry an internal degree of freedom from the finite set $\{1, \dots, d\}$, for some integer d . The consequence of the internal degree of freedom is an enhancement of the tree graph structure induced by the replacement of ordinary edges by multi-edges, indexed by the set $\{1, \dots, d\} \times \{1, \dots, d\}$. This indexing conveys the information on the internal degree of freedom of the vertices contiguous to each edge. The term *random environment* means that the jumping rates for the random walk are a family of edge-indexed random variables, independent of the natural filtration generated by the random variables entering in the definition of the random walk; their joint distribution depends on the index of each component of the multi-edges. We study the large time asymptotic behaviour of this random walk and classify it with respect to positive recurrence or transience in terms of a specific parameter of the probability distribution of the jump rates. This classifying parameter is shown to coincide with the critical value of a matrix-valued multiplicative cascade on the ordinary tree (*i.e.* the one without internal degrees of freedom attached to the vertices) having the same vertex set as the state space of the random walk. Only results are presented here since the detailed proofs will appear elsewhere.

Keywords: Markov chain, trees, random environment, recurrence criteria, matrix multiplicative cascades

1 Introduction

1.1 *On the generality of the random walk in a random environment on a tree*

Markov chains on denumerable graphs enter in the modeling of a rich variety of phenomena; among such graphs, trees play a basic and generic rôle in the sense that they can encode simultaneously

[†]Partially supported by CNPq (302981/02-0)

- the topological structure of a vast class of general (*i.e.* not necessarily tree-like) denumerable graphs,
- the combinatorial structure of paths on these general graphs, and
- the probability structure generated on the trajectory space of the Markov chain evolving on the these general graphs.

In the sequel we give a short explanation of the reason trees play such a generic rôle among denumerable graphs. We know, after Kolmogorov, that in order to describe a random variable X with space of outcomes a discrete measurable set $(\mathbb{X}, \mathcal{X})$, of law \mathbb{P}_X , one has to use an abstract probability space $(\Omega, \mathcal{F}, \mathbb{P})$ on which the random variable is defined. However, the choice of this space is not unique. Among the infinite possible choices, there exists a “minimal” one, given by $\Omega = \mathbb{X}$, realizing the random variable as the identity map $X(\omega) = \omega$ and $\mathbb{P} = \mathbb{P}_X$. Similarly, when $(X_n)_{n \in \mathbb{N}}$ is a sequence of *independent* and identically distributed random variables with space of outcomes the discrete space $(\mathbb{X}, \mathcal{X})$, the minimal realization of the abstract probability space carrying the whole sequence is the *space of trajectories or full shift* $\Omega = \mathbb{X}^{\mathbb{N}}$ and the sequence is realized by the canonical projection $X_n(\omega) = \omega_n$, for $n \in \mathbb{N}$. However for a sequence of *dependent* random variables the space of trajectories may be not minimal. The reader can easily convince herself by considering the example of $\mathbb{X} = \{a, b, \dots, z\}$ and $(X_n)_{n \in \mathbb{N}}$ being the sequence of letters appearing in a natural language. Then the occurrence $(X_n, X_{n+1}, X_{n+2}) = rzt$ never appears in a language like English or French.

For the special case of Markovian dependence, the natural space for the realization of the sequence $(X_n)_{n \in \mathbb{N}}$ is the so called unilateral or bilateral *subshift spaces* obtained from the full shift by deleting all sequences containing forbidden subwords. For Markovian sequences, defined through a stochastic matrix P , subshift spaces can also be obtained in terms of the adjacency matrix of the sequence $A : \mathbb{X} \times \mathbb{X} \rightarrow \{0, 1\}$, given by the formula $A(x, y) = 1$ whenever $P(x, y) > 0$ and zero otherwise.

Suppose now we are given an arbitrary directed graph having a finite or denumerable set of vertices and being such that only a finite number of edges is emitted out of every vertex, not having any sources or sinks, and having (complex) weights attached on its edges. Since the vertex set is at most countable, it is in bijection with an at most countable alphabet. We say that a sequence of letters from the alphabet is a path of the graph, if the corresponding sequence of vertices is such that any ordered pair of subsequent vertices is an element of the edge set. The set of paths of arbitrary length is called *path space of the graph* (see [13] for precise definitions.) Notice that the condition that two subsequent vertices must be an allowed edge prevents some sequences of vertices from being a path. The adjacency matrix defines a subshift space for the trajectories of a symbolic unilateral or bilateral Markov chain that is identified with the path space of the graph. It is easy to show that the path space of any graph has a natural tree structure [1], giving thus a first hint that trees play a prominent rôle among graphs. Their importance does not stop here however. We can in fact define a so-called *evaluation map* from the path space of a graph into some set \mathbb{A} . Depending on the precise algebraic structure of the set \mathbb{A} and of the evaluation map, a vast class of objects can be defined. Instead of giving precise definitions, let us give the following

Example 1.1 (An elementary case) Start with the finite complete graph on 4 vertices, denote them by the letters E, N, W, and S for definiteness, and consider the unilateral path space of the graph, *i.e.* the set of words of arbitrary length on the alphabet of these 4 letters. Choose for the space \mathbb{A} the Abelian group \mathbb{Z}^2 and for the evaluation map the function taking the value $n_E \mathbf{e}_1 + n_N \mathbf{e}_2 + n_W (-\mathbf{e}_1) + n_S (-\mathbf{e}_2) \in \mathbb{Z}^2$, where n_E, n_N, n_W, n_S denote the occurrences of the letters E, N, W, and S in a given word, and $\mathbf{e}_1, \mathbf{e}_2$ are

the unit vectors of \mathbb{Z}^2 . The evaluation map can be thought as the physical observable “position of the random walker” when the history of the individual directions of movement is kept in memory; individual directions define a path in the path space and the evaluation map computes the actual position of the walker which has done n_E eastward movements, n_W westward movements, etc. The graph that is generated in this way is the graph that coincides with the Cayley graph of the Abelian group \mathbb{Z}^2 . The evaluation map being many-to-one, its multiplicity encodes the combinatorics of the path space while the product of the graph weights gives the relative weight of individual paths in the path space of this graph. When the weights are probability vectors, this weight coincides with the probability of the trajectory of the simple random walk on \mathbb{Z}^2 . Notice however that the weights need not to be probability vectors, allowing us to consider both random and quantum grammars.

The above example serves as a basic paradigm; by appropriately changing the evaluation map, it can be generalised in numerous ways to produce configuration space for DNA strands (when the path space coincides with the path space of the complete graph over 4 letters), non-Abelian groups like the free group on 2 generators, Cuntz–Krieger C^* -algebras (*i.e.* non-commutative operator algebras attaching non-zero partial isometries on every edge; see [24] for additional details) naturally arising in some problems of quantum information. Infinite trees provide a rich variety of mathematical problems, particularly connected to their non-amenability; beyond their mathematical interest they arise as more or less realistic models in several applied fields like random search algorithms in large data structures, Internet traffic, random grammars and probabilistic Turing machines, DNA coding, interacting random strings and automated languages, etc. The previously exposed ideas, already lurking in [17, 19], have been exemplified in [8, 1], and will be further exploited in [24].

Random environment is a means to introduce context-dependence in the language. Again, due to the generality of trees as underlying graphs, random walks in random environment on trees provide interesting context-dependent results for random walks on a huge class of more general graphs.

Random walks in random environments on various types of graphs are known to display a behaviour dramatically differing from the one for ordinary random walks on the same graph. See [27, 26, 11, 12, 16, 4, 18, 22, 5, 2] etc. for a very partial list of known results and models.

1.2 Rough statement of the main results for bindweeds

As a simple example, consider a rooted tree with constant branching b , and let us construct a random walk in random environment on it by sampling the transition probabilities (or transition rates, for continuous time) in each vertex from a given distribution, independently. When studying the question of (positive) recurrence of such a walk, one naturally arrives (see [16, 22]) at the following model. On each edge a of the tree, we place an independent copy of a positive random variable $\hat{\xi}_a$. Then, for each vertex \mathbf{v} denote $\hat{\xi}[\mathbf{v}] = \hat{\xi}_{a_1} \dots \hat{\xi}_{a_n}$, where a_1, \dots, a_n is the (unique) path connecting the root to \mathbf{v} . Models of this type are called *multiplicative cascades* (see e.g. [10, 15]), and, to study the positive recurrence of the corresponding random walk in random environment, one has to answer the question whether the sum of $\hat{\xi}[\mathbf{v}]$ is finite. It turns out (see e.g. [16]) that the classification parameter for this problem is

$$\hat{\lambda} = \inf_{s \in [0,1]} \mathbb{E} \hat{\xi}_a^s, \quad (1)$$

which is then compared to $1/b$ (in fact, in [16] the case of general tree was considered) and the following result is established:

- if $\hat{\lambda}b < 1$ then the random walk for almost all environments is positive recurrent
- if $\hat{\lambda}b > 1$ then the random walk for almost all environments is transient.

The critical case $\hat{\lambda}b = 1$ is more complicated.

In [23], we introduced a model of random walk with internal degrees of freedom whose multiplicative cascade counterpart is expressed by a model placing random matrices on the edges in place of scalar random variables. The main classification parameter, λ , will be defined in the formula (3) below, and the main results are Theorems 3.1 and 3.2 (a lot of preliminary work is required, however, before formulating these results). This model is also equivalent to random walk in random environment on a multiplexed tree, a process we term *bindweed* in the sequel. However, for trees with average branching b , the rough statement of our result is as above with $\hat{\lambda}$ replaced by λ .

It was remarked in [22] that asymptotic properties like recurrence/transience of random walk on trees with constant branching b are intimately connected to the existence of non-trivial solutions for the so-called multiplicative chaos equation of order b , first introduced as a simple turbulence model in [20]. The simplest variant of the multiplicative chaos equation is the following: let $(\xi_i)_{i=1,\dots,b}$, with $b \in \mathbb{N}$, be a finite family of non-negative random variables having known joint distribution and $(Y'_i)_{i=1,\dots,b}$ and Y be a family of $b+1$ independent non-negative random variables distributed according the same unknown law and verifying

$$Y \stackrel{\text{law}}{=} \sum_{i=1}^b Y'_i \xi_i.$$

The multiplicative chaos problem consists in determining under which conditions on the joint distribution of the ξ 's the above equation has a non-trivial solution. This scalar problem is thoroughly studied in the literature, see e.g. [3, 6, 14]. As we remark later in this paper, the matrix multiplicative chaos equation may be an interesting problem to study as well.

2 Notation

In this section we give the formal definitions concerning trees, in particular, we define the notions of the growth rate and the branching number.

We denote $\mathbb{R}_+ = [0, \infty[$, $\mathbb{Z}_+ = \{0, 1, 2, \dots\}$, $\mathbb{N} = \{1, 2, 3, \dots\}$, and for every $n \in \mathbb{N}$, $\mathbb{N}_n = \{1, 2, \dots, n\}$, while $\mathbb{N}_0 = \emptyset$. Let $\mathcal{A} \equiv \mathcal{A}^1$ be a finite or infinite denumerable set, called the *alphabet*. Define $\mathcal{A} = \{\emptyset\}$ and for every $n \in \mathbb{N}$ denote

$$\mathcal{A}^n = \{\alpha = \alpha_1 \cdots \alpha_n : \alpha_i \in \mathcal{A} \text{ for } i \in \mathbb{N}_n\}$$

the set of words of length n (i.e. having n letters),

$$\mathcal{A}^* = \bigcup_{n \in \mathbb{Z}_+} \mathcal{A}^n$$

the set of words of arbitrary (finite) length, and

$$\partial \mathcal{A}^* \equiv \mathcal{A}^\infty = \{\alpha = \alpha_1 \alpha_2 \cdots : \alpha_i \in \mathcal{A} \text{ for } i \in \mathbb{N}\}$$

the set of infinite words. Finally, denote $\overline{\mathcal{A}^*} = \mathcal{A}^* \cup \partial \mathcal{A}^*$ and $\overset{\circ}{\mathcal{A}^*} = \mathcal{A}^* \setminus \mathcal{A}^0$.

For every $\alpha \in \mathcal{A}^*$, there exists $n \in \mathbb{Z}_+$ such that $\alpha \in \mathcal{A}^n$; in this situation $|\alpha| := n$ denotes the *length* of the word α with the convention $|\emptyset| = 0$. Consistently, for every $\alpha \in \partial \mathcal{A}^*$, we have $|\alpha| = \infty$. For $\alpha \in \overline{\mathcal{A}^*}$

with $|\alpha| \geq n$ we denote by $\alpha \downarrow_n = \alpha_1 \cdots \alpha_n \in \mathcal{A}^n$ the *restriction* of α to its n first letters with the convention $\alpha \downarrow_0 = \emptyset$. For every $\alpha \in \mathcal{A}^*$, the *ancestor* $\hat{\alpha}$ of α is defined by $\hat{\alpha} = \alpha \downarrow_{|\alpha|-1}$. For $\alpha \in \mathcal{A}^*$ and $\beta \in \overline{\mathcal{A}^*}$, the *concatenation* of α followed by β is the word $\alpha\beta = \alpha_1 \cdots \alpha_{|\alpha|}\beta_1\beta_2 \cdots$ and for $\alpha, \beta \in \mathcal{A}^*$, their *common radix* $\alpha \wedge \beta$ is the longest word $\gamma \in \mathcal{A}^*$ such that $\alpha = \gamma\alpha'$ and $\beta = \gamma\beta'$ for some words $\alpha', \beta' \in \mathcal{A}^*$. We write $\alpha \leq \beta$ if $\alpha = \alpha \wedge \beta$.

Remark: Notice that, consistently with the above notation, the symbol \mathbb{N}^* denotes the set of finite words on the alphabet \mathbb{N} , contrary to some tradition (especially the French one) where this symbol is used to denote what we call here \mathbb{N} .

Definition 2.1 A mapping $B : \mathbb{N}^* \rightarrow \mathbb{Z}_+$ is called a *branching function*.

To each branching function corresponds a uniquely determined rooted tree $\mathbb{T} = (\mathbb{V}, \mathbb{A})$ with vertex set $\mathbb{V} \equiv \mathbb{V}^*(B) \subseteq \mathbb{N}^*$ and edge set $\mathbb{A} = \overset{\circ}{\mathbb{V}}$ defined as follows: $\mathbb{V}^*(B) = \cup_{n \in \mathbb{N}} \mathbb{V}^n(B)$ where $\mathbb{V}^0(B) = \{\emptyset\} = \{\text{root}\} \equiv \{\mathbf{0}\}$ and for $n \in \mathbb{N}$,

$$\mathbb{V}^n(B) = \{\mathbf{v} = v_1 \cdots v_n : v_l \in \mathbb{N}_{B(\mathbf{v} \downarrow_{l-1})}, \text{ for } l = 1, \dots, n\}.$$

The branching function is said to be *without extinction* if the corresponding tree has non-trivial boundary $\partial\mathbb{V}$. The edge set is the subset of unordered pairs of vertices $[\mathbf{u}, \mathbf{v}] = [\mathbf{v}, \mathbf{u}]$ such that either $\mathbf{v} = \hat{\mathbf{u}}$ or $\mathbf{u} = \hat{\mathbf{v}}$. Since every vertex has a unique ancestor, every edge is indexed by its outmost vertex, *i.e.* for every $\mathbf{v} \in \overset{\circ}{\mathbb{V}}$, the corresponding edge is $a(\mathbf{v}) = [\hat{\mathbf{v}}, \mathbf{v}]$, showing thus that $\mathbb{A} \simeq \overset{\circ}{\mathbb{V}}$.

If $\mathbf{u}, \mathbf{v} \in \mathbb{V}$ and $\mathbf{u} \leq \mathbf{v}$ we define the *path* $[\mathbf{u}, \mathbf{v}]$ as the collection of the $|\mathbf{v}| - |\mathbf{u}|$ edges $[\mathbf{u}, \mathbf{v} \downarrow_{|\mathbf{u}+1}], \dots, [\hat{\mathbf{v}}, \mathbf{v}]$, and if $\mathbf{u} = \emptyset$ then we simply denote by $[\mathbf{v}]$ the path $[\emptyset, \mathbf{v}]$ for every $\mathbf{v} \in \overset{\circ}{\mathbb{V}}$. In the sequel we shall consider only *branching functions without extinction*.

Definition 2.2 Let $\kappa_n = \text{card } \mathbb{V}^n(B)$ denote the cardinality of the n^{th} generation of the tree defined by the branching function without extinction B . We call *lower growth rate* of the tree

$$\underline{\text{gr}}(\mathbb{V}) = \liminf_n \kappa_n^{1/n},$$

upper growth rate of the tree

$$\overline{\text{gr}}(\mathbb{V}) = \limsup_n \kappa_n^{1/n},$$

and, if $\underline{\text{gr}}(\mathbb{V}) = \overline{\text{gr}}(\mathbb{V})$, we call the common value *growth rate*

$$\text{gr}(\mathbb{V}) = \lim_n \kappa_n^{1/n}.$$

For $\mathbf{u}, \mathbf{v} \in \partial\mathbb{V}$, define $\delta(\mathbf{u}, \mathbf{v}) = \exp(-|\mathbf{u} \wedge \mathbf{v}|)$. It can be shown that δ is a distance on $\partial\mathbb{V}$. Moreover if $\|B\|_\infty = \sup_{\mathbf{v} \in \mathbb{V}} B(\mathbf{v}) < \infty$ then the space $(\partial\mathbb{V}, \delta)$ is compact and we can define its Hausdorff dimension $\dim_H \partial\mathbb{V}$ as usual (see [7] for instance).

Definition 2.3 For a tree \mathbb{V} generated by a branching function B with $\|B\|_\infty < \infty$, we define its *branching rate*

$$\text{br}(\mathbb{V}) = \exp(\dim_H \partial\mathbb{V}).$$

It is shown in [21, 16] that $\text{br}(\mathbb{V}) = \sup\{\lambda : \inf_{\sum_{v \in C} \lambda^{-|v|} > 0} \lambda > 0\}$ where the infimum is evaluated over all cutsets C of \mathbb{V} . We have in general that $\text{br}(\mathbb{V}) \leq \underline{\text{gr}}(\mathbb{V})$.

3 Matrix multiplicative cascades and the corresponding results

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be some abstract probability space which carries all the random variables that will be needed in the model. Let (\mathbb{V}, \mathbb{A}) be the rooted tree associated with a given branching function B . Let G be the topological group $\text{GL}(d, \mathbb{R})$, \mathcal{G} its Borel σ -algebra and μ a probability on (G, \mathcal{G}) . Denote by $\sigma_\mu = \text{supp } \mu \subset G$ the support of the measure μ and by Σ_μ the semi-group generated by σ_μ . On $(\Omega, \mathcal{F}, \mathbb{P})$, define an edge-indexed family of independent G -valued random variables $(\xi_a)_{a \in \mathbb{A}}$ identically distributed according to μ , *i.e.*

$$\mathbb{P}(\xi_a \in dg) = \mu(dg), \text{ for all } a \in \mathbb{A}.$$

For $\mathbf{u}, \mathbf{v} \in \mathbb{V}$ with $\mathbf{u} \leq \mathbf{v}$ define

$$\xi[\mathbf{u}, \mathbf{v}] \equiv \prod_{a \in [\mathbf{u}, \mathbf{v}]}^{\leftarrow} \xi_a,$$

where \prod^{\leftarrow} denotes the product in reverse order, *i.e.* if $[\mathbf{u}, \mathbf{v}] = a_1 \cdots a_k$ then $\xi[\mathbf{u}, \mathbf{v}] = \xi_{a_k} \cdots \xi_{a_1}$ with the convention $\xi[\mathbf{v}, \mathbf{v}] = e$ where e is the neutral element of G . We introduce the following G -valued random processes: the *matrix-multiplicative cascade process*

$$\Psi_n = \sum_{\mathbf{v} \in \mathbb{V}^n} \xi[\mathbf{v}], \quad n \in \mathbb{N}$$

and the *integrated matrix-multiplicative cascade process*

$$\zeta_n = \sum_{k=1}^n \Psi_k, \quad n \in \mathbb{N}.$$

For a fixed $\mathbf{v} \in \partial \mathbb{V}$ and all $n \in \mathbb{N}$

$$X_n \equiv X_n(\mathbf{v}) = \xi[\mathbf{v} \upharpoonright_n].$$

It is immediate to see (cf. [25]) that $(X_n)_{n \in \mathbb{N}}$ is a G -valued multiplicative Markov chain with stochastic kernel

$$P(g, dg') \equiv \mathbb{P}(X_{n+1} \in dg' | X_n = g) = \mu \star \delta_g(dg'), \quad n \in \mathbb{N},$$

where for two measures μ, μ' on (G, \mathcal{G}) their convolution $\mu \star \mu'$ is defined by its dual action on $L^1(G)$ via

$$\langle \mu \star \mu', f \rangle \equiv \int_G f(g) \mu \star \mu'(dg) = \int_G \int_G f(gg') \mu(dg) \mu'(dg'),$$

for all $f \in L^1(G)$. We equip $\text{End}(\mathbb{R}^d) \equiv \text{GL}(d, \mathbb{R}) = G$ with the operator norm, denoted $\|\cdot\|$, stemming from the l_1 norm of the vector space \mathbb{R}^d .

In order to be able to apply the results of [9] to our special case, we require the following conditions on μ :

Condition 1 (Integrability): For all $s \in \mathbb{R}_+$,

$$\int_G \|g\|^s \mu(dg) < \infty.$$

Condition 2 (Strong irreducibility): We assume that the set Σ_μ is strongly irreducible, *i.e.* there is no finite Σ_μ -invariant family of proper subspaces.

Condition 3 (Strict positivity): We assume that $\sigma_\mu \subseteq \overline{G}_+$ and that $\mu(\overline{G}_+ \setminus G_+) = 0$.

For $s \geq 0$, define

$$k(s) = \lim_n \left(\int_G \|g\|^s \mu^{*n}(dg) \right)^{1/n}. \tag{2}$$

(By virtue of theorem 1 of [9], under conditions 1–3 this limit exists in \mathbb{R}_+ and defines a log-convex function. As a matter of fact, in [9] a weaker condition than 3, called proximity, is needed to prove this result.) We define in the sequel the quantity λ , that turns out to be the main classification parameter for the matrix multiplicative cascades model, by

$$\lambda = \inf_{s \in [0,1]} k(s) \tag{3}$$

(compare (3) with (1).)

We are now in the position to state our main results.

Theorem 3.1 *Let (\mathbb{V}, \mathbb{A}) be some tree defined in terms of a given branching function B and $\overline{\text{gr}}(\mathbb{V})$ and λ defined as in definition 2.2 and equation (3) respectively. Under the conditions 1, 2, and 3,*

$$\lambda \overline{\text{gr}}(\mathbb{V}) < 1 \Rightarrow \zeta_{\infty,ij} < \infty \text{ almost surely, for all } i, j = 1, \dots, d.$$

Theorem 3.2 *Let $\text{br}(\mathbb{V})$ and λ be the quantities introduced in definition 2.3 and equation 3 respectively and let $\chi \in \mathbb{R}^d$ be the vector having all its components equal to 1: $\chi_i = 1$, for all $i = 1, \dots, d$. Let (\mathbb{V}, \mathbb{A}) be some tree defined in terms of a given branching function B without extinction. Under the conditions 1, 2, and 3',*

$$\lambda \text{br}(\mathbb{V}) > 1 \Rightarrow Z_\infty := (\chi, \zeta_\infty \chi) = \infty \text{ almost surely.}$$

Remark: Similarly to [16], there is a gap between Theorems 3.1 and 3.2, since in general the branching number need not be equal to the growth rate. However, this is not very important, because in most of the practical examples these quantities do coincide.

Remark: As mentioned above, the classification parameter for this problem is $\lambda = \inf_{s \in [0,1]} k(s)$. This parameter is not explicitly computable in general since it involves the infinite product of matrices. However for some particular cases this quantity can be computed explicitly as stated in the following proposition.

Proposition 3.3 *Suppose that the measure μ is such that $g_{ij} < 1/d$ almost surely for all $i, j = 1, \dots, d$. Then λ is the largest eigenvalue of the matrix $\mathbb{E}g$.*

Remark: It is interesting to consider the *chaos equation* for the case of matrix-valued random variables and constant branching b :

$$Y \stackrel{\text{law}}{=} \sum_{j=1}^b Y'_j \xi_j, \tag{4}$$

where Y, Y'_j, ξ_j are G -valued random variables, and ξ_j (which are not necessarily independent) are distributed according to μ ; $Y'_j, j = 1, \dots, b$, are i.i.d. and have the same (unknown) law as Y . Analogously to [22] we can get that (at least in the case when ξ_1 satisfies conditions 1,2,3) $\lambda d = 1$ is a necessary condition for the existence of solution of (4). It is an open problem whether this condition is sufficient.

Remark: The condition of independence of the random variables ξ_a can be relaxed; what is important is

1. if ξ_a and ξ_b are not adjacent to the same vertex then they must be independent, and
2. the ξ 's that belong to any path emanating from the root must be independent.




Fig. 1: A typical state of the bindweed model for $\mathcal{S} = \{1, 2, 3\}$, $\sigma = 2313$, and \mathbb{T} the binary tree.

4 The bindweed model

In this section we introduce a model describing an evolution of a random string in random environment on a tree (which is somewhat similar to the model studied in [4]) which we call the bindweed model. Then, we show that its classification from the point of view of positive recurrence can be obtained by using theorems 3.1 and 3.2.

Let $\mathcal{S} = \{1, \dots, d\}$ be a finite alphabet and denote, in accordance with the notations introduced in Section 2, $\mathcal{S}^{n+1} = \{\sigma = \sigma_0 \dots \sigma_n : \sigma_i \in \mathcal{S}\}$ the set of words of length $n+1$ composed from the symbols of the alphabet \mathcal{S} , \mathcal{S}^0 the set containing only the empty word and \mathcal{S}^* the set of words of arbitrary length. Suppose that a branching function B is given on \mathbb{N}^* and denote $\mathbb{V}^n \equiv \mathbb{V}^n(B)$ the corresponding generations of the tree determined by B . Therefore, the rooted tree $\mathbb{T} = (\mathbb{V}, \mathbb{A})$ is uniquely defined.

Now we are going to construct a continuous-time Markov chain with state space \mathfrak{S} , defined by

$$\mathfrak{S} = \{\hat{\theta}\} \cup \bigcup_{n=1} (\mathbb{V}^n \times \mathcal{S}^{n+1}),$$

where $\hat{\theta}$ is a special state to be defined later. In fact, what happens is the following: we place a word $\sigma = \sigma_0 \dots \sigma_n$ on the tree \mathbb{T} in such a way that the 0th symbol of the word is placed on the root $\mathbf{0}$, for any $i = 1, \dots, n$ the i^{th} symbol of the word is placed somewhere in \mathbb{V}^i , and, if the i^{th} symbol σ_i is placed on vertex \mathbf{u} , and σ_{i+1} on \mathbf{v} , then $\mathbf{u} < \mathbf{v}$ and $[\mathbf{u}, \mathbf{v}] \in \mathbb{A}$ (see figure 1). The state $\hat{\theta}$ means that nothing is placed on the tree.

Now, let us define the dynamics of the bindweed model. Suppose that for any $a \in \mathbb{A}$ two collections of positive numbers $(v_{yz}(a), y, z \in \mathcal{S})$, $(\mu_y(a), y \in \mathcal{S})$ are given. If the bindweed model is in the state (\mathbf{u}, σ) , where $\sigma = \sigma_0 \dots \sigma_{n-1}$, $\mathbf{u} \in \mathbb{V}^n$, then

- for $n \geq 0$ it jumps to the state $(\mathbf{v}, \sigma_0 \dots \sigma_{n-1}yz)$ with rate $v_{yz}(a(\mathbf{v}))$, for all $\mathbf{v} \in \mathbb{V} : \mathbf{u} = \hat{\mathbf{v}}$;
- for $n \geq 1$ it jumps to the state $(\hat{\mathbf{u}}, \sigma_0 \dots \sigma_{n-1})$ with rate $\mu_y(a(\mathbf{u}))$.

For any $\sigma_0 \in \mathcal{S}$ the transitions $\hat{\theta} \rightarrow (\mathbf{0}, \sigma_0)$ and $(\mathbf{0}, \sigma_0) \rightarrow \hat{\theta}$ occur with rate 1. Thus, we have defined a continuous-time Markov chain with state space \mathfrak{S} .

Let us describe now how to choose the transition rates. Let ρ be any probability measure on $\mathbb{R}_+^{d^2+d}$. Suppose that for any $a \in \mathbb{A}$ the vector $\Xi(a) = (v_{yz}(a), y, z \in \mathcal{S}, \mu_y(a), y \in \mathcal{S})$ is random, having distribution

ρ , and $(\Xi(a), a \in \mathbb{A})$ are independent and identically distributed. Fix a realization of that collection of random vectors and consider the bindweed model with the transition rates ruled by that realization. So, the model that we constructed is a continuous-time Markov chain in a quenched random environment.

Now, we are interested in obtaining a classification of this Markov chain with respect to positive recurrence. For $(\mathbf{v}, \sigma) \in \mathfrak{G}$ denote by $\pi(\mathbf{v}, \sigma)$ the stationary measure. For any $a \in \mathbb{A}$ let ξ_a be a $d \times d$ matrix whose matrix elements are defined in the following way: $\xi_{a,xy} = v_{xy}(a)/\mu_y(a)$, $x, y \in \mathcal{S}$. It is not difficult to see that we have a reversible Markov chain, so it is clear that $\pi(\hat{\theta}) = \pi(\mathbf{0}, x)$, for all $x \in \mathcal{S}$, and, for any $\mathbf{v} \in \mathbb{V}^n$, $n \geq 1$, and $x, y, \sigma_0, \dots, \sigma_{n-2} \in \mathcal{S}$, we can formally write

$$\begin{aligned} \pi(\mathbf{v}, \sigma_0 \dots \sigma_{n-2}xy) &= \frac{v_{xy}(a(\mathbf{v}))}{\mu_y(a(\mathbf{v}))} \pi(\hat{\mathbf{v}}, \sigma_0 \dots \sigma_{n-2}x) \\ &= \xi_{a(\mathbf{v}),xy} \pi(\hat{\mathbf{v}}, \sigma_0 \dots \sigma_{n-2}x). \end{aligned} \tag{5}$$

Then it is shown in [23] that

$$\sum_{\substack{\mathbf{v} \in \mathbb{V}^n \\ \sigma \in \mathcal{S}^n}} \pi(\mathbf{v}, \sigma) = \pi(\hat{\theta}) \left(\chi, \sum_{\mathbf{v} \in \mathbb{V}^n} \xi[\mathbf{v}]\chi \right)$$

where χ is the vector of order d with all its coordinates equal to 1. Thus $\sum_{(\mathbf{v}, \sigma) \in \mathfrak{G}} \pi(\mathbf{v}, \sigma)$ is finite if and only if Z_∞ is finite. Thus, theorems 3.1 and 3.2 allow us to obtain the classification of the bindweed model in random environment from the point of view of positive recurrence, in the following way:

Proposition 4.1 *Suppose that the distribution of the random matrix ξ_a is such that the Conditions 1, 2, and 3 are satisfied. Let λ be the quantity defined as in Section 3. Then*

- if $\lambda \overline{\text{gr}}(\mathbb{V}) < 1$, then the bindweed model is positive recurrent;
- if $\lambda \text{br}(\mathbb{V}) > 1$, then the bindweed model is not positive recurrent.

5 Open problems and further developments

We demonstrated a close relationship between matrix multiplicative cascades and random walks in random environment on multiplexed trees. In particular it is proved in [23] that both systems are classified by the same parameter. However, the critical region remains out of reach for the moment. Firstly it is not known whether, for sufficiently regular trees so that $\overline{\text{gr}}(\mathbb{V}) = \text{br}(\mathbb{V}) = b$, the walk is null recurrent for $\lambda b = 1$ or some additional condition is needed on μ as is the case for scalar multiplicative chaos [14] and for random walk [16]. Returning to the general tree where $\overline{\text{gr}}(\mathbb{V}) \neq \text{br}(\mathbb{V})$, we obtain a gap in the space of classifying parameters. It is however conjectured in [17] that in general, the set of critical values is of zero Lebesgue measure for string problems. We expect the same phenomenon to occur here. Nevertheless, whether the critical value of the parameter is $\text{br}(\mathbb{V})$ or $\overline{\text{gr}}(\mathbb{V})$ or some intermediate value is unknown for the moment.

An important step toward understanding these problems should be made if conditions for the existence of non-trivial fixed points of the functional equation (4) were obtained. This remains for the moment an open problem although under investigation.

References

- [1] M. Campanino and D. Petritis. On the physical relevance of random walks: an example of random walks on randomly oriented lattices, in "Random walks and geometry", V. Kaimanovich and W. Woess eds., in press, (2003).
- [2] M. Campanino and D. Petritis. Random walks on randomly oriented lattices, preprint 2001, eprint arXiv:math.PR/0111305, to appear in Markov Proc. Rel. Fields (2003).
- [3] P. Collet and F. Koukiou. Large deviations for multiplicative chaos. *Commun. Math. Phys.*, 147:329–342, 1992.
- [4] F. Comets, M. Menshikov, and S. Popov. Lyapunov functions for random walks and strings in random environment. *Ann. Probab.*, 26:1433–1445, 1998.
- [5] F. Comets and S. Yu. Popov. Limit law for transition probabilities and moderate deviations for Sinai's random walk in random environment, To appear in: Probab. Theory Relat. Fields, 2003.
- [6] R. Durrett and Th. Liggett. Fixed points of smoothing transformation. *Z. Wahrscheinlichkeitstheorie verw. Gebiete*, 64:275–301, 1983.
- [7] K. J. Falconer. *The geometry of fractal sets*. Cambridge University Press, Cambridge, 1986.
- [8] Ph. Flajolet and R. Sedgwick. Analytic combinatorics: functional equations, rational and algebraic functions, Research report 4103 INRIA, 2001.
- [9] Yves Guivarc'h and Émile Le Page. Simplicité de spectres de Lyapunov et propriété d'isolation spectrale pour une famille d'opérateurs sur l'espace projectif, to appear in *Random walks and geometry*, V. Kaimanovich and W. Woess, eds.
- [10] J.-P. Kahane and J. Peyrière. Sur certaines martingales de Benoît Mandelbrot. *Adv. Math.*, 22:131–145, 1976.
- [11] H. Kesten, M.V. Kozlov, and F. Spitzer. A limit law for random walk in a random environment. *Compositio Math.*, 30:145–168, 1975.
- [12] Harry Kesten. The limit distribution of Sinai's random walk in random environment. *Phys. A*, 138(1-2):299–309, 1986.
- [13] A. Kumjian, D. Pask, I. Raeburn, and J. Renault. Graphs, groupoids, and Cuntz-Krieger algebras. *J. Funct. Anal.*, 144(2):505–541, 1997.
- [14] Q. Liu. Sur une équation fonctionnelle et ses applications: une extension du théorème de Kesten-Stigun concernant les processus de branchement. *Ann. Appl. Probab.*, 29:353–373, 1997.
- [15] Quansheng Liu and Alain Rouault. Limit theorems for Mandelbrot's multiplicative cascades. *Ann. Appl. Probab.*, 10(1):218–239, 2000.
- [16] R. Lyons and R. Pemantle. Random walk in a random environment and first passage percolation on trees. *Ann. Probab.*, 20:125–136, 1991.

- [17] V. A. Malyshev. Interacting strings of symbols. *Uspekhi Mat. Nauk*, 52(2(314)):59–86, 1997.
- [18] V. A. Malyshev. Random grammars. *Uspekhi Mat. Nauk*, 53(2(320)):107–134, 1998.
- [19] V. A. Malyshev. Stochastic evolution via graph grammars, Research report 3380 INRIA, 2001.
- [20] Benoit Mandelbrot. Multiplications aléatoires itérées et distributions invariantes par moyenne pondérée aléatoire: quelques extensions. *C. R. Acad. Sci. Paris Sér. A*, 278:355–358, 1974.
- [21] R. D. Mauldin, S. Graf, and S. C. Williams. Exact Hausdorff dimension in random recursive constructions. *Proc. Nat. Acad. Sci. U.S.A.*, 84(12):3959–3961, 1987.
- [22] M. Menshikov and D. Petritis. Random walks in random environment on trees and multiplicative chaos, Mathematics and computer science II, B. Chauvin, Ph. Flajolet, D. Gardy, A. Molkadem, eds, pp. 415–422, Birkhäuser, Basel (2002).
- [23] M. Menshikov, D. Petritis, and S. Popov. Matrix multiplicative chaos and bindweeds, preprint 03-05 IRMAR 2003.
- [24] D. Petritis. Directed graphs driven grammars, in preparation 2003.
- [25] D. Revuz. *Markov chains*. North-Holland Publishing Co., Amsterdam, second edition, 1984.
- [26] Ya. G. Sinai. The limit behavior of a one-dimensional random walk in a random environment. *Teor. Veroyatnost. i Primenen.*, 27(2):247–258, 1982.
- [27] F. Solomon. Random walk in a random environment. *Ann. Probab.*, 3:1–31, 1975.

