

On Point-sets that Support Planar Graphs

Vida Dujmović, Will Evans, Sylvain Lazard, William Lenhart, Giuseppe Liotta, David Rappaport, Steve Wismath

► **To cite this version:**

Vida Dujmović, Will Evans, Sylvain Lazard, William Lenhart, Giuseppe Liotta, et al.. On Point-sets that Support Planar Graphs. 19th International Symposium on Graph Drawing, Sep 2011, Eindhoven, Netherlands. 2011. <hal-00643824>

HAL Id: hal-00643824

<https://hal.inria.fr/hal-00643824>

Submitted on 22 Nov 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

On Point-sets that Support Planar Graphs^{*}

V. Dujmović¹, W. Evans², S. Lazard³, W. Lenhart⁴, G. Liotta⁵, D. Rappaport⁶, and S. Wismath⁷

¹ Carleton University, Canada

² University of British Columbia, Canada

³ INRIA Nancy, LORIA, France

⁴ Williams University, U.S.A.

⁵ Università degli Studi di Perugia, Italy

⁶ Queen's University, Canada

⁷ University of Lethbridge, Canada

Abstract. A universal point-set supports a crossing-free drawing of any planar graph. For a planar graph with n vertices, if bends on edges of the drawing are permitted, universal point-sets of size n are known, but only if the bend-points are in arbitrary positions. If the locations of the bend-points must also be specified as part of the point-set, we prove that any planar graph with n vertices can be drawn on a universal set \mathcal{S} of $O(n^2/\log n)$ points with at most one bend per edge and with the vertices and the bend points in \mathcal{S} . If two bends per edge are allowed, we show that $O(n \log n)$ points are sufficient, and if three bends per edge are allowed, $\Theta(n)$ points are sufficient. When no bends on edges are permitted, no universal point-set of size $o(n^2)$ is known for the class of planar graphs. We show that a set of n points in balanced biconvex position supports the class of maximum degree 3 series-parallel lattices.

1 Introduction

A set of points *supports* the drawing of a graph G if there is a one-to-one mapping f of the vertices of G to the points so that for all pairs of edges $(a, b), (c, d)$ in G (where a, b, c, d are distinct), segments $\overline{f(a)f(b)}$ and $\overline{f(c)f(d)}$ do not intersect. A set of points that supports the drawing of all n -vertex graphs in some class is called *universal* for that class, or simply *universal* if the class is all planar graphs. The size of any universal point-set for planar graphs requires at least $1.235n$ points as shown by Kurowski [9] (see also Chrobak and Karloff [2]). Early graph drawing results, such as the canonical ordering technique of de Frasseix, Pach, Pollack [3] and Schnyder's embedding [10] demonstrate that an $n \times n$

^{*} This paper was initiated at the 2011 McGill/INRIA/UVictoria Bellairs workshop. Discussion with other participants is gratefully acknowledged. Research supported by NSERC, and by MIUR of Italy under project AlgoDEEP prot. 2008TFBWL4.

grid of points is a universal point-set. However, no universal point-set of size $o(n^2)$ is known.

Smaller universal point-sets for sub-classes of planar graphs are known. For example, any outerplanar graph can be drawn on *any* set of n points in general position [7]. Indeed, if the point-set is in convex position, then it supports exactly the family of outerplanar graphs. Determining other families of planar graphs for which universal point-sets of size n exist is an interesting problem. We examine a particular type of point-set, of size n , in which points are arranged in biconvex position, and show that it supports the drawing of all maximum degree 3 series-parallel lattices, a class of graphs that contains members that are not outerplanar. These notions are precisely defined in Section 2.

The main contributions in this paper are stated in Theorems 1 and 2, and pertain to universal point-sets for straight-line drawings, and drawings with bends respectively.

Theorem 1. *For all n , there exist universal point-sets of cardinality n that support the family of maximum degree 3 series-parallel lattices with n vertices.*

Suppose we relax the definition of *support* to allow edges of the graph to map to polylines composed of (at most) $k + 1$ line segments. In other words, we allow edges that “bend” at most k times. In this case, universal point-sets of size n exist for two bends [8] and even one bend [6]. However, these results assume that the bend-points can be placed in arbitrary locations and these bend-points are not included as part of the universal point-set. It is natural to ask if there exists a point-set that supports all planar graphs where each vertex *and each bend-point* occurs at a point in the set. As before, we require all pairs of edges (a, b) and (c, d) (where a, b, c, d are distinct) to map to non-intersecting polylines. Previous to this paper, no such point-set of cardinality $o(n^2)$ was known for any value of k . Extending the results of [6] and [8] in a straightforward manner imply point-sets of size $O(n^3)$. For $k = 3, 2, 1$, we present such universal point-sets of cardinality $O(n)$, $O(n \log n)$, and $O(n^2 / \log n)$ respectively.

Theorem 2. *For all n , there exist universal point-sets of cardinality $O(n)$, $O(n \log n)$, and $O(n^2 / \log n)$ that support the drawing of all n -vertex planar graphs with at most 3, 2, or 1 bend per edge, respectively.*

Table 1 summarizes our results in terms of which sets of planar graphs can be supported on point-sets of a given cardinality with a specified number of bends.

Table 1. Summary of results – cardinality of universal point-sets for classes of graphs. The first and last results are well-known. All other results are new.

Graphs	Number of Points	Number of Bends	Reference
outerplanar	n	0	[7]
3SP lattice	n	0	Thm 1
planar	$O(n)$	3	Thm 2
planar	$O(n \log n)$	2	Thm 2
planar	$O(n^2 / \log n)$	1	Thm 2
sub-Hamiltonian	$O(n)$	2	Lemma 3.1
sub-Hamiltonian	$O(n \log n)$	1	Lemma 3.2
planar	$O(n^2)$	0	[3],[10]

2 Preliminaries

We adopt standard notation from the graph drawing literature and we henceforth assume all graphs have n vertices. Many proofs are omitted for space reasons.

Two of our results rely on point-sets that have a specific form; see, for example, Fig. 2. Two non-intersecting non-linear arcs of curves λ_1 and λ_2 are defined to be *biconvex* if: each of these arcs is simple and convex, the convex hull of their 4 endpoints completely contains the two arcs, and the line segment joining any point a of λ_1 to any point b of λ_2 does not intersect either arc of curve except at a and b ; for simplicity, we refer to such arcs of curves as curves. Without loss of generality, we assume the existence of a horizontal line separating the two curves with λ_1 below λ_2 . A point-set all of whose points lie on two curves that are biconvex is in *biconvex position*. We note that point-sets in such a configuration have been used in other contexts under different names.

3 Universal Point-sets for Drawing Planar Graphs with Bends

In this section we establish Theorem 2 by constructing universal point-sets for each of the three cases: 3, 2 or 1 bend per edge allowed. A fundamental tool in our constructions for universal point-sets with bends is the following result proving the existence of a book embedding of planar graphs in which the edges are permitted to cross the spine [5]. A *monotone topological book embedding* of a planar graph G is a planar drawing such that all vertices of G are represented as distinct points on a spine (i.e. the x -axis), and each edge is either represented as an arc in the bottom page (below




Fig. 1. A graph and a monotone topological book embedding of it.

the x -axis), or as an arc in the top page (above the x -axis), or as the concatenation of two arcs: the first (leftmost) in the bottom page and the second in the top page with their common crossing point between spine points. See Fig. 1.

Theorem 3 ([5]). *Every planar graph has a proper monotone topological book embedding which can be computed in linear time.*

3.1 A Set of $\Theta(n)$ Points for Drawing Planar Graphs with Three Bends per Edge

Lemma 1. *There exists a universal set of $10n - 18$ points that supports the drawing of planar graphs with 3 bends per edge.*

Proof. Before introducing the (fixed) universal point-set, we first outline how the graph will be processed. Consider a proper monotone topological book embedding of the input graph (see Fig. 1). For each edge that intersects the spine, introduce a dummy vertex creating an *augmented two page book embedding* with the vertices of the spine drawn on a horizontal line. There are at most $n + m \leq 4n - 6$ vertices on the spine. Imagine a horizontal line slightly above the spine that intersects all arcs in the top page – call these points of intersection from left to right b_1, \dots, b_{2a} where a is the number of arcs. Note that $a \leq 3n - 6$.

Consider a point-set that lies on two curves in biconvex position and consists of $6n - 12$ points on the top curve and $4n - 6$ points on the bottom curve. Any such point-set is universal and supports the drawing of all planar graphs with at most three bends per edge. Refer to Fig. 2 for an example of the construction. For any specific graph, its augmented two page book embedding defines the drawing and requires at most $10n - 18$ points. The at most $4n - 6$ vertices on the spine (including dummy vertices) are assigned, in order, to the first points on the bottom curve.




Fig. 2. Example drawing of the graph of Fig. 1 on a biconvex point-set using three bends per edge. The curvature is exaggerated and only the first 14 of the 36 upper curve points and 11 of the 26 lower curve points in the universal point-set are shown.

The bend-points b_1, \dots, b_{2a} are assigned to the first $2a$ points of the upper curve in left to right order and then each arc in the top page is drawn using the associated bend-points. These polylines do not intersect since the upper curve is convex and any segment joining the two curves does not properly intersect these curves. The arcs in the bottom page can be drawn with no bends – they are cords of the bottom curve. Each arc in the top page uses two bend-points. Substituting a bend-point for each of the dummy vertices results in a drawing with at most three bends per edge. \square

Note that a sub-Hamiltonian planar graph corresponds exactly to a graph that has a two page (unaugmented) book embedding [5]. Since such graphs do not require dummy vertices, they can be drawn with at most two bends per edge.

3.2 A Set of $O(n \log n)$ Points for Drawing Planar Graphs with 2 Bends per Edge

The geometric idea underlying our construction is as follows. Similar to Section 3.1, we draw the spine vertices of an augmented two page book embedding on a set of points that lie on a slightly concave curve close to the x -axis. This implies that all the arcs in the bottom page of the book embedding can be drawn as straight line segments. For arcs in the top page, if the arc is from the i th to the $(i + j)$ th spine vertex, it is drawn to bend at a point at level j . We place approximately n/j bend-points approximately equally spaced in the x -dimension at level j , since only n/j top arcs can have “length” j . The bend-point that lies between the i th and $(i + j)$ th spine vertices is used by this arc. Each level is at a y -coordinate that is large enough that the drawing of an arc that uses a

bend-point at a lower level “nests” inside any drawing of an arc from the same vertex using a higher level bend-point. Of course, for each $j > n/2$, there can be only one arc of “length” j and it uses a single bend-point at level j . The total number of bend-points placed is $O(n \log n)$.

Lemma 2. *There exists a universal set of $O(n \log n)$ points that supports the drawing of planar graphs with 2 bends per edge.*

3.3 A Set of $O(n^2 / \log n)$ Points for Drawing Planar Graphs with 1 Bend per Edge

Lemma 3. *There exists a universal set of $O(n^2 / \log n)$ points that supports the drawing of planar graphs with 1 bend per edge.*

Proof. The construction is similar to that in [6]. We recall briefly this construction, referring to Figures 1 and 3. Given a planar graph G with n




Fig. 3. The one-bend drawing of the three top-page edges adjacent to $v_0 = w_1$ in the graph of Fig. 1 following the construction of [6]. The points p_6 and p_7 are not shown since the figure is to scale.

vertices, we embed the graph on vertices $p_i = (-2^i, i)$ for $i = 0, \dots, n - 1$ with at most one bend per edge, as follows. We first compute a proper monotone topological book embedding, Γ of G . We relabel the vertices of that book embedding from right to left, as v_0, \dots, v_{n-1} . We then map these vertices to p_0, \dots, p_{n-1} , respectively. All the edges below the spine are drawn as straight-line segments. The others are drawn with a bend point as follows. Consider an edge whose rightmost vertex is v_i and that intersects the spine on the interval $(v_u, v_{u+1}]$ (inclusive v_{u+1} for the case where the leftmost endpoint of the edge is v_{u+1}). Such an edge is drawn with a bend point at the same height as v_u , and in the vertical strip delimited by v_i and v_{i+1} . A universal set of points for the bend location can easily be determined in this construction. However, this construction

the right of the line p_0p_1 . Let p_{u_1} be the rightmost point of p_0, \dots, p_{4n-1} that is to the left of q_1 . The second point q_2 is infinitesimally to the right of the line $p_{u_1}p_{u_1+1}$ (at height k), and so on.

The addition of dummy points on the spine ensured that there is at most one edge that intersects the spine between two vertices on the spine. Hence, for any given graph, there is at most one bend point used on any bend line. This essentially implies that the construction of [6] works here.

How many bend points are there on a bend-line? It can be shown that there are $\Theta(k/\log k)$ candidate bend points on the bend line at height k , and $\Theta(n^2/\log n)$ candidate bend points in total. \square

4 Biconvex Point-sets and Series-Parallel Graphs

A *two terminal series-parallel digraph* (also called *TTSP-digraph*) is a planar digraph recursively defined as follows [4, 11]: A directed edge joining two vertices forms a TTSP-digraph. Let G' and G'' be two TTSP-digraphs; the digraph obtained by identifying the sink of G' with the source of G'' (*Series Composition*) is also a TTSP-digraph. Let G' and G'' be two TTSP-digraphs; the digraph obtained by identifying the source of G' with the source of G'' and the sink of G' with the sink of G'' (*Parallel Composition*) is also a TTSP-digraph.

A TTSP-digraph has one source and one sink which are called its *poles*. Also, a TTSP-digraph is always acyclic and admits a planar embedding with the poles on the same face. A TTSP-digraph is a *TTSP lattice* if for every edge (u, v) , there is no directed path from u to v that does not contain (u, v) . Note that a TTSP lattice cannot have multiple edges.

The undirected underlying graph of a TTSP-digraph (resp. lattice) is called a *TTSP-graph* (resp. *TTSP-lattice*). We further shorten these terms and refer to them as series-parallel (SP).

Any point-set in general position supports the class of outerplanar graphs [7]. Indeed a point-set in convex position supports exactly the class of outerplanar graphs, and no other planar graphs. Motivated by this insight we now consider the class of planar graphs that are supported by a point-set in which $n/2$ points are on one convex curve and the remainder are on another convex curve – in biconvex position. Clearly outerplanar graphs can be supported by this point-set and efficient algorithms such as that developed by Bose [1] exist. We show that any $(n/2, n/2)$ biconvex point-set is universal for a subclass of the series-parallel graphs. Since our purpose is to exhibit universal point-sets for classes of planar graphs, the

balancing condition is critical and since the number of vertices could be odd, the balancing must allow for one vertex to be placed arbitrarily.

A planar graph G is *biconvex* if there exists a crossing-free straight-line drawing Γ of G with all vertices located on the curves λ_1 and λ_2 . A planar graph G is *balanced biconvex* if it is biconvex with a drawing Γ in which the numbers of vertices on the two curves differ by at most one; more formally if:

for n even, $n/2$ vertices are on λ_1 and $n/2$ vertices on λ_2 (called *uniform* and denoted as $\Gamma^=$), and for n odd, either:

- $\frac{n-1}{2}$ vertices are on λ_1 and $\frac{n+1}{2}$ vertices are on λ_2 (called *top-heavy* and denoted as Γ^+) or
- $\frac{n+1}{2}$ vertices are on λ_1 and $\frac{n-1}{2}$ vertices are on λ_2 (called *bottom-heavy* and denoted as Γ_+)

Our construction is recursive and attempts to contain the drawing of the SP lattice in a box spanning the biconvex curves with a balanced number of vertices on each curve and with s and t forming a diagonal of the box. Unfortunately, such a strong invariant cannot be maintained and slightly weaker conditions must be carefully considered.

A series-parallel digraph with poles s and t is *bottom-cornered* if it is balanced biconvex with a drawing Γ (n even) or Γ_+ (n odd) such that:

1. there exists a box (i.e. a convex quadrilateral) $B(s, t)$ with s on λ_1 and t on λ_2 , \overline{st} forms one diagonal of B , and the other diagonal has one corner on λ_1 and one on λ_2 , and
2. the entire drawing lies inside B .

Similarly, a series-parallel digraph with poles s and t is *top-cornered* if it is balanced biconvex with a drawing Γ (n even), or Γ^+ (n odd) such that conditions 1 and 2 hold. If a series-parallel graph is both top-cornered and bottom-cornered, it is called *double-cornered* – i.e. if n is odd, there exist two drawings Γ^+ and Γ_+ both of which satisfy conditions 1 and 2.

In some situations, only weaker conditions on the drawings can be maintained, in which one of t or s is contained strictly inside a box rather than on the diagonal forming the box:

- 1'. there exists a box $B(s, x)$ with s on λ_1 and x on λ_2 , \overline{sx} forms one diagonal of $B(s, x)$, and the other diagonal has one corner on λ_1 and one on λ_2 and t is on λ_2 inside $B(s, x)$.
- 1''. there exists a box $B(x, t)$ with x on λ_1 and t on λ_2 , \overline{xt} forms one diagonal of $B(x, t)$, and the other diagonal has one corner on λ_1 and one on λ_2 and s is on λ_1 inside $B(x, t)$.

A series-parallel graph with source s and sink t is *bottom half-cornered* if it is balanced biconvex and conditions 1' and 2 hold; similarly, if conditions 1'' and 2 hold, then the graph is *top half-cornered*.

In the lemmas that follow, we demonstrate only the existence of biconvex drawings for the graphs that we consider. The following lemma indicates that this is sufficient to claim a *universal* biconvex point-set of suitable size.

Lemma 4. *If a graph G on n vertices has a balanced biconvex drawing, then every balanced biconvex point-set of size n supports G .*

A series-parallel graph in which every vertex is of maximum degree 3 is denoted as 3SP. We distinguish between two critical cases. If both the source and sink of a 3SP lattice have degree ≤ 2 then the graph is called *thin* and otherwise (i.e. if either pole has degree 3) it is called *thick*. It is the class of 3SP lattices that we show to be balanced biconvex. There are several cases to consider depending on whether the graph is biconnected or not, and whether the graph is thin or thick. Our proof is recursive in nature – interior components are replaced by appropriate balanced boxes. Lemmas 5 – 14 distinguish and organize these cases and Fig. 5 provides a simple example of each case, the type of drawing obtained, and the prerequisite lemmas used in the proof.

Lemma 5. *A simple path consisting of $n \geq 2$ vertices from s to t is double-cornered.*

First we present the lemmas used for subcases that are thin: biconnected and then not biconnected.

Lemma 6. *Let G be a biconnected thin 3SP lattice. Then G is double-cornered.*

Lemma 7. *Let G be a thin 3SP lattice with source s and sink t . If either $\deg(s) = \deg(t) = 1$ or $\deg(s) = \deg(t) = 2$ then G is double-cornered.*

Lemma 8. *Let G be a thin 3SP lattice with source s and sink t . If $\deg(s) = 1$ and $\deg(t) = 2$ then G is bottom-cornered.*

Lemma 9. *Let G be a thin 3SP lattice with source s and sink t . If $\deg(s) = 2$ and $\deg(t) = 1$ then G is top-cornered.*

The next sequence of lemmas pertains to the cases when the global poles have degree 3. The drawings obtained rely on the previous lemmas and are balanced biconvex, but may not be double-cornered.




Fig. 5. Roadmap of the various cases

Lemma 10. *Let G be a biconnected 3SP lattice with source s and sink t . If $\deg(s) = \deg(t) = 3$ and G consists of 3 series components combined in parallel, then G is balanced biconvex.*

Lemma 11. *Let G be a biconnected thick 3SP lattice with source s and sink t . If $\deg(s) = \deg(t) = 3$ then G is balanced biconvex.*

Lemma 12. *Let G be a biconnected thick 3SP lattice with source s and sink t . If $\deg(s) = 2$ and $\deg(t) = 3$ then G is bottom half-cornered.*

Lemma 13. *Let G be a biconnected thick 3SP lattice with source s and sink t . If $\deg(s) = 3$ and $\deg(t) = 2$ then G is top half-cornered.*

In the final case, G is not biconnected and at least one of the global poles has degree 3.

Lemma 14. *Let G be a thick 3SP lattice with source s and sink t . If $\deg(s) = 3$ or $\deg(t) = 3$ then G is balanced biconvex.*

Biconvex point-sets are the only known point-sets of size n that universally support some class of planar graphs other than the outerplanar graphs.

5 Conclusions and Open Problems

Our main contributions in this paper are stated in Theorems 1 and 2: that any balanced biconvex point-set supports the straight-line drawing of any 3SP lattice, and the specification of universal point-sets for drawings of any planar graph with a small number of bends per edge.

Closing the gap between the upper and lower bounds of the cardinality of a universal point-set for planar graphs with no bends allowed remains an open problem. When k bends per edge are permitted, universal point-sets of smaller asymptotic cardinality may be determined for $k = 1, 2$.

References

1. P. Bose. On embedding an outer-planar graph on a point set. *Computational Geometry: Theory and Applications*, 23:303–312, 2002.
2. M. Chrobak and H. Karloff. A lower bound on the size of universal sets for planar graphs. *SIGACT News*, 20(4):83–86, 1989.
3. H. de Fraysseix, J. Pach, and R. Pollack. How to draw a planar graph on a grid. *Combinatorica*, 10:41–51, 1990.
4. G. Di Battista, P. Eades, R. Tamassia, and I. G. Tollis. *Graph Drawing*. Prentice Hall, NJ, 1999.
5. E. Di Giacomo, W. Didimo, G. Liotta, and S. K. Wismath. Curve-constrained drawings of planar graphs. *Computational Geometry*, 30:1–23, 2005.
6. H. Everett, S. Lazard, G. Liotta, and S. Wismath. Universal sets of n points for one-bend drawings of planar graphs with n vertices. *Discrete and Computational Geometry*, 43(2):272–288, 2010.
7. P. Gritzmann, B. Mohar, J. Pach, and R. Pollack. Embedding a planar triangulation with vertices at specified points. *Amer. Math. Monthly*, 98(2):165–166, 1991.
8. M. Kaufmann and R. Wiese. Embedding vertices at points: Few bends suffice for planar graphs. *Journal of Graph Algorithms and Applications*, 6(1):115–129, 2002.
9. M. Kurowski. A 1.235 lower bound on the number of points needed to draw all n -vertex planar graphs. *Inf. Process. Lett.*, 92(2):95–98, 2004.
10. W. Schnyder. Embedding planar graphs on the grid. In *Proc. 1st ACM-SIAM Sympos. Discrete Algorithms (SODA '90)*, pages 138–148, 1990.
11. J. Valdes, R. E. Tarjan, and E. Lawler. The recognition of series-parallel digraphs. *SIAM J. Comput.*, 11(2):298–313, 1982.