



Tract-based statistical analyzes in dMRI in autism spectrum disorder

Demian Wassermann, Anne-Charlotte Philippe, Pablo Barttfeld, Bruno Wicker, Mariano Sigman, Rachid Deriche

► To cite this version:

Demian Wassermann, Anne-Charlotte Philippe, Pablo Barttfeld, Bruno Wicker, Mariano Sigman, et al.. Tract-based statistical analyzes in dMRI in autism spectrum disorder. 17th Annual Meeting of the Organization for Human Brain Mapping, Jun 2011, Quebec, Canada. <hal-00649766>

HAL Id: hal-00649766

<https://hal.inria.fr/hal-00649766>

Submitted on 9 Dec 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Tract-based statistical analyzes in dMRI in autism spectrum disorder

Submission No:

3854

Introduction:

Abnormal face processing is one of the hallmark features of social impairments in autism spectrum disorder (ASD). Previous neuroimaging studies showed that the fusiform gyrus is involved in face perception and is not or abnormally activated in autistic subjects [1][5]. The aim of this study was to quantify potential anatomical differences in the white matter tracts that traverse the fusiform gyrus, the prefrontal cortex and the superior temporal gyrus, and correlate them with ADOS scores in ASD subjects. We used Diffusion Tensor MRI (DT) images to assess the integrity of automatically segmented white matter bundles connecting these brain areas. Then, we perform statistical analysis on these fiber bundles using diffusivity measures calculated from DT to characterize tissue microstructure changes and correlate these changes with ADOS scores.

Methods:

Participants. 7 adults with high functioning autism or Asperger syndrome and 11 typical adults participated in the study.

Data acquisition and preparation. For each participant, diffusion-weighted images were acquired with $1 \times 1 \times 1.3$ mm voxel size using 80 non-collinear directions at $b=1000$ s/mm² and 1 image with $b=0$ s/mm². We started preparation by generating an unbiased template of the DT images and registering non-linearly all of the images to it. Then, we computed full brain tractography for every subject. Finally, we used the tools developed by Wassermann et al. [2] to cluster fiber bundles and identify the bundles traversing the fusiform gyrus. This gyrus was identified using the gyri parcellation of the JHU atlas. We clustered these bundles across subjects extracting a set of population-obtained bundles.

Data analysis. We applied the statistical analysis of Wassermann et al. [3] for each population-obtained bundle, we calculated its tract-probability map (TPM) and skeletonized this map to obtain a bidimensional representation of each bundle. For each patient, we projected the measures of diffusivity (FA, axial and radial diffusivity) around the tract to their closest points on the skeleton and averaged them with a weight according to the TPM. This produced two populations (one for patients and one for controls) of projected functions on the skeleton. For each tract and each measure, we used a cluster-based permutation hypothesis testing approach [4] to detect dissimilarities between populations. For significant clusters, we calculated correlation between the mean diffusivity measure and ADOS scores.

Results:

We found several clusters with dissimilarities between ASD and control subjects in FA measures on tracts traversing the fusiform gyrus in both hemispheres of the brain (Fig.1). We observed a significant reduction of FA values in a cluster on a bundle joining the superior temporal gyrus to the prefrontal

cortex (Fig.2). In this cluster, the mean FA was strongly correlated with the ADOS scores (correlation coefficient < -0.96 , $p < 0.004$) (Fig.3).

Conclusions:

Results revealing difference between controls and ASD subjects in clusters on fiber bundle traversing the fusiform gyrus are in agreement with current literature suggesting abnormal FFA function in ASD. The localization of the cluster where we found a correlation with ADOS social interaction scores is in agreement with current anatomical literature, the superior temporal gyrus being a classical area of face features recognition and the prefrontal cortex an area know to be involved in social information processing. This is reinforced by the strong inverse correlation (Fig. 3) showing that in this linking cluster a drop of FA value is strongly correlated with a rise of the intensity of ASD. Disruption of white matter tracts between regions implicated in social functioning and face perception may contribute to the social impairments that characterize ASD.




Fig1. One of the studied bundle traversing the fusiform gyrus. In orange significant clusters of difference between autists and controls on this bundle.





Fig2. Studied TPM and significant cluster (red voxels)



Imaging Methods

Diffusion MRI

Abstract Information

References

- [1] K.Pierce (2001), 'Face processing occurs outside the fusiform 'face area' in autism: evidence from functional MRI.' *Brain*, vol. 124, no. 10, pp. 2059-73. [2] Wassermann et al (2010). 'Unsupervised white matter fiber clustering and tract probability map generation: Applications of a Gaussian process framework for white matter fibers', *NeuroImage*, vol. 51, no. 1, pp 228-241. [3] Wassermann et al (2010), 'Diffusion-Based Population Statistics Using Tract Probability Maps', *MICCAI*, vol. 6361, pp. 631-639. [4] Thomas E. Nichols et al (2002), 'Nonparametric permutation tests for functional neuroimaging: A primer with examples', *Human Brain Mapping*, Vol. 15, no. 1, pp 1-25. [5] Robert T. Schultz et al (2005), 'Developmental deficits in social perception in autism: the role of the amygdala and fusiform face area', *International Journal of Developmental Neuroscience*, Vol. 23, no. 2-3, pp 125-141.