

On smooth time functions

Albert Fathi, Antonio Siconolfi

► **To cite this version:**

Albert Fathi, Antonio Siconolfi. On smooth time functions. Mathematical Proceedings, Cambridge University Press (CUP), 2011, 10.1017/S0305004111000661 . hal-00660452

HAL Id: hal-00660452

<https://hal.inria.fr/hal-00660452>

Submitted on 16 Jan 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

On smooth time functions

ALBERT FATHI AND ANTONIO SICONOLFI

Version after Referee's Corrections August 12, 2011

Abstract

We are concerned with the existence of smooth time functions on connected time-oriented Lorentzian manifolds. The problem is tackled in a more general abstract setting, namely in a manifold M where is just defined a field of tangent convex cones $(\mathcal{C}_x)_{x \in M}$ enjoying mild continuity properties. Under some conditions on its integral curves, we will construct a time function.

Our approach is based on the definition of an intrinsic length for curves indicating how a curve is far from being an integral trajectory of \mathcal{C}_x . We find connections with topics pertaining to Hamilton–Jacobi equations, and make use of tools and results issued from weak KAM theory.

1 Introduction

In this paper we are concerned with the existence of smooth time functions on connected time-oriented Lorentzian manifolds, space-times for short, under appropriate assumptions relative to the causal structure.

As a matter of fact our analysis takes place, and so our results are valid, in a more general environment. What we really need as underlying space is a (connected, paracompact, smooth) manifold M on which is defined a multivalued vector field $x \mapsto \mathcal{C}_x \subset T_x M$, with convex cones as values, a *cone structure* for short. Here and throughout the paper *cone* means a closed convex cone, with vertex at 0, not containing any complete affine line and whose interior is not empty.

We will require quite mild continuity properties to hold for \mathcal{C} , as well as some specific conditions on its integral curves, namely the (Lipschitz) curves ξ with $\dot{\xi}(t) \in \mathcal{C}_{\xi(t)}$. Framing the problem in this broader setting allows us to better understand the proof of existence of smooth time functions in Lorentzian geometry.

For a space-time with Lorentzian metric h of signature $(+, \dots, +, -)$, the relevant cone structure is evidently given by the family of future directed causal cones, that is to say the cones made up by timelike and lightlike vectors v , i.e.

with negative and vanishing Lorentzian norm respectively, satisfying, in addition, $h_x(X(x), v) < 0$, for any timelike vector field X providing the time orientation. They represent the physically admissible displacements pointing to the future.

Going back to our abstract setting, we need to introduce some definitions in order to present our results. The terminology is borrowed from Lorentzian geometry and General Relativity, and some explanations on it will be given later on. The cone structure \mathcal{C} is said *continuous* if it possesses such property with respect to the Hausdorff metric, in local coordinates, up to intersection of the cones with the unit closed Euclidean ball. We call \mathcal{C} -causal and \mathcal{C} -future directed the integral curves of \mathcal{C}_x , $\mathring{\mathcal{C}}_x$, respectively, where $\mathring{\mathcal{C}}_x$ is the interior of \mathcal{C}_x . We define an order relation \succ on cone structures by declaring $\mathcal{C}' \succ \mathcal{C}$ whenever $\mathcal{C}_x \setminus \{0\} \subset \mathring{\mathcal{C}}'_x$, for every $x \in M$.

The cone structure \mathcal{C} is said *causal* if it is continuous and there are no closed \mathcal{C} -causal curves; if, in addition, there is a causal cone structure \mathcal{C}' with $\mathcal{C}' \succ \mathcal{C}$, then \mathcal{C} is called *stably causal*. Finally we define \mathcal{C} *globally hyperbolic* if it is stably causal and all causal curves connecting two given points lie in a compact (possibly empty) subset of M . We establish, see Theorem 1.2, that globally hyperbolic is actually a stable property.

A function with the property of being strictly increasing on any nonconstant \mathcal{C} -causal curve is called a *global time function* for \mathcal{C} . If, in addition, it takes any value of its range when the argument varies on an inextendible (with maximal domain of definition) \mathcal{C} -causal curve, then the time function is said to be *Cauchy*. Moreover, we will say that $f : M \rightarrow \mathbb{R}$ is a smooth or C^∞ global time function for the cone structure \mathcal{C} , if its C^∞ , and

$$\forall x \in M, \forall v \in \mathcal{C}_x \setminus \{0\}, d_x f(v) > 0.$$

Our main results are the following:

Theorem 1.1. *If \mathcal{C} is a stably causal cone structure on M , then there exists a C^∞ global time function for \mathcal{C} .*

For the existence of smooth Cauchy time functions, we essentially need the already mentioned stability property for the global hyperbolicity. We believe that this result is indeed of independent interest.

Theorem 1.2. *If the cone structure \mathcal{C} is globally hyperbolic then there exists some $\mathcal{C}' \succ \mathcal{C}$ which is also globally hyperbolic.*

Theorem 1.3. *For any globally hyperbolic cone structure \mathcal{C} on M there exists a C^∞ Cauchy global time function f for \mathcal{C} , i.e. a smooth global time function such that*

$$\lim_{t \rightarrow a} f(\gamma(t)) = -\infty \quad \text{and} \quad \lim_{t \rightarrow b} f(\gamma(t)) = +\infty$$

for any inextendible \mathcal{C} -causal curve $\gamma :]a, b[\rightarrow M$.

In the context of space–times and future directed causal cone structures the time functions are clearly related to the existence of foliations of the space by surfaces of simultaneity, which is a central problem in General Relativity. If f is a global time function then all its levels $f^{-1}(c)$ are achronal in the sense that there cannot be any causal connection between different points of it. If, in addition, the function f is Cauchy time function then any level is accordingly a Cauchy surface, which means that any inextendible causal curve intersects it exactly once. The name originates from the fact that it is the natural region to pose initial conditions for Einstein’s equations.

The causality condition seems quite natural in this setting since the violation of it would apparently yield causality breakdown: one could travel into one’s past. Notice also that this requirements rules out compact space–time since it can be proved, see [2] Proposition 3.10, that compactness and causality are incompatible. The global hyperbolicity ensures that no causal curve joining two points can reach the edge of the space–time, the term is related to some special properties holding for the wave equation in this case.

The paper of Geroch [9] is seminal in the topic. Geroch establishes there the equivalence of global hyperbolicity and existence of a topological Cauchy hypersurface S . The property that the space–time is homeomorphic to $\mathbb{R} \times S$ is, in addition, derived in the so–called Splitting Theorem. A modification of Geroch’s technique allows also to show the stable causality being equivalent to the existence of a continuous global time function.

The possibility of going beyond the topological nature of these results and giving a differential version of them, has remained as an open problem for a long period and, after some fallacious attempts, has been only recently solved by Bernal and Sánchez, see [4]. Starting from Geroch’s Splitting Theorem, these authors first prove the existence of a smooth Cauchy surface and from this they construct a smooth Cauchy time function.

We tackle the issue from a new angle and propose an approach which stays independent of Geroch and Bernal–Sánchez contributions, and use instead some tools and results issued from weak KAM theory, see [8] for a comprehensive treatment of this topic. This connection is, at a first sight, quite surprising since such theory concerns the qualitative analysis of Hamilton–Jacobi equations at the critical value, i.e. at the minimum level for which a.e. subsolutions do exist, while, in the time function problem, no Hamiltonians are involved.

To bridge the gap, we first notice that what really matters in the study of a stationary Hamilton–Jacobi equations of the form $H(x, Du) = a$ is not the whole Hamiltonian but just the a –sublevels of it, namely the sets

$$\{p \mid H(x, p) \leq a\},$$

that we assume in this discussion with non-empty interior. Under the customary conditions of continuity, convexity and coercivity on H , the map associating to any x the corresponding a –sublevel is convex compact valued and continuous

with respect to the Hausdorff metric. These are the same kind of conditions that we have required for the map $x \mapsto \mathcal{C}_x$, and also hold true for $x \mapsto \mathcal{C}_x^*$, where \mathcal{C}_x^* is the polar cone of \mathcal{C}_x , apart the compactness that we recover through suitable truncations.

The property of being a strict subsolution to the equation for a smooth function can be expressed by requiring its differential to be contained in the interior of the a -sublevel for any x . In the same way a global smooth time function f for \mathcal{C} is characterized by the fact that it is solution of the partial differential inclusion $-Df(x) \in \overset{\circ}{\mathcal{C}}_x^*$. Similar ideas have already been used in [12] to construct a Lyapunov function for a multivalued dynamics enjoying suitable stability and attractiveness conditions.

In the Hamilton–Jacobi framework a smooth strict subsolution is constructed, through a partition of unity technique and mollification process, see [7], provided that for any point one can find a neighborhood U of x , $\epsilon > 0$, and an a.e. subsolution u to the equation in the whole space, which, in addition satisfies the strict inequality $H(x, Du(x)) \leq a - \epsilon$ for a.e. $x \in U$. The obstruction to this procedure is then represented by the points around which no a.e. global subsolution is strict. This set is named after Aubry and a crucial step in the analysis is to provide a metric characterization of it. This is made through the introduction of an intrinsic length for curves which again only depends on the a -sublevels of the Hamiltonian, and, more precisely, on their support functions, see [7]. In our case the intrinsic length is defined starting from the cone structure \mathcal{C} , and in a sense it indicates the deviation of a curve from being \mathcal{C} -causal, see (15), (20), so that a nonconstant trajectory possess zero intrinsic length if and only if it is \mathcal{C} -causal

The property detecting the points of the Aubry set is thus given via comparison of this intrinsic length and the natural length induced in the ground space by some reference Riemannian metric. More precisely a point belongs to the Aubry set if and only if there is a sequence of closed curves passing through it with Riemannian length bounded from below by a positive constant and arbitrarily small intrinsic length. The role of closed curves in both of definitions of Aubry set and causal cone structure, and the fact that the intrinsic length is zero whenever γ is causal, cast some light on the way where weak KAM theory enters in our topic.

The crucial point is to show that, given $\mathcal{C}' \succ \mathcal{C}$ the reference metric can be chosen in such a way that if the intrinsic length related to \mathcal{C} of a curve joining two points x and y is small compared to the natural one then x can be also connected to y through a \mathcal{C}' -future directed curve, see Lemma 2.7, Proposition 2.24, Theorem 3.4. If, in addition, \mathcal{C} is stably causal and we pick \mathcal{C}' causal, this forces $x \neq y$, which shows that no cycle can have \mathcal{C} -length small compared to the natural one, and so, under the stable causality assumption, the corresponding Aubry set is empty, which yields Theorem 1.1.

Theorem 1.2 is essentially based on some stability properties holding in the class of \mathcal{C} -causal curves, and is, in turn, crucial for proving Theorem 1.3. Here

we also exploit a density property for smooth global time functions, see Theorem 5.1.

2 Cone Structures

2.1 Cones in \mathbb{R}^N

We consider a finite dimensional Euclidean space \mathbb{R}^N and denote by $|\cdot|$ its Euclidean norm. For every $x \in \mathbb{R}^N$, we use the canonical identification of the tangent $T_x\mathbb{R}^N$ with \mathbb{R}^N . Given v_0, v_1 in \mathbb{R}^N we will denote by $v_0 \cdot v_1$ the usual Euclidean scalar product of v_0 and v_1 . We systematically identify \mathbb{R}^N with its dual $(\mathbb{R}^N)^*$, using the canonical basis on \mathbb{R}^N . Accordingly, if $p \in (\mathbb{R}^N)^*$, $v \in \mathbb{R}^N$, we will also use the scalar product notation, $p \cdot v$, to denote the action $p(v)$ of p on v .

We write \mathbb{B} for the closed unit Euclidean ball of \mathbb{R}^N centered at 0, accordingly $x + r\mathbb{B}$ will denote the closed ball of radius r centered at x , for any $x \in \mathbb{R}^N$, $r > 0$. For every closed non-empty set $K \subset \mathbb{R}^N$, the *distance* $d(x, K)$ of any x from K is defined by

$$d(x, K) = \inf_{y \in K} |x - y|.$$

By the compactness of balls in the Euclidean space, this infimum is achieved. The points in K realizing such infimum are called *projections* of x on K , and the set made up by them is denoted by $\text{proj}_K(x)$. As is well-known, the projection is unique if K is, in addition, convex, since the norm $|\cdot|$ is Euclidean. In fact, in that case, for $v \in \mathbb{R}^N$, the projection of v on K is the unique $v_0 \in K$ such that

$$(v - v_0) \cdot (w - v_0) \leq 0 \quad \text{for any } w \in K. \quad (1)$$

In the case where K is closed and convex, we will denote by $\text{proj}_K(v)$ the unique $v_0 \in K$ such that $d(v, K) = d(v, v_0) = |v - v_0|$. It follows from property (1) that for a closed and convex subset $K \subset \mathbb{R}^N$ the map $\text{proj}_K : \mathbb{R}^N \rightarrow K$ has Lipschitz constant equal to 1, i. e.

$$|\text{proj}_K(v) - \text{proj}_K(v')| \leq |v - v'| \quad \text{for all } v, v' \in \mathbb{R}^N. \quad (2)$$

For an arbitrary non-empty closed set, still denoted by K , such that $\mathbb{R}^N \setminus K \neq \emptyset$ (or equivalently $\partial K \neq \emptyset$), it is convenient to introduce the *signed distance* given by the formula

$$d^\#(\cdot, K) = 2d(\cdot, K) - d(\cdot, \partial K).$$

Proposition 2.1. *If K is a closed non-empty subset of \mathbb{R}^N with $\mathbb{R}^N \setminus K \neq \emptyset$, we have*

$$d^\#(v, K) = \begin{cases} d(v, K) = d(v, \partial K) & \text{if } v \notin \overset{\circ}{K}, \\ -d(v, \partial K) & \text{if } v \in K. \end{cases}$$

In particular, we have $|d^\#(v, K)| = d(v, \partial K)$, for every $v \in \mathbb{R}^N$.

Proof. Given the definition of $d^\#$, it suffices to show that $d(v, K) = d(v, \partial K)$, if $v \notin \overset{\circ}{K}$. We obviously have $d(v, K) \leq d(v, \partial K)$. Moreover, if $v_0 \in K$ is such that $d(v, v_0) = d(v, K)$, then, by connectedness, the segment $[v, v_0]$ must intersect ∂K in a point v_1 . Using $d(v, v_0) = d(v, K)$, we must have $v_1 = v_0$, and $d(v, \partial K) \leq d(v, v_0) = d(v, K)$. \square

Another useful result is:

Proposition 2.2. *If K is a closed convex subset of \mathbb{R}^N with $\mathbb{R}^N \setminus K \neq \emptyset$, then both the distance and the signed distance are convex functions on \mathbb{R}^N*

In fact, it is well-known that $d(\cdot, K)$ is convex when K is convex. For the signed distance this is not as well known, but still not difficult to prove. Anyway, we will use this lemma for closed convex cones with vertex at 0 (called just cones later), and this case follows from Lemma 2.4 below, where, in particular, it is shown that both the distance and the signed distance to such a cone are a supremum of linear functions, hence they are convex.

It is useful to consider positively homogeneous subset of \mathbb{R}^N . A subset $C \subset \mathbb{R}^N$ is called positively homogeneous if

$$\forall v \in C, \forall \lambda \geq 0, \lambda v \in C.$$

For such a positively homogeneous subset C , it is clear that its closure \bar{C} and its boundary ∂C are homogeneous.

The next proposition summarizes the main properties of the distance to a positively homogeneous closed subset or a cone.

Proposition 2.3. *Let C be a closed non-empty positively homogeneous subset of \mathbb{R}^N . If $v \in \mathbb{R}^N$, $\lambda \geq 0$, and $v_0 \in C$ is a projection of v on C , i.e. $d(v, v_0) = d(v, C)$ then we have*

- (i) $d(\lambda v, C) = \lambda d(v, C)$;
- (ii) λv_0 is a projection of λv on C ;
- (iii) $(v - v_0) \cdot v_0 = 0$;
- (iv) $|v_0| \leq |v|$;
- (v) if $v \in \mathbb{B}$ then $d(v, C) = d(v, C \cap \mathbb{B})$;
- (vi) if $v \in \partial \mathbb{B}$, then $d(v, C \cap \partial \mathbb{B}) \leq 2d(v, C \cap \mathbb{B}) = 2d(v, C)$.

Proof. Properties (i) and (ii) are immediate.

To prove (iii), we remark that the function $\varphi : [0, +\infty[\rightarrow |v - tv_0|^2$ has a minimum at $t = 1$. Hence $\varphi'(1) = 0$. But $\varphi'(1) = -2(v - v_0) \cdot v_0$.

To prove (iv), we note that the triangle with vertices at 0, v_0 and v has a right angle at v_0 . Therefore $|v|^2 = |v - v_0|^2 + |v_0|^2$, and $|v_0| \leq |v|$.

To prove (v), using $C \cap \mathbb{B} \subset C$, we first notice that $d(v, C) \leq d(v, C \cap \mathbb{B})$. On the other hand, since $v \in \mathbb{B}$, the projection v_0 has norm ≤ 1 by (iv). Therefore $v_0 \in C \cap \mathbb{B}$. Hence

$$d(v, C) = |v - v_0| \geq d(v, C \cap \mathbb{B}).$$

Consider now $v \in \partial\mathbb{B}$, and v_0 a projection of v on C . If $v_0 = 0$, then $d(v, C) = |v| = 1$. Since the diameter of $\partial\mathbb{B}$ is equal to 2, it follows (vi) holds if $v_0 = 0$. If $v_0 \neq 0$, then we can consider $v_0/|v_0| \in \partial\mathbb{B}$. The triangle with vertices at v, v_0 , and $v_0/|v_0|$ has a right angle at v_0 . Hence

$$\begin{aligned} d(v, C \cap \partial\mathbb{B})^2 &\leq \left| v - \frac{v_0}{|v_0|} \right|^2 \\ &= |v - v_0|^2 + \left| v_0 - \frac{v_0}{|v_0|} \right|^2 \\ &= |v - v_0|^2 + (1 - |v_0|)^2. \end{aligned}$$

Using (iv), and the triangle inequality, we note that $0 \leq |v| - |v_0| \leq |v - v_0|$. Since $|v| = 1$, squaring yields $(1 - |v_0|)^2 \leq |v - v_0|^2$. Combining with the estimate above, we obtain $d(v, C \cap \partial\mathbb{B})^2 \leq 2|v - v_0|^2 = 2d(v, C)^2$. Using (v), this proves (vi). □

Throughout the paper, we will simply say *cone* to mean a closed convex cone not containing any complete affine line, with vertex at 0, and non-empty interior. These cones are usually qualified as *complete*, see [5], in the literature. We will denote the set of such cones by $\text{Cones}(\mathbb{R}^N)$.

Obviously a cone \mathcal{C}^0 is positively homogeneous, since we assume that its vertex is 0. We can therefore apply to it Proposition 2.3. In particular, part (iii) of Proposition 2.3 and property (1) yield

$$(v - \text{proj}_{\mathcal{C}^0}(v)) \cdot w \leq 0 \quad \text{for any } w \in \mathcal{C}^0. \quad (3)$$

For a cone $\mathcal{C}^0 \subset \mathbb{R}^N$, we define its polar by

$$\mathcal{C}^{0*} = \{p \in (\mathbb{R}^N)^* \approx \mathbb{R}^N \mid p \cdot v \leq 0 \text{ for } v \in \mathcal{C}^0\}.$$

The polar \mathcal{C}^{0*} is also a (complete) cone, and is therefore in $\text{Cones}((\mathbb{R}^N)^*) = \text{Cones}(\mathbb{R}^N)$. Note that if $\mathcal{C}^0 \subset \mathcal{C}^1$ then $\mathcal{C}^{1*} \subset \mathcal{C}^{0*}$. Moreover, we have $(\mathcal{C}^0)^{**} = \mathcal{C}^0$ by the Hahn–Banach Theorem.

It is convenient to give a dual characterization of the distance $d(\cdot, \mathcal{C}^0)$ and the signed distance $d^\#(\cdot, \mathcal{C}^0)$ from a cone. Note that the convexity of these functions is an immediate consequence of this characterization.

Lemma 2.4. *Let \mathcal{C}^0 be a cone, then for any $v \in \mathbb{R}^N$*

$$d(v, \mathcal{C}^0) = \max\{p \cdot v \mid p \in \mathcal{C}^{0*} \cap \mathbb{B}\}, \quad (4)$$

$$d^\#(v, \mathcal{C}^0) = \max\{p \cdot v \mid p \in \mathcal{C}^{0*} \cap \partial\mathbb{B}\}. \quad (5)$$

Proof. To simplify notations, we set for $v \in \mathbb{R}^N$

$$\begin{aligned} \sigma(v) &= \max\{p \cdot v \mid p \in \mathcal{C}^{0*} \cap \mathbb{B}\} \\ \hat{\sigma}(v) &= \max\{p \cdot v \mid p \in \mathcal{C}^{0*} \cap \partial\mathbb{B}\}. \end{aligned}$$

Note that $\sigma \geq 0$, since $0 \in (\mathcal{C}^0)^* \cap \mathbb{B}$. Obviously $\hat{\sigma} \leq \sigma$, and both σ and $\hat{\sigma}$ are Lipschitz with Lipschitz constant 1, as supremums of families of Lipschitz functions with Lipschitz constant 1.

Let us start by (4). Pick $v \in \mathbb{R}^N$, if $\sigma(v) = 0$ then $v \in (\mathcal{C}^0)^{**} = \mathcal{C}^0$ and so $d(v, \mathcal{C}^0) = 0$. If instead $\sigma(v) > 0$ then $\sigma(v) = p \cdot v$ with $p \in (\mathcal{C}^0)^* \cap \partial\mathbb{B}$, thus $\sigma(v) = \hat{\sigma}(v)$. Moreover we have

$$\begin{aligned} \sigma(v) &= p \cdot (v - \text{proj}_{\mathcal{C}^0}(v)) + p \cdot \text{proj}_{\mathcal{C}^0}(v) \\ &\leq p \cdot (v - \text{proj}_{\mathcal{C}^0}(v)) \\ &\leq |v - \text{proj}_{\mathcal{C}^0}(v)| \\ &= d(v, \mathcal{C}^0). \end{aligned}$$

To prove the converse inequality we first derive from property (3) above that $(v - \text{proj}_{\mathcal{C}^0}(v)) \cdot w \leq 0$, for any $w \in \mathcal{C}^0$. It follows that $v - \text{proj}_{\mathcal{C}^0}(v) \in \mathcal{C}^{0*}$. Consequently taking into account part (iii) of Proposition 2.3, we obtain

$$\begin{aligned} \sigma(v) &\geq \left(\frac{v - \text{proj}_{\mathcal{C}^0}(v)}{|v - \text{proj}_{\mathcal{C}^0}(v)|} \right) \cdot v \\ &= \left(\frac{v - \text{proj}_{\mathcal{C}^0}(v)}{|v - \text{proj}_{\mathcal{C}^0}(v)|} \right) \cdot (v - \text{proj}_{\mathcal{C}^0}(v)) \\ &= d(v, \mathcal{C}^0). \end{aligned}$$

Note that if $v \notin \mathcal{C}^0 = (\mathcal{C}^0)^{**}$ then $\sigma(v) > 0$, and $\sigma(v) = \hat{\sigma}(v)$, as we have already pointed out in the first part of the proof. Summing up we have not only proved (4) but also that

$$\sigma(v) = \hat{\sigma}(v) = d(v, \mathcal{C}^0) = d^\#(v, \mathcal{C}^0) \quad \text{for } v \notin \mathcal{C}^0.$$

Since all involved function are continuous, the equality also holds on $\partial\mathcal{C}^0$. It follows that (5) is true outside of $\mathring{\mathcal{C}}^0$.

It remains to prove (5), for $v \in \mathring{\mathcal{C}}^0$. To simplify notation set $r_0 = d(v, \partial\mathcal{C}^0)$. Since the closed Euclidean ball centered at v with radius r_0 is contained in \mathcal{C}^0 , we have $p \cdot (v - r_0 w) \leq 0$ for every $p \in (\mathcal{C}^0)^* \cap \partial\mathbb{B}$ and every $w \in \mathbb{B}$. Hence, we

obtain $p \cdot v \leq r_0(p \cdot w)$. If we now take the infimum of all $w \in \mathbb{B}$, taking into account that $|p| = 1$, we obtain $p \cdot v \leq -r_0$, and finally $\hat{\sigma}(v) \leq -r_0 = d^\#(v, \mathcal{C}^0)$.

To get the converse inequality, we pick a projection v_0 of v on $\partial\mathcal{C}^0$. By what we showed above $\hat{\sigma}(v_0) = d^\#(v_0, \mathcal{C}^0) = 0$. Since we know that $\hat{\sigma}$ has Lipschitz constant 1, we obtain

$$-\hat{\sigma}(v) = \hat{\sigma}(v_0) - \hat{\sigma}(v) \leq |v - v_0| = r_0.$$

Hence $\hat{\sigma}(v) \geq -r_0 = d^\#(v, \mathcal{C}^0)$. \square

We now come to the order \prec on cones already given in the Introduction.

Definition 2.5. If $\mathcal{C}^1, \mathcal{C}^2$ are two cones in \mathbb{R}^N , we say that $\mathcal{C}^1 \prec \mathcal{C}^2$ if $\mathcal{C}^1 \setminus \{0\} \subset \overset{\circ}{\mathcal{C}}^2$.

Proposition 2.6. Let $\mathcal{C}^1, \mathcal{C}^2$ be two cones in \mathbb{R}^N . We have $\mathcal{C}^1 \prec \mathcal{C}^2$ if and only if $\mathcal{C}^{2*} \prec \mathcal{C}^{1*}$

Proof. Since $\mathcal{C}^{**} = \mathcal{C}$, we need only to prove the implication from left to right. We know that the compact set $\partial\mathbb{B} \cap \mathcal{C}^1$ is contained in $\overset{\circ}{\mathcal{C}}^2$. therefore we can find $\delta > 0$ such that for every $v \in \partial\mathbb{B} \cap \mathcal{C}^1$ the Euclidean ball $v + \delta\mathbb{B} \subset \overset{\circ}{\mathcal{C}}^2$. If $p \in \mathcal{C}^{2*}$, we then have

$$p \cdot (v + v') \leq 0 \quad \text{for all } v \in \partial\mathbb{B} \cap \mathcal{C}^1, v' \in \delta\mathbb{B}.$$

We fix v and take into account that $\sup\{p \cdot v' \mid v' \in \delta\mathbb{B}\} = \delta|p|$, to obtain

$$p \cdot v + \delta|p| \leq 0 \quad \text{for any } v \in \partial\mathbb{B} \cap \mathcal{C}^1.$$

In particular we get

$$(p + p') \cdot v \leq p \cdot v + |p'| |v| \leq p \cdot v + \delta|p| \leq 0 \quad \text{for any } p' \in \delta|p|\mathbb{B}, v \in \partial\mathbb{B} \cap \mathcal{C}^1.$$

Therefore by homogeneity $p + \delta|p|\mathbb{B} \subset \mathcal{C}^{1*}$, for every $p \in \mathcal{C}^{2*}$, which implies $\mathcal{C}^{2*} \setminus \{0\} \subset \overset{\circ}{\mathcal{C}}^{1*}$ \square

We end this subsection on cones with a Lemma that will be used later. As usual, we denote by $\ell(\gamma)$ the Euclidean length of a curve γ contained in \mathbb{R}^N .

Lemma 2.7. Let $\mathcal{C}^1, \mathcal{C}^2$ be two cones in \mathbb{R}^N with $\mathcal{C}^1 \prec \mathcal{C}^2$. There exists $\delta > 0$ such that for any Lipschitz curve $\gamma : [a, b] \rightarrow \mathbb{R}^N$, with

$$\int_a^b d(\dot{\gamma}(s), \mathcal{C}^1) ds < \delta \ell(\gamma), \tag{6}$$

one has $\gamma(b) - \gamma(a) \in \overset{\circ}{\mathcal{C}}^2$.

Proof. Note that (6) implies that $\ell(\gamma) > 0$, hence γ is not constant. By Proposition 2.6 above the compact subset $S = \partial\mathbb{B} \cap \mathcal{C}^{2*}$ is contained in the interior of \mathcal{C}^{1*} , we can thus find $\epsilon > 0$ such that $p + \epsilon\mathbb{B} \subset \mathcal{C}^{1*}$, for any $p \in S$. This implies, in view of (4)

$$d(v, \mathcal{C}^1) \geq \frac{p + p'}{1 + \epsilon} \cdot v \quad \text{for any } v \in \mathbb{R}^N, p \in S, p' \in \epsilon\mathbb{B}.$$

Multiplying by $1 + \epsilon$, and taking the supremum over $p' \in \epsilon\mathbb{B}$ yields

$$(1 + \epsilon) d(v, \mathcal{C}^1) \geq \epsilon|v| + p \cdot v \quad \text{for any } v \in \mathbb{R}^N, p \in S,$$

and consequently

$$(1 + \epsilon) \int_a^b d(\dot{\gamma}(s), \mathcal{C}^1) ds \geq 2\epsilon \ell(\gamma) + p(\gamma(b) - \gamma(a)).$$

We choose now $\delta > 0$ such that $(1 + \epsilon)\delta < \epsilon$. If (6) holds for such a δ , we therefore get

$$p \cdot (\gamma(b) - \gamma(a)) < [(1 + \epsilon)\delta - \epsilon]\ell(\gamma) < 0, \quad \text{for any } p \in S,$$

which gives $\gamma(b) - \gamma(a) \in \mathcal{C}^{\circ 2}$. □

2.2 Metric structure on the set of cones in \mathbb{R}^N

We will use the Hausdorff metric on compact sets to define a distance in $\text{Cones}(\mathbb{R}^N)$.

Before doing that, we first recall some basic facts about Hausdorff distance. If (X, d) is a metric space, one defines the Hausdorff distance d^H between compact subsets of X by

$$d^H(K_1, K_2) = \max \left\{ \max_{x \in K_1} d(x, K_2), \max_{y \in K_2} d(y, K_1) \right\}. \quad (7)$$

The fact that d^H is a metric follows from the following useful identity

$$d^H(K_1, K_2) = \sup_{x \in X} |d(x, K_1) - d(x, K_2)|. \quad (8)$$

Suppose that $f : X \rightarrow Y$ is a Lipschitz map from (X, d) to the metric space (Y, d) , with Lipschitz constant L . It is not difficult to verify that for any pair K_1, K_2 of closed subsets in X we have

$$d^H(f(K_1), f(K_2)) \leq L d^H(K_1, K_2). \quad (9)$$

We will denote by $\mathcal{K}(X)$ the set of non-empty compact subsets of X . We will always tacitly consider that $\mathcal{K}(X)$ is a metric space with d^H as a metric.

The following results will be useful at some later point.

Lemma 2.8. For $x, x' \in X$, and $K, K' \in \mathcal{K}(X)$, we have

$$|d(x, K) - d(x', K')| \leq d(x, x') + d^H(K, K').$$

Therefore the map $X \times \mathcal{K}(X), (x, K) \mapsto d(x, K)$ is Lipschitz.

Proof. The map $x \mapsto d(x, K)$ has Lipschitz constant 1, therefore

$$\begin{aligned} |d(x, K) - d(x', K')| &\leq |d(x, K) - d(x', K)| + |d(x', K) - d(x', K')| \\ &\leq d(x, x') + d^H(K, K'), \end{aligned}$$

where the last inequality was obtained using also (8) above. \square

We will need to consider Lipschitz functions on the product of two metric spaces (X, d) and (Λ, d) . Note that all the metrics that one **usually** puts on $X \times \Lambda$ are of the form $d((x_1, \lambda_1), (x_2, \lambda_2)) = \|(d(x_1, x_2), d(\lambda_1, \lambda_2))\|$, where $\|\cdot\|$ is a norm on \mathbb{R}^2 . All these metrics define the product topology, and they are all Lipschitz equivalent (like norms on \mathbb{R}^2). Therefore it makes perfect sense to speak of Lipschitz or locally Lipschitz functions on the product of two spaces. We will, for example, assume that the metric on $X \times \Lambda$ is given by

$$d((x_1, \lambda_1), (x_2, \lambda_2)) = \max[d(x_1, x_2), d(\lambda_1, \lambda_2)].$$

Lemma 2.9. Suppose that X, Λ are metric spaces, and that $f : X \times \Lambda \rightarrow \mathbb{R}$ is a continuous function. Let $\mathcal{K}(X)$ denote the set of non-empty compact subsets of X endowed with the Hausdorff distance d^H . Then the two functions $f_+, f_- : \mathcal{K}(X) \times \Lambda \rightarrow \mathbb{R}$ defined by

$$f_+(K, \lambda) = \sup_{x \in K} f(x, \lambda) \text{ and } f_-(K, \lambda) = \inf_{x \in K} f(x, \lambda)$$

are continuous. Moreover, if f is Lipschitz (resp. locally Lipschitz), so are f_+ and f_- .

Proof. We will deal with the case of f_+ . Fix $C_0 \in \mathcal{K}(X)$ and $\lambda_0 \in \Lambda$. We define $\rho : [0, +\infty[\rightarrow [0, +\infty[$ by

$$\rho(\delta) = \sup |f(x, \lambda) - f(y, \lambda_0)|,$$

where the sup is taken over all $x, y \in X$, and $\lambda \in \Lambda$ satisfying $d(x, C_0) \leq \delta, d(y, C_0) \leq \delta, d(x, y) \leq \delta$, and $d(\lambda, \lambda_0) \leq \delta$. Since C_0 is compact, and f is assumed continuous, one can show (for example by contradiction) that $\rho(\delta) \rightarrow 0$, when $\delta \rightarrow 0$. To prove the continuity of f_+ at (C_0, λ_0) it suffices to show that

$$|f_+(C, \lambda) - f_+(C_0, \lambda_0)| \leq \rho[\max(d^H(C, C_0), d(\lambda, \lambda_0))].$$

Let us set $\delta = \max(d^H(C, C_0), d(\lambda, \lambda_0))$. For $x \in C$, we can find $y \in C_0$ such that $d(x, y) \leq \delta$. Since $d(x, C_0) \leq \delta, d(y, C_0) \leq \delta, d(x, y) \leq \delta, d(\lambda, \lambda_0) \leq \delta$, we have

$$|f(x, \lambda) - f(y, \lambda_0)| \leq \rho(\delta).$$

Using $y \in C_0$, this yields

$$f(x, \lambda) \leq f(y, \lambda_0) + \rho(\delta) \leq f_+(C_0, \lambda_0) + \rho(\delta).$$

Taking the sup over all $x \in C$, we get

$$f_+(C, \lambda) - f_+(C_0, \lambda_0) \leq \rho(\delta).$$

Conversely if $x \in C_0$, we can find $y \in C$, with $d(x, y) \leq \delta$. By the same argument as above

$$f(x, \lambda_0) \leq f(y, \lambda) + \rho(\delta) \leq f_+(C, \lambda) + \rho(\delta),$$

and

$$f_+(C_0, \lambda_0) - f_+(C, \lambda) \leq \rho(\delta).$$

It remains to deal with the Lipschitz and locally Lipschitz cases. The Lipschitz case is obtained from the proof above because we can then take $\rho(\delta) = K\delta$, where K is a Lipschitz constant for f .

We will now treat the locally Lipschitz case. Fix $C_0 \in \mathcal{K}(X)$ and $\lambda_0 \in \Lambda$. By definition of locally Lipschitz, for each $x \in X$, we can find $\delta_x > 0$, and $L_x < +\infty$, such that f is Lipschitz on $B(x, 2\delta_x) \times B(\lambda_0, \delta_x)$ with Lipschitz constant $\leq L_x$. Note that f is therefore bounded on $B(x, 2\delta_x) \times B(\lambda_0, \delta_x)$.

By compactness of C_0 , we can find a finite number x_1, \dots, x_n of points in C_0 such that $C_0 \subset \cup_{i=1}^n B(x_i, \delta_{x_i})$. Setting $\delta = \min_{i=1}^n \delta_{x_i} > 0$, and $L_1 = \max_{i=1}^n L_{x_i}$, it can be checked that for $x_1, x_2 \in B_\delta(C) = \{z \in X \mid d(z, C) < \delta\}$, with $d(x_1, x_2) < \delta$, and $\lambda_1, \lambda_2 \in B(\lambda_0, \delta)$, we have

$$|f(x_1, \lambda_1) - f(x_2, \lambda_2)| \leq L \max[d(x_1, x_2), d(\lambda_1, \lambda_2)].$$

Note also that $B_\delta(C_0) \times B(\lambda_0, \delta) \subset \cup_{i=1}^n B(x_i, 3\delta_{x_i}) \times B(\lambda_0, \delta_{x_i})$, and that f is bounded on each element of that finite union. Therefore f is bounded on $B_\delta(C_0) \times B(\lambda_0, \delta)$. If M is such a bound of f , and $L = \max(L_1, 2M/\delta)$, it is easy to check that f is Lipschitz on $B_\delta(C_0) \times B(\lambda_0, \delta)$, with Lipschitz constant L . Since any $C \in \mathcal{K}(X)$ with $d^H(C, C_0) < \delta$ satisfies $C \subset B_\delta(C_0)$. From the Lipschitz case applied to $B_\delta(C_0)$ instead of X , and $B(\lambda_0, \delta)$ instead of Λ , we obtain that f_+ is Lipschitz on $\{C \in \mathcal{K}(X) \mid d^H(C, C_0) < \delta\} \times B(\lambda_0, \delta)$. \square

Lemma 2.10. *Let Λ be a metric space, and C a compact convex subset of some Euclidean space \mathbb{R}^N . Suppose $f : C \times \Lambda \rightarrow \mathbb{R}$ is a continuous function such that, for every $\lambda \in \Lambda$, the function $x \mapsto f(x, \lambda)$ is convex and continuous on C .*

For $\lambda \in \Lambda$, we set

$$C_\lambda = \{x \in C \mid f(x, \lambda) \leq 0\}.$$

Then the set $\Lambda_f = \{\lambda \in \Lambda \mid \inf_{x \in C} f(x, \lambda) < 0\}$ is open and the map $\Lambda_f \rightarrow \mathcal{K}(C), \lambda \mapsto C_\lambda$ is continuous. Moreover, if f is locally Lipschitz, then the map $\lambda \mapsto C_\lambda$ is locally Lipschitz on the open subset Λ_f .

Proof. Define $F : \Lambda \times \Lambda \rightarrow [0, +\infty[$ by

$$F(\lambda_1, \lambda_2) = \sup_{x \in C} |f(x, \lambda_1) - f(x, \lambda_2)|.$$

Note that F is finite everywhere by the compactness of C and the continuity of f . Moreover, the same continuity and compactness implies that for λ_1 fixed, we have

$$\lim_{\lambda_2 \rightarrow \lambda_1} F(\lambda_1, \lambda_2) = 0.$$

Suppose now that $\lambda_0 \in \Lambda_f$. We can find $x_0 \in C$ such that $f(x_0, \lambda_0) < 0$. Therefore by continuity of $\lambda \mapsto f(\lambda, x_0)$ we can find a neighborhood V of λ_0 and $\epsilon > 0$ such that

$$\forall \lambda \in V, f(x_0, \lambda) \leq -\epsilon.$$

Set $K = \text{diam}(C)/\epsilon$. We will first show that

$$\forall \lambda_1, \lambda_2 \in V, \forall x \in C_{\lambda_1}, d(x, C_{\lambda_2}) \leq KF(\lambda_1, \lambda_2). \quad (10)$$

We fix $\lambda_1, \lambda_2 \in V$, and $x \in C_{\lambda_1}$. Since $f(x, \lambda_1) \leq 0$, we have

$$f(x, \lambda_2) \leq f(x, \lambda_2) - f(x, \lambda_1) \leq F(\lambda_1, \lambda_2).$$

If $x \in C_{\lambda_2}$, then $d(x, C_{\lambda_2}) = 0$ and (10) is clear, since F is non-negative. Therefore we can assume $f(x, \lambda_2) > 0$. We set $x_\alpha = (1 - \alpha)x_0 + \alpha x, \alpha \in [0, 1]$. Since $f(x_0, \lambda_2) < 0 < f(x, \lambda_2)$, by continuity of f in the first argument, we can find α_0 such that $f(x_{\alpha_0}, \lambda_2) = 0$. Using the convexity of $f(\cdot, \lambda_2)$, we obtain

$$0 = f(x_{\alpha_0}, \lambda_2) \leq (1 - \alpha_0)f(x_0, \lambda_2) + \alpha_0 f(x, \lambda_2).$$

Since $f(x_0, \lambda_2) \leq -\epsilon, f(x, \lambda_2) \leq F(\lambda_1, \lambda_2)$, and F is non-negative, this implies

$$0 \leq -(1 - \alpha_0)\epsilon + \alpha_0 F(\lambda_1, \lambda_2) \leq -(1 - \alpha_0)\epsilon + F(\lambda_1, \lambda_2).$$

Therefore

$$(1 - \alpha_0) \leq \frac{F(\lambda_1, \lambda_2)}{\epsilon}.$$

Since $f(x_{\alpha_0}, \lambda_2) = 0$, we get

$$\begin{aligned} d(x, C_{\lambda_2}) &\leq |x - x_{\alpha_0}| = (1 - \alpha_0)|x - x_0| \\ &\leq \frac{F(\lambda_1, \lambda_2)}{\epsilon} \text{diam}(C) \\ &= KF(\lambda_1, \lambda_2). \end{aligned}$$

The inequality (10) implies

$$\forall \lambda_1, \lambda_2 \in V, \sup_{x \in C_{\lambda_1}} d(x, C_{\lambda_2}) \leq KF(\lambda_1, \lambda_2).$$

Exchanging the role of λ_1 and λ_2 , we obtain

$$\forall \lambda_1, \lambda_2 \in V, d^H(C_{\lambda_1}, C_{\lambda_2}) \leq KF(\lambda_1, \lambda_2).$$

Together with (2.2), this proves the continuity of the map $\lambda \mapsto C_\lambda$ on the neighborhood V of λ_0 .

To prove the last part of the Lemma, suppose f is locally Lipschitz. We fix $\lambda_0 \in \Lambda$. As in the proof of the previous lemma, we could find $\delta > 0$ and $L < +\infty$ such that f is Lipschitz on $C \times B(\lambda_0, \delta)$ with Lipschitz constant $\leq L$.

$$\forall \lambda_1, \lambda_2 \in B(\lambda_0, \delta), F(\lambda_1, \lambda_2) \leq Ld(\lambda_1, \lambda_2).$$

Hence, by what we have just obtained applied to $B(\lambda_0, \delta)$ instead of Λ , the map $\lambda \mapsto C_\lambda$ is Lipschitz on the neighborhood $B(\lambda_0, \delta)$ of λ_0 . \square

We now proceed to define a distance δ^H on $\text{Cones}(\mathbb{R}^N)$ by

$$\delta^H(\mathcal{C}^1, \mathcal{C}^2) = d^H(\mathcal{C}^1 \cap \mathbb{B}, \mathcal{C}^2 \cap \mathbb{B}),$$

for any pair $\mathcal{C}^1, \mathcal{C}^2 \in \text{Cones}(\mathbb{R}^N)$, see [3].

We will always consider $\text{Cones}(\mathbb{R}^N)$ as a metric space endowed with the metric δ^H . From the next Lemma, δ^H is equivalent to the metric $\hat{\delta}^H$ defined by

$$\hat{\delta}^H(\mathcal{C}^1, \mathcal{C}^2) = d^H(\mathcal{C}^1 \cap \partial\mathbb{B}, \mathcal{C}^2 \cap \partial\mathbb{B}).$$

Lemma 2.11. *For every $\mathcal{C}^1, \mathcal{C}^2 \in \text{Cones}(\mathbb{R}^N)$, we have*

$$\delta^H(\mathcal{C}^1, \mathcal{C}^2) \leq d^H(\mathcal{C}^1 \cap \partial\mathbb{B}, \mathcal{C}^2 \cap \partial\mathbb{B}) \leq 2\delta^H(\mathcal{C}^1, \mathcal{C}^2).$$

Proof. If $v \in \mathcal{C}^1 \cap \mathbb{B}$, and $v \neq 0$, taking into account Proposition 2.3 we get

$$\begin{aligned} d(v, \mathcal{C}^2 \cap \mathbb{B}) &= d(v, \mathcal{C}^2) = |v|d\left(\frac{v}{|v|}, \mathcal{C}^2\right) \\ &\leq d\left(\frac{v}{|v|}, \mathcal{C}^2\right) \leq d\left(\frac{v}{|v|}, \mathcal{C}^2 \cap \partial\mathbb{B}\right) \\ &\leq d^H(\mathcal{C}^1 \cap \partial\mathbb{B}, \mathcal{C}^2 \cap \partial\mathbb{B}), \end{aligned}$$

which yields

$$\max_{v \in \mathcal{C}^1 \cap \mathbb{B}} d(v, \mathcal{C}^2 \cap \mathbb{B}) \leq d^H(\mathcal{C}^1 \cap \partial\mathbb{B}, \mathcal{C}^2 \cap \partial\mathbb{B}).$$

By symmetry, we get the left hand side inequality of the lemma. For the right hand side, we use the item (vi) in the statement of Proposition 2.3 to obtain for $v \in \mathcal{C}^1 \cap \partial\mathbb{B}$

$$d(v, \mathcal{C}^2 \cap \partial\mathbb{B}) \leq 2d(v, \mathcal{C}^2 \cap \mathbb{B}) \leq 2\delta^H(\mathcal{C}^1, \mathcal{C}^2).$$

Again by symmetry, this yields the right hand side inequality of the lemma. \square

From Lemma 2.4, we first derive that the map $\mathcal{C}^0 \mapsto \mathcal{C}^{0*}$ is an isometry with respect to δ^H .

Proposition 2.12. *For every pair $\mathcal{C}^1, \mathcal{C}^2 \in \text{Cones}(\mathbb{R}^N)$, we have $\delta^H(\mathcal{C}^1, \mathcal{C}^2) = \delta^H(\mathcal{C}^{1*}, \mathcal{C}^{2*})$.*

Proof. Taking into account Lemma 2.4, we have by duality

$$d(p, \mathcal{C}^{1*}) = \max\{p \cdot v \mid v \in \mathcal{C}^1 \cap \mathbb{B}\} \quad \text{for any } p,$$

and a similar formula is true for $d(p, \mathcal{C}^{2*})$. Now, bearing in mind (7), assume

$$\delta^H(\mathcal{C}^{1*}, \mathcal{C}^{2*}) = d(p_2, \mathcal{C}^{1*}) = p_2 \cdot v_2 \quad \text{for some } p_2 \in \mathcal{C}^{2*} \cap \mathbb{B}, v_2 \in \mathcal{C}^2 \cap \mathbb{B},$$

then, exploiting again (4)

$$\delta^H(\mathcal{C}^{1*}, \mathcal{C}^{2*}) = p_2 \cdot v_2 \leq d(v_2, \mathcal{C}^1) = d(v_2, \mathcal{C}^1 \cap \mathbb{B}) \leq \delta^H(\mathcal{C}^1, \mathcal{C}^2).$$

By interchanging the role of the cones appearing in the statement, we can complete the proof through the above argument. □

We also obtain from Lemma 2.4, the Lipschitz character of the distance and signed distance functions from a cone:

Proposition 2.13. *Let $\mathcal{C}^1, \mathcal{C}^2 \in \text{Cones}(\mathbb{R}^N)$, $v \in \mathbb{R}^N$, then we have*

$$\begin{aligned} |d(v, \mathcal{C}^1) - d(v, \mathcal{C}^2)| &\leq |v| \delta^H(\mathcal{C}^1, \mathcal{C}^2) \\ |d^\#(v, \mathcal{C}^1) - d^\#(v, \mathcal{C}^2)| &\leq 2|v| \delta^H(\mathcal{C}^1, \mathcal{C}^2). \end{aligned}$$

It follows that both functions $(v, \mathcal{C}) \mapsto d(v, \mathcal{C})$ and $(v, \mathcal{C}) \mapsto d^\#(v, \mathcal{C})$ are Lipschitz on $R\mathbb{B} \times \text{Cones}(\mathbb{R}^N)$ for any $R \geq 0$.

Proof. It suffices to show that

$$\begin{aligned} d(v, \mathcal{C}^1) &\leq d(v, \mathcal{C}^2) + |v| \delta^H(\mathcal{C}^1, \mathcal{C}^2), \\ d^\#(v, \mathcal{C}^1) &\leq d^\#(v, \mathcal{C}^2) + 2|v| \delta^H(\mathcal{C}^1, \mathcal{C}^2). \end{aligned}$$

By (4), we have

$$d(v, \mathcal{C}^1) = p_1 \cdot v \quad \text{for some } p_1 \in \mathcal{C}^{1*} \cap \mathbb{B}.$$

We can find $p_2 \in \mathcal{C}^{2*} \cap \mathbb{B}$, with $|p_1 - p_2| = d(p_1, \mathcal{C}^{2*} \cap \mathbb{B})$. Since $d(p_1, \mathcal{C}^{2*} \cap \mathbb{B}) \leq \delta^H(\mathcal{C}^{1*}, \mathcal{C}^{2*}) = \delta^H(\mathcal{C}^1, \mathcal{C}^2)$, we obtain

$$\begin{aligned} d(v, \mathcal{C}^1) &= p_1 \cdot v \leq p_2 \cdot v + |p_2 - p_1| |v| \\ &\leq d(v, \mathcal{C}^2) + |v| \delta^H(\mathcal{C}^1, \mathcal{C}^2). \end{aligned}$$

This proves the part of the assertion above relative to the distances $d(\cdot, \mathcal{C}^i)$, $i = 1, 2$. By (5), we have

$$d^\#(v, \mathcal{C}^i) = \max\{p \cdot v \mid p \in \mathcal{C}^i \cap \partial\mathbb{B}\} \quad i = 1, 2$$

We adapt the same argument as above, and use Proposition 2.4 and Proposition 2.12. We have

$$d^\#(v, \mathcal{C}^1) = \tilde{p}_1 \cdot v, \text{ for some } \tilde{p}_1 \in \mathcal{C}^{1*} \cap \partial\mathbb{B}.$$

We denote by \tilde{p}_2 an element of $\mathcal{C}^{2*} \cap \partial\mathbb{B}$ with $|\tilde{p}_1 - \tilde{p}_2| = d(\tilde{p}_1, \mathcal{C}^{2*} \cap \partial\mathbb{B})$. We get

$$\begin{aligned} d^\#(v, \mathcal{C}^1) &= \tilde{p}_1 \cdot v \leq \tilde{p}_2 \cdot v + |\tilde{p}_2 - \tilde{p}_1| |v| \\ &\leq d^\#(v, \mathcal{C}^2) + |v| d(\tilde{p}_1, \mathcal{C}^{2*} \cap \partial\mathbb{B}). \end{aligned}$$

and

$$\begin{aligned} d^\#(v, \mathcal{C}^1) &\leq d^\#(v, \mathcal{C}^2) + |v| \max_{\tilde{p}_1 \in \mathcal{C}^{1*} \cap \partial\mathbb{B}} d(\tilde{p}_1, \mathcal{C}^{2*} \cap \partial\mathbb{B}) \\ &\leq d^\#(v, \mathcal{C}^2) + 2|v| \max_{\tilde{p}_1 \in \mathcal{C}^{1*} \cap \partial\mathbb{B}} d(\tilde{p}_1, \mathcal{C}^{2*} \cap \mathbb{B}) \\ &\leq d^\#(v, \mathcal{C}^2) + 2|v| \delta^H(\mathcal{C}^1, \mathcal{C}^2). \end{aligned}$$

This proves the part of the assertion above relative the distances $d^\#(\cdot, \mathcal{C}^i)$. $i = 1, 2$.

Since any distance function to a set have Lipschitz constant equal to 1, we obtain for all v_1, v_2 in $R\mathbb{B}$

$$\begin{aligned} |d(v_1, \mathcal{C}^1) - d(v_2, \mathcal{C}^2)| &\leq |d(v_1, \mathcal{C}^1) - d(v_2, \mathcal{C}^1)| + |d(v_2, \mathcal{C}^1) - d(v_2, \mathcal{C}^2)| \\ &\leq |v_1 - v_2| + R \delta^H(\mathcal{C}^1, \mathcal{C}^2). \end{aligned}$$

In the same way, since the signed distance function have Lipschitz constant equal to 1, we obtain for all v_1, v_2 in $R\mathbb{B}$

$$|d^\#(v_1, \mathcal{C}^1) - d^\#(v_2, \mathcal{C}^2)| \leq |v_1 - v_2| + 2R \delta^H(\mathcal{C}^1, \mathcal{C}^2).$$

□

We deduce:

Corollary 2.14. *The map $\mathcal{C} \mapsto \partial\mathcal{C} \cap \partial\mathbb{B}$ from $\text{Cones}(\mathbb{R}^N)$ to the space of compact subsets of \mathbb{R}^N endowed with the Hausdorff metric is Lipschitz with Lipschitz constant ≤ 2 .*

Proof. Suppose C is a closed subset of \mathbb{R}^N and $v \in \mathbb{R}^N$. By Proposition 2.1 we have $|d^\#(v, C)| = d(v, \partial C)$. In particular $v \in \partial C$ if and only if $d^\#(v, C) = 0$.

Consider now a pair of cones $\mathcal{C}^1, \mathcal{C}^2$, if $v \in \partial\mathcal{C}^1 \cap \partial\mathbb{B}$, then using Proposition 2.3 (vi) and what we said above, we obtain

$$\begin{aligned} d(v, \partial\mathcal{C}^2 \cap \partial\mathbb{B}) &\leq 2d(v, \partial\mathcal{C}^2) \\ &= 2|d^\#(v, \mathcal{C}^2)| \\ &= 2|d^\#(v, \mathcal{C}^1) - d^\#(v, \mathcal{C}^2)| \\ &\leq 2\delta^H(\mathcal{C}^1, \mathcal{C}^2), \end{aligned}$$

where the last inequality used Proposition 2.13. Therefore

$$\sup_{v \in \partial\mathcal{C}^1 \cap \partial\mathbb{B}} d(v, \partial\mathcal{C}^2 \cap \partial\mathbb{B}) \leq 2\delta^H(\mathcal{C}^1, \mathcal{C}^2).$$

By symmetry, we obtain $d^H(\partial\mathcal{C}^1 \cap \partial\mathbb{B}, \partial\mathcal{C}^2 \cap \partial\mathbb{B}) \leq 2\delta^H(\mathcal{C}^1, \mathcal{C}^2)$. \square

We proceed proving a compatibility result between the order relation \succ and the metric δ^H in $\text{Cones}(\mathbb{R}^N)$.

Theorem 2.15. *Given $\mathcal{C} \in \text{Cones}(\mathbb{R}^N)$ and $\varepsilon > 0$, there are cones $\mathcal{C}', \mathcal{C}''$ satisfying $\delta^H(\mathcal{C}, \mathcal{C}') < \varepsilon$, $\delta^H(\mathcal{C}, \mathcal{C}'') < \varepsilon$ and $\mathcal{C}' \prec \mathcal{C} \prec \mathcal{C}''$.*

We first give some preliminary notions and results.

Given $\mathcal{C} \in \text{Cones}(\mathbb{R}^N)$, we define the *inradius* $I(\mathcal{C})$ by

$$I(\mathcal{C}) = \max\{d(v, \partial\mathcal{C}) \mid v \in \mathcal{C} \cap \partial\mathbb{B}\} > 0. \quad (11)$$

Lemma 2.16. *The function $I : \text{Cones}(\mathbb{R}^N) \rightarrow]0, +\infty[$ defined above is Lipschitz.*

Proof. On $\mathbb{B} \times \text{Cones}(\mathbb{R}^N)$, the function $(v, \mathcal{C}) \mapsto d^\#(v, \mathcal{C}) = d^\#(v, \mathbb{B} \cap \mathcal{C})$ is Lipschitz by Proposition 2.13. Therefore by Lemma 2.9, the map $\mathcal{K}(\mathbb{B}) \times \text{Cones}(\mathbb{R}^N) \rightarrow \mathbb{R}$, $(K, \mathcal{C}) \mapsto \min_{v \in K} d^\#(v, \mathbb{B} \cap \mathcal{C})$ is Lipschitz. To finish the proof of the lemma it suffices to observe that the map $\text{Cones}(\mathbb{R}^N) \rightarrow \mathcal{K}(\mathbb{B})$, $\mathcal{C} \mapsto \mathbb{B} \cap \mathcal{C}$ is Lipschitz by Lemma 2.11. \square

For $\delta \geq 0$, we set

$$\mathcal{C}^{-\delta} = \{v \mid d^\#(v, \mathcal{C}) + \delta|v| \leq 0\}$$

Lemma 2.17. *Given $\mathcal{C} \in \text{Cones}(\mathbb{R}^N)$. We have:*

- (i) $\mathcal{C}^{-0} = \mathcal{C}$.
- (ii) *The subset $\mathcal{C}^{-\delta}$ of \mathbb{R}^N is a cone for any $\delta \in [0, I(\mathcal{C})[$.*
- (ii) *For $\delta_2 < \delta_1$, we have $\mathcal{C}^{-\delta_1} \prec \mathcal{C}^{-\delta_2}$.*

Moreover, the map $(\delta, \mathcal{C}) \rightarrow \mathcal{C}^{-\delta} \in \text{Cones}(\mathbb{R}^N)$ is locally Lipschitz on the open subset $\{(\delta, \mathcal{C}) \mid 0 \leq \delta < I(\mathcal{C})\} \subset [0, +\infty[\times \text{Cones}(\mathbb{R}^N)$.

Proof. The fact that $\mathcal{C}^{-0} = \mathcal{C}$ is obvious. The set $\mathcal{C}^{-\delta}$, being contained in \mathcal{C} , does not contain any affine complete line, moreover its interior is non-empty, since any $v \in \mathcal{C} \cap \partial\mathbb{B}$ whose signed distance from \mathcal{C} realizes $I(\mathcal{C})$, up to a sign, belongs to it. Finally, $\mathcal{C}^{-\delta}$ is a closed convex cone with vertex at 0 since it is the 0-sublevel of a continuous convex positively homogeneous function. The relation $\mathcal{C}^{-\delta_2} \prec \mathcal{C}^{-\delta_1}$, for $\delta_2 < \delta_1$, is immediate, since for $v \in \mathcal{C}^{-\delta_2} \setminus \{0\}$, we have

$$d^\#(v, \mathcal{C}) + \delta_1|v| = d^\#(v, \mathcal{C}) + \delta_2|v| + (d_2 - d_1)|v| \leq (d_2 - d_1)|v| < 0.$$

It remains to show that the map $(\delta, \mathcal{C}) \rightarrow \mathcal{C}^{-\delta}$ is locally Lipschitz on the open subset $\{(\delta, \mathcal{C}) \mid 0 \leq \delta < I(\mathcal{C})\}$ of $[0, +\infty[\times \text{Cones}(\mathbb{R}^N)$. For this we remark that the map $f : \mathbb{B} \times ([0, +\infty[\times \text{Cones}(\mathbb{R}^N)) \rightarrow \mathbb{R}, v \mapsto d^\#(v, \mathcal{C}) + \delta|v|$ is Lipschitz, and convex in v . Since we have $\inf_{v \in \mathbb{B}} f(v, (\delta, \mathcal{C})) < 0$, for $0 \leq \delta < I(\mathcal{C})$, it follows from Lemma 2.10 that $\mathcal{C}^{-\delta} \cap \mathbb{B} = \{v \in \mathbb{B} \mid f(v, (\delta, \mathcal{C})) \leq \delta\}$ is a locally Lipschitz function of (δ, \mathcal{C}) on the set $0 \leq \delta < I(\mathcal{C})$. \square

Proof of Theorem 2.15. It is enough to show that there is \mathcal{C}' with $\delta^H(\mathcal{C}, \mathcal{C}') < \varepsilon$ and $\mathcal{C}' \prec \mathcal{C}$. The existence of \mathcal{C}'' with the properties asserted in the statement then comes by duality taking into account Proposition 2.6 and the fact that $\mathcal{C}^0 \mapsto \mathcal{C}^{0*}$ is an isometry by Proposition 2.12.

The existence of \mathcal{C}' now follows from Lemma 2.17, since for $\delta \in]0, I(\mathcal{C})[$, we have $\mathcal{C}^{-\delta} \prec \mathcal{C}$, and $\mathcal{C}^{-\delta} \rightarrow \mathcal{C}$, as $\delta \rightarrow 0$. \square

For $\mathcal{C}, \mathcal{C}' \in \text{Cones}(\mathbb{R}^N)$, we define

$$D^\#(\mathcal{C}, \mathcal{C}') = \sup_{x \in \mathcal{C} \cap \partial\mathbb{B}} d^\#(x, \mathcal{C}').$$

Note that $D^\#(\mathcal{C}, \mathcal{C}') \leq 0$ if and only if $\mathcal{C} \subset \mathcal{C}'$, and $D^\#(\mathcal{C}, \mathcal{C}') < 0$ if and only if $\mathcal{C} \prec \mathcal{C}'$.

Lemma 2.18. *The function $D^\# : \text{Cones}(\mathbb{R}^N) \times \text{Cones}(\mathbb{R}^N), (\mathcal{C}, \mathcal{C}') \mapsto D^\#(\mathcal{C}, \mathcal{C}')$ is Lipschitz. In particular, the subset $\{(\mathcal{C}, \mathcal{C}') \mid \mathcal{C} \prec \mathcal{C}'\}$ is open in $\text{Cones}(\mathbb{R}^N) \times \text{Cones}(\mathbb{R}^N)$*

Proof. It is very similar to the proof of Lemma 2.16. In the proof of that Lemma we obtained that the map $\mathcal{K}(\mathbb{B}) \times \text{Cones}(\mathbb{R}^N) \rightarrow \mathbb{R}, (K, \mathcal{C}') \mapsto \min_{v \in K} d^\#(v, \mathbb{B} \cap \mathcal{C}')$ is Lipschitz. Again to finish the proof of the first part of the lemma it suffices to observe that the map $\text{Cones}(\mathbb{R}^N) \rightarrow \mathcal{K}(\mathbb{B}), \mathcal{C} \mapsto \mathbb{B} \cap \mathcal{C}$ is Lipschitz by Lemma 2.11.

The last part of the lemma is now clear since $\{(\mathcal{C}, \mathcal{C}') \mid \mathcal{C} \prec \mathcal{C}'\} = \{(\mathcal{C}, \mathcal{C}') \mid D^\#(\mathcal{C}, \mathcal{C}') < 0\}$ \square

We now get the continuity of the action of the linear group on cones. We use, as usual, the symbol $\text{GL}(\mathbb{R}^N)$ to denote the group of linear invertible maps from

\mathbb{R}^N to itself. We endow it with the operator norm, denoted, like the Euclidean norm in \mathbb{R}^N , by $|\cdot|$ and defined, for any $T \in \text{GL}(\mathbb{R}^N)$ as

$$|T| = \max\{|Tx| \mid x \in \mathbb{B}\}.$$

It is clear that $T\mathcal{C} \in \text{Cones}(\mathbb{R}^N)$ for any $\mathcal{C} \in \text{Cones}(\mathbb{R}^N)$, and any $T \in \text{GL}(\mathbb{R}^N)$.

Lemma 2.19. *The map $\text{GL}(\mathbb{R}^N) \times \text{Cones}(\mathbb{R}^N) \rightarrow \text{Cones}(\mathbb{R}^N)$, $(T, \mathcal{C}) \mapsto T\mathcal{C}$, is locally Lipschitz. More precisely, if we set for $K \geq 1$*

$$G_K = \{T \in \text{GL}(\mathbb{R}^N) \mid |T| \leq K, \text{ and } |T^{-1}| \leq K\},$$

then we have for all $T, T' \in G_K$, $\mathcal{C}, \mathcal{C}' \in \text{Cones}(\mathbb{R}^N)$

$$\delta^H(T\mathcal{C}, T'\mathcal{C}') \leq K|T - T'| + K^2\delta^H(\mathcal{C}, \mathcal{C}'). \quad (12)$$

Proof. We consider $T, T' \in G_K$, $\mathcal{C}, \mathcal{C}' \in \text{Cones}(\mathbb{R}^N)$, and fix $x \in \mathbb{B}$. The idea is to exploit the formula (8), therefore, since x has been arbitrarily taken in \mathbb{B} , instead of (12) we will equivalently show

$$|d(x, T\mathcal{C} \cap \mathbb{B}) - d(x, T'\mathcal{C}' \cap \mathbb{B})| \leq K|T - T'| + K^2\delta^H(\mathcal{C}, \mathcal{C}'). \quad (13)$$

There exists $y' \in \mathcal{C}'$ with $|T'y'| \leq 1$ such that $d(x, T'\mathcal{C}' \cap \mathbb{B}) = |x - T'y'|$. The fact that $T' \in G_K$ implies that $|y'| \leq K$. We have

$$|x - Ty'| \leq |x - T'y'| + |T' - T||y'| \leq d(x, T'\mathcal{C}' \cap \mathbb{B}) + K|T' - T|.$$

Since $y' \in \mathcal{C}'$ and $|y'| \leq K$, we can find $y \in \mathcal{C}$ such that $|y| \leq K$ and $|y - y'| \leq K\delta^H(\mathcal{C}, \mathcal{C}')$. Therefore we obtain

$$\begin{aligned} |x - Ty| &\leq |x - Ty'| + |Ty' - Ty| \\ &\leq d(x, T'\mathcal{C}' \cap \mathbb{B}) + K|T' - T| + |T||y - y'| \\ &\leq d(x, T'\mathcal{C}' \cap \mathbb{B}) + K|T' - T| + K^2\delta^H(\mathcal{C}, \mathcal{C}'). \end{aligned}$$

Since $|x| \leq 1$, by Proposition 2.3, we have

$$d(x, T\mathcal{C} \cap \mathbb{B}) = d(x, T\mathcal{C}) \leq |x - Ty|,$$

and consequently

$$d(x, T\mathcal{C} \cap \mathbb{B}) - d(x, T'\mathcal{C}' \cap \mathbb{B}) \leq K|T' - T| + K^2\delta^H(\mathcal{C}, \mathcal{C}').$$

By exchanging the role of \mathcal{C} and \mathcal{C}' , we in the end obtain (13). \square

2.3 Cone Structures on open subsets of \mathbb{R}^N

Definition 2.20. A cone structure (resp. Lipschitz cone structure) on an open subset U of \mathbb{R}^N is a family of cones $\mathcal{C} = (\mathcal{C}_x)_{x \in U}$ with $\mathcal{C}_x \in \text{Cones}(\mathbb{R}^N)$, and the map $U \rightarrow \text{Cones}(\mathbb{R}^N), x \mapsto \mathcal{C}_x$ continuous (resp. locally Lipschitz), where, as we said above, $\text{Cones}(\mathbb{R}^N)$ is endowed with the distance δ^H and the topology inherited from that distance.

If $\varphi : U \rightarrow V$ is a C^1 diffeomorphism between two open subsets of \mathbb{R}^N , and $\mathcal{C} = (\mathcal{C}_x)_{x \in U}$ is a cone structure on U , we define the family of cones $\varphi_*\mathcal{C}$ on V by

$$\varphi_*\mathcal{C}_y = D_x\varphi(\mathcal{C}_x), \text{ with } \varphi(x) = y.$$

The following proposition is now clearly a consequence of Lemma 2.19

Lemma 2.21. *The family of cones $\varphi_*\mathcal{C}$ is a (continuous) cone structure on V . Moreover, if $\mathcal{C} = (\mathcal{C}_x)_{x \in U}$ is a Lipschitz cone structure, so is $\varphi_*\mathcal{C}$.*

By Lemma 2.12, if \mathcal{C} is a cone structure on $U \subset \mathbb{R}^N$, the dual family \mathcal{C}^* defined by $\mathcal{C}_x^* = (\mathcal{C}_x)_{x \in U}^*$ is also a cone structure on U . Note that for a Lipschitz cone structure \mathcal{C} the dual \mathcal{C}^* is also Lipschitz by Proposition 2.12.

We recall the notions of \mathcal{C} -causal and \mathcal{C} -future directed curves given in the Introduction.

Definition 2.22. If \mathcal{C} is a cone structure on the open subset U of \mathbb{R}^N , a Lipschitz curve $\xi : [a, b] \rightarrow U$ is \mathcal{C} -causal (resp. \mathcal{C} -future directed) if $\dot{\xi}(t) \in \mathcal{C}_{\xi(t)}$ (resp. $\dot{\xi}(t) \in \mathring{\mathcal{C}}_{\xi(t)}$) for almost every $t \in [a, b]$.

We recall the a Lipschitz curve has a derivative almost everywhere, so the notions above make sense.

We now define the function $d_{\mathcal{C}} : U \times \mathbb{R}^N \rightarrow [0, +\infty[$ by

$$d_{\mathcal{C}}(x, v) = d(v, \mathcal{C}_x). \quad (14)$$

Since the function $d_{\mathcal{C}}$ is non negative and positively homogeneous, we can use it to define an intrinsic length of a curve. Namely, given a curve $\xi : [a, b] \rightarrow \mathbb{R}^N$, we set

$$\int_{\xi} d_{\mathcal{C}} = \int_a^b d_{\mathcal{C}}(\xi(t), \dot{\xi}(t)) dt = \int_a^b d(\dot{\xi}(t), \mathcal{C}_{\xi(t)}) dt. \quad (15)$$

Notice that the above integral is invariant for orientation-preserving change of parameter, as an intrinsic length should be. Obviously a curve ξ is \mathcal{C} -causal if and only if $\int_{\xi} d_{\mathcal{C}} = 0$.

We extend the relations \prec to cone structures in the obvious way. Namely if \mathcal{C} and \mathcal{C}' are cone structures on U we say that $\mathcal{C} \prec \mathcal{C}'$ if we have $\mathcal{C}_x \prec \mathcal{C}'_x$, for every $x \in U$.

Lemma 2.23. *Let $\mathcal{C}, \mathcal{C}'$ be two cone structures on the open subset $U \subset \mathbb{R}^N$ with $\mathcal{C} \prec \mathcal{C}'$. Given $x_0 \in U$, we can find a neighborhood V of x_0 in U and $\mathcal{C}_1, \mathcal{C}_2 \in \text{Cones}(\mathbb{R}^N)$ such that*

$$\mathcal{C}_x \prec \mathcal{C}_1 \prec \mathcal{C}_2 \prec \mathcal{C}'_x, \text{ for every } x \in V.$$

Proof. We will first find \mathcal{C}_1 and \mathcal{C}_2 such that

$$\mathcal{C}_{x_0} \prec \mathcal{C}_1 \prec \mathcal{C}_2 \prec \mathcal{C}'_{x_0}.$$

According Lemma 2.17 we have $(\mathcal{C}'_{x_0})^{-\delta} \prec \mathcal{C}_{x_0}$, for $\delta > 0$, and $(\mathcal{C}'_{x_0})^{-\delta} \rightarrow \mathcal{C}'_{x_0}$, for $\delta \rightarrow 0$. It therefore follows from Lemma 2.18 that $(\mathcal{C}'_{x_0})^{-\delta} \succ \mathcal{C}_{x_0}$, for δ small enough. Therefore we can take $\mathcal{C}_2 = (\mathcal{C}'_{x_0})^{-\delta}$ for some appropriate $\delta > 0$. To obtain \mathcal{C}_1 , it suffices to apply what we just did with $\mathcal{C}_{x_0}, \mathcal{C}_2$ instead of $\mathcal{C}_{x_0}, \mathcal{C}'_{x_0}$.

By the last part of Lemma 2.18 and the continuity of $x \mapsto \mathcal{C}_x, \mathcal{C}'_x$, both sets $V' = \{\mathcal{C}_x \mid \mathcal{C}_x \prec \mathcal{C}_1\}$ and $V'' = \{\mathcal{C}'_x \mid \mathcal{C}'_x \succ \mathcal{C}_2\}$ are open. Since they contain x_0 , we can then take $V = V' \cap V''$. \square

The next result is about curves with intrinsic length small with respect to the Euclidean length. It will be of crucial importance for the proof of Theorem 1.1.

Proposition 2.24. *Let $\mathcal{C}_x, \mathcal{C}'_x$ be continuous cone structures defined on a neighborhood of a compact convex subset K of \mathbb{R}^N with $\mathcal{C} \prec \mathcal{C}'$. There exist positive constants δ, ρ such that for any nonconstant curve $\gamma : [a, b] \rightarrow K$, with*

$$\int_{\gamma} d_{\mathcal{C}} < \min\{\rho, \delta \ell(\gamma)\}, \quad (16)$$

we can find a \mathcal{C}' -future directed piecewise linear curve lying in K which joins $\gamma(a)$ to $\gamma(b)$.

Proof. By Lemma 2.23 we can construct a finite covering of K by open Euclidean balls $x_i + 3\eta\mathbb{B}$, $i = 1, \dots, k$, for some $\eta > 0$, $x_i \in K$, such that for any i we can find cones $\mathcal{C}_1^i, \mathcal{C}_2^i$ satisfying

$$\mathcal{C}'_y \succ \mathcal{C}_2^i \succ \mathcal{C}_1^i \succ \mathcal{C}_y \quad \text{for any } y \in x_i + 3\eta\mathbb{B}. \quad (17)$$

We denote by δ_i a constant satisfying (6) with $\mathcal{C}_1^i, \mathcal{C}_2^i$ in place of $\mathcal{C}^1, \mathcal{C}^2$, respectively, $i = 1, \dots, k$, and set $\delta = \min_i \delta_i$, $\rho = \eta \delta$. Note that the assertion is true for any curve $\xi : [c, d] \rightarrow x_i + 3\eta\mathbb{B}$, for some i , satisfying

$$\int_{\xi} d_{\mathcal{C}} < \delta \ell(\xi),$$

because in this case we infer from (17), Lemma 2.7, and the convex character of K , that the oriented segment from $\xi(c)$ to $\xi(d)$ is a \mathcal{C}' -future directed curve.

Given $\gamma : [a, b] \rightarrow K$, the idea is to define inductively an increasing finite sequence of times

$$t_0 = a, t_1, \dots, t_{m-1}, t_m = b$$

such that $\gamma|_{[t_{j-1}, t_j]} \subset x_{i_j} + 3\eta\mathbb{B}$, for some index i_j , and then to argue on each subinterval.

We assume that $\gamma(a) \in x_{i_1} + \eta\mathring{\mathbb{B}}$ and denote by t_1 the largest $t \in [a, b]$ such that $\gamma([a, t]) \subset \gamma(a) + \eta\mathbb{B}$. If $|\gamma(a) - \gamma(t_1)| < \eta$ then $t_1 = b$, and the support of the curve γ is entirely contained in $x_{i_1} + 2\eta\mathbb{B}$, so that the assertion is obtained according to what we pointed out above. If instead $|\gamma(a) - \gamma(t_1)| = \eta$ then by the assumption

$$\int_{\gamma|_{[a, t_1]}} d_{\mathcal{C}} \leq \int_{\gamma} d_{\mathcal{C}} \leq \rho = \eta \delta \leq \delta \ell(\gamma|_{[a, t_1]}), \quad (18)$$

which implies that the oriented segment from $\gamma(a)$ to $\gamma(t_1)$ is \mathcal{C}' -future directed, since $\gamma([a, t_1]) \subset x_{i_1} + 2\eta\mathbb{B}$.

If $t_1 < b$, using the same process as above, we construct by induction a sequence $t_0 = a, t_1, \dots, t_k$ such that $|\gamma(t_j) - \gamma(t_{j+1})| = \eta$, and the oriented segment from $\gamma(t_j)$ to $\gamma(t_{j+1})$ is \mathcal{C}' -future directed continuous C^∞ curve from $\gamma(t_j)$ to $\gamma(t_{j+1})$, for $j = 0, \dots, k-1$. Since γ has finite length, there must be $m > 1$ with $t_m = b$. Some precautions must be taken for the last step to complete the proof. In fact we cannot estimate the length of $\gamma|_{[t_{m-1}, t_m]}$, and so we argue on $\gamma|_{[t_{m-2}, t_m]}$. Note that $\gamma([t_{m-2}, t_m]) \subset x_{i_{m-2}} + 3\eta\mathbb{B}$, for some i_{m-2} , and $\ell(\gamma|_{[t_{m-2}, t_m]}) \geq \eta$, and we can repeat the estimate (18), with $[t_{m-2}, t_m]$ in place of $[0, t_1]$, to conclude the argument. \square

2.4 Cone Structures on a Manifold

Let M be a connected, paracompact, C^∞ manifold of dimension N . We denote, for any $x \in M$, by $T_x M$, $T_x^* M$ the tangent and cotangent spaces of M at x , respectively. The tangent and cotangent bundle are TM and T^*M , respectively. We will assume in the sequel that M is endowed with a Riemannian metric g .

We proceed introducing some notations.

For $v, v' \in T_x M$, we will set $|v|_x = (g_x(v, v))^{1/2}$. We will also denote by $|\cdot|_x$ the norm on $T_x^* M$ dual to $|\cdot|_x$ on $T_x M$. The formula $d_x^g(v, v') = |v - v'|_x$ gives a distance on $T_x M$, we will denote the corresponding closed unit ball by \mathbb{B}_x . We will write $d_g(x, y)$ for the Riemannian distance between the points $x, y \in M$, and denote by $B_g(x, r)$ the related metric closed ball centered at x , with radius $r > 0$.

Definition 2.25. A (resp. Lipschitz) cone structure \mathcal{C} on M is a map which associates to any $x \in M$ a (complete) cone $\mathcal{C}_x \subset T_x M$, such that for any C^∞ chart $\varphi : U \rightarrow \varphi(U) \subset \mathbb{R}^N$ the family of cones $\varphi_* \mathcal{C}$ defined by

$$(\varphi_* \mathcal{C})_y = D\varphi(\varphi^{-1}(y)) \mathcal{C}_{\varphi^{-1}(y)}.$$

is a continuous (resp. locally Lipschitz) family of cones on the open subset $\varphi(U) \subset \mathbb{R}^N$.

In the definition above $D\varphi(x) : T_x M \rightarrow T_{\varphi(x)} \mathbb{R}^N = \mathbb{R}^N$, is the derivative (or tangent map) of φ at $x \in U$.

By Lemma 2.21, as usual, to verify that the family of cone \mathcal{C} is a (resp. Lipschitz) cone structure, it is sufficient to find family of charts $(U_i, \varphi_i)_{i \in I}$ such that $\varphi_{i*} \mathcal{C}$ is a (resp. Lipschitz) cone structure on the open subset $\varphi_i(U_i)$ of \mathbb{R}^N , for each $i \in I$.

In the same way we can define the notion of (resp. Lipschitz) dual cone structure on M . A dual cone structure $\tilde{\mathcal{C}}$ is a family $(\tilde{\mathcal{C}}_x)_{x \in M}$, where $\tilde{\mathcal{C}}_x \subset T_x^* M$ is a (complete) cone for each $x \in M$, such that for any C^∞ chart $\varphi : U \rightarrow \varphi(U) \subset \mathbb{R}^N$ the family of cones $\varphi^* \tilde{\mathcal{C}}$ defined by

$$(\varphi^* \tilde{\mathcal{C}})_y = [{}^t D\varphi((\varphi)^{-1}(y))]^{-1} \tilde{\mathcal{C}}_{(\varphi)^{-1}(y)}.$$

is a continuous (resp. locally Lipschitz) family of cones on the open subset $\varphi(U) \subset \mathbb{R}^N$. Here we denote by ${}^t D\varphi(x) : (\mathbb{R}^N)^* \rightarrow T_x^* M$, the transpose (or dual) of the derivative $D\varphi(x) : T_x M \rightarrow \mathbb{R}^N$, and we identified $(\mathbb{R}^N)^*$ to \mathbb{R}^N using the canonical basis (or the canonical Euclidean product) as we did since the beginning of §2.1. Again by the dual analog of Lemma 2.21, as usual, to verify that the family of cone $\tilde{\mathcal{C}}$ is a (resp. Lipschitz) dual cone structure, it is sufficient to find family of charts $(U_i, \varphi_i)_{i \in I}$ such that $\varphi_i^* \tilde{\mathcal{C}}$ is a (resp. Lipschitz) cone structure on the open subset $\varphi_i(U_i)$ of \mathbb{R}^N , for each $i \in I$.

We now extend to cone structures on M the notion of dual cone. Given $x \in M$ and a cone $\mathcal{C}^0 \subset T_x M$ we define the dual (or polar) $\mathcal{C}^{0*} \subset T_x^* M$ via

$$\mathcal{C}^{0*} = \{p \in T_x^* M \mid p(v) \leq 0 \text{ for } v \in \mathcal{C}^0\}.$$

Therefore if \mathcal{C} is a cone structure on M , the dual cone structure is \mathcal{C}^* defined by $\mathcal{C}_x^* = (\mathcal{C}_x)^*$. It is not difficult to check, with the notations above, that for a chart $\varphi : U \rightarrow \varphi(U) \subset \mathbb{R}^N$, the cones $\varphi_* \mathcal{C}_y$ and $\varphi^* \mathcal{C}_y^*$ are dual cones in \mathbb{R}^N for every $y \in \varphi(U)$. It follows that \mathcal{C}^* is a (resp. Lipschitz) dual cone structure on M if \mathcal{C} is a (resp. Lipschitz) cone structure.

The order relation \succ for cone structures is naturally extended in this setting. If $\mathcal{C}, \mathcal{C}'$ are cone structures on M , we say that $\mathcal{C} \prec \mathcal{C}'$ if $\mathcal{C}_x \prec \mathcal{C}'_x$, for every $x \in M$. Same thing for dual cone structures. Of course $\mathcal{C} \prec \mathcal{C}'$ if and only if $\mathcal{C}^* \succ \mathcal{C}'^*$.

We now introduce the function $d_{\mathcal{C}}^g, d_{\mathcal{C}}^{g\#} : TM \rightarrow \mathbb{R}$ defined for all $(x, v) \in TM$ by

$$\begin{aligned} d_{\mathcal{C}}^g(x, v) &= d_x^g(v, \mathcal{C}_x) = \inf_{v' \in \mathcal{C}_x} d_x^g(v, v'). \\ d_{\mathcal{C}}^{g\#}(x, v) &= 2d_x^g(v, \mathcal{C}_x) - d_x^g(v, \partial \mathcal{C}_x). \end{aligned}$$

By Lemma 2.4 we have

$$\begin{aligned} d_{\mathcal{C}}^g(x, v) &= \sup\{p(v) \mid p \in \mathcal{C}_x^*, |p|_x \leq 1\} \\ d_{\mathcal{C}}^{g\#}(x, v) &= \sup\{p(v) \mid p \in \mathcal{C}_x^*, |p|_x = 1\}. \end{aligned}$$

We can now introduce the function $I_g(\mathcal{C}) : M \rightarrow [0, +\infty[$ defined by

$$I_g(\mathcal{C})(x) = -\min\{d_{\mathcal{C}}^{g\#}(x, v) \mid v \in \mathcal{C}_x, \cap \mathbb{B}_x\}.$$

For $\delta \geq 0$, we also introduce the family $\mathcal{C}^{-\delta}$ defined by

$$\mathcal{C}_x^{-\delta} = \{v \in T_x M \mid d_{\mathcal{C}}^{g\#}(x, v) \leq -\delta|v|_x\}.$$

For each $\delta < I_g(\mathcal{C})(x)$, we know that $\mathcal{C}_x^{-\delta}$ is a cone in $T_x M$. If moreover $\delta > 0$, we have $\mathcal{C}_x^{-\delta} \prec \mathcal{C}_x$.

If $x \in M$, and $\mathcal{C}_1, \mathcal{C}_2$ are two cones in $T_x M$, we define a Hausdorff-type distance between them, by

$$\delta_x^{g,H}(\mathcal{C}_1, \mathcal{C}_2) = \max \left\{ \max_{w \in \mathcal{C}_2 \cap \mathbb{B}_x} d_x^g(w, \mathcal{C}_1), \max_{v \in \mathcal{C}_1 \cap \mathbb{B}_x} d_x^g(v, \mathcal{C}_2) \right\}.$$

Proposition 2.26. *The map $TM \rightarrow [0, +\infty[, (x, v) \mapsto |v|_x$ is locally Lipschitz on TM .*

Let \mathcal{C} be a (resp. Lipschitz) cone structures on the Riemannian manifold M , the following functions are continuous (resp. locally Lipschitz):

- (i) $(x, v) \mapsto d_{\mathcal{C}}^g(x, v)$, from TM to $[0, +\infty[$;
- (ii) $(x, v) \mapsto d_{\mathcal{C}}^{g\#}(x, v)$, from TM to \mathbb{R} ;
- (iii) $I_g(\mathcal{C}) : M \rightarrow [0, +\infty[$, in particular, the subset $\{(\delta, x) \in [0, +\infty[\times M \mid \delta < I_g(\mathcal{C})(x)\}$ is open in $[0, +\infty[\times M$;
- (iv) $(\delta, x) \mapsto \delta_x^{g,H}(\mathcal{C}_x^{-\delta}, \mathcal{C}_x)$ from the open subset $\{(\delta, x) \in [0, +\infty[\times M \mid \delta < I_g(\mathcal{C})(x)\}$ to $[0, +\infty[$.

If $\theta : M \rightarrow [0, +\infty[$ is continuous (resp. locally Lipschitz), with $\theta(x) < I_g(\mathcal{C})(x)$, for every $x \in M$, then $\mathcal{C}^{-\theta} = (\mathcal{C}_x^{-\theta(x)})_{x \in M}$ is a (resp. Lipschitz) cone structure on M .

If \mathcal{C}' is another cone structure on M , then the sets $\{(\delta, x) \in [0, +\infty[\times M \mid \delta < I_g(\mathcal{C})(x), \mathcal{C}'_x \prec \mathcal{C}_x^{-\delta}\}$ and $\{(\delta, x) \in [0, +\infty[\times M \mid \delta < I_g(\mathcal{C})(x), \mathcal{C}'_x \succ \mathcal{C}_x^{-\delta}\}$ are open in $[0, +\infty[\times M$.

The easiest way to prove this proposition is to use the Gram-Schmidt orthogonalization method. This method provides a way to choose an orthonormal basis for each scalar product on \mathbb{R}^N , depending smoothly on the scalar product. Define $Q^+(\mathbb{R}^N)$ as the space of positive definite bilinear forms (or scalar products) on \mathbb{R}^N .

Proposition 2.27 (Gram-Schmidt). *There exist C^∞ (even real analytic) maps $Q^+(\mathbb{R}^N) \rightarrow \mathbb{R}^N, g \mapsto e_i^g, i = 1, \dots, N$ such that (e_1^g, \dots, e_N^g) is a basis of \mathbb{R}^N orthonormal for the scalar product g .*

In particular if we define $L^g : \mathbb{R}^N \rightarrow \mathbb{R}^N$ by $L^g(x_1, \dots, x_N) = \sum_{i=1}^N x_i e_i^g$, then L^g is an isometry of \mathbb{R}^N with the usual scalar product to \mathbb{R}^N endowed with the scalar product g , and the map $Q^+(\mathbb{R}^N) \rightarrow \text{GL}(\mathbb{R}^N), g \mapsto L^g$ is C^∞ (even real analytic).

Proof. We recall the well-known argument. We will construct $e_i^g, i = 1, \dots, N$ by induction on i , with the additional property that, for each $i = 1, \dots, N$, the subset $\{e_1^g, \dots, e_i^g\}$ generates the same vector space as $\{e_1, \dots, e_i\}$, where e_1, \dots, e_N is the canonical basis of \mathbb{R}^N . It suffices to take $e_1^g = e_1 / \sqrt{g(e_1, e_1)}$. If e_1^g, \dots, e_{i-1}^g have already been defined, we denote by f_i^g the orthogonal (with respect of the scalar product g) projection of e_i , on the subspace generated by (e_1, \dots, e_{i-1}) . Since $(e_1^g, \dots, e_{i-1}^g)$ is an orthonormal basis, for g , of this last vector subspace, we have

$$f_i^g = \sum_{j=1}^{i-1} g(e_i, e_j^g) e_j^g.$$

Obviously, the projection f_i^g is, by induction, C^∞ (even real analytic) as a function of g . Since e_i is not in the space generated by $\{e_1, \dots, e_{i-1}\}$ (or $\{e_1^g, \dots, e_{i-1}^g\}$), the vector $v_i^g = e_i - f_i^g$ is never 0, therefore it suffices to set $e_i^g = v_i^g / \sqrt{g(v_i^g, v_i^g)}$. \square

Proof of Proposition 2.26. Since continuity (resp. being locally Lipschitz) is a local property, we can assume that $M = U$ is an open subset of \mathbb{R}^N . Therefore, the tangent space $TM = TU$ is $U \times \mathbb{R}^N$. If we call g_x , the Riemannian metric on $T_x U = \{x\} \times \mathbb{R}^N \simeq \mathbb{R}^N$, applying Lemma 2.27 to g_x , we can define a map $L_x = L^{g_x} \in \text{GL}(\mathbb{R}^N)$ which is an isometry of \mathbb{R}^N with the usual scalar product on \mathbb{R}^N endowed with the scalar product g_x and such that the map $U \rightarrow \text{GL}(\mathbb{R}^N), x \mapsto L_x$ is C^∞ . We have

$$|v| = |L_x(v)|_x \quad \text{for any } (x, v) \in TM,$$

which shows the first part of the proposition

Now we define $\tilde{\mathcal{C}}_x = L_x^{-1}(\mathcal{C}_x)$. By Lemma 2.19, the family of cone $\tilde{\mathcal{C}} = (\tilde{\mathcal{C}}_x)$, $x \in U$, depends continuously on $x \in U$. By construction, we have for any $(x, v) \in TM$

$$\begin{aligned} d(v, \tilde{\mathcal{C}}_x) &= d_x^g(L_x v, \mathcal{C}_x) = d_{\mathcal{C}}^g(x, L_x v), \\ d^\#(v, \tilde{\mathcal{C}}_x) &= d_x^{g^\#}(L_x v, \mathcal{C}_x) = d_{\mathcal{C}}^{g^\#}(x, L_x v), \end{aligned}$$

where $d(x, \tilde{\mathcal{C}}_x)$ and $d^\#(v, \tilde{\mathcal{C}}_x)$ are the distance and the signed distance from v to $\tilde{\mathcal{C}}_x$ measured in the usual Euclidean distance. Since the map $(x, v) \mapsto (x, L_x v)$ is a C^∞ diffeomorphism of $U \times \mathbb{R}^N$, to get the items i) and ii), it suffices to prove, that the maps $(x, v) \mapsto d(v, \tilde{\mathcal{C}}_x)$ and $(x, v) \mapsto d^\#(v, \tilde{\mathcal{C}}_x)$ are continuous (resp. locally Lipschitz). But this follows from Proposition 2.13. We pass to the proof of iii). Since $L_x \mathbb{B} = \mathbb{B}_x$, we obtain

$$\begin{aligned} I(\tilde{\mathcal{C}}_x) &= - \min_{v \in \mathbb{B} \cap \tilde{\mathcal{C}}_x} d^\#(v, \tilde{\mathcal{C}}_x) \\ &= - \min_{v \in \mathbb{B} \cap \tilde{\mathcal{C}}_x} d^{g\#}(L_x v, \mathcal{C}_x) \\ &= - \min_{w \in \mathbb{B}_x \cap \mathcal{C}_x} d_{\mathcal{C}}^{g\#}(x, w) \\ &= I_g(\mathcal{C})(x). \end{aligned}$$

The continuity of $I_g(\mathcal{C})$ is now a consequence of Lemma 2.16.

To prove iv), we first argue that $L_x[(\tilde{\mathcal{C}}_x)^{-\delta}] = \mathcal{C}_x^{-\delta}$. In fact, for $v \in \mathbb{R}^N$, we have $L_x(v) \in \mathcal{C}_x^{-\delta}$ if and only if

$$d^{g\#}(x, L_x v) \leq -\delta |L_x v|_x.$$

By the computations made above this is the same as the inequality

$$d^\#(v, \tilde{\mathcal{C}}_x) \leq -\delta |v|,$$

which precisely defines $(\tilde{\mathcal{C}}_x)^{-\delta}$.

Now if \mathcal{C}' is another cone structure on M , we obtain, arguing as above

$$\max_{v \in \tilde{\mathcal{C}}'_x \cap \mathbb{B}} d(v, \tilde{\mathcal{C}}_x) = \max_{w \in \mathcal{C}'_x \cap \mathbb{B}_x} d_x^g(w, \mathcal{C}_x) = \max_{w \in \mathcal{C}'_x \cap \mathbb{B}_x} d_{\mathcal{C}'}^g(x, w)$$

and, consequently

$$\begin{aligned} \delta^H(\tilde{\mathcal{C}}_x, \tilde{\mathcal{C}}'_x) &= \max\left\{ \max_{v \in \tilde{\mathcal{C}}'_x \cap \mathbb{B}} d(v, \tilde{\mathcal{C}}_x), \max_{u \in \tilde{\mathcal{C}}_x \cap \mathbb{B}} d(u, \tilde{\mathcal{C}}'_x) \right\} \\ &= \max\left\{ \max_{w \in \mathcal{C}'_x \cap \mathbb{B}_x} d_{\mathcal{C}}^g(x, w), \max_{z \in \tilde{\mathcal{C}}_x \cap \mathbb{B}_x} d_{\mathcal{C}'}^g(x, z) \right\} \\ &= \delta_x^{g,H}(\mathcal{C}_x, \mathcal{C}'_x). \end{aligned}$$

In particular, this shows that

$$\delta_x^{g,H}(\mathcal{C}_x^{-\delta}, \mathcal{C}_x) = \delta^H((\tilde{\mathcal{C}}_x)^{-\delta}, \tilde{\mathcal{C}}_x).$$

Since δ^H is the distance on $\text{Cones}(\mathbb{R}^N)$, we can now obtain iv) from the continuity of $(\delta, x) \mapsto (\tilde{\mathcal{C}}_x)^{-\delta}$ which follows from Lemma 2.17.

Now the continuity of $(\delta, x) \mapsto (\tilde{\mathcal{C}}_x)^{-\delta}$ also implies that $\mathcal{C}^{-\theta}$ is a (continuous) cone structure for θ is continuous because

$$\mathcal{C}_x^{-\theta(x)} = L_x \left((\tilde{\mathcal{C}}_x)^{-\theta(x)} \right).$$

Note that besides the continuity of $(\delta, x) \mapsto (\tilde{\mathcal{C}}_x)^{-\delta}$, Lemma 2.17 also shows that this map is locally Lipschitz if \mathcal{C} is a Lipschitz cone structure and θ is continuous. Therefore $\mathcal{C}^{-\theta}$ is a Lipschitz cone structure if θ is locally Lipschitz.

We now prove the last part of the Proposition. Since $\mathcal{C}'_x \prec \mathcal{C}_x^{-\delta}$ if and only if $\tilde{\mathcal{C}}'_x \prec (\tilde{\mathcal{C}}_x)^{-\delta}$, the fact that the $\{(\delta, x) \in [0, +\infty[\times M \mid \delta < I_g(\mathcal{C})(x), \mathcal{C}'_x \prec \mathcal{C}_x^{-\delta}\}$ is open follows from Lemma 2.18. \square

We now proceed to extend theorem 2.15 to cone structures on manifolds.

Theorem 2.28. *Let \mathcal{C} be a (resp. Lipschitz) cone structure on the Riemannian manifold M . If $\epsilon : M \rightarrow]0, +\infty[$ is a continuous function, we can find (resp. Lipschitz) cone structures $\mathcal{C}' \prec \mathcal{C} \prec \mathcal{C}''$ with $\delta_x^{g,H}(\mathcal{C}_x, \mathcal{C}'_x) < \epsilon(x)$, $\delta_x^{g,H}(\mathcal{C}_x, \mathcal{C}''_x) < \epsilon(x)$ for any $x \in M$.*

Proof. As in Theorem 2.15, it is enough to show the existence of a cone structure \mathcal{C}' with the properties asserted in the statement, \mathcal{C}'' is in fact obtained from this by duality (more on this below).

The subset

$$\mathcal{O} = \{(\delta, x) \in [0, +\infty[\times M \mid \delta < I_g(\mathcal{C})(x), \delta_x^{g,H}(\mathcal{C}_x^{-\delta}, \mathcal{C}_x) < \epsilon(x)\}$$

is open in $[0, +\infty[\times M$, by item (iii) and (iv) of Proposition 2.26. It obviously contains $\{0\} \times M$, we can therefore find an open cover $(U_i)_{i \in I}$ of M and a family $(\delta_i)_{i \in I}$ of (strictly) positive numbers such that $\{0\} \times M \subset \cup_{i \in I}]0, \delta_i[\times U_i \subset \mathcal{O}$. We take $(\theta_i)_{i \in I}$ a C^∞ partition of unity on M subordinated to the open cover $(U_i)_{i \in I}$. The function θ defined by

$$\theta(x) = \sum_{i \in I} \theta_i(x) \delta_i,$$

is C^∞ , and > 0 . Moreover, we have $(\theta(x), x) \in \mathcal{O}$, for every $x \in M$. It follows from Proposition 2.26, that we can take $\mathcal{C}' = \mathcal{C}^{-\theta}$.

To prove the existence of \mathcal{C}'' we argue by duality. We could just take a dual cone structure $\tilde{\mathcal{C}}'' \prec \mathcal{C}^*$ with $\delta_x^{g,H}(\tilde{\mathcal{C}}''_x, \mathcal{C}^*_x) < \epsilon(x)$, and take \mathcal{C}'' as the (pre-)dual of $\tilde{\mathcal{C}}''$. Note that instead of \mathcal{C}^* , we could use the (essentially same) cone structure \mathcal{C}^\sharp on M given by

$$\mathcal{C}_x^\sharp = \{w \in T_x M \mid g_x(w, v) \leq 0, \forall v \in \mathcal{C}_x\}.$$

This cone structure is simply \mathcal{C}^* using the identification of TM with T^*M given by the Riemannian metric g \square

We get as a consequence:

Corollary 2.29. *Given two continuous cone structures $\mathcal{C}, \mathcal{C}''$ on M with $\mathcal{C}'' \succ \mathcal{C}$, there exists a continuous cone structure \mathcal{C}' satisfying $\mathcal{C}'' \succ \mathcal{C}' \succ \mathcal{C}$. Moreover if either \mathcal{C} or \mathcal{C}'' is a Lipschitz cone structure, we could choose \mathcal{C}' also Lipschitz.*

Proof. Keeping the notation of the proof previous proposition, we consider subset

$$\mathcal{O}' = \{(\delta, x) \in [0, +\infty[\times M \mid \delta < I_g(\mathcal{C})(x), \mathcal{C}'' \prec \mathcal{C}_x^{-\delta}\}.$$

By the last part of Proposition 2.26, this set is open in $[0, +\infty[\times M$. Moreover, it contains $\{0\} \times M$ since $\mathcal{C}'' \prec \mathcal{C}$. Arguing as in the proof of the previous proposition, we can find a C^∞ function $\theta : M \rightarrow]0, +\infty[$ such that $(x, \theta(x)) \in \mathcal{O} \cap \mathcal{O}'$. We can then take $\mathcal{C}' = \mathcal{C}^{-\theta}$.

Note that when \mathcal{C} is Lipschitz, this gives a Lipschitz \mathcal{C}' . If it is rather \mathcal{C}'' that it is Lipschitz, we could use duality to exchange the role of \mathcal{C} and \mathcal{C}'' . \square

2.5 Adapted Cover and Metric

It will simplify a lot the proofs if we can impose some technical conditions on Riemannian balls of a fixed radius.

We can always find a countable atlas of C^∞ charts $\varphi_n : U_n \rightarrow \mathbb{R}^N, n \in \mathbb{N}$ such that for each n

- (i) The set U_n is open and relatively compact in M ;
- (ii) the map $\varphi_n : U_n \rightarrow \mathbb{R}^N$ is a C^∞ diffeomorphism;
- (iii) the family $\varphi_n^{-1}(\mathring{\mathbb{B}}), n \in \mathbb{N}$ covers M ;
- (iv) the family $(U_n)_{n \in \mathbb{N}}$ is a locally finite open covering of M .

It follows from the previous items that the set $\{m \in \mathbb{N} \mid U_m \cap U_n \neq \emptyset\}$ is finite for each $n \in \mathbb{N}$. We set $K_n = \varphi_n^{-1}(\mathbb{B})$ and $K'_n = \varphi_n^{-1}(2\mathbb{B})$. Therefore we get

$$\begin{aligned} \bigcup_{n \in \mathbb{N}} K_n &= M, \\ K_n &\subset \mathring{K}'_n \subset K'_n \subset U_n, \\ \varphi_n(K'_n) &\text{ is convex.} \end{aligned}$$

In the sequel of the paper, we will assume that the cover above has been chosen once and for all.

As we said before, we assume that M is endowed with a Riemannian metric g . The next result tells us that g can be modified, by multiplying it by a suitable smooth function, in order to obtain a complete Riemannian metric enjoying some additional properties.

Lemma 2.30. *We can find a positive C^∞ function $\beta : M \rightarrow]0, +\infty[$ such that the Riemannian metric $\tilde{g} := \beta^2 g$ satisfies*

$$\bar{B}_{\tilde{g}}(x, 3) = \{y \in M \mid d^{\tilde{g}}(x, y) \leq 3\} \subset K'_n \quad \text{for every } n \in \mathbb{N}, \text{ every } x \in K_n .$$

Proof. We set

$$\delta_n = \min\{d_g(x, y) \mid x \in K_n, y \in \partial K'_n\}.$$

By compactness $\delta_n > 0$. We claim that \tilde{g} will satisfy the required property whenever $\beta : M \rightarrow]0, +\infty[$ verifies

$$\inf_{K'_n} \beta \geq \frac{4}{\delta_n}. \quad (19)$$

In fact, the distance δ_n is nothing but the infimum of the lengths $\ell_g(\gamma)$ all curves $\gamma : [a, b] \rightarrow K'_n$ with $\gamma(a) \in K_n, \gamma(b) \in \partial K'_n$, and for such a curve γ we have by (19)

$$\ell_{\tilde{g}}(\gamma) \geq \left(\inf_{K'_n} \beta \right) \ell_g(\gamma) \geq \frac{4}{\delta_n} \delta_n = 4.$$

It follows that any point y in M at a distance strictly less than 4 for \tilde{g} from a point in K_n is necessarily in K'_n .

It remains to construct β such that $\inf_{K'_n} \beta \geq 4\delta_n^{-1}$, for all $n \in \mathbb{N}$. Notice that $\{m \in \mathbb{N} \mid U_m \cap U_n \neq \emptyset\}$ is finite, therefore we can set

$$\lambda_n = \max \left\{ \frac{4}{\delta_m} \mid m \in \mathbb{N}, U_m \cap U_n \neq \emptyset \right\} < +\infty.$$

We fix a C^∞ partition of unity $(\theta_n)_{n \in \mathbb{N}}$ subordinated to the open cover $(U_n)_{n \in \mathbb{N}}$, and we define

$$\beta(x) = \sum_{n \in \mathbb{N}} \lambda_n \theta_n(x).$$

We have $\inf_{U_n} \beta \geq 4\delta_n^{-1}$. In fact, if $x \in U_n$, and $\theta_m(x) > 0$, we get $x \in U_m \cap U_n$. This implies $\lambda_m \geq 4\delta_n^{-1}$, and therefore $\beta(x) \geq 4\delta_n^{-1}$. \square

From now on we will assume that we have replaced g by the Riemannian metric \tilde{g} . Therefore changing the name of \tilde{g} back to g we will assume

$$\bar{B}_g(x, 3) \subset K'_n \text{ for every } x \in K_n.$$

Note that the Riemannian metric g is necessarily complete since the closed balls $B_g(x, 3)$ are compact for any $x \in M$.

3 Intrinsic semi-distances and Aubry sets: Proof of Theorem 1.1

We will assume that \mathcal{C} is a (continuous) cone structure on M , which is endowed with the Riemannian metric g . We will also suppose that we have chosen a

continuous function $\alpha : M \rightarrow]0, +\infty[$. We define the *intrinsic length* $\sigma_{g,\alpha}^{\mathcal{C}}(\gamma)$ of the Lipschitz curve $\gamma : [a, b] \rightarrow M$ by

$$\sigma_{g,\alpha}^{\mathcal{C}}(\gamma) = \int_{\gamma} \alpha d_{\mathcal{C}}^g = \int_a^b \alpha(\gamma(t)) d_{\gamma(t)}^g(\dot{\gamma}(t), \mathcal{C}_{\gamma(t)}) dt. \quad (20)$$

The above quantity is nonnegative and invariant under orientation-preserving reparametrization of γ . Using intrinsic length of curves, we can introduce the *intrinsic semi-distance* $S_{\alpha,g}^{\mathcal{C}}$ on M , namely we set for x, y in M

$$S_{\alpha,g}^{\mathcal{C}}(x, y) = \inf_{\gamma} \sigma_{g,\alpha}^{\mathcal{C}}(\gamma), \quad (21)$$

where the infimum is taken on all Lipschitz curves γ joining x to y . It is also clear that if \mathcal{C}' is another cone structure on M with $\mathcal{C} \prec \mathcal{C}'$ then $S_{\alpha,g}^{\mathcal{C}'} \leq S_{\alpha,g}^{\mathcal{C}}$. We now check that $S_{\alpha,g}^{\mathcal{C}}$ is indeed a (not necessarily symmetric) nonnegative semi-distance.

Proposition 3.1. *The nonnegative function $S_{\alpha,g}^{\mathcal{C}}$ satisfies*

$$\begin{aligned} S_{\alpha,g}^{\mathcal{C}}(x, x) &= 0, \\ S_{\alpha,g}^{\mathcal{C}}(x, z) &\leq S_{\alpha,g}^{\mathcal{C}}(x, y) + S_{\alpha,g}^{\mathcal{C}}(y, z). \end{aligned}$$

Moreover, the function $S_{\alpha,g}^{\mathcal{C}}$ is locally Lipschitz on $M \times M$.

Proof. The fact that $S_{\alpha,g}^{\mathcal{C}}(x, x) = 0$ follows from the homogeneity in v of $d_{\mathcal{C}}^g(x, v)$ which implies that $\sigma_{g,\alpha}^{\mathcal{C}}(\gamma) = 0$ for a constant curve γ . The triangle inequality $S_{\alpha,g}^{\mathcal{C}}(x, z) \leq S_{\alpha,g}^{\mathcal{C}}(x, y) + S_{\alpha,g}^{\mathcal{C}}(y, z)$ is a routine argument of concatenation of curves.

Since by the triangle inequality

$$S_{\alpha,g}^{\mathcal{C}}(x', y') - S_{\alpha,g}^{\mathcal{C}}(x, y) \leq S_{\alpha,g}^{\mathcal{C}}(x', x) + S_{\alpha,g}^{\mathcal{C}}(y, y'),$$

it suffices to prove, to get the Lipschitz regularity of $S_{\alpha,g}^{\mathcal{C}}$, that for each $x_0 \in M$, there is a neighborhood V of x_0 a finite constant $C > 0$ such that

$$S_{\alpha,g}^{\mathcal{C}}(x', x) \leq C d^g(x, x') \quad \text{for all } x, x' \in V.$$

Since this is a local problem we can assume M to be an open set in the Euclidean space \mathbb{R}^N , and $x_0 = 0$. We pick $r > 0$ such that $V \subset r\mathbb{B} \subset M \subset \mathbb{R}^N$. By compactness

$$C' = \sup\{\alpha(x) d_{\mathcal{C}}^g(x, v) \mid x \in r\mathbb{B}, |v| = 1\} < +\infty.$$

By homogeneity

$$\alpha(x) d_{\mathcal{C}}^g(x, v) \leq C' |v| \quad \text{for any } x \in r\mathbb{B}, v \in \mathbb{R}^N.$$

Obviously, if $x, x' \in r\mathbb{B}$ we consider the Euclidean segment $\xi(t) = (1-t)x' + tx$, which satisfies $\xi(t) \in r\mathbb{B}$, and $\dot{\xi}(t) = x - x'$, for any $t \in [0, 1]$, to get

$$S_{\alpha,g}^{\mathcal{C}}(x', x) \leq \sigma_{\alpha,g}^{\mathcal{C}}(\xi) \leq K' |x - x'|.$$

This finishes the proof since d^g and the Euclidean distance are Lipschitz equivalent on $r\mathbb{B}$. \square

We recall that any locally Lipschitz function on M is differentiable for almost every (a.e. for short) $x \in M$, i.e. up to a subset of vanishing Riemannian measure. The intrinsic semi-distance is related to the dual cone structure \mathcal{C}^* by the following proposition.

Proposition 3.2. *Given $y \in M$, the locally Lipschitz functions $u_y = S_{\alpha,g}^{\mathcal{C}}(y, \cdot)$, and $v_y = S_{\alpha,g}^{\mathcal{C}}(\cdot, y)$ satisfy $Du_y(x) \in \mathcal{C}_x^*$, $-Dv_y(x) \in \mathcal{C}_x^*$, $|Du_y(x)|_x \leq \alpha(x)$, and $|Dv_y(x)|_x \leq \alpha(x)$ for a.e. $x \in M$.*

Proof. The proof is taken from [7] with minor adjustments. We will do the proof for u_y . Consider a differentiability point $x \in M$ of u_y , it suffices to prove that

$$Du_y(x) \in \mathcal{C}_x^* \quad \text{and} \quad |Du_y(x)|_x \leq \alpha(x).$$

Again this is a local result, therefore we can assume that M is an open subset of \mathbb{R}^N . For $v \in \mathbb{R}^n$, and $\varepsilon > 0$ small enough, the path $\gamma(t) = x + t\varepsilon v$, $t \in [0, 1]$, is included in M , therefore

$$\begin{aligned} u_y(x + \varepsilon v) - u_y(x) &= S_{\alpha,g}^{\mathcal{C}}(y, x + \varepsilon v) - S_{\alpha,g}^{\mathcal{C}}(y, x) \\ &\leq S_{\alpha,g}^{\mathcal{C}}(x, x + \varepsilon v) \\ &\leq \sigma_{\alpha,g}^{\mathcal{C}}(\gamma) = \int_0^1 \alpha(x + t\varepsilon v) d_{\mathcal{C}}^g(x + t\varepsilon v, \varepsilon v) dt \\ &= \varepsilon \int_0^1 \alpha(x + t\varepsilon v) d_{\mathcal{C}}^g(x + t\varepsilon v, v) dt. \end{aligned}$$

Therefore dividing by $\varepsilon > 0$ and letting $\varepsilon \rightarrow 0$, we get

$$Du_y(x)[v] \leq \alpha(x) d_{\mathcal{C}}^g(x, v).$$

Since $d_{\mathcal{C}}^g(x, v) = d_x^g(v, \mathcal{C}_x) \leq |v|_x$, we obtain $|Du_y(x)|_x \leq \alpha(x)$. Moreover $d_{\mathcal{C}}^g(x, v) \leq 0$ for $v \in \mathcal{C}_x$. Hence $Du_y(x)[v] \leq 0$ for $v \in \mathcal{C}_x$, which means $Du_y(x) \in \mathcal{C}_x^*$ \square

The proof of Theorem 1.1 is based on the notion of Aubry set that we now introduce. Given a positive smooth function α on M , we define $\mathcal{A}_{\alpha,g}^{\mathcal{C}}$ as the set of points x of M such that there is a sequence of closed curves γ_n based at x with

$$\inf_n \ell_g(\gamma_n) > 0 \quad \text{and} \quad \inf_n \sigma_{\alpha,g}^{\mathcal{C}}(\gamma_n) = 0.$$

The main point will be to show that α can be chosen in such a way that the corresponding Aubry set is empty.

The connection of this issue with the global time function we are looking for, is given by the following theorem which is a direct consequence of what is proved in [7], see in particular Theorem 6.2 there, and go back to the Introduction for more explanation.

Theorem 3.3. *If $\mathcal{A}_{\alpha,g}^{\mathcal{C}}$ is empty then there exists a C^∞ function $f : M \rightarrow \mathbb{R}$ such that $-Df(x) \in \mathring{\mathcal{C}}_x^*$ and $|Df(x)|_x \leq \alpha(x)$ for every $x \in M$.*

In other terms, f is the sought global time function.

We refer to the Introduction for the definitions of causal and stably causal cone structure. Let \mathcal{C} be a stably causal cone structure on M , we fix \mathcal{C}' a causal cone structure with $\mathcal{C}' \succ \mathcal{C}$.

In the sequel we will assume that $(U_n, \varphi_n), K_n, K'_n$ are the ones obtained at the beginning of section 2.5. We will further assume that the Riemannian metric g on M satisfies

$$\bar{B}_g(x, 3) \subset K'_n \quad \text{for all } n \geq 0, x \in K_n,$$

as this was shown to be possible in Lemma 2.30.

The next result, which essentially exploits Proposition 2.24, ensures that, for a suitable choice of α , if the intrinsic length $\int_\gamma \alpha d_g^{\mathcal{C}}$ of a non-constant curve $\gamma : [a, b] \rightarrow M$ is small compared to its Riemannian length, then we can find a \mathcal{C}' -future directed curve $\xi : [a, b] \rightarrow M$ with $\xi(a) = \gamma(a)$, and $\xi(b) = \gamma(b)$. In view of the causality assumption on \mathcal{C}' , this implies that γ cannot be a closed curve. Looking at the very definition of Aubry set above, we can, in turn, deduce that $\mathcal{A}_{\alpha,g}^{\mathcal{C}} = \emptyset$ and conclude the proof of Theorem 1.1 using Theorem 3.3. It is then left to show:

Theorem 3.4. *We can find a C^∞ positive function $\alpha : M \rightarrow]0, +\infty[$ such that if $\gamma : [a, b] \rightarrow M$ is a curve with*

$$\sigma_{\alpha,g}^{\mathcal{C}}(\gamma) = \int_\gamma \alpha d_g^{\mathcal{C}} < \min \left\{ \ell_g(\gamma), \frac{1}{2} \right\},$$

then there exists a \mathcal{C}' -future directed piecewise C^∞ curve $\tilde{\gamma} : [a, b] \rightarrow M$ with $\tilde{\gamma}(a) = \gamma(a)$ and $\tilde{\gamma}(b) = \gamma(b)$. Moreover, at each $t \in [a, b]$, both left and right derivatives $\tilde{\gamma}_-(t), \tilde{\gamma}_+(t)$ are contained in $\mathring{\mathcal{C}}'_{\tilde{\gamma}(t)}$.

Proof. Note that the curve γ cannot be constant, since it would have length 0, and this would contradict $\sigma_{\alpha,g}^{\mathcal{C}}(\gamma) \geq 0$. To avoid heavy notations, we identify K'_n , for any n , to a convex compact subset of \mathbb{R}^N through the diffeomorphism φ_n . Using the function $d_{\mathcal{C}}$ obtained from the Euclidean distance in (14), by Proposition 2.24 we can find suitable positive constants $\tilde{\rho}_n, \tilde{\delta}_n$ such that for any Lipschitz curve $\xi : [a, b] \rightarrow K'_n$ which satisfies

$$\int_\xi d_{\mathcal{C}} < \min\{\tilde{\rho}_n, \tilde{\delta}_n \ell(\xi)\}, \tag{22}$$

where $\ell(\xi)$ is the Euclidean length of ξ , there exists a piecewise linear \mathcal{C}' -future directed curve $\tilde{\xi}$ contained in K'_n , joining $\xi(a)$ to $\xi(b)$. Notice that such a piecewise linear curve satisfies automatically the last part of the statement of the theorem concerning the right and left derivative.

By compactness of K'_n we can find a positive number a_n such that

$$\frac{1}{a_n}|v| \leq |v|_x \leq a_n|v| \quad \text{for all } x \in K'_n, v \in \mathbb{R}^N. \quad (23)$$

Since $d_{\mathcal{C}}(x, \cdot)$ and $d_{\mathcal{C}}^g(x, \cdot)$ are distance functions to the same \mathcal{C}_x , this implies that

$$\frac{1}{a_n}d_{\mathcal{C}}(x, v) \leq d_{\mathcal{C}}^g(x, v) \leq a_n d_{\mathcal{C}}(x, v) \quad \text{for all } x \in K'_n, v \in \mathbb{R}^N. \quad (24)$$

It also implies that for any curve ξ with support contained in K'_n , we have

$$\frac{1}{a_n} \ell(\xi) \leq \ell_g(\xi) \leq a_n \ell(\xi), \quad (25)$$

where $\ell_g(\xi)$ is the Riemannian length of ξ .

We claim that for any Lipschitz curve $\xi : [a, b] \rightarrow K'_n$ which satisfies

$$\int_{\xi} d_{\mathcal{C}}^g < \min \left\{ \frac{1}{a_n} \tilde{\rho}_n, \frac{1}{a_n^2} \tilde{\delta}_n \ell_g(\xi) \right\}, \quad (26)$$

there exists a piecewise linear \mathcal{C}' -future directed curve $\tilde{\xi}$, with support contained in K'_n , joining $\xi(a)$ to $\xi(b)$. This is true simply because (26) implies (22), as we now show in the following computation which exploits (24), (25)

$$\begin{aligned} \int_{\xi} d_{\mathcal{C}} &\leq a_n \int_{\xi} d_{\mathcal{C}}^g < a_n \min \left\{ \frac{1}{a_n} \tilde{\rho}_n, \frac{1}{a_n^2} \tilde{\delta}_n \ell_g(\xi) \right\} \\ &\leq a_n \min \left\{ \frac{1}{a_n} \tilde{\rho}_n, \frac{1}{a_n} \tilde{\delta}_n \ell(\xi) \right\} = \min \{ \tilde{\rho}_n, \tilde{\delta}_n \ell(\xi) \}. \end{aligned}$$

We set $\rho_n = \frac{\tilde{\rho}_n}{a_n}$, $\delta_n = \frac{\tilde{\delta}_n}{a_n^2}$, and we choose a C^∞ function $\alpha : M \rightarrow]0, +\infty[$ with $\alpha > \max \left\{ \frac{1}{\delta_n}, \frac{2}{\rho_n} \right\}$ on K'_n , for each $n \in \mathbb{N}$. Such a function can be constructed arguing as in Lemma 2.30.

The rest of the proof goes along the same lines as that of Proposition 2.24, so we just give the main points of it. We assume $\gamma(a) \in K_{n_1}$, if $\gamma([a, b])$ is entirely contained in $B_g(\gamma(a), 1) \subset K'_{n_1}$ then

$$\begin{aligned} \frac{1}{\delta_{n_1}} \int_{\gamma} d_g^{\mathcal{C}} &< \int_{\gamma} \alpha d_g^{\mathcal{C}} \leq \ell_g(\gamma), \\ \frac{2}{\rho_{n_1}} \int_{\gamma} d_g^{\mathcal{C}} &< \int_{\gamma} \alpha d_g^{\mathcal{C}} \leq \frac{1}{2}, \end{aligned}$$

hence

$$\int_{\gamma} \alpha d_g^{\mathcal{C}} \leq \min \left\{ \frac{1}{a_{n_1}} \tilde{\rho}_{n_1}, \frac{1}{a_{n_1}^2} \tilde{\delta}_{n_1} \right\}.$$

and the curve $\tilde{\gamma}$ exists by what we saw above. If instead the curve goes out from $B_g(\gamma(a), 1)$ then its first exit time from this set, say t_1 , satisfies $d_g(\gamma(a), \gamma(t_1)) = 1$ and we have

$$\max \left\{ \frac{2}{\rho_1}, \frac{1}{\delta_1} \right\} \int_{\gamma|_{[a, t_1]}} d_{\mathcal{C}}^g \leq \int_{\gamma} \alpha d_{\mathcal{C}}^g \leq \frac{1}{2} \leq d_g(\gamma(a), \gamma(t_1)) \leq \ell_g(\gamma|_{[a, t_1]}),$$

from which we deduce, from what was obtained before, that we can find a piecewise linear \mathcal{C}' -future directed curve $\tilde{\gamma} : [a, t_1] \rightarrow K'_{n_1}$, with $\tilde{\gamma}(a) = \gamma(a)$ and $\tilde{\gamma}(t_1) = \gamma(t_1)$.

Then, still following the argument of Proposition 2.24, we construct by induction a finite sequence $t_0 = a, t_1, \dots, t_k = b$, and piecewise smooth \mathcal{C}' -future directed curves $\tilde{\gamma} : [t_j, t_{j+1}] \rightarrow K'_{n_j}$, with $\tilde{\gamma}(t_j) = \gamma(t_j)$ and $\tilde{\gamma}(t_{j+1}) = \gamma(t_{j+1})$. In the final step, like in the proof of Proposition 2.24, we argue on $\gamma|_{[t_{k-2}, t_k]}$, since the Riemannian length of $\gamma|_{[t_{k-1}, t_k]}$ cannot be estimated from below. \square

We deduce from Theorem 1.1 a corollary which will be used in the proof of Theorems 1.2 and 1.3.

Corollary 3.5. *Let \mathcal{C} be a stably causal cone structure. For any compact subset $K \subset M$ there are two positive constants ρ_K, l_K such that if \mathcal{C}' is any cone structure on M satisfying $\max_{x \in K} \delta_x^{g, H}(\mathcal{C}_x, \mathcal{C}'_x) \leq \rho_K$ then any \mathcal{C}' -causal curve contained in K has Riemannian length bounded from above by l_K .*

Proof. By Theorem 1.1 there is C^∞ global time function f for \mathcal{C} . Therefore, we can find $\theta > 0$ such that

$$Df(x)[v] \geq \theta \quad \text{for any } x \in K, \text{ and any } v \in \mathcal{C}_x \text{ with } |v|_x = 1.$$

We define $m = \max_K f - \min_K f < +\infty$, and $A = \sup_{x \in K} |Df(x)|_x < +\infty$. We choose $\delta_K > 0$ such that $\eta = \theta - 2A\rho_K > 0$. Assume that $\max_{x \in K} \delta_x^{g, H}(\mathcal{C}_x, \mathcal{C}'_x) \leq \rho_K$. For $x \in K$, and $v' \in \mathcal{C}'_x \cap \partial\mathbb{B}_x$, by Lemma 2.11, applied to the Euclidean space $(T_x M, \|\cdot\|_x)$, we can find $v \in \mathcal{C}_x \cap \partial\mathbb{B}_x$ such that

$$\|v' - v\|_x \leq 2\delta_x^{g, H}(\mathcal{C}_x, \mathcal{C}'_x) \leq 2\rho_K.$$

It follows that

$$\begin{aligned} Df(x)(v') &\geq Df(x)(v) - |Df(x)|_x \|v' - v\|_x \\ &\geq \theta - 2A\rho_K \\ &= \eta > 0. \end{aligned}$$

In particular, if $\gamma : [0, \ell] \rightarrow K$ is a \mathcal{C}' -causal curve parametrized by arc-length, we have

$$Df(\gamma(t))(\dot{\gamma}(t)) \geq \eta, \text{ for almost every } t \in [0, \ell].$$

Integrating on $[0, \ell]$ yields $f(\gamma(\ell)) - f(\gamma(0)) \geq \eta\ell$. Since $\gamma(\ell), \gamma(0) \in K$. This implies $\eta\ell \leq m = \max_K f - \min_K f$. Therefore if we choose $l_K = m/\eta$, we indeed have $\ell_g(\gamma) = \ell \leq l_K$. \square

4 Proof of Theorem 1.2

We recall from the introduction that a cone structure \mathcal{C} defined in M is called globally hyperbolic if it is stably causal and all \mathcal{C} -causal curves connecting two given points have support contained in a compact (possibly empty) subset of M .

We start with some lemmas.

The first one is a particular version of a quite well-known semi-continuity of action for convex functional. For the convenience of the reader we give a more elementary proof of the particular version we need using only Jensen's inequality.

Lemma 4.1. *Let $K, \mathcal{C}, \mathcal{C}^n$ be a compact subset of M and continuous cone structures, respectively, satisfying*

$$\max_{x \in K} \delta_x^H(\mathcal{C}_x, \mathcal{C}_x^n) \rightarrow 0, \text{ as } n \rightarrow +\infty. \quad (27)$$

If $\gamma_n : [a, b] \rightarrow K$ is a sequence of equi-Lipschitz \mathcal{C}^n -causal curves uniformly converging in $[a, b]$ to some curve γ , then γ is \mathcal{C} -causal.

Proof. Since the \mathcal{C} -causality is a local property, we can assume K to be a compact subset of \mathbb{R}^N , and δ^H is the Hausdorff distance on cones obtained from the usual Euclidean norm. Since the sequence γ_n is equi-Lipschitz, we can find a constant $L < +\infty$ such that

$$|\dot{\gamma}_n(t)| \leq L \quad (28)$$

for any n , and any $t \in [a, b]$ for which $\dot{\gamma}_n(t)$ exists.

The Lipschitz path γ is differentiable almost everywhere. We will prove that at a point $t_0 \in]a, b[$, at which the derivative $\dot{\gamma}(t_0)$ exists we have $\dot{\gamma}(t_0) \in \mathcal{C}_{\gamma(t_0)}$.

For this we fix $\epsilon > 0$. By continuity of γ and the cone structure \mathcal{C} we can find $\delta > 0$ such that

$$\delta^H(\mathcal{C}_{\gamma(t)}, \mathcal{C}_{\gamma(t_0)}) \leq \epsilon, \text{ for every } t \in [t_0 - \delta, t_0 + \delta].$$

Since $\max_{x \in K} \delta^H(\mathcal{C}_x, \mathcal{C}_x^n) \rightarrow 0$, as $n \rightarrow +\infty$, we can find n_ϵ such that

$$\max_{x \in K} \delta^H(\mathcal{C}_x, \mathcal{C}_x^n) \leq \epsilon, \text{ for all } n \geq n_\epsilon.$$

By the continuity of $x \mapsto \mathcal{C}_x$ and the fact the γ_n converges uniformly to γ , taking n_ϵ larger if necessary we can also assume that

$$\delta^H(\mathcal{C}_{\gamma_n(t)}, \mathcal{C}_{\gamma(t)}) \leq \epsilon, \text{ for every } n \geq n_\epsilon, \text{ and every } t \in [a, b].$$

It follows that

$$\delta^H(\mathcal{C}_{\gamma_n(t)}, \mathcal{C}_{\gamma(t_0)}) \leq 3\epsilon, \text{ for every } n \geq n_\epsilon, \text{ and every } t \in [t_0 - \delta, t_0 + \delta].$$

Since by (28), we have $|\dot{\gamma}_n(t)| \leq L$, using Proposition 2.13 we get

$$|d(\dot{\gamma}_n(s), \mathcal{C}_{\gamma_n(s)}) - d(\dot{\gamma}_n(s), \mathcal{C}_{\gamma(t_0)})| \leq 3L\epsilon$$

for every $n \geq n_\epsilon$, and every $s \in [t_0 - \delta, t_0 + \delta]$ ant which $\dot{\gamma}_n(s)$ exists. Since $\dot{\gamma}_n(t) \in \mathcal{C}_{\gamma_n(t)}^n$, for almost every $t \in [a, b]$, we obtain

$$d(\dot{\gamma}_n(s), \mathcal{C}_{\gamma(t_0)}) \leq 3L\epsilon,$$

for every $n \geq n_\epsilon$, and almost every $s \in [t_0 - \delta, t_0 + \delta]$. Using that $v \mapsto d(v, \mathcal{C}_{\gamma(t_0)})$ is convex, we obtain from Jensen's inequality that

$$d\left(\frac{\int_t^{t'} \dot{\gamma}_n(s) ds}{t' - t}, \mathcal{C}_{\gamma(t_0)}\right) \leq \frac{\int_t^{t'} d(\dot{\gamma}_n(s), \mathcal{C}_{\gamma(t_0)}) ds}{t' - t} \leq 3L\epsilon,$$

for every $n \geq n_\epsilon$, and every $t, t' \in [t_0 - \delta, t_0 + \delta]$, with $t < t'$. Since γ_n is Lipschitz, we know that $\int_t^{t'} \dot{\gamma}_n(s) ds = \gamma_n(t') - \gamma_n(t)$. It follows that

$$d\left(\frac{\gamma_n(t') - \gamma_n(t)}{t' - t}, \mathcal{C}_{\gamma(t_0)}\right) \leq 3L\epsilon,$$

for every $n \geq n_\epsilon$, and every $t, t' \in [t_0 - \delta, t_0 + \delta]$, with $t < t'$. We can now let $n \rightarrow +\infty$ to obtain

$$d\left(\frac{\gamma(t') - \gamma(t)}{t' - t}, \mathcal{C}_{\gamma(t_0)}\right) \leq 3L\epsilon,$$

for every $t, t' \in [t_0 - \delta, t_0 + \delta]$, with $t < t'$. If we set $t = t_0$, and let $t' \rightarrow t_0$, we obtain

$$d(\dot{\gamma}(t_0), \mathcal{C}_{\gamma(t_0)}) \leq 3L\epsilon.$$

Since $\epsilon > 0$ was arbitrary, and L is a constant that does not depend on ϵ , we can let $\epsilon \rightarrow 0$ to obtain $d(\dot{\gamma}(t_0), \mathcal{C}_{\gamma(t_0)}) = 0$, and therefore $\dot{\gamma}(t_0) \in \mathcal{C}_{\gamma(t_0)}$. \square

Lemma 4.2. *Let \mathcal{C} be a stably causal cone structure. For any compact subset $K \subset M$ and $\varepsilon > 0$ there is a positive constant $\rho = \rho_{K, \varepsilon}$ such that if $\gamma : [a, b] \rightarrow K$ is a \mathcal{C}' -causal curve and $\max_{x \in K} \delta_x^H(\mathcal{C}_x, \mathcal{C}'_x) \leq \rho$ then there exists a \mathcal{C} -causal curve $\hat{\gamma} : [a, b] \rightarrow K$ with $\max_{[a, b]} d_g(\gamma(t), \hat{\gamma}(t)) \leq \varepsilon$.*

Proof. The argument is by contradiction. We assume the existence of a $\varepsilon_0 > 0$, of a sequence \mathcal{C}^n of cone structures with

$$\max_{x \in K} \delta_x^H(\mathcal{C}_x, \mathcal{C}_x^n) \leq \frac{1}{n} \quad \text{for any } n,$$

and of a sequence $\gamma_n : [a_n, b_n] \rightarrow K$ of \mathcal{C}^n -causal curves such that for any n , any \mathcal{C} -causal curve $\gamma : [a_n, b_n] \rightarrow K$

$$\max_{[a_n, b_n]} d_g(\gamma_n(t), \gamma(t)) > \varepsilon_0. \quad (29)$$

We know from Lemma 3.5 that for n large enough, say $n > n_0$, the length of the \mathcal{C}^n -causal curves with support contained in K are equibounded from above by

some constant l_K . We therefore have, up to an orientation preserving change of parameters which does not affect the property of being a \mathcal{C}^n -causal curve, that all such curves, for $n > n_0$, are parametrized by a common interval $[a, b] = [a_n, b_n]$ and they are in addition equi-Lipschitz on it. Therefore, we can apply Ascoli's Theorem to the sequence $\gamma_n, n > n_0$, in order to obtain, up to a subsequence, a uniform limit $\hat{\gamma}$ in $[a, b]$. We derive from Lemma 4.1 that $\hat{\gamma}$ is a \mathcal{C} -causal curve, which is in contradiction with (29). \square

We now give a characterization of the global hyperbolicity.

Proposition 4.3. *The global hyperbolicity property of \mathcal{C} can be equivalently expressed by requiring that the union of the supports of all \mathcal{C} -causal curves with endpoints in a given compact subset of M is compact.*

This result is well known in Lorentzian Geometry, see for instance [10], and can be demonstrated along the same lines in our setting. We supply a proof for reader's convenience. Some definitions and two lemmata are preliminary. Given $x \in M$, we define

$$\begin{aligned} J_{\mathcal{C}}^+(x) &= \{y \in M \mid \text{there exists a } \mathcal{C}\text{-causal curve joining } x \text{ to } y\} \\ J_{\mathcal{C}}^-(x) &= \{y \in M \mid \text{there exists a } \mathcal{C}\text{-causal curve joining } y \text{ to } x\} \end{aligned}$$

Using these sets, the definition of global hyperbolicity of \mathcal{C} can be expressed by requiring that $J_{\mathcal{C}}^+(x) \cap J_{\mathcal{C}}^-(y)$ has compact closure for all pair of points x, y of M . Proposition 4.3 then asserts that this is equivalent to $J_{\mathcal{C}}^+(K) \cap J_{\mathcal{C}}^-(K)$ being compact for any compact subset K of M , where $J_{\mathcal{C}}^+(K) = \cup_{x \in K} J_{\mathcal{C}}^+(x)$ and $J_{\mathcal{C}}^-(K)$ is defined similarly.

Before proving Proposition 4.3, we establish two lemmas.

Lemma 4.4. *Let \mathcal{C} be a globally hyperbolic cone structure. For any y, z in M the set $J_{\mathcal{C}}^+(y) \cap J_{\mathcal{C}}^-(z)$ is compact, possibly empty.*

Proof. Let $J_{\mathcal{C}}^+(y) \cap J_{\mathcal{C}}^-(z)$ be non-empty, and take a sequence x_n belonging to it. There consequently exist \mathcal{C} -causal curves ξ_n defined in $[a_n, b_n]$ with $\xi_n(a_n) = y$, $\xi_n(b_n) = z$, $\xi_n(t_n) = x_n$ for some t_n . By the hyperbolicity assumption all the ξ_n have support contained in some compact set K , therefore, thanks to Corollary 3.5, they have equibounded Riemannian length.

We can therefore assume, up to a change of parameter which does not affect the property of being \mathcal{C} -causal, that such curves are equi-Lipschitz and are defined in a common interval $[a, b]$. By applying Ascoli Theorem we find that the ξ_n uniformly converge, up to a subsequence, in $[a, b]$ to a curve ξ which is still \mathcal{C} -causal by Lemma 4.1. Consequently the x_n converge to some x_0 belonging to the support of ξ , since $\xi(a) = y$ and $\xi(b) = z$, we deduce that the limit point x_0 belongs to $J_{\mathcal{C}}^+(y) \cap J_{\mathcal{C}}^-(z)$. This ends the proof. \square

Lemma 4.5. *Given $x_0 \in M$, there are sequences z_n and y_n converging to x_0 such that the sets $J_{\mathcal{C}}^-(z_n)$, $J_{\mathcal{C}}^+(y_n)$ are both nonincreasing with respect to the inclusion as n goes to $+\infty$, and $x_0 \in J_{\mathcal{C}}^+(y_n) \cap J_{\mathcal{C}}^-(z_n)$ for n large enough.*

Proof. The argument is local in x_0 , so we can assume to be in \mathbb{R}^N . By continuity of \mathcal{C} , we can determine a convex neighborhood W of x_0 and an Euclidean unit vector v_0 with

$$\sup_{x \in W} d^\#(v_0, \mathcal{C}_x) < 0. \quad (30)$$

Therefore $t \mapsto x_0 + tv_0$ is a \mathcal{C} -future directed curve for its part contained in W . We set $y_n = x_0 - 1/n v_0$ and $z_n = x_0 + 1/n v_0$ for n so large that y_n and z_n are in W . By (30), up to further increasing n , the segments starting at (resp. ending to) points suitably close to x_0 and other endpoint at z_n (resp. at y_n) are still \mathcal{C} -future directed, which proves that $x_0 \in J_{\mathcal{C}}^+(y_n) \cap J_{\mathcal{C}}^-(z_n)$, when n is large enough. Finally, the claimed monotonicity of $J_{\mathcal{C}}^-(z_n)$, $J_{\mathcal{C}}^+(y_n)$ is apparent. This concludes the proof. \square

Proof of Proposition 4.3. One implication is clear. It is therefore enough to show that any globally hyperbolic cone structure possesses the property in the statement.

We first deduce from Lemmas 4.4 and 4.5 that $J_{\mathcal{C}}^+(x)$ and $J_{\mathcal{C}}^-(x)$ are closed for any x . Let a point x_0 be the limit of a sequence x_n of $J_{\mathcal{C}}^+(x)$, since $x_0 \in J_{\mathcal{C}}^-(z)$ for a suitable z , then $x_n \in J_{\mathcal{C}}^+(x) \cap J_{\mathcal{C}}^-(z)$ when n is large enough. Taking into account that $J_{\mathcal{C}}^+(x) \cap J_{\mathcal{C}}^-(z)$ is compact by Lemma 4.4, we infer that $x_0 \in J_{\mathcal{C}}^+(x)$, as claimed. Same argument works for $J_{\mathcal{C}}^-(x)$.

The second step is to show that $J_{\mathcal{C}}^+(K)$ and $J_{\mathcal{C}}^-(K)$ are closed for any compact subset K of M . We consider a point x_0 in the closure of $J_{\mathcal{C}}^+(K)$. Bearing in mind Lemma 4.5 and the fact that any neighborhood of x_0 intersects $J_{\mathcal{C}}^+(K)$, we can find a sequence z_n converging to x_0 with $J_{\mathcal{C}}^-(z_n) \cap J_{\mathcal{C}}^+(K) \neq \emptyset$ and nonincreasing in n with respect to the inclusion. This implies that $J_{\mathcal{C}}^-(z_n) \cap K$ are non-empty and, clearly, nonincreasing with respect to the inclusion. By the first part of the proof these sets are, in addition, compact and so possess a non-empty intersection.

If $z \in \bigcap_n J_{\mathcal{C}}^-(z_n) \cap K$ then $z_n \in J_{\mathcal{C}}^+(z)$ for any n and, taking into account that $J_{\mathcal{C}}^+(z)$ is closed and $z \in K$, we find that $x_0 \in J_{\mathcal{C}}^+(z) \subset J_{\mathcal{C}}^+(K)$. Similarly we prove the claim for $J_{\mathcal{C}}^-(K)$.

We can determine, by Lemma 4.5, for any $x \in K$ two points y_x, z_x and an open neighborhood V_x of x with $V_x \subset J_{\mathcal{C}}^-(z_x) \cap J_{\mathcal{C}}^+(y_x)$. Since the V_x make up an open covering of K , we can extract a finite subcover, and consequently determine two finite sets of points $\{y_1, \dots, y_h\}$, $\{z_1, \dots, z_h\}$, for some h in \mathbb{N} , such that $K \subset \bigcup_{i=1}^h J_{\mathcal{C}}^+(y_i)$, $K \subset \bigcup_{j=1}^h J_{\mathcal{C}}^-(z_j)$, which implies $J_{\mathcal{C}}^+(K) \subset \bigcup_{i=1}^h J_{\mathcal{C}}^+(y_i)$, $J_{\mathcal{C}}^-(K) \subset \bigcup_{j=1}^h J_{\mathcal{C}}^-(z_j)$ and

$$J_{\mathcal{C}}^-(K) \cap J_{\mathcal{C}}^+(K) \subset \bigcup_{i,j} (J_{\mathcal{C}}^-(z_j) \cap J_{\mathcal{C}}^+(y_i)).$$

Hence $J_{\mathcal{C}}^-(K) \cap J_{\mathcal{C}}^+(K)$ is contained in a compact subset of M by Lemma 4.4 and so, being in addition closed by the previous step, it is in the end compact. \square

We come to the proof of Theorem 1.2.

We exploit Proposition 4.3 and define inductively an increasing sequence of compact domains $M_n \subset M$ with $\cup M_n = M$, by taking M_1 arbitrarily. We construct $M_n, n > 1$ by induction. Suppose M_{n-1} has been constructed, by Proposition 4.3, we can find a compact subset K_n , such that any \mathcal{C} -causal curve with endpoints in the compact subset $\bar{V}_1(M_{n-1}) = \{x \in M \mid d_g(x, M_{n-1}) \leq 1\}$ is entirely contained in K_n , we then set $M_n = \bar{V}_1(K_n) = \{x \in M \mid d_g(x, K_n) \leq 1\}$.

We denote by N_n the compact subset

$$\begin{aligned} N_n &= M_n && \text{for } n = 1 \text{ or } 2, \\ N_n &= M_n \setminus \overset{\circ}{M}_{n-2} && \text{for } n > 2. \end{aligned}$$

It is clear that the N_n are compact and constitute a locally finite covering of M . By Lemmas 3.5 and 4.2 we can determine, for any n , positive constants l_n, ρ_n , with ρ_n decreasing, such that if a cone structure \mathcal{C}' satisfies $\sup_{x \in N_n} \delta_x^H(\mathcal{C}_x, \mathcal{C}'_x) \leq \rho_n$ then:

- (i) any \mathcal{C}' -causal curve with support contained in N_n has length estimated from above by l_n ;
- (ii) for any \mathcal{C}' -causal curve $\gamma : [a, b] \rightarrow N_n$ there is a \mathcal{C} -causal curve $\hat{\gamma} : [a, b] \rightarrow N_n$ with $\sup_{[a, b]} d_g(\gamma(t), \hat{\gamma}(t)) < 1$.

We can construct through Urysohn's Lemma a continuous function $\rho : M \rightarrow \mathbb{R}^+$ satisfying

$$\max_{x \in N_n} \rho(x) \leq \rho_n. \quad (31)$$

We claim that any cone structure \mathcal{C}' satisfying

$$\mathcal{C}' \succ \mathcal{C} \quad \text{and} \quad \delta_x^H(\mathcal{C}_x, \mathcal{C}'_x) < \rho(x) \quad \text{for any } x \quad (32)$$

is globally hyperbolic. We know from Theorem 2.28 that the family of cone structures satisfying both these properties is non-empty.

We proceed to prove that, for any n , any \mathcal{C}' satisfying (32), a \mathcal{C}' -causal curve joining two points of M_n is contained in M_{n+1} . In fact, if this is not the case, we find an \bar{n} and one of such curves γ joining two points of $\tilde{M}_{\bar{n}}$ with

$$n_0 := \max\{n \mid \gamma \subset M_n\} \geq \bar{n} + 2,$$

so that there is a portion of γ , say γ_0 , defined on the $[a, b]$, contained in $N_{n_0} = M_{n_0} \setminus \overset{\circ}{M}_{n_0-2}$ with non-empty intersection with $M_{n_0} \setminus M_{n_0-1}$, and with its endpoints $\gamma_0(a), \gamma_0(b)$ in ∂M_{n_0-2} .

Since

$$\max_{x \in N_{n_0}} \rho(x) \leq \rho_{n_0}$$

we find a \mathcal{C} -causal curve $\hat{\gamma} : [a, b] \rightarrow N_{n_0}$ with $\sup_{t \in [a, b]} d_g(\gamma_0(t), \hat{\gamma}(t)) < 1$. Therefore the endpoints of $\hat{\gamma}$ are in $\bar{V}_1(M_{n_0-2}) = \{x \in M \mid d_g(x, M_{n_0-2}) \leq 1\}$, since $\gamma_0(a)$ and $\gamma_0(b)$ belong to ∂M_{n_0-2} . Therefore by the choice of K_{n_0-1} , the curve $\hat{\gamma}$ is contained in K_{n_0-1} , and γ_0 is contained in $\bar{V}_1(K_{n_0-1}) = \{x \in M \mid d_g(x, K_{n_0-1}) \leq 1\} = M_{n_0-1}$. This is impossible since γ_0 has a point in $M_{n_0} \setminus M_{n_0-1}$.

We now claim that if \mathcal{C}' satisfies (32) then any \mathcal{C}' -causal curve with support contained in a compact subset of M has finite length. This implies that any such cone structures are causal and consequently stably causal since for any \mathcal{C}' it is possible to find \mathcal{C}'' still satisfying (32) and with $\mathcal{C}'' \succ \mathcal{C}'$.

To prove the claim it suffices to show that for any n , any \mathcal{C}' -causal curve γ contained in M_n there exists a positive constant l'_n with $\ell_g(\gamma) \leq l'_n$. The argument is by induction on n . The property is true for M_1 with $l'_1 = l_1$. Let γ , defined in $[a, b]$, be a \mathcal{C}' -causal curve contained in M_n , if γ is contained in N_n then $\ell_g(\gamma) \leq l_n$ by (31). Otherwise there are points of γ in \tilde{M}_{n-2} , and we set

$$\begin{aligned} a' &= \min\{s \in [a, b] : \gamma(s) \in M_{n-2}\} \\ b' &= \max\{s \in [a, b] : \gamma(s) \in \tilde{M}_{n-2}\} \end{aligned}$$

Since $\gamma|_{[a, a']}$ and $\gamma|_{[b', b]}$, if not reduced to a point, are both contained in N_n , their length is estimated from above by l_n , thanks to (31). The \mathcal{C}' -causal curve $\gamma|_{[a', b']}$ connects two points of M_{n-2} and so it is contained in \tilde{M}_{n-1} by what we proved above, and, by inductive assumption, we know that $\ell_g(\gamma|_{[a', b]}) \leq l'_{n-1}$. We get in any case the estimate

$$\ell_g(\gamma) \leq 2l_n + l'_{n-1} =: l'_n,$$

which concludes the proof of the claim and of the theorem.

5 Proof of Theorem 1.3

Throughout the section we still denote by g and α the Riemannian metric and the C^∞ positive function introduced in Section 3, respectively. We indicate by \mathcal{C}' a globally hyperbolic cone structure with $\mathcal{C}' \succ \mathcal{C}$. The existence of such an object is guaranteed by Theorem 1.2. Theorem 3.4 holds true and, in particular, $\mathcal{A}_{\alpha, g}^{\mathcal{C}'}$ is empty. We will use a result following from [6, Theorem 8.5] or the proof of [7, Theorem 6.6]. For a cone structure, we can introduce, in the same way as we did for \mathcal{C} , the function $d_{\mathcal{C}^*}^{\#g} : \mathcal{T}^*M \rightarrow \mathbb{R}$ defined by

$$\begin{aligned} d_{\mathcal{C}^*}^{\#g}(x, p) &= 2d_x^g(p, \mathcal{C}_x^*) - d_x^g(v, \partial \mathcal{C}_x^*). \\ &= \sup\{p(v) \mid v \in \mathcal{C}_x, |v|_x = 1\}. \end{aligned}$$

This function is locally Lipschitz and convex in p . Moreover, for any $(x, p) \in T^*M$, we have $p \in \mathring{C}_x^*$ if and only if $d_{\mathring{C}^*}^{\#g}(x, p) < 0$.

Theorem 5.1. *Consider a cone structure \mathcal{C} on the manifold M . Assume $\mathcal{A}_{\alpha, g}^{\mathcal{C}}$ to be empty. Suppose that $\epsilon : M \rightarrow]0, +\infty[$ is a continuous function. If $h : M \rightarrow \mathbb{R}$ is a locally Lipschitz function with $d_{\mathring{C}^*}^{\#g}(x, Dh(x)) \leq -\epsilon(x)$ for a.e. $x \in M$, we can construct a C^∞ function \tilde{h} with $d_{\mathring{C}^*}^{\#g}(x, D\tilde{h}(x)) \leq -\epsilon(x)/2$ for all $x \in M$ and, in addition, satisfying $\sup_{x \in M} |h(x) - \tilde{h}(x)| \leq 1$.*

Proof. We introduce the Hamiltonian $H : T^*M \rightarrow \mathbb{R}$ defined by

$$H(x, p) = \frac{d_{\mathring{C}^*}^{\#g}(x, p)}{\epsilon(x)}.$$

It is not difficult to see that H is continuous and convex in p . Moreover, we have $H(x, Dh(x)) \leq -1$, for a.e. $x \in M$. Therefore by [6, Theorem 8.5] or the proof of [7, Theorem 6.6], we can find a C^∞ function $\tilde{h} : M \rightarrow \mathbb{R}$ such that $H(x, D\tilde{h}(x)) \leq -1/2$, and $|h(x) - \tilde{h}(x)| \leq 1$, for every $x \in M$. \square

We will say that a \mathcal{C} -causal curve γ defined in some interval $]a, b[$ is *inextendible* if both $\lim_{t \rightarrow a} \gamma(t)$, $\lim_{t \rightarrow b} \gamma(t)$ do not exist. The first step is a characterization of inextendible \mathcal{C} -causal curves exploiting that \mathcal{C} is stably causal.

Lemma 5.2. *A \mathcal{C} -causal curve γ defined on some interval $]a, b[$ is inextendible if and only if it does not admit any limit point for $t \rightarrow a$ and $t \rightarrow b$.*

In particular, since we are assuming that the Riemannian metric on M is complete, if the \mathcal{C} -causal curve $\gamma : I \rightarrow M$ is inextendible and parametrized by arc-length, then $I =]-\infty, +\infty[$.

Proof. Let γ be \mathcal{C} -causal and inextendible. If the statement is false for $t \rightarrow b$, there are, by the very definition of inextendible curve, two distinct limit points x and y , and one can select times $t_1 < t_2 < t_3$ in such a way that $\gamma(t_1)$, $\gamma(t_3)$ are arbitrarily close to x and $\gamma(t_2)$ is arbitrarily close to y . We can form a closed loop ξ based at x by juxtaposition of the Riemannian geodesic between x and $\gamma(t_1)$, $\gamma|_{[t_1, t_3]}$, and the Riemannian geodesic between $\gamma(t_3)$ and x . Without loss of generality, we can assume that the points $\gamma(t_1)$ and $\gamma(t_3)$ are contained in the compact Riemannian $\bar{B}(x, 1)$. It follows that we can find a constant K such that

$$\int_{\xi} \alpha d_{\mathcal{C}}^g \leq K[d_g(x, \gamma(t_1)) + d_g(x, \gamma(t_3))] + \int_{\gamma|_{[t_1, t_3]}} \alpha d_{\mathcal{C}}^g.$$

Since $\gamma|_{[t_1, t_3]}$ is causal, the last term in the right hand side above is 0, and

$$\int_{\xi} \alpha d_{\mathcal{C}}^g \leq K[d_g(x, \gamma(t_1)) + d_g(x, \gamma(t_3))].$$

Therefore by choosing appropriately, we can assume $\int_{\xi} \alpha d_{\mathcal{C}}^g$ as small as we want. On the other hand $\ell_g(\xi) \geq 2d_g(x, \gamma(t_3))$. Since $\gamma(t_3)$ is as close as we want to y , we can assume that $\ell_g(\xi) \geq d_g(x, y)$. It follows that we can arrange ξ to satisfy

$$\int_{\xi} \alpha d_{\mathcal{C}}^g < \min \left\{ \ell_g(\xi), \frac{1}{2} \right\}.$$

This entails, in view of Theorem 3.4, that there exists a \mathcal{C}' -future directed cycle based at x , which is impossible since \mathcal{C}' is causal.

The converse implication of the statement is immediate. □

We break the body of the proof of Theorem 1.3 in two lemmas and a final part. We use Proposition 4.3 applied to \mathcal{C}' in the following construction of an increasing sequence of compact domains W_n .

The compact domains $W_n \subset M$, with $\cup_i W_i = M$, are defined taking W_1 arbitrarily but non-empty, and choosing, for $n > 1$, $W_n = \bar{V}_1(K_n) = \{x \in M \mid d_g(x, K_n) \leq 1\}$, where K_n is a compact subset containing $\bar{V}_1(W_{n-1}) = \{x \in M \mid d_g(x, W_{n-1}) \leq 1\}$, and such that any \mathcal{C}' -causal curve connecting two points of W_{n-1} is contained in K_n .

We will essentially use that $\mathcal{C}' \succ \mathcal{C}$ is globally hyperbolic to show that the W_n enjoy the crucial property given in the following:

Lemma 5.3. *Let y_1, y_2 be in W_{n-1} , for some $n \geq 2$, and take $x \notin W_n$, then*

$$\max\{S_{\alpha,g}^{\mathcal{C}}(y_1, x), S_{\alpha,g}^{\mathcal{C}}(x, y_2)\} \geq 1/2$$

Proof. We assume by contradiction

$$S_{\alpha,g}^{\mathcal{C}}(y_1, x), S_{\alpha,g}^{\mathcal{C}}(x, y_2) < 1/2, \tag{33}$$

by the very definition of W_n we have

$$d_g(y_1, x), d_g(x, y_2) > 1. \tag{34}$$

The inequalities (33), (34) imply, by Theorem 3.4, that there are \mathcal{C}' -future directed curves from y_1 to x and from x to y_2 , so that there exists a \mathcal{C}' -causal curve connecting y_1 to y_2 and passing through x . From this we, in turn, derive that $x \in W_n$, in contradiction with the assumption. □

We now consider a smooth function $\psi : M \rightarrow \mathbb{R}$ satisfying

$$\begin{aligned} \psi &\geq 2\alpha \quad \text{on } W_1; \\ \psi &\geq 2n\alpha \quad \text{on } W_n \setminus W_{n-1}, \text{ for } n \geq 2. \end{aligned}$$

We define $S_{\psi,g}^{\mathcal{C}}$ through formula (21) with ψ in place of α . Note that, since $\psi > \alpha$ on M , the Aubry set $\mathcal{A}_{\psi,g}^{\mathcal{C}}$ is empty, being $\mathcal{A}_{\alpha,g}^{\mathcal{C}} = \emptyset$ in force of Theorem 3.4. We deduce from the very definition of ψ and Lemma 5.3

Lemma 5.4. *Let $\gamma :] - \infty, +\infty[$ be a \mathcal{C} -causal inextendible curve parametrized by the Riemannian arc-length, then*

$$\lim_{t \rightarrow +\infty} S_{\psi,g}^{\mathcal{C}}(\gamma(t), x) = \lim_{t \rightarrow -\infty} S_{\psi,g}^{\mathcal{C}}(x, \gamma(t)) = +\infty \quad (35)$$

for any fixed $x \in M$.

Proof. It suffices to show (35) for $x = \gamma(0)$, in fact by the triangle inequality

$$S_{\psi,g}^{\mathcal{C}}(\gamma(t), x) \leq S_{\psi,g}^{\mathcal{C}}(\gamma(t), \gamma(0)) + S_{\psi,g}^{\mathcal{C}}(\gamma(0), x),$$

for any t , and a similar inequality applies to $S_{\psi,g}^{\mathcal{C}}(x, \gamma(t))$.

We select an n so large that $\gamma(0) \in W_{n-1}$. By Lemma 5.2, $\gamma(t) \notin W_n$, whenever $|t|$ is sufficiently large. Given a $t_0 > 0$ enjoying such a property, we claim that

$$S_{\psi,g}^{\mathcal{C}}(\gamma(t_0), \gamma(0)) \geq n. \quad (36)$$

To prove it, we fix $\varepsilon > 0$, and select an ε -optimal curve $\xi : [a, b] \rightarrow M$ for $S_{\psi,g}^{\mathcal{C}}(\gamma(t_0), \gamma(0))$. We set

$$t_1 = \min\{t \in [a, b] \mid \xi(t) \in W_{n-1}\} \quad \text{and} \quad \xi_1 = \xi|_{[a, t_1]}.$$

We derive from Lemma 5.3 and the fact that γ is \mathcal{C} -causal

$$\frac{1}{2} \leq \max\{S_{\alpha,g}^{\mathcal{C}}(\gamma(0), \gamma(t_0)), S_{\alpha,g}^{\mathcal{C}}(\gamma(t_0), \xi(t_1))\} = S_{\alpha,g}^{\mathcal{C}}(\gamma(t_0), \xi(t_1)). \quad (37)$$

Since ξ_1 lies outside W_{n-1} , we have $\int_{\xi_1} \psi \sigma_g^{\mathcal{C}} \geq 2n \int_{\xi_1} \alpha \sigma_g^{\mathcal{C}}$, and we deduce from this inequality, the ε -optimality of ξ and (37)

$$\begin{aligned} \frac{1}{2} &\leq S_{\alpha,g}^{\mathcal{C}}(\gamma(t_0), \xi(t_1)) \leq \int_{\xi_1} \alpha \sigma_g^{\mathcal{C}} \leq \frac{1}{2n} \int_{\xi_1} \psi \sigma_g^{\mathcal{C}} \\ &\leq \frac{1}{2n} \int_{\xi} \psi \sigma_g^{\mathcal{C}} \leq \frac{1}{2n} (S_{\psi,g}^{\mathcal{C}}(\gamma(t_0), \gamma(0)) + \varepsilon), \end{aligned}$$

which gives (36), since ε has been arbitrarily chosen. This, in turn, implies

$$\lim_{t \rightarrow +\infty} S_{\psi,g}^{\mathcal{C}}(\gamma(t), x) = +\infty.$$

The same argument applies, with obvious adaptation, to prove the limit relation for t going to $-\infty$. □

We come to the final part of the proof of Theorem 1.3. To simplify notation, we fix some x_0 , and we introduce the locally Lipschitz function $f_1 : M \rightarrow \mathbb{R}$ defined by

$$f_1(x) = S_{\psi,g}^{\mathcal{C}}(x, x_0) - S_{\psi,g}^{\mathcal{C}}(x_0, x).$$

By Proposition 3.2, we have $-DS_{\alpha,g}^{\mathcal{C}}(x, x_0) \in \mathcal{C}_x^*$, $DS_{\alpha,g}^{\mathcal{C}}(x_0, x) \in \mathcal{C}_x^*$ for a.e. x . It follows that $-Df_1(x) \in \mathcal{C}_x^*$ for a.e. $x \in M$. Moreover for any \mathcal{C} -causal curve $\gamma : I \rightarrow M$, the $t \mapsto f_1(\gamma(t))$ is non-decreasing, since this is the case for both $S_{\psi,g}^{\mathcal{C}}(\gamma(t), x_0)$ and $-S_{\psi,g}^{\mathcal{C}}(x_0, \gamma(t))$. It therefore follows from Lemma 5.4, that for any inextendible \mathcal{C} -causal curve $\gamma :]-\infty, +\infty[\rightarrow M$, we have

$$\lim_{t \rightarrow +\infty} f_1(\gamma(t)) = +\infty \text{ and } \lim_{t \rightarrow -\infty} f_1(\gamma(t)) = -\infty.$$

We now pick a C^∞ global time function f_0 for \mathcal{C} provided by Theorem 1.1 in correspondence to α . We set

$$f(x) = f_0(x) + f_1(x),$$

Because $t \mapsto f_0(\gamma(t))$ is also non-decreasing along a \mathcal{C} -causal curve γ , we obtain that

$$\lim_{t \rightarrow +\infty} f(\gamma(t)) = +\infty \text{ and } \lim_{t \rightarrow -\infty} f(\gamma(t)) = -\infty, \quad (38)$$

for any inextendible \mathcal{C} -causal curve $\gamma :]-\infty, +\infty[\rightarrow M$. We have that $-Df_0(x) \in \mathring{\mathcal{C}}_x^*$ for any $x \in M$ and $-Df_1(x) \in \mathcal{C}_x^*$, or a.e. $x \in M$. Since $(x, p) \rightarrow d_{\mathcal{C}^*}^{g\#}(x, p)$ is convex and positively homogeneous in p , it is subadditive in p . We therefore get for a.e. $x \in M$

$$\begin{aligned} d_{\mathcal{C}^*}^{g\#}(x, -Df(x)) &\leq d_{\mathcal{C}^*}^{g\#}(x, -Df_0(x)) + d_{\mathcal{C}^*}^{g\#}(x, -Df_1(x)) \\ &\leq d_{\mathcal{C}^*}^{g\#}(x, -Df_0(x)). \end{aligned}$$

Note that since f_0 is C^1 and $-Df_0(x) \in \mathring{\mathcal{C}}_x^*$ for every $x \in M$, the function $\epsilon(x) = -d_{\mathcal{C}^*}^{g\#}(x, -Df_0(x))$ is continuous and (strictly) positive everywhere on M .

Starting from $h = -f$, Theorem 5.1 finally allows us to find a smooth Cauchy time function going from $-\infty$ to $+\infty$ on any inextendible \mathcal{C} -causal curve, as claimed.

The proof of Theorem 1.3 is therefore concluded.

References

- [1] J.P. AUBIN, A. CELLINA *Differential Inclusion*. Grundlehren der mathematischen Wissenschaften **264** Springer-Verlag, Berlin, 1984.
- [2] J.K. BEEM, P.E. EHRLICH & K.L. EASLEY *Global Lorentzian geometry*. Monograph and Textbooks in Pure Appl. Math. **202** Marcel Dekker Inc., New York, 1996.
- [3] G. BEER, *Topologies on closed and closed convex sets*, Kluwer Academic Publisher, Dordrecht, 1993

- [4] A.N BERNAL, M. SÁNCHEZ “On smooth Cauchy hypersurfaces and Geroch’s splitting theorem”. *Comm. Math Phys*, **243** (2003), 461–470.
- [5] J. DATTORRO, *Convex optimization and Euclidean distance geometry*, Meboo Publishing, USA, 2005.
- [6] A. FATHI, E. MADERNA “Weak KAM theorem on non compact manifolds”, *Nonlinear differ. equ. appl.*, **14** (2007), 1–27.
- [7] A. FATHI, A. SICONOLFI, “PDE aspects of Aubry-Mather theory for quasi-convex Hamiltonians”, *Calc. Var. Partial Differential Equations*, **22** (2005), 185–228.
- [8] A. FATHI, *Weak KAM Theorem in Lagrangian Dynamics*, Cambridge University Press, to appear.
- [9] R. GEROCH, “Domain of dependence”. *J. Math. Phys* **11** (1970), 437–449.
- [10] S.W. HAWKING, G.F.R ELLIS *The large scale structure of space-time*. Cambridge Monographs on Mathematical Physics, Cambridge University Press, London–New York, 1973.
- [11] B. O’NEILL *Semi-Riemannian Geometry*. Academic Press London–New York, 1983.
- [12] A. SICONOLFI, G. TERRONE, “A Metric Approach to the Converse Lyapunov Theorem for Continuous Multivalued Dynamics”, *Nonlinearity*, **20** (2007), 1077–1093.

Albert Fathi, Unité de mathématiques pures et appliquées, CNRS UMR 5669 & École Normale Supérieure de Lyon, 46 allée d’Italie, 69364 Lyon, France
afathi@umpa.ens-lyon.fr

Antonio Siconolfi, Dipartimento di Matematica, Università di Roma “La Sapienza”. Piazzale A. Moro, 2, 00185 Roma, Italy
siconolf@mat.uniroma1.it