

Application of timestepping schemes based on time discontinuous Galerkin methods to multi-dimensional examples

Thorsten Schindler, Vincent Acary

► **To cite this version:**

Thorsten Schindler, Vincent Acary. Application of timestepping schemes based on time discontinuous Galerkin methods to multi-dimensional examples. *Euromech 514 - New trends in Contact Mechanics*, Mar 2012, Cargèse, France. 2012. <hal-00691477>

HAL Id: hal-00691477

<https://hal.inria.fr/hal-00691477>

Submitted on 26 Apr 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Application of timestepping schemes based on time discontinuous Galerkin methods to multi-dimensional examples

Thorsten Schindler¹, Vincent Acary²

¹ *Institute of Applied Mechanics, Technische Universität München, Boltzmannstraße 15, 85748 Garching, Germany, thorsten.schindler@mytum.de*

² *INRIA Grenoble, 655 avenue de l'Europe, Montbonnot, 38334 Saint Ismier Cedex, France, vincent.acary@inrialpes.fr*

Summary: Classic timestepping schemes for nonsmooth dynamical systems are locally of integration order one both in smooth and nonsmooth periods. A consistent enhancement based on time discontinuous Galerkin methods and splitting of smooth and nonsmooth force propagation is presented. Convergence and general behaviour are discussed with different multi-dimensional examples with and without considering friction.

Introduction

In nonsmooth mechanical systems, the velocity function contains smooth and nonsmooth propagation episodes. With classic function derivatives, one has to distinguish these two ranges. Measure theory is the correct mathematical tool which yields a uniform description. The interpretation in the sense of distributions permits the following problem formulation.

Problem 1 (Distribution differential inclusion) *Solve*

$$q(0) := q_0, \quad (1)$$

$$v(0) := v_0, \quad (2)$$

$$\langle \dot{q}, \varphi_q \rangle_{\mathcal{D}^*, \mathcal{D}} = \langle v, \varphi_q \rangle_{\mathcal{D}^*, \mathcal{D}}, \quad \forall \varphi_q \in \mathcal{D}(I), \quad (3)$$

$$\langle D^1 v, \varphi_v \rangle_{\mathcal{D}^*, \mathcal{D}} = \langle m^{-1} f, \varphi_v \rangle_{\mathcal{D}^*, \mathcal{D}} + \langle m^{-1} di, \varphi_v \rangle_{\mathcal{D}^*, \mathcal{D}}, \quad \forall \varphi_v \in \mathcal{D}(I), \quad (4)$$

$$\text{contact and impact relations } (q, v, di, t) \in \mathcal{N}. \quad (5)$$

The initial values of position q and velocity v are given by q_0 and v_0 , respectively. The position offers a classical time derivative almost everywhere. On the other hand, non-regular reaction measures di require to use the distributional derivative $D^1 v$ of v . The space of all real-valued \mathcal{C}^∞ -functions with compact support contains the test functions for position φ_q and velocity φ_v . It is denoted by $\mathcal{D}(I)$. Its dual space $\mathcal{D}^*(I)$ contains distributions. Inverse mass m^{-1} , external force f as well as time t can be interpreted in this general setting using the notion of the primal-dual pairing $\langle *, * \rangle_{\mathcal{D}^*, \mathcal{D}}$.

Classical timestepping schemes discretise the equations of motion of Problem 1 including the constraints with integration order one. This avoids impact detections: a large number of contact transitions can be handled efficiently. Event-driven schemes resolve the exact contact transition times. Between the events, the motion of the system is computed by a classic integration method for differential algebraic equations. This is very accurate but the detection of events can be time consuming and is not possible for Zeno phenomena.

Time Discontinuous Galerkin Methods

To consistently improve the behaviour during smooth episodes, classical timestepping schemes are embedded in time discontinuous Galerkin methods. In Problem 1, one assumes that ansatz

and test functions might have jumps across discretisation intervals, and that they are of higher order inside discretisation intervals. The second assumption states that there is not an instantaneous influence of the analytic nonsmooth dynamics on the numerical solution: the exact time of discontinuity is not resolved. Figure 1 shows that one can define the location of discontinuities of velocity and interaction impulses at the left or right interval border. Piecewise constant




Figure 1: Piecewise smooth propagation with left and right discontinuity.

functions and right interaction impulse discontinuity yield classic explicit (right velocity discontinuity) and implicit (left velocity discontinuity) timestepping schemes.

For each interval $[t_i, t_{i+1}]$, we use splitting techniques

$$di := rdt + p_{i+1}\delta_{t_{i+1}}$$

to take care of contact reactions r separated from the single impact reaction p_{i+1} always evaluated at t_{i+1} with the Dirac measure $\delta_{t_{i+1}}$ because of stability reasons. Hence, contacts benefit from higher order ansatz functions and impacts yield local integration order one.

Numerical Experiments

We analyse convergence and general behaviour of a D^{1-} timestepping scheme (right velocity discontinuity, linear velocity ansatz function) and the Moreau-Jean timestepping scheme (constant velocity ansatz function) with different examples with and without considering friction (e.g. bouncing ball example, slider-crank mechanism [2], circuit breaker, cam follower).




Figure 2: Convergence diagram (bouncing ball): free flight and finite accumulation of impacts.

References

- [1] Thorsten Schindler and Vincent Acary. Timestepping schemes for nonsmooth dynamics based on discontinuous Galerkin methods: definition and outlook. Research Report RR-7625, INRIA, 2011. <http://hal.inria.fr/inria-00595460/en/>.
- [2] Paulo Flores, Remco Leine, and Christoph Glocker. Modeling and analysis of planar rigid multibody systems with translational clearance joints based on the non-smooth dynamics approach. *Multibody System Dynamics*, 23:165–190, 2010.