



**HAL**  
open science

## Finding an induced subdivision of a digraph.

Jørgen Bang-Jensen, Frédéric Havet, Nicolas Trotignon

► **To cite this version:**

Jørgen Bang-Jensen, Frédéric Havet, Nicolas Trotignon. Finding an induced subdivision of a digraph.. Theoretical Computer Science, 2012, 443, pp.10–24. hal-00749187

**HAL Id: hal-00749187**

**<https://inria.hal.science/hal-00749187>**

Submitted on 23 Oct 2016

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Finding an induced subdivision of a digraph\*

Jørgen Bang-Jensen<sup>†</sup>      Frédéric Havet<sup>‡</sup>      Nicolas Trotignon<sup>§</sup>

March 10, 2012

## Abstract

We consider the following problem for oriented graphs and digraphs: Given an oriented graph (digraph)  $G$ , does it contain an induced subdivision of a prescribed digraph  $D$ ? The complexity of this problem depends on  $D$  and on whether  $G$  must be an oriented graph or is allowed to contain 2-cycles. We give a number of examples of polynomial instances as well as several NP-completeness proofs.

**Keywords:** NP-completeness, induced paths and cycles, linkings, 3-SAT.

## 1 Introduction

Many interesting classes of graphs are defined by forbidding induced subgraphs, see [4] for a survey. This is why the detection of several kinds of induced subgraphs is interesting, see [7] where several such problems are surveyed. In particular, the problem of deciding whether a graph  $G$  contains, as an induced subgraph, some graph obtained after possibly subdividing prescribed edges of a prescribed graph  $H$  has been studied. This problem can be polynomial or NP-complete depending on  $H$  and to the set of edges that can be subdivided. The aim of the present work is to investigate various similar problems in digraphs, focusing only on the following problem: given a digraph  $H$ , is there a polynomial algorithm to decide whether an input digraph  $G$  contains a subdivision of  $H$ ?

Of course the answer depends heavily on what we mean by “contain”. Let us illustrate this by surveying what happens in the realm of non-oriented graphs. If the containment relation is the subgraph containment, then for any fixed  $H$ , detecting a subdivision of  $H$  in an input graph  $G$  can be performed in polynomial time by the Robertson and Seymour linkage algorithm [9] (for a short explanation of this see e.g. [2]). But if we want to detect an *induced* subdivision of  $H$  then the answer depends on  $H$  (assuming  $P \neq NP$ ). It is proved in [7] that detecting an induced subdivision of  $K_5$  is NP-complete, and the argument can be reproduced for any  $H$  whose minimum degree is at least 4. Polynomial-time solvable instances trivially exist, such as detecting an induced subdivision of  $H$  when  $H$  is a path, or a graph on at most 3 vertices. But non-trivial polynomial-time solvable instances also exist, such as detecting an induced subdivision of  $K_{2,3}$  that can be performed in time

---

\*This work was done while the first author was on sabbatical at Team Mascotte, INRIA, Sophia Antipolis France whose hospitality is gratefully acknowledged. Financial support from the Danish National Science research council (FNU) (under grant no. 09-066741) is gratefully acknowledged.

<sup>†</sup>Department of Mathematics and Computer Science, University of Southern Denmark, Odense DK-5230, Denmark (email: jbj@imada.sdu.dk).

<sup>‡</sup>Projet Mascotte, I3S (CNRS, UNSA) and INRIA, Sophia Antipolis, France. Partly supported by ANR Blanc AGAPE ANR-09-BLAN-0159. (email:Frederic.Havet@sophia.inria.fr).

<sup>§</sup>CNRS, LIP – ENS Lyon, France. Partially supported by the French *Agence Nationale de la Recherche* under reference ANR 10 JCJC 0204 01. (email: nicolas.trotignon@ens-lyon.fr).

1  $O(n^{11})$  by the Chudnovsky and Seymour's three-in-a-tree algorithm, see [5]. Note that for many  
 2 graphs  $H$ , nothing is known about the complexity of detecting an induced subdivision of  $H$ : when  $H$   
 3 is cubic (in particular when  $H = K_4$ ) or when  $H$  is a disjoint union of 2 triangles, and in many other  
 4 cases.

5 When we move to digraphs, the situation becomes more complicated, even for the subdigraph  
 6 containment relation. All the digraphs we will consider here are *simple*, i.e. they have no loops nor  
 7 multiple arcs. We rely on [1] for classical notation and concepts. A *subdivision of a digraph  $D$* , also  
 8 called a  *$D$ -subdivision*, is a digraph obtained from  $D$  by replacing each arc  $ab$  of  $D$  by a directed  
 9  $(a, b)$ -path. From the NP-completeness of the 2-linkage problem, proved by Fortune, Hopcroft and  
 10 Wyllie [6], it is straightforward to construct an oriented graph  $H$  such that deciding whether a given  
 11 oriented graph  $G$  contains a subdivision of  $H$  as a subgraph is NP-complete. See Theorem 33.

12 Let us now think about the induced subdigraph relation. An induced subdigraph of a digraph  $G$   
 13 which is a subdivision of  $D$  is called an *induced subdivision* of  $D$ . When  $D$  is a digraph, we define:

14 **PROBLEM  $\Pi_D$**

15 Input: A digraph  $G$ .

16 Question: Does  $G$  contain an induced subdivision of  $D$ ?

17 In  $\Pi_D$ , the instance digraph  $G$  may have (directed) 2-cycles, where the 2-cycle is the digraph  
 18  $C_2$  on 2 vertices  $a, b$  with 2 arcs  $ab$  and  $ba$ . Because of these 2-cycles, NP-completeness results  
 19 are often quite easy to obtain, because no induced directed path can go through a 2-cycle (which  
 20 by itself contains a chord). Hence 2-cycles are very convenient to force an induced directed path  
 21 to go through many places of a large digraph that models an instance of 3-SAT. This yields NP-  
 22 completeness results that cover large classes of detection problems. See Section 4. In fact, it can be  
 23 easily shown (see Section 2) that if  $D$  is the disjoint union of *spiders* (trees obtained from disjoint  
 24 directed paths by identifying one end of each path into a vertex) and at most one 2-cycle, then  $\Pi_D$   
 25 is polynomial-time solvable. However, except from those digraphs, we are not aware of any  $D$  for  
 26 which  $\Pi_D$  is polynomial time solvable. We indeed conjecture that there are none. As an evidence, we  
 27 show that if  $D$  is an *oriented graph*, i.e. a digraph with no 2-cycles, then  $\Pi_D$  is NP-complete unless it  
 28 is the disjoint union of spiders (see Corollary 13).

29 It seems that allowing or not allowing 2-cycles is an essential distinction. Hence we also consider  
 30 the restricted problem  $\Pi'_D$  in which the input graph  $G$  is an oriented graph.

31 **PROBLEM  $\Pi'_D$**

32 Input: An oriented graph  $G$ .

33 Question: Does  $G$  contain an induced subdivision of  $D$ ?

34 Observe that if  $\Pi_D$  is polynomial-time solvable then  $\Pi'_D$  is also polynomial-time solvable. Con-  
 35 versely, if  $\Pi'_D$  is NP-complete then  $\Pi_D$  is also NP-complete. Hence, NP-completeness results cover  
 36 less cases for  $\Pi'_D$ .

37 Similarly to  $\Pi_D$ , for several  $D$ 's,  $\Pi'_D$  is solvable by very simple polynomial-time algorithms (See  
 38 Section 2). However, in this case they are not the only ones. We could obtain several digraphs for  
 39 which  $\Pi'_D$  is solvable in polynomial time with non-trivial algorithms.

40 We denote by  $TT_3$  the transitive tournament on 3 vertices  $a, b, c$  and arcs  $ab, ac, bc$ . In Subsec-  
 41 tion 5.1, we use a variant of Breadth First Search that computes only induced trees to solve  $\Pi'_{TT_3}$  in  
 42 polynomial time.

43 We also study oriented paths in Subsection 5.2. An *oriented path* is an orientation of a path. The  
 44 *length* of an oriented path  $P$  is its number of arcs and is denoted  $l(P)$ . Its first vertex is called its *origin*  
 45 and its last vertex its *terminus*. The *blocks* of an oriented paths are its maximal directed subpaths. We

1 denote by  $A_k^-$  the path on vertices  $s_1, s_2, \dots, s_k, s_{k+1}$  and arcs  $s_2s_1, s_2s_3, s_4s_3, s_4s_5, \dots$  and  $A_k^+$  the path  
2 on vertices  $s_1, s_2, \dots, s_k, s_{k+1}$  and arcs  $s_1s_2, s_3s_2, s_3s_4, s_5s_4, \dots$ . These two paths are the *antidirected*  
3 *paths* of length  $k - 1$ . Observe that  $A_k^-$  is the converse of  $A_k^+$  (i.e. it is obtained from  $A_k^+$  by reversing  
4 all the arcs); if  $k$  is odd they are isomorphic but the origin and terminus are exchanged. Clearly, an  
5 oriented path with  $k$ -blocks can be seen as a subdivision of  $A_k^-$  or  $A_k^+$ . In particular, paths with one  
6 block are the directed paths. We show that if  $P$  is an oriented path with three blocks such that the  
7 last one has length one then  $\Pi_P$  is polynomial-time solvable. We also use classical flow algorithms to  
8 prove that  $\Pi'_{A_4^-}$  is polynomial-time solvable.

9 If  $D$  is any of the two tournaments on 3 vertices, namely the directed 3-cycle  $C_3$  and the transitive  
10 tournament  $TT_3$ , then  $\Pi'_D$  is polynomial time solvable. Hence it is natural to study the complexity  
11 of larger tournaments. In Section 6, it is shown that if  $D$  is a transitive tournament on more than 3  
12 vertices or the strong tournament on 4 vertices, then  $\Pi'_D$  is NP-complete.

13 Finally, in Section 7, we point out several open questions.

## 14 2 Easily polynomial-time solvable problems

15 There are digraphs  $D$  for which  $\Pi_D$  or  $\Pi'_D$  can be easily proved to be polynomial-time solvable. For  
16 example, it is the case for the directed  $k$ -path  $P_k$  on  $k$  vertices. Indeed, a  $P_k$ -subdivision is a directed  
17 path of length at least  $k - 1$  and an induced directed path of length at least  $k - 1$  contains an induced  
18  $P_k$ . Hence a digraph has a  $P_k$ -subdivision if and only if it has  $P_k$  as an induced subdigraph. This can  
19 be checked in time  $O(n^k)$  by checking for every set of  $k$  vertices whether or not it induces a  $P_k$ .

20 A vertex of a digraph is a *leaf* if its degree is one, a *node* if its out-degree or its in-degree is at least  
21 2, and a *continuity* otherwise, that is if both its out- and in-degree equal 1. A *spider* is a tree having  
22 at most one node.

23 **Proposition 1.** *If  $D$  is the disjoint union of spiders then  $\Pi_D$  is polynomial-time solvable.*

24 *Proof.* A digraph  $G$  contains an induced  $D$ -subdivision if and only if it contains  $D$  as an induced  
25 subdigraph. This can be checked in time  $O(n^{|V(D)|})$ .  $\square$

26 It is also not difficult to see that  $\Pi_{C_2}$  is polynomial-time solvable.

27 **Proposition 2.**  *$\Pi_{C_2}$  is polynomial-time solvable.*

28 *Proof.* A subdivision of the directed 2-cycle is a directed cycle. In a digraph, a shortest cycle is  
29 necessarily induced, hence a digraph has a  $C_2$ -subdivision if and only if it is not acyclic. Since  
30 one can check in linear time if a digraph is acyclic or not [1, Section 2.1],  $\Pi_{C_2}$  is polynomial-time  
31 solvable.  $\square$

32 Since an oriented graph contains no 2-cycle, then  $\Pi'_{C_2} = \Pi'_{C_3}$ . Similarly to  $\Pi_{C_2}$ , this problem is  
33 polynomial-time solvable.

34 **Proposition 3.**  *$\Pi'_{C_3}$  is polynomial-time solvable.*

35 *Proof.* An oriented graph contains an induced subdivision of  $C_3$  if and only if it is not acyclic.  $\square$

36 Moreover, the following is polynomial-time solvable.

37 **Proposition 4.** *If  $D$  is the disjoint union of spiders and a  $C_2$  then  $\Pi_D$  is polynomial-time solvable.*

38 *Proof.*  $D' = D - C_2$  is a collection of spiders. Let  $p$  be its order. For each set  $A$  of  $p$  vertices, we  
39 check if the digraph  $G\langle A \rangle$  induced by  $A$  is  $D'$  and if yes we check if  $G - (A \cup N(A))$  has a directed  
40 cycle.  $\square$

1 Similarly,

2 **Proposition 5.** *If  $D$  is the disjoint union of spiders and a  $C_3$  then  $\Pi'_D$  is polynomial-time solvable.*

### 3 NP-completeness results for oriented graphs

4 In all proofs below it should be clear that the reductions can be performed in polynomial time and  
5 hence we omit saying this anymore. Before starting with the NP-completeness proofs, we state a  
6 proposition.

7 **Proposition 6.** *Let  $D$  be a digraph and  $C$  a connected component<sup>1</sup> of  $D$ . If  $\Pi_C$  is NP-complete then  
8  $\Pi_D$  is NP-complete. Similarly, if  $\Pi'_C$  is NP-complete then  $\Pi'_D$  is NP-complete.*

9 *Proof.* Let  $D_1, \dots, D_k$  be the components of  $D$  and assume that  $\Pi_{D_1}$  is NP-complete. To show that  
10  $\Pi_D$  is NP-complete, we will give a reduction from  $\Pi_{D_1}$  to  $\Pi_D$ .

11 Let  $G_1$  be an instance of  $\Pi_{D_1}$  and  $G$  be the digraph obtained from  $D$  by replacing  $D_1$  by  $G_1$ . We  
12 claim that  $G$  has an induced  $D$ -subdivision if and only if  $G_1$  has an induced  $D_1$ -subdivision.

13 Clearly, if  $G_1$  has an induced  $D_1$ -subdivision  $S_1$  then the disjoint union of  $S_1$  and the  $D_i$ ,  $2 \leq i \leq k$   
14 is an induced  $D$ -subdivision in  $G$ .

15 Reciprocally, assume that  $G$  contains an induced  $D$ -subdivision  $S$ . Let  $S_i$ ,  $1 \leq i \leq k$  be the con-  
16 nected components of  $S$  such that  $S_i$  is an induced  $D_i$ -subdivision. Set  $G_i = D_i$  if  $i \geq 2$ . Then the  
17  $G_i$ 's are the connected components of  $G$ . Thus  $S_1$  is contained in one of the  $G_i$ 's. If it is  $G_1$  then  
18 we have the result. Otherwise, it is contained in some other component say  $G_2 = D_2$ . In turn,  $S_2$   
19 is contained in some  $G_j$ . Hence  $G_j$  contains a  $D_1$ -subdivision because  $S_2$  contains a  $D_1$ -subdivision  
20 since  $D_2$  contains  $S_1$ . Thus  $G_j$  cannot be  $G_2$  since  $G_2$  already contains  $D_1$  and  $|S_2| \geq |G_2|$ . If  $j = 1$   
21 then we have the result. If not we may assume that  $j = 3$ . And so on, for every  $i \geq 3$ , applying the  
22 same reasoning, we show that one of the following occurs:

- 23 •  $S_i$  is contained in  $G_1$  and thus  $G_1$  contains a  $D_1$ -subdivision because  $S_i$  did.
- 24 •  $S_i$  is contained in  $G_j$  which cannot be any of the  $G_i$ ,  $1 \leq l \leq i$ , for cardinality reasons. Hence  
25 we may assume that  $G_j = G_{i+1}$  and that  $G_{i+1}$  and hence  $S_{i+1}$  contains a  $D_1$ -subdivision.

26 Since the number of components is finite, the process must stop, so  $G_1$  contains an induced  $D_1$ -  
27 subdivision. □

#### 28 3.1 Induced $(a, b)$ -path in an oriented graph

29 Our first result is an easy modification of Bienstock's proof [3] that finding an induced cycle through  
30 two given vertices is NP-complete for undirected graphs.

31 **Lemma 7.** *It is NP-complete to decide whether an oriented graph contains an induced  $(a, b)$ -path  
32 for prescribed vertices  $a$  and  $b$ .*

33 *Proof.* Given an instance  $I$  of 3-SAT with variables  $x_1, x_2, \dots, x_n$  and clauses  $C_1, C_2, \dots, C_m$  we first  
34 create a variable gadget  $V_i^1$  for each variable  $x_i$ ,  $i = 1, 2, \dots, n$  and a clause gadget  $C_j^1$  for each clause  
35  $C_j$ ,  $j = 1, 2, \dots, m$  as shown in Figure 1. Then we form the digraph  $G_1(I)$  as follows (see Figure 2):  
36 Form a chain  $U$  of variable gadgets by adding the arcs  $b_i a_{i+1}$  for  $i = 1, 2, \dots, n - 1$  and a chain  $W$   
37 of clause gadgets by adding the arcs  $d_j c_{j+1}$ ,  $j = 1, 2, \dots, m - 1$ . Add the arcs  $aa_1, b_n c_1, c_m b$  to get  
38 a chain from  $a$  to  $b$ . For each clause  $C$ , we connect the three literal vertices of the gadget for  $C$  to  
39 the variable gadgets for variables occurring as literals in  $C$  in the way indicated in the figure. To be  
40 precise, suppose  $C_p = (x_i \vee \bar{x}_j \vee x_k)$ , then we add the following three 3-cycles  $l_p^1 x_i v_i l_p^1$ ,  $l_p^2 \bar{x}_j \bar{v}_j l_p^2$  and  
41  $l_p^3 x_k v_k l_p^3$ . This concludes the construction of  $G_1(I)$ .

<sup>1</sup>A connected component of a digraph  $H$  is a connected component in the underlying undirected graph of  $H$ .




Figure 1: The variable gadget  $V_i^1$  (left) and the clause gadget  $C_j^1$  (right).




Figure 2: The digraph  $G_1(I)$  when  $I$  has variables  $x_1, x_2, x_3$  and three clauses  $C_1, C_2, C_3$  where  $C_1 = (\bar{x}_1 \vee x_2 \vee \bar{x}_3)$  and  $C_3 = (x_1 \vee \bar{x}_2 \vee x_3)$  (for clarity we do not show the arcs corresponding to  $C_2$ )

1 We claim that there is an induced directed  $(a, b)$ -path in  $G_1(I)$  if and only if  $I$  is satisfiable.  
2 Suppose first that  $I$  is satisfiable and consider a truth assignment  $T$  which satisfies  $I$ . Now form a  
3 directed  $(a, b)$ -path  $P$  by taking the arcs  $aa_1, c_m b$  and the following subpaths: for each variable  $x_i$  take  
4 the subpath  $a_i \bar{x}_i \bar{v}_i b_i$  if  $T$  sets  $x_i$  true and otherwise take the subpath  $a_i x_i v_i b_i$ . For each clause  $C_j$  we fix  
5 a literal  $l'_j$  which is satisfied by  $T$  and take the subpath  $c_j l'_j d_j$ . It is easy to check that  $P$  is induced  
6 as we navigate it to avoid each of the arcs between the variable chain  $U$  and the clause chain  $W$ .  
7 Suppose now that  $Q$  is an induced directed  $(a, b)$ -path in  $G_1(I)$ . It follows from the construction that  
8  $Q$  starts by a directed  $(a_1, b_n)$ -path through all variable gadgets which contains no vertices from  $W$   
9 and continues with a directed  $(c_1, d_m)$ -path through all clause gadgets which contains no vertices from  
10  $U$ . This follows from the presence of the directed 3-cycles that prevent  $Q$  from using any of the arcs  
11 going from a variable gadget to a clause gadget other than the arc  $b_n c_1$ . Similarly there is no induced  
12 directed  $(c_1, d_m)$ -path which contains any vertex from  $U$ . Now form a truth assignment by setting  
13  $x_i$  true if and only if  $Q$  uses the subpath  $a_i \bar{x}_i \bar{v}_i b_i$  and false otherwise. Since  $Q$  is induced, for each  
14 clause  $C_j$  if  $Q$  uses the subpath  $c_j l'_j d_j$ , then we claim that  $l'_j$  will be true with the truth assignment just  
15 described: if  $l'_j = x_k$  for some  $k$  then since  $Q$  is induced the presence of the arc  $l'_j x_k$  implies that  $Q$  uses  
16 the path  $a_k \bar{x}_k \bar{v}_k b_k$  and similarly, if  $l'_j = \bar{x}_k$  then  $Q$  uses the path  $a_k x_k v_k b_k$  and again  $C_j$  is satisfied.  $\square$

### 17 3.2 Induced subdivisions of directed cycles

18 We first show that for any  $k \geq 4$ , the problem  $\Pi'_{C_k}$  is NP-complete.

19 **Theorem 8.** *It is NP-complete to decide whether an oriented graph contains an induced subdivision*  
20 *of a fixed directed cycle of length at least 4.*

21 *Proof.* Given an instance  $I$  of 3-SAT with variables  $x_1, x_2, \dots, x_n$  and clauses  $C_1, C_2, \dots, C_m$  we form  
22 the digraph  $G_1^*(I)$  from  $G_1(I)$  which we defined above by adding the arc  $ba$ .

23 Let  $C$  be an induced cycle of  $G_1^*(I)$ . Since the variable chain  $U$  and the clause chain  $W$  are both  
24 acyclic,  $C$  must contain an arc with tail  $l$  in  $W$  and head  $y$  in  $U$ . If  $ly \neq ba$ , then there exists  $i$  such that  
25  $y \in \{x_i, \bar{x}_i\}$  and so  $C = lx_i v_i l$  or  $C = l\bar{x}_i \bar{v}_i l$  by construction of  $G_1^*(I)$ . Hence every induced directed  
26 cycle of length at least 4 contains the arc  $ba$ . Thus  $G_1^*(I)$  has an induced cycle of length at least 4 if  
27 and only if  $G_1(I)$  has an induced directed  $(a, b)$ -path. As shown in the proof of Lemma 7 this is if  
28 and only if  $I$  is satisfiable.  $\square$

29 **Theorem 9.** *Let  $D$  be an oriented graph containing an induced directed cycle of length at least 4*  
30 *with a vertex of degree<sup>2</sup> 2. It is NP-complete to decide whether a given oriented graph contains an*  
31 *induced subdivision of  $D$ .*

32 *Proof.* Let  $D$  be given and let  $I$  be an arbitrary instance of 3-SAT. Fix an induced directed cycle  $C$  of  
33 length at least 4 in  $D$  and fix an arc  $uv$  on  $C$  such that  $u$  is of degree 2. Let  $G'_1(I)$  be the oriented graph  
34 that we obtain by replacing the arc  $uv$  by a copy of  $G_1(I)$  and the arcs  $ua, bv$ . We claim that  $G'_1(I)$   
35 contains an induced subdivision of  $D$  if and only if  $I$  is satisfiable (which is if and only if  $G_1(I)$   
36 contains an induced directed  $(a, b)$ -path).

37 Clearly, if  $G_1(I)$  has an induced directed  $(a, b)$ -path, then we may use the concatenation of this  
38 path with  $ua$  and  $bv$  instead of the deleted arc  $uv$  to obtain an induced  $D$ -subdivision in  $G'_1(I)$  (the  
39 only subdivided arc will be  $uv$ ).

40 Conversely, suppose that  $G'_1(I)$  contains an induced subdivision  $D'$  of  $D$ . Clearly  $D'$  has at least  
41 as many vertices as  $D$  and thus must contain at least one vertex  $z$  of  $V(G_1(I))$ . Since  $u$  is of degree  
42 2, the digraph  $D \setminus uv$  has fewer induced directed cycles of length at least 4 than  $D$ . (Note that the fact

<sup>2</sup>The degree of a vertex  $v$  in a digraph is the number of arcs with one end in  $v$ , that is, the sum of the in- and out-degree of  $v$ .

1 that  $u$  is of degree 2 is important: if  $u$  has degree more than 2, deleting  $uv$  could create new induced  
 2 directed cycles. ) Thus  $z$  must be on a cycle of length at least 4 in  $D'$ . But this and the fact that  $G_1(I)$   
 3 has no induced directed cycle of length at least 4 implies that  $G'_1(I)$  contains an induced directed  
 4  $(a, b)$ -path (which passes through  $z$ ).  $\square$

5 We move now to the detection of induced subdivisions of digraphs  $H$  when  $H$  is the disjoint  
 6 union of one or more directed cycles, all of length 3. If there is just one cycle in  $H$ , the problem  
 7 is polynomial-time solvable by Proposition 3. But from two on, it becomes NP-complete. We need  
 8 results on the following problem.

9 **PROBLEM DIDPP**

10 Input: An acyclic digraph  $G$  and two vertex pairs  $(s_1, t_1), (s_2, t_2)$ . Moreover, there is no directed path  
 11 from  $\overline{\{s_2, t_2\}}$  to  $\{s_1, t_1\}$ .

12 Question: Does  $G$  have two paths  $P_1, P_2$  such that  $P_i$  is a directed  $(s_i, t_i)$ -path,  $i = 1, 2$ , and  $G \setminus (V(P_1) \cup$   
 13  $V(P_2))$  is the disjoint union of  $P_1$  and  $P_2$ ?

14 Problem  $k$ -DIDPP was shown to be NP-complete by Kobayashi [8] using a proof similar to Bien-  
 15 stock's proof in [3].

16 **Theorem 10.** *Let  $D$  be the disjoint union of two directed cycles with no arcs between them. Then  $\Pi'_D$*   
 17 *is NP-complete.*

18 *Proof.* Let  $G$  be an instance of DIDPP and  $H$  the oriented graph obtained from it by adding new  
 19 vertices  $u_1, u_2$  and the arcs  $t_1 u_1, u_1 s_1, t_2 u_2$  and  $u_2 s_2$ . Since  $G$  was acyclic it is not difficult to see that  
 20  $H$  is a yes-instance of  $\Pi'_D$  if and only if  $G$  is a yes-instance of DIDPP.  $\square$

21 **4 NP-completeness results for digraphs**




Figure 3: The variable gadget  $V_i^2$  (left) and the clause gadget  $C_j^2$  (right). Unoriented bold edges represent 2-cycles.

22 **Theorem 11.** *Let  $k \geq 3$  be an integer. Then  $\Pi_{C_k}$  is NP-complete.*

23 *Proof.* Reduction from 3-SAT. Let  $I$  be an instance of 3-SAT with variables  $x_1, x_2, \dots, x_n$  and clauses  
 24  $C_1, C_2, \dots, C_m$ . We first create a variable gadget  $V_i^2$  for each variable  $x_i$ ,  $i = 1, 2, \dots, n$  and a clause  
 25 gadget  $C_j^2$  for each clause  $C_j$ ,  $j = 1, 2, \dots, m$  as shown in Figure 3. Then we form the digraph  $G_2(I)$   
 26 as follows (see Figure 4): Form a chain  $U$  of variable gadgets by adding the arcs  $b_i a_{i+1}$  for  $i =$

1  $1, 2, \dots, n-1$  and a chain  $W$  of clause gadgets by adding the arcs  $d_j c_{j+1}$ ,  $j = 1, 2, \dots, m-1$ . Add the  
2 arcs  $aa_1, b_n c_1, c_m b$  to get a chain from  $a$  to  $b$ . For each clause  $C$ , we connect the three literal vertices  
3 of the gadget for  $C$  to the variable gadgets for variables occurring as literals in  $C$  in the following way.  
4 Suppose  $C_p = (x_i \vee \bar{x}_j \vee x_k)$ , then we add the following three 2-cycles  $l_p^1 x_i l_p^1$ ,  $l_p^2 \bar{x}_j l_p^2$  and  $l_p^3 x_k l_p^3$ . This  
5 concludes the construction of  $G_2(I)$ . See Figure 4.




Figure 4: The digraph  $G_2(I)$  when  $I$  has variables  $x_1, x_2, x_3$  and three clauses  $C_1, C_2, C_3$  where  $C_1 = (\bar{x}_1 \vee x_2 \vee \bar{x}_3)$  and  $C_3 = (x_1 \vee \bar{x}_2 \vee x_3)$  (for clarity we do not show the arcs corresponding to  $C_2$ )

6 Similarly to the proof of Lemma 7, one can show that there is an induced directed  $(a, b)$ -path in  
7  $G_2(I)$  if and only if  $I$  is satisfiable.

8 Let  $G_2^k(I)$  be the digraph obtained from  $C_k$  by replacing one arc  $ab$  by  $G_2(I)$ . It is easy to check  
9 that  $G_2(I)$  has no induced cycle of length at least 3. Hence  $G_2^k(I)$  has an induced directed cycle of  
10 length  $k$  if and only if  $G_2(I)$  has an induced directed  $(a, b)$ -path. Hence by Lemma 7,  $G_2^k(I)$  has an  
11 induced  $D$ -subdivision if and only if  $I$  is satisfiable.  $\square$

12 A *branch* is a directed walk such that all the vertices are distinct except possibly its ends, its ends  
13 are nodes or leaves and all its internal vertices are continuities. A branch is *central* if its two ends are  
14 nodes.

15 The *skeleton* of a multidigraph  $D$  is the digraph whose vertices are the nodes and leaves in  $D$  and  
16 in which  $ab$  is an arc if and only if there is a directed  $(a, b)$ -branch in  $D$ . Observe that a skeleton may  
17 have loops and multiple arcs. Clearly, any subdivision of  $D$  has the same skeleton as  $D$ .

18 **Theorem 12.** *Let  $D$  be an oriented graph. If  $D$  contains a central branch, then  $\Pi_D$  is NP-complete.*

1 *Proof.* Reduction from 3-SAT. Let  $I$  be an instance of 3-SAT. Let  $B$  be a central branch with origin  $a$   
2 and terminus  $c$ . Let  $G_2^D(I)$  be the digraph obtained from  $D$  by replacing the first arc  $ab$  of  $B$  by  $G_2(I)$ .

3 Clearly if  $G_2(I)$  has an induced directed  $(a, b)$ -path  $P$ , then the union of  $P$  and  $D \setminus ab$  is a  $D$ -  
4 subdivision (in which only  $ab$  is subdivided) in  $G_2^D(I)$ .

5 Conversely, assume that  $G_2^D(I)$  contains an induced  $D$ -subdivision  $S$ . It is easy to check that no  
6 vertex in  $V(G_2(I)) \setminus \{a, b\}$  can be a node of  $S$  (the 2-cycles prevent this). Then since  $S$  has the same  
7 skeleton as  $D$ ,  $a$  and  $b$  are nodes of  $S$ . In addition, since the number of central branches in  $D \setminus ab$  is  
8 one less than the number of central branches in  $D$ , one central branch of  $D$  must use vertices of  $G_2(I)$ .  
9 Thus, there is an induced directed  $(a, b)$ -path in  $G_2(I)$ .

10 Hence  $G_2^D(I)$  has an induced  $D$ -subdivision if and only if  $G_2(I)$  has an induced directed  $(a, b)$ -  
11 path and thus if and only if  $I$  is satisfiable.  $\square$

12 **Corollary 13.** *Let  $D$  be an oriented graph. Then  $\Pi_D$  is NP-complete unless  $D$  is the disjoint union of*  
13 *spiders.*

14 *Proof.* Let  $D$  be an oriented graph. If one of its connected components is neither a directed cycle nor  
15 a spider, then it contains at least one central branch. So  $\Pi_D$  is NP-complete by Theorem 12.

16 If one of the components is directed cycle of length at least 3, then  $\Pi_D$  is NP-complete by Theo-  
17 rem 11 and Proposition 6.

18 Finally, if all its connected components are spiders then  $\Pi_D$  is polynomial-time solvable according  
19 to Theorem 5.  $\square$

20 We believe that Corollary 13 can be generalized to digraphs.

21 **Conjecture 14.** *Let  $D$  be a digraph. Then  $\Pi_D$  is NP-complete unless  $D$  is the disjoint union of spiders*  
22 *and at most one 2-cycle.*

23 As support for this conjecture, we give some other digraphs  $D$  (which are not oriented graphs),  
24 for which  $\Pi_D$  is NP-complete. In particular, when  $D$  is the *lollipop*, that is the digraph  $L$  with vertex  
25 set  $\{x, y, z\}$  and arc set  $\{xy, yz, zy\}$ . Note that the lollipop seems to be the simplest digraph that is not  
26 an oriented graph nor a  $C_2$ . So it should be an obvious candidate for a further polynomial case if one  
27 existed.

28 **Theorem 15.** *Deciding if a digraph contains an induced subdivision of the lollipop is NP-complete.*

29 *Proof.* Reduction from 3-SAT. Let  $I$  be an instance of 3-SAT with variables  $x_1, x_2, \dots, x_n$  and clauses  
30  $C_1, C_2, \dots, C_m$ . We first create a variable gadget  $V_i^3$  for each variable  $x_i$ ,  $i = 1, 2, \dots, n$  and a clause  
31 gadget  $C_j^3$  for each clause  $C_j$ ,  $j = 1, 2, \dots, m$  as shown in Figure 5. Then we form the digraph  $G_3(I)$   
32 as follows: Form a chain  $U$  of variable gadgets by adding the arcs  $b_i a_{i+1}$  for  $i = 1, 2, \dots, n - 1$  and a  
33 chain  $W$  of clause gadgets by adding the arcs  $d_j c_{j+1}$ ,  $j = 1, 2, \dots, m - 1$ . Add the arcs  $aa_1, b_n c_1, c_m b$   
34 to get a chain from  $a$  to  $b$ . For each clause  $C$ , we connect the three literal vertices of the gadget for  $C$   
35 to the variable gadgets for variables occurring as literals in  $C$  in the way indicated in the figure.

36 Similarly to the proof of Lemma 7, one can check that there is an induced directed  $(a, b)$ -path in  
37  $G_3(I)$  if and only if  $I$  is satisfiable.

38 The digraph  $G_3^L(I)$  is obtained from  $L$  and  $G_3(I)$  by deleting the arc  $yz$  and adding the arcs  $ya$   
39 and  $bz$ .

40 It is easy to see that  $G_3(I)$  has no induced directed cycle of length 3 and that no 2-cycle is  
41 contained in an induced lollipop. Hence if  $G_3^L(I)$  contains an  $L$ -subdivision, the induced directed  
42 cycle in it is the concatenation of the path  $bzya$  and a induced directed  $(a, b)$ -path in  $G_3(I)$ . Thus  $I$  is  
43 satisfiable. The other direction is (as usual) clear.  $\square$




Figure 5: The variable gadget  $V_i^3$ , (top left), the clause gadget  $C_i^3$  (bottom left) and the way to connect them in  $G_3(I)$  (right). Bold unoriented edges represent 2-cycles. Only the connection for one variable gadget and one clause gadget is shown and the general strategy for connecting variable and clause gadgets is the same as in  $G_1(I)$  (Figure 2).

1 **Remark 16.** The *cone* is the digraph  $C$  with vertex set  $\{x, y, z\}$  and arc set  $\{xy, xz, yz, zy\}$ . In the very  
 2 same way as Theorem 15, one can show that finding an induced subdivision of the cone in a digraph  
 3 is NP-complete.

## 4 5 Polynomial-time algorithms for induced subdivisions in oriented graphs

5 According to Conjecture 14, the only digraphs for which  $\Pi_D$  is polynomial-time solvable are disjoint  
 6 unions of spiders and possibly one 2-cycle. For such digraphs, easy polynomial-time algorithms exist  
 7 (See Section 2).

8 In this section, we show that the picture is more complicated for  $\Pi'_D$  than for  $\Pi_D$ . We show some  
 9 oriented graphs  $D$  for which  $\Pi'_D$  is polynomial-time solvable. For all these oriented graphs,  $\Pi_D$  is  
 10 NP-complete by Corollary 13.

### 11 5.1 Induced subdivision of cherries in oriented graphs

12 Let  $s, u, v$  be three vertices such that  $s \neq v$  and  $u \neq v$  (so  $s = u$  is possible). A *cherry on*  $(s, u, v)$  is any  
 13 oriented graph made of three induced directed paths  $P, Q, R$  such that:

- 14 •  $P$  is directed from  $s$  to  $u$  (so when  $s = u$  it has length 0);
- 15 •  $Q$  and  $R$  are both directed from  $u$  to  $v$  (so they both have length at least 1 and since we do not  
 16 allow parallel edges, at least one of them has length at least 2);
- 17 •  $u, v$  are the only vertices in more than one of  $P, Q, R$ ;
- 18 • there are no other arcs than those from  $P, Q, R$ .

19 The cherry is *rooted at*  $s$ .

20 An induced cherry contains an induced  $TT_3$ -subdivision (made of  $Q$  and  $R$ ) and a  $TT_3$ -subdivision  
 21 is a cherry (with  $u = s$ ). Hence detecting an induced cherry is equivalent to detecting an induced  $TT_3$ -  
 22 subdivision.

1 In order to give an algorithm that detects a cherry rooted at a given vertex, we use a modification  
 2 of the well-known Breadth First Search algorithm (BFS), see e.g. [1, Section 3.3]. Given an oriented  
 3 graph  $G$  and a vertex  $s \in V(D)$ , BFS returns an out-tree rooted at  $s$  and spanning all the vertices  
 4 reachable from  $s$ . It proceeds as follows:

**BFS( $G, s$ )**

Create a queue  $Q$  consisting of  $s$ ; Initialize  $T = (\{s\}, \emptyset)$

**while**  $Q$  is not empty **do**

    Consider the head  $u$  of  $Q$  and visit  $u$ , that is

**foreach** out-neighbour  $v$  of  $u$  in  $D$  **do**

**if**  $v \notin V(T)$  **then**

$V(T) := V(T) \cup \{v\}$  and  $A(T) := A(T) \cup \{uv\}$

            Put  $v$  to the end of  $Q$

        Delete  $u$  from  $Q$

6  
 7 Note that the arc-set of the out-branching produced by BFS depends on the order in which the vertices  
 8 are visited, but the vertex-set is always the same: it is the set of the vertices reachable from  $s$ . See [1]  
 9 p. 92 for more details on BFS. We need the following variant:

**IBFS( $G, s$ )**

Create a queue  $Q$  consisting of  $s$ ; Initialize  $T = (\{s\}, \emptyset)$

**while**  $Q$  is not empty **do**

    Consider the head  $u$  of  $Q$  and visit  $u$ , that is

**foreach** out-neighbour  $v$  of  $u$  in  $G$  **do**

**if**  $N_G(v) \cap V(T) = \{u\}$  **then**

$V(T) := V(T) \cup \{v\}$  and  $A(T) := A(T) \cup \{uv\}$

            Put  $v$  to the end of  $Q$

        Delete  $u$  from  $Q$

11  
 12 Observe that IBFS (which we also call *induced-BFS*) is the same as BFS except that we add the  
 13 out-neighbour  $v$  of  $u$  to  $T$  only if it has no other neighbour already in  $T$ , hence ensuring that the  
 14 resulting out-tree is an induced subdigraph of  $G$ . Contrary to BFS, the vertex-set of a tree obtained  
 15 after IBFS may depend on the order in which the vertices are visited.

16 IBFS can easily be implemented to run in time  $O(n^2)$ . When  $T$  is an oriented tree, we denote by  
 17  $T[x, y]$  the unique oriented path from  $x$  to  $y$  in  $T$ .

18 **Theorem 17.** *Let  $G$  be an oriented graph,  $s$  a vertex and  $T$  a tree obtained after running IBFS( $G, s$ ).  
 19 Then exactly one of the following outcomes is true:*

- 20 (i)  $D$  contains an induced subdigraph that is a cherry rooted at  $s$ ;  
 21 (ii) for every vertex  $x$  of  $T$ , any out-neighbour of  $x$  not in  $T$  has an out-neighbour that is an ancestor  
 22 of  $x$  in  $T$ .

23 *This is algorithmic in the sense that there is an  $O(n^2)$  algorithm that either outputs the cherry of  
 24 (i) or checks that (ii) holds.*

25 *Proof.* Suppose that  $T$  does not satisfy (ii). Then some vertex  $x$  of  $T$  has an out-neighbour  $y$  not in  
 26  $T$  and no out-neighbour of  $y$  is an ancestor of  $x$ . Without loss of generality, we assume that  $x$  is the  
 27 first vertex added to  $T$  when running IBFS with such a property. In particular,  $T[s, x]y$  is an induced  
 28 directed path because a chord would contradict (ii) or the choice of  $x$ . Let  $v$  be the neighbour of  $y$  in  
 29  $T$ , different from  $x$ , that was first added to  $T$  when running IBFS. Note that  $v$  exists for otherwise  $y$   
 30 would have been added to  $T$  when visiting  $x$ . If  $x$  is the parent of  $v$  in  $T$  then  $T[s, x]y$  together with  $v$

1 form a cherry rooted at  $s$  (whatever the orientation of the arc between  $y$  and  $v$ ). So we may assume  
2 that  $x$  is not the parent of  $v$ . When visiting  $x$ , vertex  $y$  was not added to  $T$ , hence  $v$  was already visited  
3 (because  $x$  is not the parent of  $v$ ). In addition, when  $v$  was visited, it was the unique neighbour of  $y$   
4 in the current out-tree, so  $y$  is an in-neighbour of  $v$ , for otherwise it would have been added to  $T$ . Let  
5  $u$  be the common ancestor of  $x$  and  $v$  in  $T$ , chosen closest to  $x$ . Since  $T$  does not satisfy (ii) by the  
6 choice of  $x$  and  $y$ ,  $u \neq v$ . Now the directed paths  $sTu$ ,  $T[u,x]yv$  and  $T[u,v]$  form an induced cherry  
7 rooted at  $s$ . Indeed since  $T$  is an induced out-tree, it suffices to prove that  $y$  has no neighbour in these  
8 three paths except  $x$  and  $v$ . By definition of  $v$ , there is no neighbour of  $y$  in  $T[s,u]$  and  $T[u,v]$  except  
9  $v$ . Moreover,  $y$  has no out-neighbour in  $T[u,x]$  by the assumption that (ii) does not hold for  $y$  and  $x$   
10 and it has no in-neighbour in  $T[u,x]$  except  $x$  by the choice of  $x$ .

11 Conversely, let us assume that  $T$  satisfies (ii) and suppose by contradiction that  $G$  contains an  
12 induced cherry  $C$  rooted at  $s$ . Since  $T$  is an induced out-branching, some vertices of  $C$  are not in  $T$ .  
13 So, let  $y$  be a vertex of  $V(C) \setminus V(T)$  as close to  $s$  as possible in the cherry. Let  $x$  be an in-neighbour of  
14  $y$  in  $C \cup T$ . From the choice of  $y$ ,  $x$  and all its ancestors along the cherry are in  $T$ . Since  $T$  is induced,  
15 the ancestors of  $x$  along the cherry are in fact the ancestors of  $x$  along  $T$ . Hence,  $x$  is a vertex of  $T$   
16 with an out-neighbour  $y$  not in  $T$  having no out-neighbour among the ancestors of  $x$  along  $T$ . This  
17 contradicts  $T$  satisfying (ii).

18 All this may be turned in an  $O(n^2)$ -algorithm that finds a cherry rooted at  $s$  if it exists or answer no  
19 otherwise. Indeed we first run IBFS and then check in time  $O(n^2)$  if the obtained tree  $T$  satisfies (ii).  
20 If not, then we can find the cherry following the first paragraph of the proof.  $\square$

21 **Remark 18.** Since a digraph contains an induced  $TT_3$ -subdivision if and only if it contains an in-  
22 duced cherry, Theorem 17 implies directly that  $\Pi'_{TT_3}$  is solvable in time  $O(n^3)$  (because we need to  
23 enumerate all potential roots).

24 We can slightly extend our result. A *tiny cherry* is a cherry such that the path  $Q$  and  $R$  as in the  
25 definition form a  $TT_3$ .

26 **Corollary 19.** *For any tiny cherry  $D$ , the problem  $\Pi'_D$  is solvable in time  $O(n^{|V(D)|})$ .*

27 *Proof.* Let  $P$  be the path of  $D$  as in the definition of cherry. Let  $G$  be the input oriented graph. Enu-  
28 merate by brute force all induced directed paths of order  $|P|$  by checking all the possible subdigraphs  
29 of order  $|P|$ . For each such path  $P'$  with terminus  $x$ , look for a cherry rooted at  $x$  in the graph  $G'$   
30 obtained by deleting all the vertices of  $P - x$  and their neighbourhoods except  $x$ . If there is such a  
31 cherry  $C$  then the union of  $P$  and  $C$  is an induced  $D$ -subdivision.  $\square$

32 Similarly to Propositions 4 and 5, we have the following.

33 **Corollary 20.** *If  $D$  is the disjoint union of spiders and a tiny cherry then  $\Pi'_D$  is polynomial-time  
34 solvable.*

## 35 5.2 Induced subdivision of oriented paths with few blocks in oriented graphs.

36 By Proposition 1, for any oriented path  $P$  with at most two blocks  $\Pi_P$  and thus  $\Pi'_P$  are polynomial-  
37 time solvable. In this section, we shall prove that  $\Pi'_P$  is polynomial-time solvable for some oriented  
38 paths with three or four blocks. In contrast,  $\Pi_P$  is NP-complete for every oriented path with at least  
39 three blocks as shown in Corollary 13.

### 40 5.2.1 Oriented path with three blocks

41 **Theorem 21.** *There exists an algorithm of complexity  $O(m^2)$  that given a connected oriented graph  
42 on  $n$  vertices and  $m$  arcs with a specified vertex  $s$  returns an induced  $A_2^+$ -subdivision with origin  $s$  if  
43 one exists, and answer 'no' if not.*

1 *Proof.* Observe that any induced  $A_2^+$ -subdivision with origin  $s$  contains an induced  $A_2^+$ -subdivision  
 2 with origin  $s$  such that the directed path corresponding to the arc  $s_3s_2$  is some arc  $f$ . Such a subdivision  
 3 is called *f-led*.

4 Given an oriented graph  $G$ , we enumerate all arcs  $f = s'_3s'_2$ . For each arc in turn we either show  
 5 that there is no *f-led* induced  $A_2^+$ -subdivision with origin  $s$  or give an induced subdivision of  $A_2^+$   
 6 with origin  $s$ , (but not necessarily *f-led*). This will detect the  $A_2^+$ -subdivision since if some exists,  
 7 it is *f-led* for some  $f$ .

8 We do this as follows. We delete all in-neighbours of  $s$  and all neighbours of  $s'_3$  except  $s'_2$ . Let us  
 9 denote by  $G'$  the resulting graph. Then we compute by BFS a shortest directed path  $P$  from  $s$  to  $s'_2$ . If  
 10 it is induced, together with  $s'_3s'_2$ , it forms the desired  $A_2^+$  subdivision. So, as  $P$  has no forward chord  
 11 (since it is a shortest path), there is an arc  $uv$  in  $G' \setminus \langle V(P) \rangle$  such that  $u$  occurs after  $v$  on  $P$ . Take such an  
 12 arc  $b_3b_2$  such that  $b_2$  is as close as possible to  $s$  (in  $P$ ). Observe that since we deleted all in-neighbours  
 13 of  $s$  we have  $b_2 \neq s$ . Now,  $P[s_1, b_2]$  together with  $b_3b_2$  forms the desired  $A_2^+$ -subdivision.

14 There are  $O(m)$  arcs and for each of them we must find a shortest path in  $G'$  which can be done  
 15 in  $O(m)$ . Hence the complexity of the algorithm is  $O(m^2)$ .  $\square$

16 From this theorem, one can show that finding an induced  $A_3^-$ -subdivision is polynomial-time  
 17 solvable. It is enough to enumerate all arcs  $s'_2s'_1$ , to delete  $s'_1$  and its neighbours except  $s'_2$ , and to  
 18 decide whether there exists in what remains an  $A_2^+$ -subdivision with origin  $s_2$ . One can also de-  
 19 rive polynomial-time algorithms for finding induced subdivisions of other oriented paths with three  
 20 blocks.

21 **Corollary 22.** *Let  $P$  be a path with three blocks such that the last one has length 1. One can check in*  
 22 *time  $O(n^{|P|-2}m^2)$  whether a given oriented graph contains an induced  $P$ -subdivision.*

23 *Proof.* By directional duality, we may assume that  $P$  is an  $A_3^-$ -subdivision. Let  $Q$  be the subdigraph  
 24 of  $P$  formed by the first block of  $P$  and the second block of  $P$  minus the last arc. Let  $s$  be the terminus  
 25 of  $Q$ . For each induced oriented path  $Q'$  in the instance graph, isomorphic to  $Q$  (there are at most  
 26  $O(n^{|P|-2})$  of them), we delete  $Q' - s$  and all vertices that have neighbours in  $Q - s$  except  $s$ . We then  
 27 detect an  $A_2^+$ -subdivision rooted at  $s$  in the resulting graph. This will detect a  $P$ -subdivision if there is  
 28 one.  $\square$

### 29 5.2.2 Induced subdivision of $A_4^-$ in an oriented graph

30 We show how to check the presence of an induced copy of  $A_4^-$  by using flows (for definitions and  
 31 algorithms for flows see e.g. [1, Chapter 4]).

32 **Theorem 23.** *There exists an algorithm of complexity  $O(nm^2)$  that given an oriented graph on  $n$*   
 33 *vertices and  $m$  arcs with a specified vertex  $s$  returns an induced  $A_3^+$ -subdivision rooted at  $s$ , if one*  
 34 *exists, and answer 'no' if not.*

35 *Proof.* The general idea is close to the one of the proof of Theorem 21. Observe that any induced  
 36  $A_3^+$ -subdivision with origin  $s = a_1$  contains an induced subdivision of  $A_3^+$  with origin  $s = a_1$  such that  
 37 the directed path corresponding to the arc  $s_3s_4$  is some arc  $f$ . If, in addition, the vertex corresponding  
 38 to  $s_2$  is  $v$ , such a subdivision is called *(v, f)-led*.

39 Given an oriented graph  $G$ , we enumerate all pairs  $(a_2, a_3a_4)$  such that  $a_2, a_3, a_4$  are distinct ver-  
 40 tices and  $a_3a_4 \in E(G)$ . For each such pair in turn we either show that there is no  $(a_2, a_3a_4)$ -led  
 41 induced  $A_3^+$ -subdivision with origin  $a_1$  or give an induced subdivision of  $A_3^+$  with origin  $a_1$  (but not  
 42 necessarily  $(a_2, a_3a_4)$ -led).

43 We do this as follows. We first delete all the neighbours of  $a_4$  except  $a_3$ , all in-neighbours of  $a_1$   
 44 and  $a_3$  and finally all out-neighbours of  $a_2$ . If this results in one or more of the vertices  $a_1, \dots, a_4$  to

1 be deleted, then there cannot be any  $(a_2, a_4a_3)$ -leaded induced  $A_3^+$ -subdivision with origin  $a_1$  because  
2 there is an arc in  $G\langle\{a_1, \dots, a_4\}\rangle$  which is not in  $\{a_1a_2, a_3a_2, a_3a_4\}$ . So we skip this pair and proceed  
3 to the next one. Otherwise we delete  $a_4$  and we use a flow algorithm to check in the resulting digraph  
4  $G'$  the existence of two internally-disjoint directed paths  $P, Q$  such that the origin of  $P$  and  $Q$  are  $a_1$   
5 and  $a_3$  respectively and such that  $a_2$  is the terminus of both  $P$  and  $Q$ . Moreover, we suppose that  
6 these two paths have no forward chord (this can easily be ensured by running BFS on the graphs  
7 induced by each of them). If no such paths exist, then we proceed to the next pair because there is  
8 no  $(a_2, a_3a_4)$ -leaded induced  $A_3^+$ -subdivision. If we find such a pair of directed paths  $P, Q$ , then we  
9 shall provide an induced subdivision of  $A_3^+$  with origin  $a_1$ . If  $P$  and  $Q$  are induced and have no arcs  
10 between them, then these paths together with the arc  $a_3a_4$  form the desired induced subdivision of  
11  $A_3^+$ .

12 Suppose that  $P$  is not induced. As  $P$  has no forward chord, there is an arc  $uv$  in  $G'\langle V(P)\rangle$  such that  
13  $u$  occurs after  $v$  on  $P$ . Take such an arc  $b_3b_2$  such that  $b_2$  is as close as possible to  $a_1$  (in  $P$ ), and subject  
14 to this, such that  $b_3$  is as close as possible to  $a_2$ . Observe that since we deleted all in-neighbours of  $a_1$   
15 and all out-neighbours of  $a_2$  before, we must have  $b_2 \neq a_1$  and  $b_3 \neq a_2$ . Let  $b_4$  be the successor of  $b_3$   
16 on  $P$ . Now  $P[a_1, b_2]$  and the arcs  $b_3b_2, b_3b_4$  form the desired induced subdivision of  $A_3^+$ . From here  
17 on, we suppose that  $P$  is induced.

18 Suppose now that there is an arc  $e$  with an end  $x \in V(P)$  and the other  $y \in V(Q)$ . Choose such an  
19 arc so that the sum of the lengths of  $P[a_1, x]$  and  $Q[a_3, y]$  is as small as possible. If  $e$  is from  $x$  to  $y$   
20 have  $y \neq a_3$  because we removed all the in-neighbours of  $a_3$ , else  $e$  is from  $y$  to  $x$  and we have  $x \neq a_1$   
21 because we removed all the in-neighbours of  $a_1$ . In all cases, we get an induced subdivision of  $A_3^+$  by  
22 taking the paths  $P[a_1, x]$  and  $Q[a_3, y]$  and the arcs  $a_3a_4, e$ . From here on, we suppose that there are no  
23 arcs with an end in  $V(P)$  and the other in  $V(Q)$ .

24 The last case is when  $Q$  is not induced. Since  $Q$  has no forward chord, there is an arc  $uv$  in  
25  $G'\langle V(Q)\rangle$  such that  $u$  occurs after  $v$  on  $Q$ . Take such an arc  $b_3b_4$  such that  $b_3$  is as close as possible to  
26  $a_2$  (in  $Q$ ). Observe that since we deleted all out-neighbours of  $a_2$  before, we must have  $b_3 \neq a_2$ . Now  
27  $P, Q[b_3, a_2]$  and the arc  $b_3b_4$  form the desired induced subdivision of  $A_3^+$ .

28 There are  $O(nm)$  pairs  $(a_2, a_3a_4)$  and for each of them, we run an  $O(m)$  flow algorithm (we just  
29 need to find a flow of value 2, say, by the Ford-Fulkerson method [1, Section 4.5.1]) and do some  
30 linear-time operations. Hence the complexity of the algorithm is  $O(nm^2)$ .  $\square$

31 One can check in polynomial time if there is an induced  $A_4^-$ -subdivision: it is enough to enumerate  
32 all arcs  $t_2t_1$ , to delete  $t_1$  and its neighbours except  $t_2$ , and to decide whether there exists in what remains  
33 an  $A_3^+$  subdivision with origin  $t_2$ . One can also derive polynomial-time algorithm for finding induced  
34 subdivision of other oriented paths with four blocks.

35 **Corollary 24.** *Let  $P$  be an oriented path that can be obtained from  $A_4^-$  by subdividing the first arc  
36 and the second arc. One can check in time  $O(n^{|P|-1}m^2)$  whether a given oriented graph contains an  
37 induced subdivision of  $P$ .*

38 *Proof.* Let  $R$  be the subdigraph of  $P$  formed by the first block of  $P$  and its second block minus the last  
39 arc. Let  $s$  be the last vertex of  $R$ . For each induced oriented path  $Q$  in the instance graph, isomorphic  
40 to  $R$  (there are  $O(n^{|P|-3})$  of them), we delete  $Q - s$ , all vertices that have neighbours in  $Q - s$  except  $s$   
41 and detect an  $A_3^-$ -subdivision with origin  $s$ . This will detect a  $P$ -subdivision if there is one.  $\square$




Figure 6: Left: clause gadget of  $G_1(I)$ . Right: clause gadget of  $G_4(I)$ .

## 6 Induced subdivisions of tournaments in oriented graphs

### 6.1 Induced subdivision of transitive tournaments

The transitive tournament on  $k$  vertices is denoted  $TT_k$ . We saw in Section 5.1 that  $\Pi'_{TT_3}$  is polynomial. The next result shows that  $\Pi'_{TT_k}$  is NP-complete for all  $k \geq 4$ .

**Theorem 25.** *For all  $k \geq 4$ ,  $\Pi'_{TT_k}$  is NP-complete*

*Proof.* For a given instance  $I$  of 3-SAT, let  $G_4(I)$  be the digraph we obtain from  $G_1(I)$  by replacing each clause gadget  $C_j^1$  by the modified one  $C_j^4$  from Figure 6. Also for each variable, modify the gadget  $V_i^1$  as follows: replace the path  $a_i x_i v_i b_i$  by a path  $a_i x_i^1 v_i^1 x_i^2 v_i^2 \dots x_i^m v_i^m b_i$ , and similarly the path  $a_i \bar{x}_i \bar{v}_i b_i$  by a path  $a_i \bar{x}_i^1 \bar{v}_i^1 \bar{x}_i^2 \bar{v}_i^2 \dots \bar{x}_i^m \bar{v}_i^m b_i$ . Then in  $G_4(I)$  the links representing a variable  $x_i$  and a clause  $C_j$  that uses this variable are represented by arcs between vertices from the variable gadget with superscript  $j$  (as in Figure 2).

Recall that  $G_1(I)$  has an induced directed  $(a, b)$ -path if and only if  $I$  is satisfiable. It is easy to see that the same holds for  $G_4(I)$ . Note that in  $G_4(I)$  no vertex has in- or out-degree larger than 2.

Given an instance  $I$  of 3-SAT we form the digraph  $G_4^k(I)$  from  $G_4(I)$  and a copy of  $TT_k$  (with vertices  $v_1, v_2, \dots, v_k$  and arcs  $v_i v_j$ ,  $1 \leq i < j \leq k$ ) by deleting the arc  $v_1 v_k$  and adding the arcs  $v_1 a, b v_k$ . We claim that  $G_4^k(I)$  contains an induced subdivision of  $TT_k$  if and only if  $G_4(I)$  has an induced directed  $(a, b)$ -path which is if and only if  $I$  is satisfiable.

Clearly, if  $I$  is satisfiable, we may use the concatenation of an induced directed  $(a, b)$ -path in  $G_4(I)$  with  $v_1 a$  and  $b v_k$  in place of  $v_1 v_k$  to obtain an induced  $TT_4$ -subdivision in  $G_4^k(I)$ .

Conversely, suppose that  $G_4^k(I)$  contains an induced subdivision of  $TT_k$  and let  $h(v_i)$ ,  $1 \leq i \leq k$ , denote the image of  $v_i$  in some fixed induced subdivision  $H$  of  $TT_k$ . Then we must have  $h(v_1) = v_1$  and  $h(v_k) = v_k$ , because  $G_4(I)$  does not contain any vertex of out-degree  $k-1$  or in-degree  $k-1$  because  $k \geq 4$ . For all  $i$ ,  $1 < i < k$ , the vertex  $h(v_i)$  could not be in  $V(G_4(I))$  since otherwise there must be either two disjoint directed  $(v_i, v_k)$ -paths to  $v_k$  or two disjoint directed  $(v_1, v_i)$ -paths. This is impossible because there is no directed  $(v_i, v_k)$ -path in  $G_4^k(I) \setminus b v_k$  and no directed  $(v_1, v_i)$ -path in  $G_4^k(I) \setminus v_1 a$ . Hence  $h(v_i) = v_i$  for all  $1 \leq i \leq k$  and so it is clear that we have an induced directed  $(a, b)$ -path in  $G_4(I)$ , implying that  $I$  is satisfiable.  $\square$

In the proof above we used that the two vertices  $v_1, v_k$  cannot be mapped to vertices of  $G_4(I)$ , the fact that the connectivity between these and the other vertices is too high to allow any of these

1 to be mapped to vertices of  $G_4(I)$  and finally we could appeal to the fact that  $G_4(I)$  has an induced  
 2 directed  $(a,b)$ -path if and only if  $I$  is satisfiable. Refining this argument it is not difficult to see  
 3 that the following holds where a  $(z,X)$ -path is a path whose initial vertex is  $z$  and whose last vertex  
 4 belongs to  $X$ .

5 **Theorem 26.** *Let  $D = (V,A)$  be a digraph and let  $X$  (resp.  $Y$ ) be the subset of vertices with out-  
 6 degree (resp. in-degree) at least 3 and let  $Z = V \setminus (X \cup Y)$  (note that  $X \cap Y \neq \emptyset$  is possible and also  
 7  $Z = \emptyset$  is possible). Suppose that for every  $z \in Z$  the digraph  $D$  contains either two internally disjoint  
 8  $(X,z)$ -paths or two internally disjoint  $(z,Y)$ -paths. Then  $\Pi'_D$  is NP-complete.*

9 *Proof.* (Sketch) For a given instance  $I$  of 3-SAT we form the digraph  $G'_4(I)$  from  $D$  by replacing  
 10 one arc  $uv$  with at least one of its endvertices in  $X \cup Y$  by  $G_4(I)$  and the arcs  $ua, bv$ . Again it is  
 11 clear how to obtain an induced subdivision of  $D$  in  $G'_4(I)$  when  $I$  is satisfiable. Let us now assume  
 12 that  $G'_4(I)$  contains an induced subdivision  $D'$  of  $D$ . Let  $\{h(v)|v \in V\}$  be the vertices corresponding  
 13 to the vertices of  $V$  in the subdivision. For degree reasons, none of the vertices in  $X \cup Y$  can have  
 14  $h(v) \in V(G_4(I))$  and because of connectivity, none of the vertices of  $Z$  can have  $h(z) \in V(G_4(I))$   
 15 because there is only one arc entering and leaving  $V(G_4(I))$  in  $G'_4(I)$ . Thus  $\{h(v)|v \in V\} = V$   
 16 (possibly with  $h(v) \neq v$  for several vertices). However, since we deleted the arc  $uv$  and replaced it by  
 17  $G_4(I)$  and the arcs  $ua, bv$ , it follows that  $G'_4(I)$  and so  $G_4(I)$  contains an induced directed  $(a,b)$ -path,  
 18 implying that  $I$  is satisfiable.  $\square$

## 19 6.2 Induced subdivision of the strong tournament on 4 vertices

20 Let  $ST_4$  be the unique strong tournament of order 4. It can be seen has a directed cycle  $\alpha\gamma\beta\delta\alpha$  together  
 21 with two chords  $\alpha\beta$  and  $\gamma\delta$ . The aim of this section is to show that  $\Pi'_{ST_4}$  is NP-complete.

22 An  $(x, y_1, y_2)$ -switch is the digraph with vertex set  $\{x, z, y_1, y_2\}$  and edge set  $\{xz, xy_1, zy_1, zy_2, y_2y_1\}$ .  
 23 See Figure 7.




Figure 7: An  $(x, y_1, y_2)$ -switch.

24 A good  $(x, y_1, y_2)$ -switch in a digraph  $D$  is an induced switch  $Y$  such that all the arcs entering  $Y$   
 25 have head  $x$  and all arcs leaving  $Y$  have tail in  $\{y_1, y_2\}$ .

26 **Lemma 27.** *Let  $Y$  be a good  $(x, y_1, y_2)$ -switch in a digraph  $D$ . Then every induced subdivision  $S$  of  
 27  $ST_4$  in  $D$  intersects  $Y$  on either the path  $(x, y_1)$ , the path  $(x, z, y_2)$ , or the empty set.*

28 *Proof.* Suppose for a contradiction, that  $y_2y_1 \in E(S)$ . Then  $S$  must contain the unique in-neighbour  $z$   
 29 of  $y_2$  and the unique in-neighbour  $x$  of  $z$ . Hence  $y_1$  has in-degree 3 in  $S$ , a contradiction.

30 Suppose for a contradiction, that  $zy_1 \in E(S)$ . Then  $S$  must contain  $x$  the unique in-neighbour of  $z$ .  
 31 Hence  $xy_1$  is a chord of  $S$  and so  $z$  must have degree 3 in  $S$ . Thus  $y_2 \in V(S)$  and  $y_1$  has in-degree 3 in  
 32  $S$ , a contradiction.  $\square$

33 **Theorem 28.**  $\Pi'_{ST_4}$  is NP-complete.

1 *Proof.* Reduction from 3-SAT. Let  $I$  be an instance of 3-SAT with variables  $x_1, x_2, \dots, x_n$  and clauses  
2  $C_1, C_2, \dots, C_m$ . We first create a variable gadget  $V_i^5$  for each variable  $x_i$ ,  $i = 1, 2, \dots, n$  and a clause  
3 gadget  $C_j^5$  for each clause  $C_j$ ,  $j = 1, 2, \dots, m$  as shown in Figure 8. Then we form the digraph  $G_5(I)$  as  
4 follows: Form a chain  $U$  of variable gadgets by adding the arcs  $b_i a_{i+1}$  for  $i = 1, 2, \dots, n - 1$  and a chain  
5  $W$  of clause gadgets by adding the arcs  $d_j c_{j+1}$ ,  $j = 1, 2, \dots, m - 1$ . Add the arcs  $aa_1, b_n b, cc_1, t_m d$ .  
6 For each clause  $C$ , we connect the three literal vertices of the gadget for  $C$  to the variable gadgets for  
7 variables occurring as literals in  $C$  in the way indicated in Figure 9.




Figure 8: The variable gadget  $V_i^5$  (left) and the clause gadget  $C_j^5$  (right).




Figure 9: Connections between a clause gadget and a variable gadget in  $G_5(I)$ . Only the connection for one variable gadget and one clause gadget is shown and the general strategy for connecting variable and clause gadgets is the same as in  $G_1(I)$  (Figure 2).

8 We denote by  $X_i$  the path  $a_i a'_i x_i^0 x_i^1 x_i^2 x_i^3 x_i^4 b_i$ , by  $\bar{X}_i$  the path  $a_i \bar{x}_i^0 \bar{x}_i^1 \bar{x}_i^2 \bar{x}_i^3 \bar{x}_i^4 b'_i b_i$ , by  $P_j$  the path  $c_j p_j p_j^0 p_j^1 p_j^2 p_j^3 d_j$ ,  
9 by  $Q_j$  the path  $c_j q_j q'_j q_j^0 q_j^1 q_j^2 q_j^3 d_j$ , and by  $R_j$  the path  $c_j q_j r_j^0 r_j^1 r_j^2 r_j^3 d_j$ .

10 Similarly to the proof of Lemma 7, one can check that  $I$  is satisfiable if and only if there are two  
11 induced disjoint directed  $(a, b)$ - and  $(c, d)$ -paths in  $G_5(I)$ .

12 Let  $G_5^*(I)$  be the digraph obtained from  $G_5(I)$  by adding the edges  $ac, cb, bd$  and  $da$ . Observe  
13 that  $G_5^*(I) \setminus da$  is acyclic.

14 Let us prove that  $G_5^*(I)$  contains an induced  $ST_4$ -subdivision if and only if  $I$  is satisfiable.

1 If  $I$  is satisfiable, then in  $G_5(I)$  there are two induced disjoint directed  $(a, b)$ - and  $(c, d)$ -paths.  
 2 The union of these paths and the directed cycle  $abcd$  is an induced  $ST_4$ -subdivision in  $G_5^*(I)$ .

3 Conversely, assume that  $G_5^*(I)$  contains an induced subdivision  $S$  of  $ST_4$ . For sake of simplicity  
 4 (and with a slight abuse of notation), we will denote the vertices of  $S$  corresponding to  $\alpha, \beta, \gamma$  and  $\delta$   
 5 by the same names. Let  $T_1$  and  $T_2$  be the paths corresponding to the chord  $\alpha\beta$  and  $\gamma\delta$  respectively in  
 6  $S$  and let  $C$  be the directed cycle corresponding to  $\alpha\gamma\beta\delta\alpha$ . Observe that the ends of  $T_1$  and  $T_2$  must  
 7 alternate on  $C$ .

8 Notice that the subdigraphs induced by the sets  $\{a_i, a'_i, x_i^0, \bar{x}_i^0\}$ ,  $1 \leq i \leq n$ ,  $\{c_j, p_j, p_j^0, q_j\}$  and  
 9  $\{q_j, q'_j, q_j^0, r_j^0\}$  are good switches. In addition, the subdigraphs induced by the sets  $\{b_i, b'_i, x_i^4, \bar{x}_i^4\}$ ,  
 10  $1 \leq i \leq n$ , are the converse of good switches. Hence Lemma 27 (and its converse) imply the following  
 11 proposition.

12 **Claim 28.1.**

13 (i) For  $1 \leq i \leq n$ , if  $a_i \in V(S)$ , then exactly one of the two paths  $(a_i, a'_i, x_i^0)$  and  $(a_i, \bar{x}_i^0)$  is in  $S$ .

14 (ii) For  $1 \leq i \leq n$ , if  $b_i \in V(S)$ , then exactly one of the two paths  $(\bar{x}_i^4, b'_i, b_i)$  and  $(x_i^4, b_i)$  is in  $S$ .

15 (iii) For  $1 \leq j \leq m$ , if  $c_j \in V(S)$ , then exactly one of the three paths  $(c_j, p_j, p_j^0)$ ,  $(c_j, q_j, q'_j, q_j^0)$  and  
 16  $(c_j, q_j, r_j^0)$  is in  $S$ .

17 Since  $G_5^*(I) \setminus da$  is acyclic,  $C$  must contain the arc  $da$ . Moreover since there is no arc with tail  
 18 in some clause gadget and head in some variable gadget,  $C$  contains at most one arc with tail in some  
 19 variable gadget and head in some clause gadget.

20 **Claim 28.2.** For any  $1 \leq i \leq n$  and any  $1 \leq j \leq m$ , the cycle  $C$  contains no arc with tail in  $\{x_i^3, \bar{x}_i^3\}$   
 21 and head in  $\{p_j^1, q_j^1, r_j^1\}$ .

22 *Proof.* Assume for a contradiction that  $C$  contains such an arc  $y_i^3 l_j^1$ . Then since  $l_j^1$  and  $l_j^2$  have out-  
 23 degree 1 then  $C$  must also contain  $l_j^2$  and  $l_j^3$ . Thus, in  $S$ ,  $y_i^3$  has out-degree 3 in  $S$ , a contradiction.  $\square$

24 **Claim 28.3.** For any  $1 \leq i \leq n$  and any  $1 \leq j \leq m$  the cycle  $C$  contains no arc with tail in  $\{x_i^3, \bar{x}_i^3\}$   
 25 and head in  $\{p_j^3, q_j^3, r_j^3\}$ .

26 *Proof.* Assume for a contradiction that  $C$  contains such an arc  $y_i^3 l_j^3$ . Then since  $y_i^3$  and  $y_i^2$  have in-  
 27 degree 1 then  $C$  must also contain  $y_i^2$  and  $y_i^1$ . Thus, in  $S$ ,  $l_j^3$  has in-degree 3 in  $S$ , a contradiction.  $\square$

28 **Claim 28.4.** For any  $1 \leq i \leq n$  and any  $1 \leq j \leq m$  then  $C$  contains no arc with tail in  $\{x_i^3, \bar{x}_i^3\}$  and  
 29 head in  $\{p_j^2, q_j^2, r_j^2\}$ .

30 *Proof.* Assume for a contradiction that  $C$  contains such an arc  $y_i^3 l_j^2$ . The vertex  $l_j^2$  has a unique out-  
 31 neighbour  $l_j^3$  which must be in  $C$ . It follows that  $y_i^3 l_j^3$  corresponds to one of the chords  $\alpha\beta$  or  $\gamma\delta$ .  
 32 Thus  $l_j^2$  must have degree 3 in  $S$ . It follows that  $l_j^1$  is in  $V(S)$  and so  $y_i^3$  has out-degree 3 in  $S$ , a  
 33 contradiction.  $\square$

34 **Claim 28.5.** For any  $1 \leq i \leq n$  and any  $1 \leq j \leq m$  the cycle  $C$  contains no arc with tail in  $\{x_i^2, \bar{x}_i^2\}$   
 35 and head in  $\{p_j^3, q_j^3, r_j^3\}$ .

36 *Proof.* Assume for a contradiction that  $C$  contains such an arc  $y_i^2 l_j^3$ . The vertex  $y_i^2$  has a unique in-  
 37 neighbour  $y_i^1$  which must be in  $C$ . It follows that  $y_i^1 l_j^3$  corresponds to one of the chords  $\alpha\beta$  or  $\gamma\delta$ .  
 38 Thus  $y_i^2$  must have degree 3 in  $S$ . It follows that  $y_i^3$  is in  $V(S)$  and so  $l_j^3$  has in-degree 3 in  $S$ , a  
 39 contradiction.  $\square$

1 **Claim 28.6.** For any  $1 \leq i \leq n$  and any  $1 \leq j \leq m$  the cycle  $C$  contains no arc with tail in  $\{x_i^1, \bar{x}_i^1\}$   
2 and head in  $\{p_j^3, q_j^3, r_j^3\}$ .

3 *Proof.* Assume for a contradiction that  $C$  contains such an arc  $y_i^1 l_j^3$ . Without loss of generality  $y_i^1 = x_i^1$ .

4 By the remark after Claim 28.1 this is the only arc from a variable gadget to a clause gadget.  
5 Furthermore, we have that  $b$  is not on  $C$ .

6 Thus, by Claim 28.1, for every  $1 \leq k < i$ , the intersection of  $C$  and  $V_k^4$  is either  $X_k$  or  $\bar{X}_k$ , and for  
7 every  $j < l \leq m$ , the intersection of  $C$  and  $C_j^5$  is either  $P_j$ ,  $Q_j$  or  $R_j$ .

8 Consider  $y \in \{\alpha, \beta\}$ . It is on  $C$  and has outdegree 2. On the other hand, applying Claim 28.1 we  
9 see that the following must hold as none of these vertices can belong to  $S$  and at the same time have  
10 two of their out-neighbours in  $S$ :

- 11 •  $y \notin \cup_{1 \leq j \leq m} \{c_j, p_j, q_j, q'_j, q_j^0\}$ ,
- 12 •  $y \notin \cup_{k \neq i} \{a_k, a'_k, x_k^0, x_k^4\}$  and
- 13 •  $y \notin \{a_i, a'_i, x_i^0, x_i^4\}$ .

14 By Claims 28.2-28.5, we have  $y \notin \{x_i^2, x_i^3\}$  and since  $b$  is not on  $C$  we also have  $y \neq b$ . If  $y = x_i^1$ ,  
15 then using that  $y l_j^3$  is an arc of  $C$  we get a contradiction because  $x_i^2 l_j^3$  is an arc (so we cannot obtain  
16 an induced copy of  $S$  using both arcs  $y l_j^3, x_i^2 l_j^3$ ). Hence (as  $y$  was any of  $\alpha, \beta$ ) we have  $a = \alpha = \beta$ , a  
17 contradiction.  $\square$

18 **Claim 28.7.**  $C = acbda$ .

19 *Proof.* Suppose not. Then by the above claims,  $C$  either does not intersect the clause gadget and  
20 intersect all the variable ones or does not intersect the variable gadget and intersect all the gadget ones.  
21 In both cases, similarly to the proof of Claim 28.5, one shows that  $a = \alpha = \beta$ , a contradiction.  $\square$

22 Since  $C = acbda$  and by construction of  $G_5^*(I)$ ,  $T_1$  and  $T_2$  are two induced disjoint path in  $G_5(I)$   
23 and so  $I$  is satisfiable.  $\square$

## 24 7 Remarks and open problems

25 It would be nice to have results proving a full dichotomy between the digraphs  $D$  for which  $\Pi_D$   
26 (resp.  $\Pi'_D$ ) is NP-complete and the ones for which it is polynomial-time solvable. Regarding  $\Pi_D$ ,  
27 Conjecture 14 gives us what the dichotomy should be. But for  $\Pi'_D$  we do not know yet.

28 A useful tool to prove such a dichotomy would be the following conjecture.

29 **Conjecture 29.** If  $D$  is a digraph such that  $\Pi_D$  (resp.  $\Pi'_D$ ) is NP-complete, then for any digraph  $D'$   
30 that contains  $D$  as an induced subdigraph,  $\Pi_{D'}$  (resp.  $\Pi'_{D'}$ ) is NP-complete.

31 We were able to settle the complexity of  $\Pi'_D$  when  $D$  is a directed cycle, a directed path, or some  
32 paths with at most four blocks. The following problems are perhaps the natural next steps.

33 **Problem 30.** What is the complexity of  $\Pi'_D$  when  $D$  is an oriented cycle which is not directed?

34 **Problem 31.** What is the complexity of  $\Pi'_D$  when  $D$  is an oriented path which is not directed?

1 Note that the approach used above to find an induced subdivision of  $A_4^-$  relied on the fact that  
 2 one can check in polynomial time (using flows) whether a digraph contains internally disjoint  $(x,z)$ -,  
 3  $(y,z)$ -paths for prescribed distinct vertices  $x,y,z$ . If we want to apply a similar approach for  $A_5^-$ , then  
 4 for prescribed vertices  $x,y,z,w$  we need to be able to check the existence of internally disjoint paths  
 5  $P,Q,R$  such that  $P$  is an  $(x,y)$ -path,  $Q$  is a  $(z,y)$ -path and  $R$  is a  $(z,w)$ -path such that these paths are  
 6 induced and have no arcs between them. However, the problem of deciding just the existence of  
 7 internally disjoint paths  $P,Q,R$  with these prescribed ends is NP-complete by the result of Fortune et  
 8 al. [6]. Thus we need another approach to obtain a polynomial-time algorithm (if one exists).




Figure 10: The digraph  $H$  with specified vertices  $u_H, v_H, x_H, y_H$ .

9 It seems that little is known about detecting a subdivision of some given digraph  $D$  as a subgraph  
 10 (possibly non-induced). This leads us to the following problem:

11 **Problem 32.** When  $D$  is fixed directed graph, what is the complexity of deciding whether a given  
 12 digraph  $G$  contains a  $D$ -subdivision as a subgraph?

13 The following shows that the problem above can be NP-complete.

14 **Theorem 33.** Let  $H$  be the digraph in Figure 10. It is NP-complete to decide whether a given digraph  
 15  $G$  contains an  $H$ -subdivision.

16 *Proof.* By the classical result of Fortune, Hopcroft and Wyllie [6], the so-called 2-linkage problem  
 17 (given a digraph and four distinct vertices  $u,v,x,y$ ; does  $G$  contain a pair of vertex-disjoint paths  $P,Q$   
 18 so that  $P$  is a directed  $(u,v)$ -path and  $Q$  is a directed  $(x,y)$ -path?) is NP-complete. By inspecting  
 19 the proof (see [1, Section 10.2]) it can be seen that the problem is NP-complete even when  $G$  has  
 20 maximum in- and out-degree at most 3. Given an instance  $G$  of the 2-linkage problem with maximum  
 21 in- and out-degree at most 3 and a copy of  $H$  we form a new digraph  $G_H$  by identifying the vertices  
 22  $\{u,v,x,y\}$  with  $\{u_H, v_H, x_H, y_H\}$  in that order. Clearly, if  $G$  has disjoint directed  $(u,v)$ ,  $(x,y)$ -paths,  
 23 then we can use these to realize the needed paths from  $u_H$  to  $v_H$  and from  $x_H$  to  $y_H$  (and all other  
 24 paths are the original arcs of  $H$ ). Conversely, suppose there is a subdivision  $H^*$  of  $H$  in  $G_H$ . For  
 25 every  $v \in \{u_H, v_H, x_H, y_H\}$ , let us denote by  $v^*$  the vertex corresponding to  $v$  in  $H^*$ . Since  $d^-(u_H) =$   
 26  $4, d^+(v_H) = 5, d^-(x_H) = 5, d^+(y_H) = 4$  in  $G_H$ , we have  $u_H^* = u_H, v_H^* = v_H, x_H^* = x_H$ , and  $y_H^* = y_H$ .  
 27 Thus the two disjoint directed  $(u_H^*, v_H^*)$ - $(x_H^*, y_H^*)$ -paths in  $H^*$  are disjoint directed  $(u,v)$ ,  $(x,y)$ -paths in  
 28  $G$  implying that  $G$  is a 'yes'-instance.  $\square$

29 Finally, we would like to point out that in all detection problems about induced digraphs, back-  
 30 ward arcs of paths play an important role, especially in NP-completeness proofs. Also, these back-  
 31 ward arcs make all "connectivity-flavoured" arguments fail: when two vertices  $x,y$  are given, it is not

1 possible to decide whether  $x$  can be linked to  $y$ . So, maybe another notion of induced subdigraph  
2 containment would make sense: chords should be kept forbidden between the different directed paths  
3 that arise from subdividing arcs, but backward arcs inside the paths should be allowed.

## 4 **Acknowledgement**

5 The authors would like to thank Joseph Yu for stimulating discussions.

## 6 **References**

- 7 [1] J. Bang-Jensen and G. Gutin. *Digraphs: Theory, Algorithms and Applications*. Springer Mono-  
8 graphs in Mathematics. Springer Verlag, London, 2008.
- 9 [2] J. Bang-Jensen and M. Kriesell. Disjoint directed and undirected paths and cycles in digraphs  
10 *Theoretical Computer Science* 410, 5138–5144, 2009.
- 11 [3] D. Bienstock. On the complexity of testing for odd holes and induced odd paths. *Discrete*  
12 *Math.* 90:85–92, 1991. See also Corrigendum by B. Reed, *Discrete Math.* 102:109, 1992.
- 13 [4] M. Chudnovsky and P. Seymour. Excluding induced subgraphs. In *Surveys in Combinatorics*,  
14 volume 346, pages 99–119. London Mathematical Society Lecture Notes Series, 2007.
- 15 [5] M. Chudnovsky and P. Seymour. The three-in-a-tree problem. *Combinatorica*, 2007. To  
16 appear.
- 17 [6] S. Fortune, J.E. Hopcroft, and J. Wyllie. The directed subgraph homeomorphism problem.  
18 *Theoretical Computer Science*, 10:111–121, 1980.
- 19 [7] B. Lévêque, D. Lin, F. Maffray, and N. Trotignon. Detecting induced subgraphs. *Discrete*  
20 *Applied Mathematics*, 157:3540–3551, 2009.
- 21 [8] Y. Kobayashi. Induced disjoint paths problem in a planar digraph *Discrete Applied Mathemat-*  
22 *ics* 157, 3231–3238, 2009.
- 23 [9] N. Robertson and P.D. Seymour. Graph minors. XIII. The disjoint paths problem. *Journal of*  
24 *Combinatorial Theory, Series B*, 63:65–110, 1995.