



Geometrical optics formulaes for Helmholtz equation

Francois Cuvelier

► To cite this version:

| Francois Cuvelier. Geometrical optics formulaes for Helmholtz equation. 2013. <hal-00785774>

HAL Id: hal-00785774

<https://hal.inria.fr/hal-00785774>

Submitted on 7 Feb 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

GEOMETRICAL OPTICS FORMULAE FOR HELMHOLTZ EQUATION

FRANÇOIS CUVELIER

ABSTRACT. The present work deals with high frequency Helmholtz equation resolution using geometrical optics. We give formulae in dimension 2 and 3 for mixed Dirichlet, Neumann and Robin boundaries conditions.

1. INTRODUCTION

Let $(K_j)_{j=1,\dots,N}$ be a set of regular, disjoint and strictly convex compacts in \mathbb{R}^d $d = 2$ or 3 . We suppose that for $j \neq l$, $K_j \cap K_l = \emptyset$. Let $\Omega_j = K_j^c$ and $\Omega = \bigcap_{j=1}^N \Omega_j$. We note Γ_j the boundary of K_j and

$$\Gamma = \bigcup_{j=1}^N \Gamma_j.$$

Let u^{inc} be the incident wave given by

$$u^{inc}(x) = e^{-ik\varphi^{inc}(x)} a^{inc}(x) \quad (1.1)$$

which satisfy Helmholtz equation in \mathbb{R}^d . We want to solve, at *high frequency*, Helmholtz equation

$$\Delta u + k^2 u = 0, \text{ in } \Omega \quad (1.2)$$

with boundaries conditions

$$u(\gamma) = -u^{inc}(\gamma), \quad \forall \gamma \in \Gamma_D, \quad (1.3)$$

$$\frac{\partial u}{\partial n}(\gamma) = -\frac{\partial u^{inc}}{\partial n}(\gamma), \quad \forall \gamma \in \Gamma_N, \quad (1.4)$$

$$\alpha(k) \frac{\partial u}{\partial n}(\gamma) + \beta(k) u(\gamma) = -\left(\alpha(k) \frac{\partial u^{inc}}{\partial n}(\gamma) + \beta(k) u^{inc}(\gamma) \right), \quad \forall \gamma \in \Gamma_R, \quad (1.5)$$

and outgoing Sommerfeld radiation condition

$$r^2 \left(\frac{\partial u}{\partial r} - iku \right) \text{ bound for } r = |x| \rightarrow \infty. \quad (1.6)$$

We assume that

$$\alpha(k) = \sum_{j=0}^{\infty} \frac{\alpha_j}{k^j}, \quad \alpha_j \in \mathbb{C}, \quad (1.7)$$

$$\beta(k) = \sum_{j=-1}^{\infty} \frac{\beta_j}{k^j}, \quad \beta_j \in \mathbb{C}, \quad (1.8)$$

and

$$\beta_{-1} + i\alpha_0 \mu \neq 0 \quad \forall \mu \in [0; 1] \quad (1.9)$$

Here we take Γ_D, Γ_N and Γ_R such that $\Gamma = \Gamma_D \cup \Gamma_N \cup \Gamma_R$, $\overset{o}{\Gamma}_D \cap \overset{o}{\Gamma}_N = \emptyset$, $\overset{o}{\Gamma}_D \cap \overset{o}{\Gamma}_R = \emptyset$ and $\overset{o}{\Gamma}_N \cap \overset{o}{\Gamma}_R = \emptyset$.

Date: September 12, 2012.




FIGURE 1. Exterior Helmholtz problem

We assume that, Σ , the front of the incident wave is given by

$$\Sigma = \{x \in \mathbb{R}^d \mid \varphi^{inc}(x) = \varphi_\Sigma\}$$

is a regular and orientable surface ($d = 3$) or curve ($d = 2$). We chose the orientation $N(\sigma) = \frac{\nabla \varphi^{inc}(\sigma)}{\|\nabla \varphi^{inc}(\sigma)\|}$.

Let $\Sigma_+ = \{x \in \mathbb{R}^d \mid \exists \sigma \in \Sigma, \exists t > 0, x = \sigma + t \nabla \varphi^{inc}(\sigma)\}$, we also made following assumption :

- (i) $\bigcup_{j=1}^N K_j \subset \Sigma_+$
- (ii) curvature radius of Σ are negatives

At *high frequency* (means that wave length is small with respect to the obstacle boundary curvature radius), geometrical optics approximation for this problem is based on k^{-1} asymptotic expansion of $v(x) = u(x) + u^{inc}(x)$, where u is solution of problem (1.2,1.3,1.4,1.5,1.6). The first term of the k^{-1} expansion is the geometrical optics ansatz.

2. NOTATIONS AND RESULTS

Definition 1 (Geometrical optic ray). Let $\rho = (\sigma, \gamma_1, \dots, \gamma_l) = (\gamma_0, \gamma_1, \dots, \gamma_l) \in \Sigma \times \Gamma^l$, we say that ρ is a geometrical optic ray coming from Σ , going through $x = \gamma_{l+1}$ and reflected l times if

- (1) $\gamma_j \neq \gamma_{j-1}, \forall j \in \{1, \dots, l+1\}$,
- (2) $\left(\bigcup_{j=1}^{l+1} [\gamma_{j-1}; \gamma_j] \right) \cap \Gamma = \emptyset$,
- (3) $\frac{\gamma_1 - \gamma_0}{|\gamma_1 - \gamma_0|} = \nabla \varphi^{inc}(\gamma_0)$,
- (4) **reflections conditions** : $\forall j \in \{1, \dots, l\}$

$$\frac{\gamma_{j+1} - \gamma_j}{|\gamma_{j+1} - \gamma_j|} = \frac{\gamma_j - \gamma_{j-1}}{|\gamma_j - \gamma_{j-1}|} - 2 \left\langle \frac{\gamma_j - \gamma_{j-1}}{|\gamma_j - \gamma_{j-1}|}, \mathbf{n}(\gamma_j) \right\rangle \mathbf{n}(\gamma_j),$$

- (5) **non grazing conditions** : $\forall j \in \{1, \dots, l\}$

$$\left\langle \frac{\gamma_j - \gamma_{j-1}}{|\gamma_j - \gamma_{j-1}|}, \mathbf{n}(\gamma_j) \right\rangle \neq 0,$$

where $\mathbf{n}(\gamma)$ note the exterior normal at Γ in γ .

We note $\mathcal{R}_l(x)$ a such set of ρ and $\mathcal{R}(x) = \bigcup_{l \in \mathbb{N}^*} \mathcal{R}_l(x)$.

Definition 2 (Grazing ray). Let $\rho = (\sigma, \gamma_1, \dots, \gamma_l) = (\gamma_0, \gamma_1, \dots, \gamma_l) \in \Sigma \times \Gamma^l$, we say that ρ is a grazing ray coming from Σ , going through $x = \gamma_{l+1}$ and reflected l times if conditions (1) to (4) of definition 1 are satisfied and not condition (5). We note $\mathcal{G}_l(x)$ a such set of ρ and $\mathcal{G}(x) = \bigcup_{l \in \mathbb{N}^*} \mathcal{G}_l(x)$.

Definition 3 (Propagation operator). Let $t > 0$, we note $S_t^d \subset \mathbb{R}^{d \times d}$ the set of matrix \mathbb{A} such that $\mathbb{I} + t\mathbb{A}$ is regular. We note \mathcal{S}_t the following application :

$$\begin{aligned} \mathcal{S}_t & : S_t^d \longrightarrow \mathbb{R}^{d \times d} \\ \mathbb{A} & \longmapsto \mathbb{A}(\mathbb{I} + t\mathbb{A})^{-1} \end{aligned}$$

Definition 4 (Reflection operator). Let $\mathbb{B} \in \mathbb{R}^{d \times d}$ a symmetric matrix. Let $\eta \in \mathbb{R}^d$ and $\zeta \in \mathbb{R}^d$ such that $\langle \zeta, \eta \rangle \neq 0$. We note $\mathcal{T}_{\mathbb{B}, \eta, \zeta}^d : \mathbb{R}^{d \times d} \rightarrow \mathbb{R}^{d \times d}$ given by :
 $\forall \mathbb{A} \in \mathbb{R}^{d \times d}, \forall x \in \mathbb{R}^d$

$$\begin{aligned} (\mathcal{T}_{\mathbb{B}, \eta, \zeta}^d(\mathbb{A}))x &= (\mathbb{A} - 2\langle \zeta, \eta \rangle \mathbb{B})x - 2\langle \eta, x \rangle (\mathbb{A}\eta + \mathbb{B}\zeta) \\ &\quad - 2\langle \mathbb{A}\eta + \mathbb{B}\zeta, x \rangle \eta + 2\left[2\langle \mathbb{A}\eta, \eta \rangle - \frac{\langle \mathbb{B}\zeta, \zeta \rangle}{\langle \zeta, \eta \rangle}\right]\langle \eta, x \rangle \eta \end{aligned}$$

Definition 5 (Reflection coefficient). Let $\eta \in \mathbb{R}^d$, we note b_η the function defined on Γ by

$$b_\eta(\gamma) = \begin{cases} -1, & \text{if } \gamma \in \Gamma_D, \\ 1, & \text{if } \gamma \in \Gamma_N, \\ i\alpha_0 \langle \eta, \mathbf{n}(\gamma) \rangle + \beta_{-1}, & \text{if } \gamma \in \Gamma_R, \\ i\alpha_0 \langle \eta, \mathbf{n}(\gamma) \rangle - \beta_{-1}, & \text{if } \gamma \in \Gamma_R. \end{cases}$$

Theorem 1. Let $x \in \Sigma_+$, such that $\mathcal{G}(x) = \emptyset$, and $\rho = (\gamma_0, \gamma_1, \dots, \gamma_l) \in \mathcal{R}_l(x)$, $l \in \mathbb{N}^*$. We note $\mathbb{B}(\gamma)$ the curvature matrix of Γ at γ and $\mathbf{n}(\gamma)$ the exterior normal to Γ at γ . Let $u_\rho^{G.O.}(x)$ be the ray contribution :

$$u_\rho^{G.O.}(x) = a_\rho(x)e^{-ik\varphi(x; \rho)}$$

with

$$\varphi(x; \rho) = \varphi_\rho^{(l)}(\gamma_l) + \|x - \gamma_l\|$$

$$a_\rho(x) = \frac{a_\rho^{(l)}(\gamma_l)}{\sqrt{\det(\mathbb{I} + \|x - \gamma_l\| \mathbb{A}_\rho^{(l)}(\gamma_l))}}$$

and for $j = l$ to 1,

$$\begin{cases} \varphi_\rho^{(j)}(\gamma_j) &= \varphi_\rho^{(j-1)}(\gamma_{j-1}) + \|\gamma_j - \gamma_{j-1}\| \\ a_\rho^{(j)}(\gamma_j) &= b_{\frac{\gamma_j - \gamma_{j-1}}{\|\gamma_j - \gamma_{j-1}\|}}(\gamma_j) \frac{a_\rho^{(j-1)}(\gamma_{j-1})}{\sqrt{\det(\mathbb{I} + \|\gamma_j - \gamma_{j-1}\| \mathbb{A}_\rho^{(j-1)}(\gamma_{j-1}))}} \\ \mathbb{A}_\rho^{(j)}(\gamma_j) &= \mathcal{T}_{\mathbb{B}(\gamma_j), \mathbf{n}(\gamma_j), \frac{\gamma_j - \gamma_{j-1}}{\|\gamma_j - \gamma_{j-1}\|}}^d \circ \mathcal{S}_{\|\gamma_j - \gamma_{j-1}\|} \left(\mathbb{A}_\rho^{(j-1)}(\gamma_{j-1}) \right) \end{cases}$$

where $\varphi_\rho^{(0)}(\gamma_0) = \varphi^{inc}(\gamma_0)$, $\mathbb{A}_\rho^{(0)}(\gamma_0) = \text{Hess } \varphi^{inc}(\gamma_0)$ and $a_\rho^{(0)}(\gamma_0) = a_0^{inc}(\gamma_0)$.

Then, we have

$$u^{inc}(x) = \sum_{\rho \in \mathcal{R}_0(x)} u_\rho^{G.O.}(x) + O\left(\frac{1}{k}\right)$$

and

$$u(x) = \sum_{\rho \in \mathcal{R}(x)} u_\rho^{G.O.}(x) + O\left(\frac{1}{k}\right)$$

3. PROOF

To obtain theorem 1, we first give geometrical optics formulae without obstacle, then with a stricly convex compact and finally with an union of stricly convex compacts.

3.1. Without obstacles. We want to solve, at high frequency, Helmholtz equation

$$\Delta u + k^2 u = 0 \text{ in } \mathbb{R}^d \quad (3.1)$$

using an asymptotic expansion in k^{-1} of the solution

$$u(x; k) = \sum_{j \in \mathbb{N}} \frac{a_j(x)}{k^j} e^{-ik\varphi(x)} \quad (3.2)$$

where we suppose that u is known on Σ .

By substituting expansion (3.2) in Helmholtz equation, we obtain

$$\begin{aligned} & k^2(1 - \|\nabla \varphi(x)\|^2)a_0(x) \\ & + k \left[-i\{2(\nabla \varphi(x), \nabla a_0(x)) + a_0(x)\Delta \varphi(x)\} + (1 - \|\nabla \varphi(x)\|^2)a_1(x) \right] \\ & + \sum_{j \in \mathbb{N}} k^{-j} (-i\{2(\nabla \varphi(x), \nabla a_{k+1}(x)) + a_{k+1}(x)\Delta \varphi(x)\} + \Delta a_k(x)) = 0 \end{aligned}$$

Equating the coefficients of different powers of k , we have :

- **Eikonal equation**

$$\|\nabla \varphi(x)\|^2 = 1 \quad (3.3)$$

- **Transport equation for a_0**

$$\langle \nabla \varphi(x), \nabla a_0(x) \rangle + \frac{1}{2}a_0(x)\Delta \varphi(x) = 0 \quad (3.4)$$

- **Transport equation for a_j in function of a_{j-1}**

$$\langle \nabla \varphi(x), \nabla a_j(x) \rangle + \frac{1}{2}a_j(x)\Delta \varphi(x) = \frac{i}{2}\Delta a_{j-1}(x) \quad (3.5)$$

3.1.1. *Solution of Eikonal equation (3.3).* Derivating eikonal equation (3.3) give

$$\text{Hess } \varphi(x) \nabla \varphi(x) = 0 \quad (3.6)$$

This equation can be solved by the characteristics method :

Let $X'(t) = \nabla \varphi(X(t))$ with $X(0) = \sigma \in \Sigma$ then we have :

$$\begin{aligned} \frac{d}{dt}(\nabla \varphi(X(t))) &= \text{Hess } \varphi(X(t))X'(t) \\ &= \text{Hess } \varphi(X(t))\nabla \varphi(X(t)) \\ &= 0 \end{aligned}$$

That is to say

$$\nabla \varphi(X(t)) = \nabla \varphi(X(0))$$

hence

$$X(t) = X(0) + t\nabla \varphi(X(0)).$$

and $\nabla \varphi(X(t))$ is constant along the line $X(t) = X(0) + t\nabla \varphi(X(0))$.

Owing to the construction of Σ , we have

$$X(t) = \sigma + t\mathbf{N}(\sigma)$$

with $\sigma \in \Sigma$ and $\mathbf{N}(\sigma) = \nabla \varphi(\sigma)$ unitary normal vector to Σ at σ .

We also have :

$$\begin{aligned} \frac{d}{dt}(\varphi(X(t))) &= \langle \nabla \varphi(X(t)), X'(t) \rangle \\ &= \langle \nabla \varphi(X(0)), \nabla \varphi(X(0)) \rangle \\ &= 1 \end{aligned}$$

Thus

$$\varphi(X(t)) = \varphi(X(0)) + t.$$

We have proved following lemma :

Lemma 1. (1) *Characteristic curves of Eikonal equation are geometrical optic rays,*
 (2) *Phase is linear along geometrical optic rays :*

$$\forall \sigma \in \Sigma, \forall t \geq 0 \quad \varphi(\sigma + t\mathbf{N}(\sigma)) = \varphi(\sigma) + t. \quad (3.7)$$

- (3) *We have $\forall \sigma \in \Sigma$*

$$\nabla \varphi(\sigma) = \mathbf{N}(\sigma) \quad (3.8)$$

and

$$\text{Hess } \varphi(\sigma)\mathbf{N}(\sigma) = 0. \quad (3.9)$$

3.1.2. *Solution of transport equation (3.4)*. We want to solve

$$\begin{cases} \langle \nabla \varphi(x), \nabla a_0(x) \rangle + \frac{1}{2} a_0(x) \Delta \varphi(x) = 0 \\ a_0 \text{ given on } \Sigma \end{cases} \quad (3.10)$$

In fact, we only have to solve this equation along the ray $\sigma + t\mathbf{N}(\sigma)$ where $\sigma \in \Sigma$ and $\mathbf{N}(\sigma) = \nabla \varphi(\sigma)$.

Thus, transport equation for a_0 becomes

$$\begin{cases} \frac{d}{dt}(a_0(\sigma + t\mathbf{N}(\sigma))) + \frac{1}{2} \Delta \varphi(\sigma + t\mathbf{N}(\sigma)) a_0(\sigma + t\mathbf{N}(\sigma)) = 0, \\ a_0(\sigma) \text{ given on } \Sigma. \end{cases} \quad (3.11)$$

We now have to compute $\Delta \varphi(\sigma + t\mathbf{N}(\sigma)) = \text{Tr} \text{Hess } \varphi(\sigma(t))$.

Let $\sigma \in \Sigma$, we note for $d = 3$ (resp. $d = 2$) $\mathcal{B}_\Sigma(\sigma) = \{\mathbf{u}, \mathbf{v}, \mathbf{N}\}(\sigma)$ (resp. $\mathcal{B}_\Sigma(\sigma) = \{\mathbf{u}, \mathbf{N}\}(\sigma)$) the **direct orthonormal curvature basis** of Σ at σ . Here \mathbf{u} and \mathbf{v} are the direction of maximum and minimal principal curvature $k_1^{(0)}(\sigma)$ and $k_2^{(0)}(\sigma)$ (resp. \mathbf{u} is the tangent vector and $k^{(0)}(\sigma)$ the curvature). By hypothesis on Σ , we have

$$k_2^{(0)}(\sigma) \leq k_1^{(0)}(\sigma) \leq 0 \quad (\text{resp. } k^{(0)}(\sigma) \leq 0).$$

With these notations, we have in basis $\mathcal{B}_\Sigma(\sigma)$

$$\text{Hess } \varphi(\sigma) = \begin{pmatrix} -k_1^{(0)}(\sigma) & 0 & 0 \\ 0 & -k_2^{(0)}(\sigma) & 0 \\ 0 & 0 & 0 \end{pmatrix} \quad (\text{resp. } \text{Hess } \varphi(\sigma) = \begin{pmatrix} -k^{(0)}(\sigma) & 0 \\ 0 & 0 \end{pmatrix})$$

Lemma 2. Let $\sigma \in \Sigma$. We note $\sigma(t) = \sigma + t\mathbf{N}(\sigma)$ then

$$\forall t \geq 0, (\text{Hess } \varphi)(\sigma(t)) = \mathcal{S}_t(\text{Hess } \varphi(\sigma)) \quad (3.12)$$

In basis $\mathcal{B}_\Sigma(\sigma)$ for $d = 3$ (resp. $d = 2$)

$$\text{Hess } \varphi(\sigma(t)) = \begin{pmatrix} -k_1^{(0)}(\sigma(t)) & 0 & 0 \\ 0 & -k_2^{(0)}(\sigma(t)) & 0 \\ 0 & 0 & 0 \end{pmatrix} \quad (\text{resp. } \text{Hess } \varphi(\sigma(t)) = \begin{pmatrix} -k^{(0)}(\sigma(t)) & 0 \\ 0 & 0 \end{pmatrix})$$

with

$$\begin{aligned} 0 \geq k_j^{(0)}(\sigma(t)) &= \frac{k_j^{(0)}(\sigma)}{1 - tk_j^{(0)}(\sigma)} \geq k_j^{(0)}(\sigma) \quad j = 1, 2 \\ (\text{resp. } 0 \geq k^{(0)}(\sigma(t)) &= \frac{k^{(0)}(\sigma)}{1 - tk^{(0)}(\sigma)} \geq k^{(0)}(\sigma)) \end{aligned}$$

Proof. To simplify notations, we note $\forall t \geq 0$

$$\mathbb{A}_t(\sigma) = (\text{Hess } \varphi)(\sigma + t\mathbf{N}(\sigma)).$$

Let $\mathcal{V}(\sigma) \subset \mathbb{R}^d$ be a neighborhood of $\sigma \in \Sigma$. We have, $\forall x \in \mathcal{V}(\sigma)$

$$\nabla \varphi(x) = \nabla \varphi(\sigma) + \mathbb{A}_0(\sigma)(x - \sigma) + O(\|x - \sigma\|) \quad (3.13)$$

Substituting $\nabla \varphi(x)$ by (3.13) in Eikonal equation (3.3) give, $\forall x \in \mathcal{V}(\sigma)$:

$$\begin{aligned} 1 &= \left\| \nabla \varphi(\sigma) + \mathbb{A}_0(\sigma)(x - \sigma) + O(\|x - \sigma\|^2) \right\|^2 \\ &= \|\nabla \varphi(\sigma)\|^2 + 2 \langle \mathbb{A}_0(\sigma)(x - \sigma), \nabla \varphi(\sigma) \rangle + O(\|x - \sigma\|^2) \\ &= 1 + 2 \langle \mathbb{A}_0(\sigma)(x - \sigma), \mathbf{N}(\sigma) \rangle + O(\|x - \sigma\|^2). \end{aligned}$$

Using $\mathbb{A}_0(\sigma)$ symmetry, we obtain

$$\forall x \in \mathcal{V}(\sigma), \langle \mathbb{A}_0(\sigma)\mathbf{N}(\sigma), x - \sigma \rangle = 0$$

Thus

$$\mathbb{A}_0(\sigma)\mathbf{N}(\sigma) = 0. \quad (3.14)$$

We now want to obtain a similar formula for $\mathbb{A}_t(\sigma)$.

We know that $\mathbb{A}_0(\sigma)$ is the curvature matrix of Σ at σ and in basis $\mathcal{B}_\Sigma(\sigma)$ we have for $d = 3$ (resp. $d = 2$)

$$\mathbb{A}_0(\sigma) = \begin{pmatrix} -k_1^\Sigma(\sigma) & 0 & 0 \\ 0 & -k_2^\Sigma(\sigma) & 0 \\ 0 & 0 & 0 \end{pmatrix} \quad (\text{resp. } \mathbb{A}_0(\sigma) = \begin{pmatrix} -k^\Sigma(\sigma) & 0 \\ 0 & 0 \end{pmatrix}).$$

By hypothesis, curvature radius of Σ are negative and

$$k_2^\Sigma(\sigma) \leq k_1^\Sigma(\sigma) \leq 0 \quad (\text{resp. } k^\Sigma(\sigma) \leq 0)$$

Thus, $\forall t \geq 0$, $\mathbb{I} + t\mathbb{A}_0(\sigma)$ is regular. Let $x = \sigma + t\mathbf{N}(\sigma)$ with $t > 0$, then, there exists a neighborhood $\mathcal{V}(x) \subset \mathbb{R}^d$ of x such that

$$\forall x' \in \mathcal{V}(x), \exists \sigma' \in \mathcal{V}(\sigma) \text{ and } t' \text{ in a neighborhood of } t \text{ verifying } x' = \sigma' + t'\mathbf{N}(\sigma')$$

So we obtain by Taylor's developpement

$$\nabla \varphi(x') = \nabla \varphi(x) + \mathbb{A}_t(\sigma)(x' - x) + O(\|x' - x\|) \quad (3.15)$$

Substituting $\nabla \varphi(x')$ for (3.15) in eikonal equation (3.3) give

$$\mathbb{A}_t(\sigma)\mathbf{N}(\sigma) = 0 \quad (3.16)$$

To prove formula (3.12) we write

$$\begin{aligned} x' - x &= \sigma' + t'\mathbf{N}(\sigma') - \sigma - t\mathbf{N}(\sigma) \\ &= \sigma' - \sigma + (t' - t)\mathbf{N}(\sigma) + t'(\mathbf{N}(\sigma') - \mathbf{N}(\sigma)). \end{aligned}$$

But, due to (3.13)

$$\mathbf{N}(\sigma') - \mathbf{N}(\sigma) = \mathbb{A}_0(\sigma)(\sigma' - \sigma) + O(\|\sigma' - \sigma\|) \quad (3.17)$$

and, thus

$$x' - x = \sigma' - \sigma + (t' - t)\mathbf{N}(\sigma) + t'\mathbb{A}_0(\sigma)(\sigma' - \sigma) + O(\|\sigma' - \sigma\|). \quad (3.18)$$

Substituing $x' - x$ for (3.18) in (3.15) and using (3.16) give

$$\mathbf{N}(\sigma') = \mathbf{N}(\sigma) + \mathbb{A}_t(\sigma)(\sigma' - \sigma) + t'\mathbb{A}_t(\sigma)\mathbb{A}_0(\sigma)(\sigma' - \sigma) + O(\|\sigma' - \sigma\| + \|t' - t\|).$$

Now, with formula (3.17) we found

$$\begin{aligned} &\mathbb{A}_0(\sigma)(\sigma' - \sigma) + O(\|\sigma' - \sigma\|) \\ &= \mathbb{A}_t(\sigma)(\sigma' - \sigma) + t'\mathbb{A}_t(\sigma)\mathbb{A}_0(\sigma)(\sigma' - \sigma) + O(\|\sigma' - \sigma\| + \|t' - t\|) \end{aligned}$$

That is to say

$$\begin{aligned} \mathbb{A}_0(\sigma) &= \mathbb{A}_t(\sigma) + t\mathbb{A}_t(\sigma)\mathbb{A}_0(\sigma) \\ &= \mathbb{A}_t(\sigma)(\mathbb{I} + t\mathbb{A}_0(\sigma)). \end{aligned}$$

But, $\forall t \geq 0$, the matrix $(\mathbb{I} + t\mathbb{A}_0(\sigma))$ is regular and formula (3.12) is proved. \square

Lemma 3. *Let $\sigma \in \Sigma$. Along the ray $\sigma(t) = \sigma + t\mathbf{N}(\sigma)$, $\forall t \geq 0$, we have*

$$a_0(\sigma(t)) = \frac{a_0(\sigma)}{\sqrt{\det(\mathbb{I} + t \operatorname{Hess} \varphi(\sigma))}} \quad (3.19)$$

and $\forall t > 0$, $\forall t' \geq 0$ such that $t > t'$

$$|a_0(\sigma(t))| \leq |a_0(\sigma(t'))|. \quad (3.20)$$

The previous inequality becomes strict if, at least, one of the curvature radius of Σ is strictly negative.

Proof. By definition of $\mathbb{A}_t(\sigma)$ (see proof of lemma 2)

$$\Delta \varphi(\sigma(t)) = \operatorname{Tr}(\mathbb{A}_t(\sigma))$$

Then we solve equation (3.11):

$$\begin{aligned} a_0(\sigma(t)) &= a_0(\sigma)e^{-\frac{1}{2}\int_0^t \operatorname{Tr} \mathbb{A}_\tau(\sigma)d\tau} \\ &= a_0(\sigma)e^{-\frac{1}{2}\int_0^t \operatorname{Tr}[\mathbb{A}_0(\sigma)(I + \tau\mathbb{A}_0(\sigma))^{-1}]d\tau} \\ &= a_0(\sigma)e^{-\frac{1}{2}[\log \det(\mathbb{I} + t\mathbb{A}_0(\sigma))]_0^t} \\ &= \frac{a_0(\sigma)}{\sqrt{\det(\mathbb{I} + t\mathbb{A}_0(\sigma))}}. \end{aligned}$$

We have seen, in proof of lemma 2, that eigenvalues of $\mathbb{A}_0(\sigma)$ are positive, thus

$$\det(\mathbb{I} + t'\mathbb{A}_0(\sigma)) \geq \det(\mathbb{I} + t\mathbb{A}_0(\sigma)) \geq 1.$$

and inequality (3.20) is immediatly prooved.

We can remark that if, at least, one of the curvature of Σ at σ is strictly negative then, for $d = 3$ (resp. $d = 2$)

$$k_2^\Sigma(\sigma) < k_1^\Sigma(\sigma) \leq 0 \quad (\text{resp. } k^\Sigma(\sigma) < 0)$$

and thus inequality (3.20) become strict. \square

3.1.3. *Conclusion.* The knowledge of a_0 , φ , $\nabla \varphi$ and $\text{Hess } \varphi$ on Σ is sufficient to compute them for all x in Σ_+ using formulae along geometrical optic rays comming through x and we have




FIGURE 2. Wave Propagation

Theorem 2. Let $x \in \Sigma_+$, then there exist an unique $\sigma \in \Sigma$ such that

$$x = \sigma + \|x - \sigma\| N(\sigma) \quad (3.21)$$

and solution of (3.1-3.2) is given by

$$u(x; k) = \frac{a_0(\sigma)}{\sqrt{\det(\mathbb{I} + \|x - \sigma\| \text{Hess } \varphi(\sigma))}} e^{-ik(\varphi(\sigma) + \|x - \sigma\|)} + O\left(\frac{1}{k}\right) \quad (3.22)$$

Proof. By definition of Σ_+ , we have existence of $\sigma \in \Sigma$ satisfying (3.21).

For unicity, we suppose there exists $\sigma_1 \in \Sigma$ and $\sigma_2 \in \Sigma$ satisfying (3.21). By convexity hypothesis on Σ , we have

$$\langle \sigma_2 - \sigma_1, N(\sigma_1) \rangle \leq 0$$

and then

$$\langle \sigma_2 - \sigma_1, x - \sigma_1 \rangle \leq 0.$$

But

$$\langle \sigma_2 - \sigma_1, x - \sigma_1 \rangle = \langle \sigma_2 - \sigma_1, x - \sigma_2 \rangle + \|\sigma_2 - \sigma_1\|^2$$

so we obtain

$$\langle \sigma_2 - \sigma_1, x - \sigma_2 \rangle \leq 0. \quad (3.23)$$

By convexity hypothesis on Σ , we have

$$\langle \sigma_1 - \sigma_2, N(\sigma_2) \rangle \leq 0$$

and then

$$\langle \sigma_1 - \sigma_2, x - \sigma_2 \rangle \leq 0. \quad (3.24)$$

Thus, inequalities (3.23) and (3.24) give us $\sigma_1 \equiv \sigma_2$.

Formula (3.22) is just an application of propagation Lemmas (lemma 1 and lemma 3) along the ray $\sigma \in \mathcal{R}_0(x)$. \square

3.2. Outside a strictly convex compact. Let $K \subset \mathbb{R}^d$ be a regular and strictly convex compact, $\Omega = K^c$ and $\Gamma = \partial K$ be its boundary.

Remark 1. Owing to the strict convexity of K and the hypotheses on Σ , we have

$$\forall x \in \Sigma_+ \cap \Omega, \forall l > 1, \mathcal{R}_l(x) = \emptyset.$$

There is only one reflexion on K .

We denote by

$$\Gamma^s = \{\gamma \in \Gamma \mid \mathcal{R}_0(\gamma) = \emptyset\} \text{ and } \Gamma^e = (\Gamma^s)^c$$

Let $\gamma \in \Gamma^e$ and $\sigma \in \mathcal{R}_0(\gamma)$. Using previous formulas give us $a_0^{(0)}(\sigma)$, $\varphi^{(0)}(\sigma)$, $\nabla \varphi^{(0)}(\sigma)$ and $\text{Hess } \varphi^{(0)}(\sigma)$ on γ .

If we can compute reflected wave on γ (i.e. $a_0^{(1)}(\sigma')$, $\varphi^{(1)}(\sigma')$, $\nabla \varphi^{(1)}(\sigma')$ and $\text{Hess } \varphi^{(1)}(\sigma')$ on γ) then we can use propagation formulae along the reflected ray given by $\gamma + t \nabla \varphi^{(1)}(\gamma)$, $t > 0$




FIGURE 3. Wave reflection

So, the local problem is only to find how to compute the reflected wave in γ . That is to say :

$$\begin{cases} a_0^{(0)}(\gamma) \\ \varphi^{(0)}(\gamma) \\ \nabla \varphi^{(0)}(\gamma) \\ \text{Hess } \varphi^{(0)}(\gamma) \end{cases} \quad \text{given, how to compute} \quad \begin{cases} a_0^{(1)}(\gamma) \\ \varphi^{(1)}(\gamma) \\ \nabla \varphi^{(1)}(\gamma) \\ \text{Hess } \varphi^{(1)}(\gamma) \end{cases} \quad ?$$

3.2.1. Computation of $\varphi^{(1)}$ and $\nabla \varphi^{(1)}$ in γ .

Lemma 4. Let $\gamma \in \Gamma$, then

$$\varphi^{(1)}(\gamma) = \varphi^{(0)}(\gamma) \quad (3.25)$$

Proof. Due to boundary condition :

- If $\gamma \in \Gamma_D$, we have

$$a^{(1)}(\gamma) e^{-ik\varphi^{(1)}(\gamma)} = -a^{(0)}(\gamma) e^{-ik\varphi^{(0)}(\gamma)} \quad (3.26)$$

- If $\gamma \in \Gamma_N$, we have

$$\begin{aligned} & \left(\frac{\partial a^{(1)}}{\partial n}(\gamma) - ika^{(1)}(\gamma) \frac{\partial \varphi^{(1)}}{\partial n}(\gamma) \right) e^{-ik\varphi^{(1)}(\gamma)} \\ &= \\ & - \left(\frac{\partial a^{(0)}}{\partial n}(\gamma) - ika^{(0)}(\gamma) \frac{\partial \varphi^{(0)}}{\partial n}(\gamma) \right) e^{-ik\varphi^{(0)}(\gamma)} \end{aligned} \quad (3.27)$$

- If $\gamma \in \Gamma_R$, we have

$$\begin{aligned} & \left[\alpha(k) \left(\frac{\partial a^{(1)}}{\partial n}(\gamma) - ika^{(1)}(\gamma) \frac{\partial \varphi^{(1)}}{\partial n}(\gamma) \right) + \beta(k)a^{(1)}(\gamma) \right] e^{-ik\varphi^{(1)}(\gamma)} \\ &= \\ & - \left[\alpha(k) \left(\frac{\partial a^{(0)}}{\partial n}(\gamma) - ika^{(0)}(\gamma) \frac{\partial \varphi^{(0)}}{\partial n}(\gamma) \right) + \beta(k)a^{(0)}(\gamma) \right] e^{-ik\varphi^{(0)}(\gamma)} \end{aligned} \quad (3.28)$$

By identification, we immediately obtain (3.25). \square

Lemma 5. Let $\gamma \in \Gamma$, and $\mathbf{n}(\gamma)$ the exterior normal to Γ at γ . We have

$$\nabla \varphi^{(1)}(\gamma) = \nabla \varphi^{(0)}(\gamma) - 2 \langle \nabla \varphi^{(0)}(\gamma), \mathbf{n}(\gamma) \rangle \mathbf{n}(\gamma) \quad (3.29)$$

Proof. Let $\mathcal{B}_\Gamma(\gamma) = \{\mathbf{u}(\gamma), \mathbf{v}(\gamma), \mathbf{n}(\gamma)\}$ be the **direct orthonormal curvature basis** of Γ at γ . Then we have the local parametrization of Γ at γ :

$$\gamma(u, v) = \gamma + (u, v, g(u, v))$$

with $g(u, v) = \frac{1}{2}(k_1^\Gamma(\gamma)u^2 + k_2^\Gamma(\gamma)v^2) + o(u^2 + v^2)$

Taylor's expansion at order 1 of $\varphi^{(0)}$ and $\varphi^{(1)}$ on Γ at point γ are

$$\begin{aligned} \varphi^{(0)}(\gamma(u, v)) &= \varphi^{(0)}(\gamma) + \langle \nabla \varphi^{(0)}(\gamma), u\mathbf{u}(\gamma) + v\mathbf{v}(\gamma) \rangle + O(u^2 + v^2) \\ \varphi^{(1)}(\gamma(u, v)) &= \varphi^{(1)}(\gamma) + \langle \nabla \varphi^{(1)}(\gamma), u\mathbf{u}(\gamma) + v\mathbf{v}(\gamma) \rangle + O(u^2 + v^2) \end{aligned}$$

Due to relation (3.25) we obtain

$$\langle \nabla \varphi^{(1)}(\gamma) - \nabla \varphi^{(0)}(\gamma), u\mathbf{u}(\gamma) + v\mathbf{v}(\gamma) \rangle + O(u^2 + v^2)$$

that is to say

$$\langle \nabla \varphi^{(1)}(\gamma) - \nabla \varphi^{(0)}(\gamma), \mathbf{u}(\gamma) \rangle = \langle \nabla \varphi^{(1)}(\gamma) - \nabla \varphi^{(0)}(\gamma), \mathbf{v}(\gamma) \rangle = 0$$

So, exists $\lambda \in \mathbb{R}$ such that

$$\nabla \varphi^{(1)}(\gamma) = \nabla \varphi^{(0)}(\gamma) + \lambda \mathbf{n}(\gamma)$$

Taking the norm of previous relation and using eikonal equation (3.6) give

$$\lambda = -2 \langle \nabla \varphi^{(0)}(\gamma), \mathbf{n}(\gamma) \rangle.$$

A similar proof will act in dimension 2. \square

3.2.2. Computation of $a_0^{(1)}$ in γ .

Lemma 6. Let $\gamma \in \Gamma^e$. If we know $a_0^{(0)}(\gamma)$ and $\nabla \varphi^{(0)}(\gamma)$ then

$$a_0^{(1)}(\gamma) = b_{\nabla \varphi^{(0)}(\gamma)}(\gamma) a_0^{(0)}(\gamma). \quad (3.30)$$

where the function $b_{\nabla \varphi^{(0)}(\gamma)}$ is given in definition 5.

Proof. • If $\gamma \in \Gamma_D \cap \Gamma^e$, formula (3.26) give

$$a^{(1)}(\gamma) = -a^{(0)}(\gamma)$$

Using asymptotic expansions of $a^{(0)}$ and $a^{(1)}$

$$\sum_{j \in \mathbb{N}} k^{-j} a_j^{(1)}(\gamma) = - \sum_{j \in \mathbb{N}} k^{-j} a_j^{(0)}(\gamma)$$

With high frequency hypothesis, we obtain the leading term in power of k :

$$a_0^{(1)}(\gamma) = -a_0^{(0)}(\gamma)$$

- If $\gamma \in \Gamma_N \cap \Gamma^e$, formula (3.27) give

$$\frac{\partial a^{(1)}}{\partial n}(\gamma) - \iota k a^{(1)}(\gamma) \frac{\partial \varphi^{(1)}}{\partial n}(\gamma) = - \left(\frac{\partial a^{(0)}}{\partial n}(\gamma) - \iota k a^{(0)}(\gamma) \frac{\partial \varphi^{(0)}}{\partial n}(\gamma) \right)$$

Using asymptotic expansions of $a^{(0)}$ and $a^{(1)}$

$$\begin{aligned} & \sum_{j \in \mathbb{N}} \left(k^{-j} \frac{\partial a_j^{(1)}}{\partial n}(\gamma) - \iota k^{-j+1} a_j^{(1)}(\gamma) \frac{\partial \varphi^{(1)}}{\partial n}(\gamma) \right) \\ &= \\ & - \sum_{j \in \mathbb{N}} \left(k^{-j} \frac{\partial a_j^{(0)}}{\partial n}(\gamma) - \iota k^{-j+1} a_j^{(0)}(\gamma) \frac{\partial \varphi^{(0)}}{\partial n}(\gamma) \right) \end{aligned}$$

The leading term in power of k is

$$a_0^{(1)}(\gamma) \frac{\partial \varphi^{(1)}}{\partial n}(\gamma) = -a_0^{(0)}(\gamma) \frac{\partial \varphi^{(0)}}{\partial n}(\gamma)$$

But lemma 5 give $\frac{\partial \varphi^{(1)}}{\partial n}(\gamma) = -\frac{\partial \varphi^{(0)}}{\partial n}(\gamma)$. Like $\gamma \in \Gamma^e$, we have $\langle \nabla \varphi^{(0)}(\gamma), \mathbf{n}(\gamma) \rangle \neq 0$. So we obtain

$$a_0^{(1)}(\gamma) = a_0^{(0)}(\gamma)$$

- If $\gamma \in \Gamma_R \cap \Gamma^e$, formula (3.28) give

$$\begin{aligned} & \alpha(k) \left(\frac{\partial a^{(1)}}{\partial n}(\gamma) - \iota k a^{(1)}(\gamma) \frac{\partial \varphi^{(1)}}{\partial n}(\gamma) \right) + \beta(k) a^{(1)}(\gamma) \\ &= - \left[\alpha(k) \left(\frac{\partial a^{(0)}}{\partial n}(\gamma) - \iota k a^{(0)}(\gamma) \frac{\partial \varphi^{(0)}}{\partial n}(\gamma) \right) + \beta(k) a^{(0)}(\gamma) \right] \end{aligned}$$

Using asymptotic expansions of $a^{(0)}$, $a^{(1)}$, $\alpha(k)$ and $\beta(k)$

$$\begin{aligned} & \left(\sum_{j \in \mathbb{N}} k^{-j} \alpha_j \right) \times \sum_{j \in \mathbb{N}} \left(k^j \frac{\partial a_j^{(1)}}{\partial n}(\gamma) - \iota k^{-j+1} a_j^{(1)}(\gamma) \frac{\partial \varphi^{(1)}}{\partial n}(\gamma) \right) \\ &+ \left(\sum_{j \in \mathbb{N}} k^{-j+1} \beta_{j-1} \right) \times \sum_{j \in \mathbb{N}} k^{-j} a_j^{(1)}(\gamma) \\ &= - \left(\sum_{j \in \mathbb{N}} k^{-j} \alpha_j \right) \times \sum_{j \in \mathbb{N}} \left(k^j \frac{\partial a_j^{(0)}}{\partial n}(\gamma) - \iota k^{-j+1} a_j^{(0)}(\gamma) \frac{\partial \varphi^{(0)}}{\partial n}(\gamma) \right) \\ &- \left(\sum_{j \in \mathbb{N}} k^{-j+1} \beta_{j-1} \right) \times \sum_{j \in \mathbb{N}} k^{-j} a_j^{(0)}(\gamma) \end{aligned}$$

The leading term in power of k is

$$-\iota \alpha_0 a_0^{(1)}(\gamma) \frac{\partial \varphi^{(1)}}{\partial n}(\gamma) + \beta_{-1} a_0^{(1)}(\gamma) = - \left(-\iota \alpha_0 a_0^{(0)}(\gamma) \frac{\partial \varphi^{(0)}}{\partial n}(\gamma) + \beta_{-1} a_0^{(0)}(\gamma) \right)$$

But $\frac{\partial \varphi^{(1)}}{\partial n}(\gamma) = -\frac{\partial \varphi^{(0)}}{\partial n}(\gamma)$, so

$$a_0^{(1)}(\gamma) \left(\beta_{-1} + \iota \alpha_0 \frac{\partial \varphi^{(0)}}{\partial n} \right) = -a_0^{(0)}(\gamma) \left(\beta_{-1} - \iota \alpha_0 \frac{\partial \varphi^{(0)}}{\partial n} \right)$$

with hypothesis $|\beta_{-1}| > |\alpha_0|$ we have

$$\beta_{-1} + \iota \alpha_0 \frac{\partial \varphi^{(0)}}{\partial n} \neq 0$$

So we obtain

$$a_0^{(1)}(\gamma) = - \frac{\beta_{-1} - \iota \alpha_0 \frac{\partial \varphi^{(0)}}{\partial n}}{\beta_{-1} + \iota \alpha_0 \frac{\partial \varphi^{(0)}}{\partial n}} a_0^{(0)}(\gamma)$$

□

We notice that

$$\forall \gamma \in \Gamma^e \quad |b_{\nabla \varphi^{(0)}(\gamma)}(\gamma)| = 1. \quad (3.31)$$

This the case when there is no absorption in the boundary condition.

3.2.3. Calculus of $\text{Hess } \varphi^{(1)}$ at $\gamma \in \Gamma^e$.

Lemma 7. Let $\gamma \in \Gamma^e$. We note $\mathbb{B}(\gamma)$ the curvature matrix of Γ at γ . Then, in dimension $d = 2$ or 3 , we have

$$\text{Hess } \varphi^{(1)}(\gamma) = \mathcal{T}_{\mathbb{B}(\gamma), \mathbf{n}(\gamma), \nabla \varphi^{(0)}(\gamma)}^d(\text{Hess } \varphi^{(0)}(\gamma)). \quad (3.32)$$

Proof. We have seen (lemma 4) that

$$\varphi^{(1)}(\gamma) = \varphi^{(0)}(\gamma).$$

Let $\mathcal{V}_\Gamma(\gamma)$ be a neigborhood of γ in Γ . Taylor's expansion at order 1 of $\nabla \varphi^{(0)}$ and $\nabla \varphi^{(1)}$ are, for all $\gamma' \in \mathcal{V}_\Gamma(\gamma)$,

$$\nabla \varphi^{(0)}(\gamma') = \nabla \varphi^{(0)}(\gamma) + \text{Hess } \varphi^{(0)}(\gamma)(\gamma' - \gamma) + o(|\gamma' - \gamma|) \quad (3.33)$$

and

$$\nabla \varphi^{(1)}(\gamma') = \nabla \varphi^{(1)}(\gamma) + \text{Hess } \varphi^{(1)}(\gamma)(\gamma' - \gamma) + o(|\gamma' - \gamma|) \quad (3.34)$$

Let $\gamma(u, v)$ be the local parametrization of Γ at γ define in section ?? We compute now, Taylor's expansion at order 1 of $\mathbf{n}(\gamma(u, v))$ in dimension 2 and 3. For that, we first evaluate

$$\begin{aligned} \left(\frac{\partial \gamma}{\partial u} \wedge \frac{\partial \gamma}{\partial v} \right)(u, v) &= \begin{pmatrix} -k_1^\Gamma(\gamma)u \\ -k_2^\Gamma(\gamma)v \\ 1 \end{pmatrix}_{\mathcal{B}_\Gamma(\gamma)} + O(u^2 + v^2), \\ &= \mathbf{n}(\gamma) + \mathbb{B}(\gamma)(\gamma(u, v) - \gamma) + O(u^2 + v^2). \end{aligned}$$

We obtain

$$\mathbf{n}(\gamma(u, v)) = \frac{\mathbf{n}(\gamma) + \mathbb{B}(\gamma)(\gamma(u, v) - \gamma)}{(1 + (k_1^\Gamma(\gamma)u)^2 + (k_2^\Gamma(\gamma)v)^2)^{\frac{1}{2}}} + O(u^2 + v^2).$$

But

$$(1 + (k_1^\Gamma(\gamma)u)^2 + (k_2^\Gamma(\gamma)v)^2)^{-\frac{1}{2}} = 1 + O(u^2 + v^2)$$

so, we have

$$\mathbf{n}(\gamma') = \mathbf{n}(\gamma) + \mathbb{B}(\gamma)(\gamma' - \gamma) + o(|\gamma' - \gamma|) \quad (3.35)$$

In similar way, we obtain the same formula in dimension $d = 2$.

Using formulas (3.33) and (3.35) in equation (3.29) gives

$$\begin{aligned} \nabla \varphi^{(1)}(\gamma') &= \nabla \varphi^{(1)}(\gamma) + (\text{Hess } \varphi^{(0)}(\gamma) - 2 \langle \nabla \varphi^{(0)}(\gamma), \mathbf{n}(\gamma) \rangle \mathbb{B}(\gamma))(\gamma' - \gamma) \\ &\quad - 2 \langle \nabla \varphi^{(0)}(\gamma), \mathbb{B}(\gamma)(\gamma' - \gamma) \rangle \mathbf{n}(\gamma) \\ &\quad - 2 \langle \text{Hess } \varphi^{(0)}(\gamma)(\gamma' - \gamma), \mathbf{n}(\gamma) \rangle \mathbf{n}(\gamma) + o(|\gamma' - \gamma|) \end{aligned}$$

So we obtain with formula (3.34), for all $\gamma' \in \mathcal{V}_\Gamma(\gamma)$,

$$\begin{aligned} \text{Hess } \varphi^{(1)}(\gamma)(\gamma' - \gamma) &= (\text{Hess } \varphi^{(0)}(\gamma) - 2 \langle \nabla \varphi^{(0)}(\gamma), \mathbf{n}(\gamma) \rangle \mathbb{B}(\gamma))(\gamma' - \gamma) \\ &\quad - 2 \langle \nabla \varphi^{(0)}(\gamma), \mathbb{B}(\gamma)(\gamma' - \gamma) \rangle \mathbf{n}(\gamma) \\ &\quad - 2 \langle \text{Hess } \varphi^{(0)}(\gamma)(\gamma' - \gamma), \mathbf{n}(\gamma) \rangle \mathbf{n}(\gamma) + o(|\gamma' - \gamma|) \end{aligned}$$

Now, we want to extend the previous formula for all x in $\mathcal{V}(\gamma) \subset \mathbb{R}^d$, a neighborhood of γ . So we must add to previous formula a function from \mathbb{R}^d to \mathbb{R}^d which vanishes on Γ . That is to say at order 2, there exists a constant vector $C(\gamma) \in \mathbb{R}^d$ such that :

for all $x \in \mathcal{V}(\gamma)$

$$\begin{aligned} \text{Hess } \varphi^{(1)}(\gamma)(x - \gamma) &= (\text{Hess } \varphi^{(0)}(\gamma) - 2 \langle \nabla \varphi^{(0)}(\gamma), \mathbf{n}(\gamma) \rangle \mathbb{B}(\gamma))(x - \gamma) \\ &\quad - 2 \langle \nabla \varphi^{(0)}(\gamma), \mathbb{B}(\gamma)(x - \gamma) \rangle \mathbf{n}(\gamma) \\ &\quad - 2 \langle \text{Hess } \varphi^{(0)}(\gamma)(x - \gamma), \mathbf{n}(\gamma) \rangle \mathbf{n}(\gamma) \\ &\quad + \langle x - \gamma, \mathbf{n}(\gamma) \rangle C(\gamma) + o(|x - \gamma|) \end{aligned} \quad (3.36)$$

To compute $C(\gamma)$, we take $x \in \mathcal{V}(\gamma)$ such that $x - \gamma = \varepsilon \nabla \varphi^{(1)}(\gamma)$ with $\varepsilon > 0$ and use (3.6) for $\varphi^{(0)}$ and $\varphi^{(1)}$

$$\text{Hess } \varphi^{(0)}(x) \nabla \varphi^{(0)}(x) = 0$$

and

$$\text{Hess } \varphi^{(1)}(x) \nabla \varphi^{(1)}(x) = 0$$

So, formula (3.36) become

$$\begin{aligned} & (\text{Hess } \varphi^{(0)}(\gamma) - 2 \langle \nabla \varphi^{(0)}(\gamma), \mathbf{n}(\gamma) \rangle \mathbb{B}(\gamma)) \nabla \varphi^{(1)}(\gamma) \\ & - 2 \langle \nabla \varphi^{(0)}(\gamma), \mathbb{B}(\gamma) (\nabla \varphi^{(1)}(\gamma)) \rangle \mathbf{n}(\gamma) \\ & - 2 \langle \text{Hess } \varphi^{(0)}(\gamma) (\nabla \varphi^{(1)}(\gamma)), \mathbf{n}(\gamma) \rangle \mathbf{n}(\gamma) \\ & + \langle \nabla \varphi^{(1)}(\gamma), \mathbf{n}(\gamma) \rangle C(\gamma) = 0 \end{aligned}$$

Using (3.29) and $\mathbb{B}(\gamma) \mathbf{n}(\gamma) = 0$ we obtain

$$\mathbb{B}(\gamma) \nabla \varphi^{(1)}(\gamma) = \mathbb{B}(\gamma) \nabla \varphi^{(0)}(\gamma)$$

and so

$$\begin{aligned} C(\gamma) = & 2 \left[2 \langle \text{Hess } \varphi^{(0)}(\gamma) \mathbf{n}(\gamma), \mathbf{n}(\gamma) \rangle - \frac{\langle \mathbb{B}(\gamma) \nabla \varphi^{(0)}(\gamma), \nabla \varphi^{(0)}(\gamma) \rangle}{\langle \nabla \varphi^{(0)}(\gamma), \mathbf{n}(\gamma) \rangle} \right] \mathbf{n}(\gamma) \\ & - 2 [\text{Hess } \varphi^{(0)}(\gamma) \mathbf{n}(\gamma) + \mathbb{B}(\gamma) \nabla \varphi^{(0)}(\gamma)] \end{aligned}$$

Replacing $C(\gamma)$ by previous formula in (3.36) immediately give (3.32). \square

These results are given in [eC89]

3.2.4. Properties of $\text{Hess } \varphi^{(1)}(\gamma)$.

Lemma 8. *The matrix $\text{Hess } \varphi^{(1)}(\gamma)$ is symmetric and*

$$\text{Hess } \varphi^{(1)}(\gamma) \nabla \varphi^{(1)}(\gamma) = 0. \quad (3.37)$$

In dimension $d = 3$, eigenvalues of $\text{Hess } \varphi^{(1)}(\gamma)$ are $(-k_1^{(1)}(\gamma), -k_2^{(1)}(\gamma), 0)$ and

$$k_2^{(1)}(\gamma) \leq k_1^{(1)}(\gamma) < 0, \quad k_1^{(1)}(\gamma) + k_2^{(1)}(\gamma) < k_1^{(0)}(\gamma) + k_2^{(0)}(\gamma) \quad \text{and} \quad k_1^{(1)}(\gamma) k_2^{(1)}(\gamma) < k_1^{(0)}(\gamma) k_2^{(0)}(\gamma).$$

In dimension $d = 2$, eigenvalues of $\text{Hess } \varphi^{(1)}(\gamma)$ are $(-k^{(1)}(\gamma), 0)$ and

$$k^{(1)}(\gamma) < k^{(0)}(\gamma) \leq 0.$$

Proof. As we have seen in section 3.1.2, in dimension $d = 3$ (resp. $d = 2$), eigenvalues of $\text{Hess } \varphi^{(0)}(\gamma)$ are $(-k_1^{(0)}(\gamma), -k_2^{(0)}(\gamma), 0)$ (resp. $-k^{(0)}(\gamma), 0$) where $k_2^{(0)}(\gamma) \leq k_1^{(0)}(\gamma) \leq 0$ (resp. $k^{(0)}(\gamma) \leq 0$). So we only have to apply lemma 12 in dimension 2 or lemma 13 in dimension 3 to end the proof. \square

3.2.5. Conclusion. Outside a strictly convex compact, we proved following result

Theorem 3. *Let $x \in \Sigma_+ \cap \Omega$, such that $\mathcal{G}(x) = \emptyset$. Then $\#\mathcal{R}(x) = \#\mathcal{R}_1(x) \leq 1$, $\#\mathcal{R}_0(x) \leq 1$, and*

$$u^{inc}(x) = \sum_{\rho \in \mathcal{R}_0(x)} u_\rho^{G.O.}(x) + O\left(\frac{1}{k}\right)$$

and

$$u(x) = \sum_{\rho \in \mathcal{R}_1(x)} u_\rho^{G.O.}(x) + O\left(\frac{1}{k}\right)$$

3.3. Outside an union of strictly convex compacts. To prove theorem 1, we just have to verify that we can use propagation and reflection lemmas along each ray coming through x . For that, we have

Lemma 9. *Let $x \in \Sigma_+ \cap \Omega$, such that $\mathcal{G}(x) = \emptyset$. Let $l \in \mathbb{N}^*$ such that $\mathcal{R}_l(x) \neq \emptyset$ and $\rho = (\gamma_0, \gamma_1, \dots, \gamma_l) \in \mathcal{R}_l(x)$. In dimension $d = 3$ (resp. $d = 2$), if eigenvalues of the symmetric matrix $\mathbb{A}_\rho^{(0)}(\gamma_0)$ are $(\lambda_1^{(0)}(\gamma_0), \lambda_2^{(0)}(\gamma_0), 0)$ (resp. $(\lambda^{(0)}(\gamma_0), 0)$) where*

$$0 \leq \lambda_1^{(0)}(\gamma_0) \leq \lambda_2^{(0)}(\gamma_0) \quad (\text{resp. } 0 \leq \lambda^{(0)}(\gamma_0))$$

then $\forall j \in \{1, \dots, l\}$ eigenvalues of $\mathbb{A}_\rho^{(j)}(\gamma_j)$ are $(\lambda_1^{(j)}(\gamma_j), \lambda_2^{(j)}(\gamma_j), 0)$ (resp. $(\lambda^{(j)}(\gamma_j), 0)$) where

$$0 < \lambda_1^{(j)}(\gamma_j) \leq \lambda_2^{(j)}(\gamma_j) \quad (\text{resp. } 0 < \lambda^{(j)}(\gamma_j))$$

Proof. Due to hypothesis we can apply propagation lemma 2 to obtain that

$$\mathbb{A}_\rho^{(0)}(\gamma_1) = \mathcal{S}_{\|\gamma_1 - \gamma_0\|}(\mathbb{A}_\rho^{(0)}(\gamma_0))$$

is well defined. and its eigenvalues are $(\lambda_1^{(0)}(\gamma_1), \lambda_2^{(0)}(\gamma_1), 0)$ where

$$0 \leq \lambda_1^{(0)}(\gamma_1) \leq \lambda_2^{(0)}(\gamma_1).$$

By hypothesis $\rho \in \mathcal{R}_l(x)$ so $\langle \gamma_1 - \gamma_0, \mathbf{n}(\gamma_1) \rangle < 0$ and we can apply reflection lemma 7 to obtain that

$$\mathbb{A}_\rho^{(1)}(\gamma_1) = \mathcal{T}_{\mathbb{B}(\gamma_1), \mathbf{n}(\gamma_1), \frac{\gamma_1 - \gamma_0}{\|\gamma_1 - \gamma_0\|}}(\mathbb{A}_\rho^{(0)}(\gamma_1))$$

is well defined.

Furthermore, lemma 8 give us that matrix $\mathbb{A}_\rho^{(1)}(\gamma_1)$ is symmetric and its eigenvalues are $(\lambda_1^{(1)}(\gamma_1), \lambda_2^{(1)}(\gamma_1), 0)$ where

$$0 < \lambda_1^{(1)}(\gamma_1) \leq \lambda_2^{(1)}(\gamma_1).$$

Then a simply recurrence proof give us lemma :

Let $j \in \{1, \dots, l-1\}$ and suppose that $\mathbb{A}_\rho^{(j)}(\gamma_j)$ is symmetric and its eigenvalues are $(\lambda_1^{(j)}(\gamma_j), \lambda_2^{(j)}(\gamma_j), 0)$ where

$$0 < \lambda_1^{(j)}(\gamma_j) \leq \lambda_2^{(j)}(\gamma_j)$$

we can apply propagation lemma 2 to obtain that

$$\mathbb{A}_\rho^{(j)}(\gamma_{j+1}) = \mathcal{S}_{\|\gamma_{j+1} - \gamma_j\|}(\mathbb{A}_\rho^{(j)}(\gamma_j))$$

is well defined. and its eigenvalues are $(\lambda_1^{(j)}(\gamma_{j+1}), \lambda_2^{(j)}(\gamma_{j+1}), 0)$ where

$$0 < \lambda_1^{(j)}(\gamma_{j+1}) \leq \lambda_2^{(j)}(\gamma_{j+1}).$$

By hypothesis $\rho \in \mathcal{R}_l(x)$ so $\langle \gamma_{j+1} - \gamma_j, \mathbf{n}(\gamma_{j+1}) \rangle < 0$ and we can apply reflection lemma 7 to obtain that

$$\mathbb{A}_\rho^{(j+1)}(\gamma_{j+1}) = \mathcal{T}_{\mathbb{B}(\gamma_{j+1}), \mathbf{n}(\gamma_{j+1}), \frac{\gamma_{j+1} - \gamma_j}{\|\gamma_{j+1} - \gamma_j\|}}(\mathbb{A}_\rho^{(j)}(\gamma_{j+1}))$$

is well defined.

Furthermore, lemma 8 give us that matrix $\mathbb{A}_\rho^{(j+1)}(\gamma_{j+1})$ is symmetric and its eigenvalues are $(\lambda_1^{(j+1)}(\gamma_{j+1}), \lambda_2^{(j+1)}(\gamma_{j+1}), 0)$ where

$$0 < \lambda_1^{(j+1)}(\gamma_{j+1}) \leq \lambda_2^{(j+1)}(\gamma_{j+1}).$$

A similar proof will act in dimension $d = 2$. □

With theorem 1 hypothesis and this lemma, we immediatly have

$$\forall j \in \{1, \dots, l\}, \det \left(\mathbb{I} + \|\gamma_j - \gamma_{j-1}\| \mathbb{A}_\rho^{(j-1)}(\gamma_{j-1}) \right) > 0$$

and

$$\det \left(\mathbb{I} + \|x - \gamma_l\| \mathbb{A}_\rho^{(l)}(\gamma_l) \right) > 0.$$

So along each ray comming throug x we can apply propagation and reflection lemmas. Then, by adding the contribution of each ray we obtain theorem 1 results.

Lemma 10. Let $x \in \Sigma_+ \cap \Omega$, such that $\mathcal{G}(x) = \emptyset$. Let $l > 1$ such that $\mathcal{R}_l(x) \neq \emptyset$ and $\rho = (\gamma_0, \gamma_1, \dots, \gamma_l) \in \mathcal{R}_l(x)$. We have

$$|a_\rho(x)| < \frac{|\mathbb{A}_\rho^{(0)}(\gamma_0)|}{\left[(1 + 2d(x)\lambda_{\min}^\Gamma) (1 + 2d_{\min}\lambda_{\min}^\Gamma)^{l-1} \right]^{\frac{d-1}{2}}} \quad (3.38)$$

where

$$0 < d(x) = \min_{\gamma \in \Gamma} \|x - \gamma\|,$$

$$0 < \lambda_{\min}^\Gamma = \begin{cases} \min_{\gamma \in \Gamma} \lambda_1(\gamma) & \text{if } d = 3, \\ \min_{\gamma \in \Gamma} \lambda(\gamma) & \text{if } d = 2, \end{cases},$$

and

$$0 < d_{\min} = \min_{i \neq j} \min_{\gamma_i \in \Gamma_i, \gamma_j \in \Gamma_j} \|\gamma_i - \gamma_j\|.$$

Proof. Using (3.31) and definition of $a_\rho^{(j)}(\gamma_j)$, we obtain

$$\forall j \in \{1, \dots, l\}, |a_\rho^{(j+1)}(\gamma_{j+1})| = \frac{|a_\rho^j(\gamma_j)|}{\sqrt{\det(\mathbb{I} + \|\gamma_{j+1} - \gamma_j\| \mathbb{A}_\rho^{(j)}(\gamma_j))}} \quad (3.39)$$

where $t_j = \|\gamma_{j+1} - \gamma_j\|$ and $\gamma_{j+1} = x$.

In dimension $d = 2$, we can use equation (4.1) of lemma 12 to obtain

$$\lambda^{(j)}(\gamma_j) = \lambda^{(j-1)}(\gamma_j) - 2t_{j-1} \frac{\lambda^\Gamma(\gamma_j)}{\langle \mathbf{n}(\gamma_j), \gamma_j - \gamma_{j-1} \rangle}$$

But we have $\frac{1}{t_{j-1}} \langle \mathbf{n}(\gamma_j), \gamma_j - \gamma_{j-1} \rangle \in [-1; 0[$ and then

$$\lambda^{(j)}(\gamma_j) \geq \lambda^{(j-1)}(\gamma_j) + 2\lambda_{\min}^\Gamma > 0.$$

From proposition (1) we get

$$\lambda^{(j-1)}(\gamma_j) = \frac{\lambda^{(j-1)}(\gamma_{j-1})}{1 + t_{j-1}\lambda^{(j-1)}(\gamma_{j-1})}$$

and so

$$\begin{aligned} \lambda^{(j-1)}(\gamma_j) &> 0 & \forall j \in \{2, \dots, l\} \\ \lambda^{(0)}(\gamma_1) &\geq 0 \end{aligned}.$$

From equation (3.39), we immediately obtain

$$\begin{aligned} |a_\rho^{(1)}(\gamma_1)| &\leq |a_\rho^{(0)}(\gamma_0)| \\ |a_\rho^{(j+1)}(\gamma_{j+1})| &< |a_\rho^{(j)}(\gamma_j)| (1 + 2d_{\min} \lambda_{\min}^\Gamma)^{-1/2} \quad \forall j \in \{1, \dots, l-1\} \\ |a_\rho^{(l)}(x)| = |a_\rho(x)| &< |a_\rho^{(l)}(\gamma_l)| (1 + 2d(x) \lambda_{\min}^\Gamma)^{-1/2} \end{aligned}$$

and thus

$$|a_\rho(x)| < \frac{|a_\rho^{(0)}(\gamma_0)|}{\left[(1 + 2d(x) \lambda_{\min}^\Gamma) (1 + 2d_{\min} \lambda_{\min}^\Gamma)^{l-1}\right]^{\frac{1}{2}}}.$$

In dimension $d = 3$, We have

$$\det(\mathbb{I} + t_j \mathbb{A}_\rho^{(j)}(\gamma_j)) = 1 + (\lambda_1^{(j)}(\gamma_j) + \lambda_2^{(j)}(\gamma_j))t_j + \lambda_1^{(j)}(\gamma_j)\lambda_2^{(j)}(\gamma_j)t_j^2$$

and we can use equations (4.5) and (4.6) of lemma 13 to obtain

$$\begin{aligned} \lambda_1^{(j)}(\gamma_j) + \lambda_2^{(j)}(\gamma_j) &\geq \lambda_1^{(j-1)}(\gamma_j) + \lambda_2^{(j-1)}(\gamma_j) + 4\lambda_{\min}^\Gamma > 0, \\ \lambda_1^{(j)}(\gamma_j)\lambda_2^{(j)}(\gamma_j) &\geq \lambda_1^{(j-1)}(\gamma_j)\lambda_2^{(j-1)}(\gamma_j) + 4(\lambda_{\min}^\Gamma)^2 > 0. \end{aligned}$$

So, with these inequalities, we have

$$\begin{aligned} \det(\mathbb{I} + t_j \mathbb{A}_\rho^{(j)}(\gamma_j)) &\geq 1 + 4\lambda_{\min}^\Gamma t_j + 4(\lambda_{\min}^\Gamma t_j)^2 + (\lambda_1^{(j-1)}(\gamma_j) + \lambda_2^{(j-1)}(\gamma_j))t_j \\ &\quad + (\lambda_1^{(j-1)}(\gamma_j)\lambda_2^{(j-1)}(\gamma_j))t_j^2 \end{aligned}$$

From proposition (1) we get

$$\lambda_\alpha^{(j-1)}(\gamma_j) = \frac{\lambda_\alpha^{(j-1)}(\gamma_{j-1})}{1 + t_{j-1}\lambda_\alpha^{(j-1)}(\gamma_{j-1})}, \quad \alpha \in \{1, 2\}$$

and, as $\lambda_2^{(j-1)}(\gamma_{j-1}) \geq \lambda_1^{(j-1)}(\gamma_{j-1}) > 0$ for $j \in \{2, \dots, l+1\}$ and $\lambda_2^{(0)}(\gamma_0) \geq \lambda_1^{(0)}(\gamma_0) \geq 0$ (see lemma 13) we have

$$\begin{aligned} \lambda_2^{(j-1)}(\gamma_j) &\geq \lambda_1^{(j-1)}(\gamma_j) &> 0 & \forall j \in \{2, \dots, l\} \\ \lambda_2^{(0)}(\gamma_1) &\geq \lambda_1^{(0)}(\gamma_1) &\geq 0 \end{aligned}.$$

With these inequalities, we have

$$\det(\mathbb{I} + t_j \mathbb{A}_\rho^{(j)}(\gamma_j)) > (1 + 2\lambda_{\min}^\Gamma t_j)^2 \quad \text{if } j > 1$$

$$\det(\mathbb{I} + t_1 \mathbb{A}_\rho^{(1)}(\gamma_1)) \geq (1 + 2\lambda_{\min}^\Gamma t_1)^2$$

From equation (3.39), we immediately obtain

$$\begin{aligned} |a_\rho^{(1)}(\gamma_1)| &\leq |a_\rho^{(0)}(\gamma_0)| \\ |a_\rho^{(j+1)}(\gamma_{j+1})| &< |a_\rho^{(j)}(\gamma_j)| (1 + 2d_{\min} \lambda_{\min}^\Gamma)^{-1} \quad \forall j \in \{1, \dots, l-1\} \\ |a_\rho^{(l)}(x)| = |a_\rho(x)| &< |a_\rho^{(l)}(\gamma_l)| (1 + 2d(x) \lambda_{\min}^\Gamma)^{-1} \end{aligned}$$

and thus

$$|a_\rho(x)| < \frac{|a_\rho^{(0)}(\gamma_0)|}{(1 + 2d(x) \lambda_{\min}^\Gamma)^{(l-1)}}.$$

□

4. TECHNICAL RESULTS

Proposition 1. Let $t > 0$ and $\mathbb{A} \in S_t^d$. Assume that λ is an eigenvalue of \mathbb{A} with corresponding eigenvector ζ then $\frac{\lambda}{1+t\lambda}$ is an eigenvalue of $\mathcal{S}_t(\mathbb{A})$ with corresponding eigenvector ζ .

Proof. By definition $\mathcal{S}_t(\mathbb{A}) = \mathbb{A}(\mathbb{I} + t\mathbb{A})^{-1}$ and by hypothesis

$$(\mathbb{I} + t\mathbb{A})\zeta = (1 + t\lambda)\zeta$$

with $1 + t\lambda \neq 0$. So $(1 + t\lambda)(\mathbb{I} + t\mathbb{A})^{-1}\zeta = \zeta$ and we obtain

$$(1 + t\lambda)\mathbb{A}(\mathbb{I} + t\mathbb{A})^{-1}\zeta = \mathbb{A}\zeta.$$

Thus

$$\mathbb{A}(\mathbb{I} + t\mathbb{A})^{-1}\zeta = \frac{\lambda}{1 + t\lambda}\zeta.$$

□

Lemma 11. Let $\eta \in \mathbb{R}^d$ and $\zeta \in \mathbb{R}^d$ such that $\|\eta\| = 1$ and $\langle \eta, \zeta \rangle \neq 0$. Let \mathbb{A} and \mathbb{B} symmetric matrices in $\mathbb{R}^{d \times d}$. Assume $\mathbb{A}\zeta = 0$ and $\mathbb{B}\eta = 0$ then

$$(\mathcal{T}_{\mathbb{B}, \eta, \zeta}^d(\mathbb{A}))(\zeta - 2\langle \zeta, \eta \rangle \eta) = 0$$

Proof. Note $\xi = \zeta - 2\langle \zeta, \eta \rangle \eta$ then, by definition,

$$\begin{aligned} (\mathcal{T}_{\mathbb{B}, \eta, \zeta}^d(\mathbb{A}))\xi &= \mathbb{A}\xi - 2\langle \zeta, \eta \rangle \mathbb{B}\xi - 2\langle \eta, \xi \rangle (\mathbb{A}\eta + \mathbb{B}\zeta) \\ &\quad - 2\langle \mathbb{A}\eta + \mathbb{B}\zeta, \xi \rangle \eta + 2\left[2\langle \mathbb{A}\eta, \eta \rangle - \frac{\langle \mathbb{B}\zeta, \zeta \rangle}{\langle \zeta, \eta \rangle}\right]\langle \eta, \xi \rangle \eta \end{aligned}$$

Due to hypothesis, we have

$$\mathbb{A}\xi = -2\langle \zeta, \eta \rangle \mathbb{A}\eta, \quad \mathbb{B}\xi = \mathbb{B}\zeta, \quad \langle \eta, \xi \rangle = -\langle \zeta, \eta \rangle$$

and

$$\langle \mathbb{A}\eta + \mathbb{B}\zeta, \xi \rangle = -2\langle \zeta, \eta \rangle \langle \mathbb{A}\eta, \eta \rangle + \langle \mathbb{B}\zeta, \zeta \rangle.$$

So, we obtain

$$\begin{aligned} (\mathcal{T}_{\mathbb{B}, \eta, \zeta}^d(\mathbb{A}))\xi &= -2\langle \zeta, \eta \rangle (\mathbb{A}\eta + \mathbb{B}\zeta) + 2\langle \zeta, \eta \rangle (\mathbb{A}\eta + \mathbb{B}\zeta) + 4\langle \zeta, \eta \rangle \langle \mathbb{A}\eta, \eta \rangle \eta \\ &\quad - 2\langle \mathbb{B}\zeta, \zeta \rangle \eta - 2\left[2\langle \mathbb{A}\eta, \eta \rangle - \frac{\langle \mathbb{B}\zeta, \zeta \rangle}{\langle \zeta, \eta \rangle}\right]\langle \eta, \zeta \rangle \eta \\ &= 0. \end{aligned}$$

□

Lemma 12. Let $\mathcal{B} = (\mathbf{u}, \mathbf{w})$ and $\mathcal{B}^{(I)} = (\mathbf{u}^{(I)}, \mathbf{w}^{(I)})$ two direct orthonormal basis of \mathbb{R}^2 such that $\langle \mathbf{w}, \mathbf{w}^{(I)} \rangle < 0$. Let $\mathbb{B} = \text{diag}(\lambda, 0)$ in \mathcal{B} basis and $\mathbb{A}^{(I)} = \text{diag}(\lambda^{(I)}, 0)$ in $\mathcal{B}^{(I)}$ basis. Then $\mathbb{A}^{(R)} = \mathcal{T}_{\mathbb{B}, \mathbf{w}, \mathbf{w}^{(I)}}^2(\mathbb{A}^{(I)})$ is a symmetric matrix having for eigenvalues $(\lambda^{(R)}, 0)$ where

$$\lambda^{(R)} = \lambda^{(I)} - 2\frac{\lambda}{\langle \mathbf{w}, \mathbf{w}^{(I)} \rangle} \tag{4.1}$$

and

$$\mathbb{A}^{(R)} \left(\mathbf{w}^{(I)} - 2 \langle \mathbf{w}, \mathbf{w}^{(I)} \rangle \mathbf{w} \right) = 0 \tag{4.2}$$

If $\lambda > 0$ and $\lambda^{(I)} \geq 0$ then

$$\lambda^{(R)} > \lambda^{(I)}. \tag{4.3}$$

Proof. In basis \mathcal{B} , we note $\mathbf{u}^{(I)} = (u_j)_{j=1}^2$, $\mathbf{w}^{(I)} = (w_j)_{j=1}^2$ and $\mathbb{A}^{(I)} = (a_{ij})_{i,j=1}^2$. By hypothesis, $\mathbb{A}^{(I)}$ is symmetric and $\mathbb{A}^{(I)}\mathbf{w}^{(I)} = 0$.

As $\langle \mathbf{w}, \mathbf{w}^{(I)} \rangle \neq 0$, we can compute $\mathbb{A}^{(R)} : \forall x \in \mathbb{R}^2$

$$\begin{aligned}\mathbb{A}^{(R)}x &= (\mathbb{A}^{(I)} - 2\langle \mathbf{w}, \mathbf{w}^{(I)} \rangle \mathbb{B})x \\ &\quad - 2\langle \mathbf{w}, x \rangle (\mathbb{A}^{(I)}\mathbf{w} + \mathbb{B}\mathbf{w}^{(I)}) \\ &\quad - 2\langle \mathbb{A}^{(I)}\mathbf{w} + \mathbb{B}\mathbf{w}^{(I)}, x \rangle \mathbf{w} \\ &\quad + 2 \left[2\langle \mathbb{A}^{(I)}\mathbf{w}, \mathbf{w} \rangle - \frac{\langle \mathbb{B}\mathbf{w}^{(I)}, \mathbf{w}^{(I)} \rangle}{\langle \mathbf{w}, \mathbf{w}^{(I)} \rangle} \right] \langle \mathbf{w}, x \rangle \mathbf{w}\end{aligned}$$

in basis \mathcal{B} , $x = (x_j)_{j=1}^2$ and

$$\begin{aligned}\mathbb{A}^{(R)}x &= \begin{pmatrix} a_{11} & a_{12} \\ a_{12} & a_{22} \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} - 2w_2 \begin{pmatrix} \lambda x_1 \\ 0 \end{pmatrix} - 2x_2 \begin{pmatrix} a_{12} + \lambda w_1 \\ a_{22} \end{pmatrix} \\ &\quad - 2 \left\langle \begin{pmatrix} a_{12} + \lambda w_1 \\ a_{22} \end{pmatrix}, \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} \right\rangle \begin{pmatrix} 0 \\ 1 \end{pmatrix} + 2 \left[2a_{22} - \frac{\lambda w_1^2}{w_2} \right] x_2 \begin{pmatrix} 0 \\ 1 \end{pmatrix}\end{aligned}$$

That is to say,

$$\mathbb{A}^{(R)} = \begin{pmatrix} a_{11} - 2\lambda w_2 & -a_{12} - 2\lambda w_1 \\ -a_{12} - 2\lambda w_1 & a_{22} - 2\lambda \frac{w_1^2}{w_2} \end{pmatrix}$$

So we have $\mathbb{A}^{(R)}$ symmetry. We can remark that formula (4.2) is a direct application of Lemma 11.

Now, we study eigenvalues of $\mathbb{A}^{(R)}$. Let \mathbb{H} be the normal matrix given by

$$\mathbb{H} = \begin{pmatrix} u_1 & w_1 \\ -u_2 & -w_2 \end{pmatrix}$$

Then $\mathbb{H}\mathbb{A}^{(R)}\mathbb{H}^t$ is a similar to $\mathbb{A}^{(R)}$.

Using $\mathbb{A}^{(I)}\mathbf{u}^{(I)} = \lambda^{(I)}\mathbf{u}^{(I)}$ and $\mathbb{A}^{(I)}\mathbf{w}^{(I)} = 0$ give

$$\begin{aligned}\mathbb{A}^{(R)}\mathbb{H} &= \begin{pmatrix} (a_{11} - 2\lambda w_2)u_1 + (a_{12} + 2\lambda w_1)u_2 & (a_{11} - 2\lambda w_2)w_1 + (a_{12} + 2\lambda w_1)w_2 \\ (a_{12} + 2\lambda w_1)u_1 - (a_{22} - 2\lambda \frac{w_1^2}{w_2})u_2 & (a_{12} + 2\lambda w_1)w_1 - (a_{22} - 2\lambda \frac{w_1^2}{w_2})w_2 \end{pmatrix} \\ &= \begin{pmatrix} \lambda^{(I)}u_1 + 2\lambda(w_1u_2 - w_2u_1) & 0 \\ -\lambda^{(I)}u_2 + 2\lambda(\frac{w_1^2}{w_2}u_2 - w_1u_1) & 0 \end{pmatrix}\end{aligned}$$

But $(\mathbf{u}^{(I)}, \mathbf{w}^{(I)})$ is a direct and orthonormal basis, so

$$w_2u_1 - w_1u_2 = 1, \quad w_1u_1 + w_2u_2 = 0 \text{ and } u_1^2 + u_2^2 = 1.$$

Then

$$\begin{aligned}\mathbb{H}^t\mathbb{A}^{(R)}\mathbb{H} &= \begin{pmatrix} \lambda^{(I)}(u_1^2 + u_2^2) + 2\lambda \left[(w_1u_2 - w_2u_1)u_1 - \frac{w_1^2}{w_2}u_2^2 + w_1u_1u_2 \right] & 0 \\ \lambda^{(I)}(u_1w_1 + u_2w_2) + 2\lambda(w_1^2u_2 - w_2u_1w_1 - w_1^2u_2 + w_2u_1w_1) & 0 \end{pmatrix} \\ &= \begin{pmatrix} \lambda^{(I)} - 2\frac{\lambda}{w_2}[w_2u_1 - w_1u_2]^2 & 0 \\ 0 & 0 \end{pmatrix} \\ &= \begin{pmatrix} \lambda^{(I)} - 2\frac{\lambda}{w_2} & 0 \\ 0 & 0 \end{pmatrix}\end{aligned}$$

□

Lemma 13. Let $\mathcal{B} = (\mathbf{u}, \mathbf{v}, \mathbf{w})$ and $\mathcal{B}^{(I)} = (\mathbf{u}^{(I)}, \mathbf{v}^{(I)}, \mathbf{w}^{(I)})$ two direct orthonormal basis of \mathbb{R}^3 such that $\langle \mathbf{w}, \mathbf{w}^{(I)} \rangle < 0$. Let $\mathbb{B} = \text{diag}(\lambda_1, \lambda_2, 0)$ in \mathcal{B} basis and the symmetric matrix $\mathbb{A}^{(I)} = \text{diag}(\lambda_1^{(I)}, \lambda_2^{(I)}, 0)$ in $\mathcal{B}^{(I)}$ basis. Then $\mathbb{A}^{(R)} = \mathcal{T}_{\mathbb{B}, \mathbf{w}, \mathbf{w}^{(I)}}^3(\mathbb{A}^{(I)})$ is a symmetric matrix having for eigenvalues $(\lambda_1^{(R)}, \lambda_2^{(R)}, 0)$ and

$$\mathbb{A}^{(R)} \left(\mathbf{w}^{(I)} - 2\langle \mathbf{w}, \mathbf{w}^{(I)} \rangle \mathbf{w} \right) = 0. \quad (4.4)$$

Under the hypothesis

$$0 < \lambda_1 \leq \lambda_2 \text{ and } 0 \leq \lambda_1^{(I)} \leq \lambda_2^{(I)} \quad (\text{H})$$

we have

$$\lambda_1^{(R)} + \lambda_2^{(R)} \geq \lambda_1^{(I)} + \lambda_2^{(I)} + 4\lambda_1 > 0, \quad (4.5)$$

$$\lambda_1^{(R)}\lambda_2^{(R)} \geq \lambda_1^{(I)}\lambda_2^{(I)} + 4(\lambda_1)^2 > 0 \quad (4.6)$$

and thus

$$\lambda_1^{(R)} > 0, \lambda_2^{(R)} > 0 \quad (4.7)$$

Proof. In basis \mathcal{B} , $\mathbf{u}^{(I)} = (u_j)_{j=1}^3$, $\mathbf{v}^{(I)} = (v_j)_{j=1}^3$, $\mathbf{w}^{(I)} = (w_j)_{j=1}^3$ and $\mathbb{A}^{(I)} = (a_{ij})_{i,j=1}^3$. By hypothesis, $\mathbb{A}^{(I)}$ is symmetric and $\mathbb{A}^{(I)}\mathbf{w}^{(I)} = 0$.

As $\langle \mathbf{w}, \mathbf{w}^{(I)} \rangle \neq 0$, we can compute $\mathbb{A}^{(R)} : \forall x \in \mathbb{R}^3$

$$\begin{aligned} \mathbb{A}^{(R)}x &= (\mathbb{A}^{(I)} - 2\langle \mathbf{w}, \mathbf{w}^{(I)} \rangle \mathbb{B})x \\ &\quad - 2\langle \mathbf{w}, x \rangle (\mathbb{A}^{(I)}\mathbf{w} + \mathbb{B}\mathbf{w}^{(I)}) \\ &\quad - 2\langle \mathbb{A}^{(I)}\mathbf{w} + \mathbb{B}\mathbf{w}^{(I)}, x \rangle \mathbf{w} \\ &\quad + 2 \left[2\langle \mathbb{A}^{(I)}\mathbf{w}, \mathbf{w} \rangle - \frac{\langle \mathbb{B}\mathbf{w}^{(I)}, \mathbf{w}^{(I)} \rangle}{\langle \mathbf{w}, \mathbf{w}^{(I)} \rangle} \right] \langle \mathbf{w}, x \rangle \mathbf{w} \end{aligned}$$

in basis \mathcal{B} , $x = (x_j)_{j=1}^3$ and

$$\begin{aligned} \mathbb{A}^{(R)}x &= \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{12} & a_{22} & a_{23} \\ a_{13} & a_{23} & a_{33} \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} - 2w_3 \begin{pmatrix} \lambda_1 x_1 \\ \lambda_2 x_2 \\ 0 \end{pmatrix} \\ &\quad - 2x_3 \begin{pmatrix} a_{13} + \lambda_1 w_1 \\ a_{23} + \lambda_2 w_2 \\ a_{33} \end{pmatrix} \\ &\quad - 2 \left\langle \begin{pmatrix} a_{13} + \lambda_1 w_1 \\ a_{23} + \lambda_2 w_2 \\ a_{33} \end{pmatrix}, \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} \right\rangle \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \\ &\quad + 2 \left[2a_{33} - \frac{\lambda_1 w_1^2 + \lambda_2 w_2^2}{w_3} \right] x_3 \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \end{aligned}$$

That is to say,

$$\mathbb{A}^{(R)} = \begin{pmatrix} a_{11} - 2\lambda_1 w_3 & a_{12} & -a_{13} - 2\lambda_1 w_1 \\ a_{12} & a_{22} - 2\lambda_2 w_3 & -a_{23} - 2\lambda_2 w_2 \\ -a_{13} - 2\lambda_1 w_1 & -a_{23} - 2\lambda_2 w_2 & a_{33} - \frac{2}{w_3}(\lambda_1 w_1^2 + \lambda_2 w_2^2) \end{pmatrix}$$

So we have $\mathbb{A}^{(R)}$ symmetry.

To prove formula (4.4), we make computation in basis \mathcal{B} where

$$\mathbf{w}^{(I)} - 2\langle \mathbf{w}, \mathbf{w}^{(I)} \rangle \mathbf{w} = \begin{pmatrix} w_1 \\ w_2 \\ -w_3 \end{pmatrix}$$

So

$$\mathbb{A}^{(R)}\mathbf{w}^{(R)} = \begin{pmatrix} a_{11}w_1 - 2\lambda_1 w_3 w_1 + a_{12}w_2 + a_{13}w_3 + 2\lambda_1 w_1 w_3 \\ a_{12}w_1 + a_{22}w_2 - 2\lambda_2 w_3 w_2 + a_{23}w_3 + 2\lambda_2 w_2 w_3 \\ -a_{13}w_1 - 2\lambda_1 w_1^2 - a_{23}w_2 - 2\lambda_2 w_2^2 - a_{33}w_3 + 2(\lambda_1 w_1^2 + \lambda_2 w_2^2) \end{pmatrix}$$

Then, using $\mathbb{A}^{(I)}\mathbf{w}^{(I)} = 0$ give immediately formula (4.4).

To establish the formulas (4.5)-(4.6)-(4.7) under the assumptions (H), we will study the eigenvalues of matrix $\mathbb{A}^{(R)}$. For that, let \mathbb{H} be the normal matrix given by

$$\mathbb{H} = \begin{pmatrix} u_1 & v_1 & w_1 \\ u_2 & v_2 & w_2 \\ -u_3 & -v_3 & -w_3 \end{pmatrix}$$

Then $\mathbb{H}\mathbb{A}^{(R)}\mathbb{H}^t$ is similar to $\mathbb{A}^{(R)}$. Using that $(\mathbf{u}^{(I)}, \mathbf{v}^{(I)}, \mathbf{w}^{(I)})$ is a direct and orthonormal basis, we have

$$\mathbf{u}^{(I)} \wedge \mathbf{v}^{(I)} = \mathbf{w}^{(I)} \Leftrightarrow \begin{pmatrix} u_2v_3 - u_3v_2 \\ -u_1v_3 + u_3v_1 \\ u_1v_2 - u_2v_1 \end{pmatrix} = \begin{pmatrix} w_1 \\ w_2 \\ w_3 \end{pmatrix} \quad (4.8)$$

$$\mathbf{v}^{(I)} \wedge \mathbf{w}^{(I)} = \mathbf{u}^{(I)} \Leftrightarrow \begin{pmatrix} v_2w_3 - v_3w_2 \\ -v_1w_3 + v_3w_1 \\ v_1w_2 - v_2w_1 \end{pmatrix} = \begin{pmatrix} u_1 \\ u_2 \\ u_3 \end{pmatrix} \quad (4.9)$$

$$\mathbf{w}^{(I)} \wedge \mathbf{u}^{(I)} = \mathbf{v}^{(I)} \Leftrightarrow \begin{pmatrix} w_2 u_3 - w_3 u_2 \\ -w_1 u_3 + w_3 u_1 \\ w_1 u_2 - w_2 u_1 \end{pmatrix} = \begin{pmatrix} v_1 \\ v_2 \\ v_3 \end{pmatrix} \quad (4.10)$$

$$\|\mathbf{u}^{(I)}\| = 1 \Leftrightarrow u_1^2 + u_2^2 + u_3^2 = 1 \quad (4.11)$$

$$\|\mathbf{v}^{(I)}\| = 1 \Leftrightarrow v_1^2 + v_2^2 + v_3^2 = 1 \quad (4.12)$$

$$\|\mathbf{w}^{(I)}\| = 1 \Leftrightarrow w_1^2 + w_2^2 + w_3^2 = 1 \quad (4.13)$$

Using $\mathbb{A}^{(I)}\mathbf{u}^{(I)} = \lambda_1^{(I)}\mathbf{u}^{(I)}$, $\mathbb{A}^{(I)}\mathbf{v}^{(I)} = \lambda_1^{(I)}\mathbf{v}^{(I)}$ and $\mathbb{A}^{(I)}\mathbf{w}^{(I)} = 0$ give

$$\begin{aligned} & \mathbb{A}^{(R)}\mathbb{H} \\ &= \\ & \left(\begin{array}{ccc} \lambda_1^{(I)}u_1 + 2\lambda_1(w_1u_3 - w_3u_1) & \lambda_2^{(I)}v_1 + 2\lambda_1(w_1v_3 - w_3v_1) & 0 \\ \lambda_1^{(I)}u_2 + 2\lambda_2(w_2u_3 - w_3u_2) & \lambda_2^{(I)}v_2 + 2\lambda_2(w_2v_3 - w_3v_2) & 0 \\ \left\{ \begin{array}{l} -\lambda_1^{(I)}u_3 - 2\lambda_1(w_1u_1 - \frac{u_3w_1^2}{w_3}) \\ -2\lambda_2(w_2u_2 - \frac{u_3w_2^2}{w_3}) \end{array} \right\} & \left\{ \begin{array}{l} -\lambda_2^{(I)}v_3 - 2\lambda_1(w_1v_1 - \frac{v_3w_1^2}{w_3}) \\ -2\lambda_2(w_2v_2 - \frac{v_3w_2^2}{w_3}) \end{array} \right\} & 0 \end{array} \right) \end{aligned}$$

To obtain $\mathbb{H}^t\mathbb{A}^{(R)}\mathbb{H}$, we compute $\langle \mathbb{H}^t\mathbb{A}^{(R)}\mathbb{H}\mathbf{x}, \mathbf{y} \rangle \quad \forall x, y \in \{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$ We can remark that

$$\langle \mathbb{H}^t\mathbb{A}^{(R)}\mathbb{H}\mathbf{x}, \mathbf{w} \rangle = 0 \quad \forall x \in \{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$$

Now, we compute the six leading terms of matrix $\mathbb{H}^t\mathbb{A}^{(R)}\mathbb{H}$

- Calculus of $\langle \mathbb{H}^t\mathbb{A}^{(R)}\mathbb{H}\mathbf{u}, \mathbf{u} \rangle$

$$\begin{aligned} \langle \mathbb{H}^t\mathbb{A}^{(R)}\mathbb{H}\mathbf{u}, \mathbf{u} \rangle &= \left(\lambda_1^{(I)}u_1 + 2\lambda_1(w_1u_3 - w_3u_1) \right) u_1 \\ &\quad + \left(\lambda_1^{(I)}u_2 + 2\lambda_2(w_2u_3 - w_3u_2) \right) u_2 \\ &\quad - \left(-\lambda_1^{(I)}u_3 - 2\lambda_1(w_1u_1 - \frac{u_3w_1^2}{w_3}) - 2\lambda_2(w_2u_2 - \frac{u_3w_2^2}{w_3}) \right) u_3 \end{aligned}$$

By hypothesis $\|\mathbf{u}^{(I)}\| = 1$ so

$$\begin{aligned} \langle \mathbb{H}^t\mathbb{A}^{(R)}\mathbb{H}\mathbf{u}, \mathbf{u} \rangle &= \lambda_1^{(I)} + 2\frac{\lambda_1}{w_3}(2w_3w_1u_3u_1 - w_3^2u_1^2 - w_1^2u_3^2) \\ &\quad + 2\frac{\lambda_2}{w_3}(2w_3w_2u_2u_3 - w_3^2u_2^2 - w_2^2u_3^2) \\ &= \lambda_1^{(I)} - 2\frac{\lambda_1}{w_3}(w_3u_1 - w_1u_3)^2 - 2\frac{\lambda_2}{w_3}(w_3u_2 - w_2u_3)^2 \end{aligned}$$

Using formula (4.10) give

$$\langle \mathbb{H}^t\mathbb{A}^{(R)}\mathbb{H}\mathbf{u}, \mathbf{u} \rangle = \lambda_1^{(I)} - \frac{2}{w_3}(\lambda_1v_2^2 + \lambda_2v_1^2)$$

- Calculus of $\langle \mathbb{H}^t\mathbb{A}^{(R)}\mathbb{H}\mathbf{u}, \mathbf{v} \rangle$

$$\begin{aligned} \langle \mathbb{H}^t\mathbb{A}^{(R)}\mathbb{H}\mathbf{u}, \mathbf{v} \rangle &= \left(\lambda_1^{(I)}u_1 + 2\lambda_1(w_1u_3 - w_3u_1) \right) v_1 \\ &\quad + \left(\lambda_1^{(I)}u_2 + 2\lambda_2(w_2u_3 - w_3u_2) \right) v_2 \\ &\quad - \left(-\lambda_1^{(I)}u_3 - 2\lambda_1(w_1u_1 - \frac{u_3w_1^2}{w_3}) - 2\lambda_2(w_2u_2 - \frac{u_3w_2^2}{w_3}) \right) v_3 \end{aligned}$$

By hypothesis $\langle \mathbf{u}^{(I)}, \mathbf{v}^{(I)} \rangle = 0$ so

$$\begin{aligned}\langle \mathbb{H}^t \mathbb{A}^{(R)} \mathbb{H} \mathbf{u}, \mathbf{v} \rangle &= 2 \frac{\lambda_1}{w_3} (w_3 w_1 u_3 v_1 - w_3^2 u_1 v_1 + w_3 w_1 u_1 v_3 - w_1^2 u_3 v_3) \\ &\quad + 2 \frac{\lambda_2}{w_3} (w_3 w_2 u_3 v_2 - w_3^2 u_2 v_2 + w_3 w_2 u_2 v_3 - w_2^2 u_3 v_3) \\ &= 2 \frac{\lambda_1}{w_3} (w_3 u_1 - w_1 u_3) (w_1 v_3 - w_3 v_1) + 2 \frac{\lambda_2}{w_3} (w_3 u_2 - w_2 u_3) (w_2 v_3 - w_3 v_2)\end{aligned}$$

Using formulas (4.9) and (4.10) give

$$\langle \mathbb{H}^t \mathbb{A}^{(R)} \mathbb{H} \mathbf{u}, \mathbf{v} \rangle = \frac{2}{w_3} (\lambda_1 u_2 v_2 + \lambda_2 u_1 v_1)$$

- Calculus of $\langle \mathbb{H}^t \mathbb{A}^{(R)} \mathbb{H} \mathbf{u}, \mathbf{w} \rangle$

$$\begin{aligned}\langle \mathbb{H}^t \mathbb{A}^{(R)} \mathbb{H} \mathbf{u}, \mathbf{w} \rangle &= \left(\lambda_1^{(I)} u_1 + 2\lambda_1 (w_1 u_3 - w_3 u_1) \right) w_1 \\ &\quad + \left(\lambda_1^{(I)} u_2 + 2\lambda_2 (w_2 u_3 - w_3 u_2) \right) w_2 \\ &\quad - \left(-\lambda_1^{(I)} u_3 - 2\lambda_1 (w_1 u_1 - \frac{u_3 w_1^2}{w_3}) - 2\lambda_2 (w_2 u_2 - \frac{u_3 w_2^2}{w_3}) \right) w_3\end{aligned}$$

By hypothesis $\langle \mathbf{u}^{(I)}, \mathbf{w}^{(I)} \rangle = 0$ so

$$\langle \mathbb{H}^t \mathbb{A}^{(R)} \mathbb{H} \mathbf{u}, \mathbf{w} \rangle = 0$$

- Calculus of $\langle \mathbb{H}^t \mathbb{A}^{(R)} \mathbb{H} \mathbf{v}, \mathbf{u} \rangle$

$$\begin{aligned}\langle \mathbb{H}^t \mathbb{A}^{(R)} \mathbb{H} \mathbf{v}, \mathbf{u} \rangle &= \left(\lambda_2^{(I)} v_1 + 2\lambda_1 (w_1 v_3 - w_3 v_1) \right) u_1 \\ &\quad + \left(\lambda_2^{(I)} v_2 + 2\lambda_2 (w_2 v_3 - w_3 v_2) \right) u_2 \\ &\quad - \left(-\lambda_2^{(I)} v_3 - 2\lambda_1 (w_1 v_1 - \frac{v_3 w_1^2}{w_3}) - 2\lambda_2 (w_2 v_2 - \frac{v_3 w_2^2}{w_3}) \right) u_3\end{aligned}$$

By hypothesis $\langle \mathbf{u}^{(I)}, \mathbf{v}^{(I)} \rangle = 0$ so

$$\begin{aligned}\langle \mathbb{H}^t \mathbb{A}^{(R)} \mathbb{H} \mathbf{v}, \mathbf{u} \rangle &= 2 \frac{\lambda_1}{w_3} (w_1 w_3 u_1 v_3 - w_3^2 u_1 v_1 + w_1 w_3 u_3 v_1 - w_1^2 u_3 v_3) \\ &\quad + 2 \frac{\lambda_2}{w_3} (w_2 w_3 u_2 v_3 - w_3^2 u_2 v_2 + w_2 w_3 u_3 v_2 - w_2^2 u_3 v_3) \\ &= 2 \frac{\lambda_1}{w_3} (w_3 u_1 - w_1 u_3) (w_1 v_3 - w_3 v_1) + 2 \frac{\lambda_2}{w_3} (w_3 u_2 - w_2 u_3) (w_2 v_3 - w_3 v_2)\end{aligned}$$

Using formulas (4.9) and (4.10) give

$$\langle \mathbb{H}^t \mathbb{A}^{(R)} \mathbb{H} \mathbf{v}, \mathbf{u} \rangle = \frac{2}{w_3} (\lambda_1 u_2 v_2 + \lambda_2 u_1 v_1)$$

- Calculus of $\langle \mathbb{H}^t \mathbb{A}^{(R)} \mathbb{H} \mathbf{v}, \mathbf{v} \rangle$

$$\begin{aligned}\langle \mathbb{H}^t \mathbb{A}^{(R)} \mathbb{H} \mathbf{v}, \mathbf{v} \rangle &= \left(\lambda_2^{(I)} v_1 + 2\lambda_1 (w_1 v_3 - w_3 v_1) \right) v_1 \\ &\quad + \left(\lambda_2^{(I)} v_2 + 2\lambda_2 (w_2 v_3 - w_3 v_2) \right) v_2 \\ &\quad - \left(-\lambda_2^{(I)} v_3 - 2\lambda_1 (w_1 v_1 - \frac{v_3 w_1^2}{w_3}) - 2\lambda_2 (w_2 v_2 - \frac{v_3 w_2^2}{w_3}) \right) v_3\end{aligned}$$

By hypothesis $\|\mathbf{v}^{(I)}\| = 1$ so

$$\begin{aligned}\left\langle \mathbb{H}^t \mathbb{A}^{(R)} \mathbb{H} \mathbf{v}, \mathbf{v} \right\rangle &= \lambda_2^{(I)} + 2 \frac{\lambda_1}{w_3} (2w_1 w_3 v_1 v_3 - w_3^2 v_1^2 - w_1^2 v_3^2) \\ &\quad + 2 \frac{\lambda_2}{w_3} (2w_2 w_3 v_2 v_3 - w_3^2 v_2^2 - w_2^2 v_3^2) \\ &= \lambda_2^{(I)} - 2 \frac{\lambda_1}{w_3} (w_3 v_1 - w_1 v_3)^2 - 2 \frac{\lambda_2}{w_3} (w_3 v_2 - w_2 v_3)^2\end{aligned}$$

Using formula (4.9) give

$$\left\langle \mathbb{H}^t \mathbb{A}^{(R)} \mathbb{H} \mathbf{v}, \mathbf{v} \right\rangle = \lambda_2^{(I)} - \frac{2}{w_3} (\lambda_1 u_2^2 + \lambda_2 u_1^2)$$

- Calculus of $\langle \mathbb{H}^t \mathbb{A}^{(R)} \mathbb{H} \mathbf{v}, \mathbf{w} \rangle$

$$\begin{aligned}\left\langle \mathbb{H}^t \mathbb{A}^{(R)} \mathbb{H} \mathbf{v}, \mathbf{w} \right\rangle &= \left(\lambda_2^{(I)} v_1 + 2\lambda_1 (w_1 v_3 - w_3 v_1) \right) w_1 \\ &\quad + \left(\lambda_2^{(I)} v_2 + 2\lambda_2 (w_2 v_3 - w_3 v_2) \right) w_2 \\ &\quad - \left(-\lambda_2^{(I)} v_3 - 2\lambda_1 (w_1 v_1 - \frac{v_3 w_1^2}{w_3}) - 2\lambda_2 (w_2 v_2 - \frac{v_3 w_2^2}{w_3}) \right) w_3\end{aligned}$$

By hypothesis $\langle \mathbf{v}^{(I)}, \mathbf{w}^{(I)} \rangle = 0$ so

$$\left\langle \mathbb{H}^t \mathbb{A}^{(R)} \mathbb{H} \mathbf{u}, \mathbf{w} \right\rangle = 0$$

With all these formulas we obtain :

$$\mathbb{H}^t \mathbb{A}^{(R)} \mathbb{H} = \begin{pmatrix} \lambda_1^{(I)} - \frac{2}{w_3} (\lambda_1 v_2^2 + \lambda_2 v_1^2) & \frac{2}{w_3} (\lambda_1 u_2 v_2 + \lambda_2 u_1 v_1) & 0 \\ \frac{2}{w_3} (\lambda_1 u_2 v_2 + \lambda_2 u_1 v_1) & \lambda_2^{(I)} - \frac{2}{w_3} (\lambda_1 u_2^2 + \lambda_2 u_1^2) & 0 \\ 0 & 0 & 0 \end{pmatrix} \quad (4.14)$$

As $\mathbb{H}^t \mathbb{A}^{(R)} \mathbb{H}$ is similar to $\mathbb{A}^{(R)}$, we refind that 0 is an eigenvalue of $\mathbb{A}^{(R)}$ and the two others eigenvalues $\lambda_1^{(R)}$ and $\lambda_2^{(R)}$ are also eingenvales of

$$\mathbb{A} = \begin{pmatrix} \lambda_1^{(I)} - \frac{2}{w_3} (\lambda_1 v_2^2 + \lambda_2 v_1^2) & \frac{2}{w_3} (\lambda_1 u_2 v_2 + \lambda_2 u_1 v_1) \\ \frac{2}{w_3} (\lambda_1 u_2 v_2 + \lambda_2 u_1 v_1) & \lambda_2^{(I)} - \frac{2}{w_3} (\lambda_1 u_2^2 + \lambda_2 u_1^2) \end{pmatrix}$$

Furthermore, we have that

$$\text{Tr}(\mathbb{A}) = \lambda_1^{(R)} + \lambda_2^{(R)} \quad (4.15)$$

$$\det(\mathbb{A}) = \lambda_1^{(R)} \lambda_2^{(R)} \quad (4.16)$$

To establish inequality 4.5, we use equation (4.15) under hypothesis (H) and $w_3 < 0$ to get

$$\lambda_1^{(R)} + \lambda_2^{(R)} \geq \lambda_1^{(I)} + \lambda_2^{(I)} - \frac{2}{w_3} \lambda_1 ((u_1^2 + u_2^2) + (v_1^2 + v_2^2)).$$

We remark that

$$\begin{aligned}((u_1^2 + u_2^2) + (v_1^2 + v_2^2)) &= (u_1 + v_2)^2 + (u_2 - v_1)^2 - 2(u_1 v_2 - u_2 v_1) \\ &= (u_1 + v_2)^2 + (u_2 - v_1)^2 - 2w_3\end{aligned}$$

and so, we obtain

$$\begin{aligned}\lambda_1^{(R)} + \lambda_2^{(R)} &\geq \lambda_1^{(I)} + \lambda_2^{(I)} - \frac{2}{w_3} \lambda_1 ((u_1 + v_2)^2 + (u_2 - v_1)^2 - 2w_3) \\ &\geq \lambda_1^{(I)} + \lambda_2^{(I)} + 4\lambda_1 + 4\lambda_1 ((u_1 + v_2)^2 + (u_2 - v_1)^2) \\ &\geq \lambda_1^{(I)} + \lambda_2^{(I)} + 4\lambda_1.\end{aligned}$$

To establish inequality 4.6, we use equation (4.16) to get

$$\begin{aligned}
\lambda_1^{(R)} \lambda_2^{(R)} &= \left(\lambda_1^{(I)} - \frac{2}{w_3} (\lambda_1 v_2^2 + \lambda_2 v_1^2) \right) \left(\lambda_2^{(I)} - \frac{2}{w_3} (\lambda_1 u_2^2 + \lambda_2 u_1^2) \right) \\
&\quad - \frac{4}{w_3^2} (\lambda_1 u_2 v_2 + \lambda_2 u_1 v_1)^2 \\
&= \lambda_1^{(I)} \lambda_2^{(I)} + \frac{4}{w_3^2} \left[(\lambda_1 v_2^2 + \lambda_2 v_1^2) (\lambda_1 u_2^2 + \lambda_2 u_1^2) - (\lambda_1 u_2 v_2 + \lambda_2 u_1 v_1)^2 \right] \\
&\quad - \frac{2}{w_3} \left((\lambda_1 v_2^2 + \lambda_2 v_1^2) \lambda_2^{(I)} + (\lambda_1 u_2^2 + \lambda_2 u_1^2) \lambda_1^{(I)} \right) \\
&= \lambda_1^{(I)} \lambda_2^{(I)} - \frac{2}{w_3} \left((\lambda_1 v_2^2 + \lambda_2 v_1^2) \lambda_2^{(I)} + (\lambda_1 u_2^2 + \lambda_2 u_1^2) \lambda_1^{(I)} \right) \\
&\quad + \frac{4}{w_3^2} [\lambda_1^2 u_2^2 v_2^2 + \lambda_1 \lambda_2 u_1^2 v_2^2 + \lambda_1 \lambda_2 v_1^2 u_2^2 - \lambda_1^2 u_2^2 v_2^2 - 2\lambda_1 \lambda_2 u_1 v_1 u_2 v_2 - \lambda_1 \lambda_2 v_1^2 u_2^2] \\
&= \lambda_1^{(I)} \lambda_2^{(I)} - \frac{2}{w_3} \left((\lambda_1 v_2^2 + \lambda_2 v_1^2) \lambda_2^{(I)} + (\lambda_1 u_2^2 + \lambda_2 u_1^2) \lambda_1^{(I)} \right) \\
&\quad + \frac{4}{w_3^2} \lambda_1 \lambda_2 [u_1^2 v_2^2 - 2u_1 v_1 u_2 v_2 + v_1^2 u_2^2] \\
&= \lambda_1^{(I)} \lambda_2^{(I)} - \frac{2}{w_3} \left((\lambda_1 v_2^2 + \lambda_2 v_1^2) \lambda_2^{(I)} + (\lambda_1 u_2^2 + \lambda_2 u_1^2) \lambda_1^{(I)} \right) \\
&\quad + \frac{4}{w_3^2} \lambda_1 \lambda_2 (u_1 v_2 - v_1 u_2)^2
\end{aligned}$$

From formula (4.8), we have $u_1 v_2 - v_1 u_2 = w_3$ and so we obtain

$$\begin{aligned}
\lambda_1^{(R)} \lambda_2^{(R)} &= \lambda_1^{(I)} \lambda_2^{(I)} - \frac{2}{w_3} \left((\lambda_1 v_2^2 + \lambda_2 v_1^2) \lambda_2^{(I)} + (\lambda_1 u_2^2 + \lambda_2 u_1^2) \lambda_1^{(I)} \right) \\
&\quad + 4\lambda_1 \lambda_2
\end{aligned}$$

Under hypothesis (H) and $w_3 < 0$, we clearly have

$$\lambda_1^{(R)} \lambda_2^{(R)} \geq \lambda_1^{(I)} \lambda_2^{(I)} + 4\lambda_1^2 > 0$$

and we obtain formula (4.6). □

REFERENCES

[eC89] M.Balabane et C.Bardos. *Equation des ondes : solutions asymptotiques et singularités*. Compte Rendus INRIA de la session électromagnétisme, 1989.

UNIVERSITÉ PARIS XIII, INSTITUT GALILIÉE, LAGA CNRS UMR 7539,, 99, Av. J.-B. CLÉMENT F-93430 VILLETTANEUSE, FRANCE

E-mail address: cuvelier@math.univ-paris13.fr