

Exact relaxation for polynomial optimization on semi-algebraic sets

Marta Abril Bucero, Bernard Mourrain

► **To cite this version:**

Marta Abril Bucero, Bernard Mourrain. Exact relaxation for polynomial optimization on semi-algebraic sets. 2014. hal-00846977v4

HAL Id: hal-00846977

<https://hal.inria.fr/hal-00846977v4>

Preprint submitted on 1 Jul 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

EXACT RELAXATION FOR POLYNOMIAL OPTIMIZATION ON SEMI-ALGEBRAIC SETS

MARTA ABRIL BUCERO, BERNARD MOURRAIN

ABSTRACT. In this paper, we study the problem of computing the infimum of a real polynomial function f on a closed basic semialgebraic set S and the points where this infimum is reached, if they exist. We show that when the infimum is reached, a Semi-Definite Program hierarchy constructed from the Karush-Kuhn-Tucker ideal is always exact and that the vanishing ideal of the KKT minimizer points is generated by the kernel of the associated moment matrix in that degree, even if this ideal is not zero-dimensional. We also show that this relaxation allows to detect when there is no KKT minimizer. Analysing the properties of the Fritz John variety, we show how to find all the minimizers of f . We prove that the exactness of the relaxation depends only on the real points which satisfy these constraints. This exploits representations of positive polynomials as elements of the preordering modulo the KKT ideal, which only involves polynomials in the initial set of variables. The approach provides a uniform treatment of different optimization problems considered previously. Applications to global optimization, optimization on semialgebraic sets defined by regular sets of constraints, optimization on finite semialgebraic sets and real radical computation are given.

1. INTRODUCTION

The problem we consider in this paper is the following:

$$(1) \quad \begin{aligned} \inf_{\mathbf{x} \in \mathbb{R}^n} \quad & f(\mathbf{x}) \\ \text{s.t.} \quad & g_1^0(\mathbf{x}) = \dots = g_{n_1}^0(\mathbf{x}) = 0 \\ & g_1^+(\mathbf{x}) \geq 0, \dots, g_{n_2}^+(\mathbf{x}) \geq 0 \end{aligned}$$

where $f, g_1^0, \dots, g_{n_1}^0, g_1^+, \dots, g_{n_2}^+ \in \mathbb{R}[\mathbf{x}]$ are polynomial functions in n variables x_1, \dots, x_n .

Hereafter, we fix the set of constraints $\mathbf{g} = \{g_1^0, \dots, g_{n_1}^0; g_1^+, \dots, g_{n_2}^+\} = \{\mathbf{g}^0; \mathbf{g}^+\}$ and denoted by S the basic semi-algebraic set defined by these constraints.

The points $\mathbf{x}^* \in S$ which satisfy $f(\mathbf{x}^*) = \inf_{\mathbf{x} \in S} f(\mathbf{x})$ are called the *minimizers points* of f on S . If the set of minimizers is not empty, we say that the *minimization problem is feasible*.

The objectives of the method we consider are to detect if the minimization problem is feasible and to compute the minimum value of f and the minimizer points where this minimum value is reached, when the problem is feasible. Though this global minimization problem is known to be NP-hard (see e.g. [24]), a practical challenge is to devise methods which can approximate or compute efficiently the solutions of the problem.

About a decade ago, a relaxation approach has been proposed in [13] (see also [29], [35]) to solve this difficult problem. Instead of searching points where the polynomial f reaches its minimum f^* , a probability measure which minimizes the function f is searched. This problem is relaxed into a hierarchy of finite dimensional convex minimization problems, which can be solved by Semi-Definite Programming (SDP) techniques. The sequence of SDP minima converges to the minimum f^* [13]. This hierarchy of SDP problems can be formulated in terms of linear matrix inequalities on moment matrices associated to the

set of monomials of degree t or less, for increasing values of t . The dual hierarchy can be described as a sequence of maximization problems over the cone of polynomials that are Sums of Squares (SoS). A feasibility condition is needed to prove that this dual hierarchy of maximization problems also converges to the minimum f^* , i.e. that there is no duality gap.

This approach provides a very interesting way to approximate a global optimum of a polynomial function on S . But one may wonder if using this approach, it is possible to compute in a finite number of steps, this minimum and the minimizer points when the problem is feasible. From a computational point of view, the following issues need to be addressed:

- (1) Is it possible to use an *exact* SDP hierarchy, i.e. which converges in a finite number of steps?
- (2) How can we recover all the points where the optimum is achieved if the optimization problem is feasible?

To address the first issue, the following strategy has been considered: add polynomial inequalities or equalities satisfied by the points where the function f is minimum.

A first family of methods are used when the set S is compact or when the minimizer set can be bounded easily. By adding an inequality constraint, one can then transform S into a compact subset of \mathbb{R}^n , for which exact hierarchies can be used [13], [22]. It is shown in [17] that if the complex variety defined by the equalities $\mathbf{g}^0 = 0$ is finite (and thus S is compact), then the hierarchy of relaxation problems introduced by Lasserre in [13] is exact. It is also proved that there is no duality gap if the generators of this ideal satisfy some regularity conditions. In [27], it is proved that if the real variety defined by the equalities $\mathbf{g}^0 = 0$ is finite, then the hierarchy of relaxation problems introduced by Lasserre is exact, this answers an open question in [18].

In a second family of methods, equality constraints which are naturally satisfied by the minimizer points are added. These constraints are for instance the gradient of f when $S = \mathbb{R}^n$ or the Karush-Kuhn-Tucker (KKT) constraints, obtained by introducing Lagrange multipliers. In [28], it is proved that a relaxation hierarchy using the gradient constraints is exact when the gradient ideal is radical. In [23], it is shown that this gradient hierarchy is exact, when the global minimizers satisfy the Boundary Hessian condition. In [5], it is proved that a relaxation hierarchy which involves the KKT constraints is exact when the KKT ideal is radical. In [9], a relaxation hierarchy obtained by projection of the KKT constraints is proved to be exact under a regularity condition on the *real* minimizer points¹. In [25], a similar relaxation hierarchy is shown to be exact under a stronger regularity condition for the *complex* points of associated KKT varieties. These regularity conditions require that the gradient of the active constraints evaluated at the points of S or of some complex varieties are linearly independent. Thus they cannot be used for general semi algebraic sets S , for instance when S is a real non-complete intersection variety.

Moreover, the assumption that the minimum is reached at a KKT point is required. Unfortunately, in some cases the set of KKT points of S can be empty. As we shall see, this obstacle can be removed using Fritz John variety (see [11, 21]). There is not much work dedicated to this issue (see [14]).

The case where the infimum value is not reached has also been studied. In [33], relaxation techniques are studied for functions for which the minimum is not reached and which satisfy some special properties “at infinity”. In [8], tangency constraints are used in a relaxation hierarchy which converges to the global minimum of a polynomial, when the polynomial

¹The results of this paper are true but a problem appears in the proof which we fix in the present paper.

is bounded by below over \mathbb{R}^n . In [7], generic changes of coordinates and a partial gradient ideal are used in a relaxation hierarchy which also converges to the global minimum of f on \mathbb{R}^n .

In the cases studied so far, the exactness of the relaxation is proved under a genericity condition or a compactness property. From an algorithmic point of view, the flat extension condition of Curto-Fialkow [4] is used in most of the works [10, 17, 16, 18] to detect the exactness of the hierarchy, when the number of minimizers is finite. In [26], it is proved that the Curto-Fialkow flat extension criterion is eventually satisfied on truncated moment matrices under some regularity conditions or archimedean conditions. In [12], a sparse extension [19] of this flat extension condition is used to compute zero-dimensional real radical ideals.

The second issue is related to the problem of computing all the minimizer points, which is also important from a practical point of view. In [16], the kernel of moment matrices is used to compute generators of the real radical of an ideal. This method is improved in [12] to compute a border basis of the real radical, involving SDP problems of significantly smaller size, when the real radical ideal is zero-dimensional. The case of positive dimensional real radical ideal is analysed in [31] and [20]. The problem of computing the minimizer ideal for general optimization problems from exact relaxation hierarchies has not been addressed, though it is mentioned in [26] for zero-dimensional minimizer ideals.

Notice that Problem (1) can be attacked from a purely algebraic point of view. It reduces to the computation of a (minimal) critical value and polynomial system solvers can be used to tackle it (see e.g. [30], [6]). But in this case, the complex solutions of the underlying algebraic system come into play and additional computation efforts should be spent to remove these extraneous solutions. Semi-algebraic techniques such as Cylindrical Algebraic Decomposition or extensions [32] may also be considered here, providing algorithms to solve Problem (1), but suffering from similar issues.

Contributions. Our aim is to show that for the general polynomial optimization problem (1), exact SDP relaxations can be constructed, which either detect that the problem is infeasible or compute the minimal value and the ideal associated to the minimizer points. The main contributions are the following:

- We prove that exact relaxation hierarchies depending on the variables \mathbf{x} can be constructed for solving the optimization problem (1) (see Theorem 6.3 and Theorem 5.10).
- We show that even if the minimizer points are not KKT points, we can find them using the Fritz John variety (see Section 3.3 and Section 3.4). We describe an approach, which splits this minimizer set into the KKT minimizer set and the singular minimizer set which can be recursively computed using the same method.
- We prove that if the set of KKT minimizers is empty, the SDP relaxation will eventually be empty (Theorem 6.3).
- We prove that the KKT minimizer ideal can be constructed from the moment matrix of an optimal linear form, when the corresponding relaxation is exact, even if the ideal is not zero-dimensional (Theorem 5.10).
- We prove that the exactness of the relaxation depends only on the real points which satisfy these constraints (Theorem 5.10).
- We provide a general approach which allows us to treat in a uniform way and to extend results on the representation of polynomials which are positive (resp. non-negative) on the critical points (see [5] and Theorem 4.9) and on the exactness of relaxation hierarchies (see [28], [8], [16], [25], [12], [27] and Theorem 6.2, Theorem 6.4, Theorem 6.5, Theorem 6.6).

Content. The paper is organized as follows. In Section 2, we recall algebraic concepts and describe the hierarchy of finite dimensional convex optimization problems considered. In Section 3, we analyse the varieties associated to the critical points of the minimization problem. Section 4, is devoted to the representation of positive and non-negative polynomials on the critical points as sum of squares modulo the gradient ideal. In Section 5, we prove that when the order of relaxation is big enough, the sequence of finite dimensional convex optimization problems attains its limit and the minimizer ideal can be generated from the solution of our relaxation problem. In Section 6, we analyse some consequences of these results. Finally, Section 7 contains several examples which illustrate the approach.

2. IDEALS, VARIETIES, OPTIMIZATION AND RELAXATION

In this section, we recall some algebraic concepts as ideals and varieties and we set our notation.

2.1. Ideals and varieties. Let $\mathbb{K}[\mathbf{x}]$ be the set of the polynomials in the variables $\mathbf{x} = (x_1, \dots, x_n)$, with coefficients in the field \mathbb{K} . Hereafter, we choose² $\mathbb{K} = \mathbb{R}$ or \mathbb{C} . Let $\overline{\mathbb{K}}$ denotes the algebraic closure of \mathbb{K} . For $\alpha \in \mathbb{N}^n$, $\mathbf{x}^\alpha = x_1^{\alpha_1} \cdots x_n^{\alpha_n}$ is the monomial with exponent α and degree $|\alpha| = \sum_i \alpha_i$. The set of all monomials in \mathbf{x} is denoted $\mathcal{M} = \mathcal{M}(\mathbf{x})$.

For $t \in \mathbb{N} \cup \{\infty\}$ and $B \subseteq \mathbb{K}[\mathbf{x}]$, we introduce the following sets:

- B_t is the set of elements of B of degree $\leq t$,
- $\langle B \rangle = \{ \sum_{f \in B} \lambda_f f \mid f \in B, \lambda_f \in \mathbb{K} \}$ is the linear span of B ,
- $(B) = \{ \sum_{f \in B} p_f f \mid p_f \in \mathbb{K}[\mathbf{x}], f \in B \}$ is the ideal in $\mathbb{K}[\mathbf{x}]$ generated by B ,
- $B_{\langle t \rangle} = \{ \sum_{f \in B_t} p_f f \mid p_f \in \mathbb{K}[\mathbf{x}]_{t - \deg(f)} \}$ is the vector space spanned by $\{ \mathbf{x}^\alpha f \mid f \in B_t, |\alpha| \leq t - \deg(f) \}$,
- $\mathcal{Q}_t^+ = \{ \sum_{i=1}^l p_i^2 \mid l \in \mathbb{N}, p_i \in \mathbb{R}[\mathbf{x}]_t \}$ is the set of finite sums of squares of polynomials of degree $\leq t$; $\mathcal{Q}^+ = \mathcal{Q}_\infty^+$.

By definition $B_{\langle t \rangle} \subseteq (B) \cap \mathbb{K}[\mathbf{x}]_t = (B)_t$, but the inclusion may be strict.

By convention, a set of constrains $C = \{c_1^0, \dots, c_{n_1}^0; c_1^+, \dots, c_{n_2}^+\} \subset \mathbb{R}[\mathbf{x}]$ is a finite set of polynomials composed of a subset $C^0 = \{c_1^0, \dots, c_{n_1}^0\}$ corresponding to the equality constraints and a subset $C^+ = \{c_1^+, \dots, c_{n_2}^+\}$ corresponding to the non-negativity constraints. For two set of constraints $C, C' \subset \mathbb{R}[\mathbf{x}]$, we say that $C \subset C'$ if $C^0 \subset C'^0$ and $C^+ \subset C'^+$.

Definition 2.1. For $t \in \mathbb{N} \cup \{\infty\}$ and a set of constraints $C = \{c_1^0, \dots, c_{n_1}^0; c_1^+, \dots, c_{n_2}^+\} \subset \mathbb{R}[\mathbf{x}]$, we define the (truncated) quadratic module of C by

$$\mathcal{Q}_t(C) = \left\{ \sum_{i=1}^{n_2} c_i^0 h_i + s_0 + \sum_{j=1}^{n_2} c_j^+ s_j \mid h_i \in \mathbb{R}[\mathbf{x}]_{2t - \deg(c_i^0)}, s_0 \in \mathcal{Q}_t^+, s_i \in \mathcal{Q}_{t - \lfloor \deg(c_i^+) / 2 \rfloor}^+ \right\}.$$

If \tilde{C} is such that $\tilde{C}^0 = C^0$ and $\tilde{C}^+ = \{ \prod (c_1^+)^{\epsilon_1} \cdots (c_{n_2}^+)^{\epsilon_{n_2}} \mid \epsilon_i \in \{0, 1\} \}$, $\mathcal{Q}_t(\tilde{C})$ is also called the (truncated) preordering of C and denoted $\mathcal{P}_t(C)$. When $t = \infty$, $\mathcal{P}(C) := \mathcal{P}_\infty(C)$ is the preordering of C . The (truncated) preordering generated by the positive constraints is denoted $\mathcal{P}^+(C) = \mathcal{P}(C^+)$.

Definition 2.2. Given $t \in \mathbb{N} \cup \{\infty\}$ and a set of constraints $C \subset \mathbb{R}[\mathbf{x}]$, we define

$$\mathcal{N}_t(C) := \{ \Lambda \in (\mathbb{R}[\mathbf{x}]_{2t})^* \mid \Lambda(p) \geq 0, \forall p \in \mathcal{Q}_t(C), \Lambda(1) = 1 \}.$$

When we replace $\mathcal{Q}_t(C)$ by $\mathcal{P}_t(C)$ in this definition, we denote the corresponding set by $\mathcal{L}_t(C)$.

²For notational simplicity, we consider only these two fields, but \mathbb{R} and \mathbb{C} can be replaced respectively by any real closed field and any field containing its algebraic closure.

Given a set $I \subseteq \mathbb{K}[\mathbf{x}]$ and a field $\mathbb{L} \supseteq \mathbb{K}$, we denote by

$$\mathcal{V}^{\mathbb{L}}(I) := \{x \in \mathbb{L}^n \mid f(x) = 0 \forall f \in I\}$$

its associated variety in \mathbb{L}^n . By convention $\mathcal{V}(I) = \mathcal{V}^{\overline{\mathbb{K}}}(I)$, where $\overline{\mathbb{K}}$ is the algebraic closure of \mathbb{K} . We also consider sets of homogeneous equations I and the varieties $\mathbb{P}\mathcal{V}(I)$ (resp. $\mathbb{P}\mathcal{V}^{\mathbb{R}}(I)$) defined in the projective space \mathbb{P}^n (resp. the real projective space $\mathbb{R}\mathbb{P}^n$).

For a set $V \subseteq \mathbb{K}^n$, we define its vanishing ideal

$$\mathcal{I}(V) := \{p \in \mathbb{K}[\mathbf{x}] \mid p(v) = 0 \forall v \in V\}.$$

For a set $V \subset \mathbb{L}^n$ with $\mathbb{L} \supseteq \mathbb{K}$, $V^{\mathbb{K}} = V \cap \mathbb{K}^n$. Hereafter, we take $\mathbb{K} = \mathbb{R}$ and $\mathbb{L} = \mathbb{C}$, so that $\mathcal{V}(I) = \mathcal{V}^{\mathbb{C}}(I)$, $\mathcal{V}^{\mathbb{R}}(I) = \mathcal{V}(I)^{\mathbb{R}} = \mathcal{V}(I) \cap \mathbb{R}^n$.

Definition 2.3. For a set of constraints $C = (C^0; C^+) \subset \mathbb{R}[\mathbf{x}]$,

$$\begin{aligned} \mathcal{S}(C) &:= \{\mathbf{x} \in \mathbb{R}^n \mid c^0(\mathbf{x}) = 0 \forall c^0 \in C^0, c^+(\mathbf{x}) \geq 0 \forall c^+ \in C^+\}, \\ \mathcal{S}^+(C) &:= \{\mathbf{x} \in \mathbb{R}^n \mid c^+(\mathbf{x}) \geq 0 \forall c^+ \in C^+\}. \end{aligned}$$

To describe the vanishing ideal of these sets, we introduce the following ideals:

Definition 2.4. For a set of constraints $C = (C^0; C^+) \subset \mathbb{R}[\mathbf{x}]$,

$$\begin{aligned} \sqrt{C^0} &= \{p \in \mathbb{R}[\mathbf{x}] \mid p^m \in (C^0) \text{ for some } m \in \mathbb{N} \setminus \{0\}\} \\ \sqrt[{\mathbb{R}}]{C^0} &= \{p \in \mathbb{R}[\mathbf{x}] \mid p^{2m} + q \in (C^0) \text{ for some } m \in \mathbb{N} \setminus \{0\}, q \in \mathcal{Q}^+\} \\ \sqrt[{}^+]{C^0} &= \{p \in \mathbb{R}[\mathbf{x}] \mid p^{2m} + q \in (C^0) \text{ for some } m \in \mathbb{N} \setminus \{0\}, q \in \mathcal{P}^+(C)\} \end{aligned}$$

These ideals are called respectively the radical of C^0 , the real radical of C^0 , the C^+ -radical of C^0 .

Remark 2.5. If $C^+ = \emptyset$, then $\sqrt[{}^+]{C^0} = \sqrt[{\mathbb{R}}]{C^0}$.

The following three famous theorems relate vanishing and radical ideals:

Theorem 2.6. Let $C = (C^0; C^+)$ be a set of constraints of $\mathbb{R}[\mathbf{x}]$.

- (i) **Hilbert's Nullstellensatz** (see, e.g., [3, §4.1]) $\sqrt{C^0} = \mathcal{I}(\mathcal{V}^{\mathbb{C}}(C^0))$.
- (ii) **Real Nullstellensatz** (see, e.g., [1, §4.1]) $\sqrt[{\mathbb{R}}]{C^0} = \mathcal{I}(\mathcal{V}^{\mathbb{R}}(C^0))$.
- (iii) **Positivstellensatz** (see, e.g., [1, §4.4]) $\sqrt[{}^+]{C^0} = \mathcal{I}(\mathcal{S}(C)) = \mathcal{I}(\mathcal{V}^{\mathbb{R}}(C^0) \cap \mathcal{S}^+(C))$.

2.2. Relaxation hierarchy. The approach proposed by Lasserre in [13] to solve Problem (1) consists in approximating the optimization problem by a sequence of finite dimensional convex optimization problems, which can be solved efficiently by Semi-Definite Programming tools. This sequence is called Lasserre hierarchy of relaxation problems. Let C be a set of constraints in $\mathbb{R}[\mathbf{x}]$ such that $\mathcal{S}(C) = \mathcal{S}(\mathbf{g})$. Hereafter, we consider the relaxation hierarchy associated to preordering sequences:

$$\cdots \subset \mathcal{L}_{t+1}(C) \subset \mathcal{L}_t(C) \subset \cdots \text{ and } \cdots \subset \mathcal{P}_t(C) \subset \mathcal{P}_{t+1}(C) \subset \cdots$$

These convex sets are used to define extrema that approximate the solution of the minimization problem (1).

Definition 2.7. Let $t \in \mathbb{N}$ and let C be the set of constraints in $\mathbb{R}[\mathbf{x}]$. We define the following extrema:

- $f_C^* = \inf_{\mathbf{x} \in \mathcal{S}(C)} f(\mathbf{x})$,
- $f_{t,C}^{\mu} = \inf \{\Lambda(f) \text{ s.t. } \Lambda \in \mathcal{L}_t(C)\}$,
- $f_{t,C}^{sos} = \sup \{\gamma \in \mathbb{R} \text{ s.t. } f - \gamma \in \mathcal{P}_t(C)\}$.

By convention if the corresponding sets are empty, $f_C^* = -\infty$, $f_{t,C}^{sos} = -\infty$ and $f_{t,C}^\mu = +\infty$.

Remark 2.8. We have $f_{t,C}^{sos} \leq f_{t,C}^\mu \leq f_C^*$.

Indeed, if there exists $\gamma \in \mathbb{R}$ such that $f - \gamma = q \in \mathcal{P}_t(C)$ then $\forall \Lambda \in \mathcal{L}_t(C)$, $\Lambda(f - \gamma) = \Lambda(f) - \gamma = \Lambda(q) \geq 0$, which proves the first inequality.

Since for any $\mathbf{s} \in S$, the evaluation $\mathbf{1}_\mathbf{s} : p \in \mathbb{R}[\mathbf{x}] \mapsto p(\mathbf{s})$ is in $\mathcal{L}_t(C)$, we have $\mathbf{1}_\mathbf{s}(f) = f(\mathbf{s}) \geq f_{t,C}^\mu$. This proves the second inequality.

As $\mathcal{L}_{t+1}(C) \subset \mathcal{L}_t(C)$ and $\mathcal{P}_t(C) \subset \mathcal{P}_{t+1}(C)$ we have the following increasing sequences for $t \in \mathbb{N}$:

$$\cdots f_{t,C}^\mu \leq f_{t+1,C}^\mu \leq \cdots \leq f_C^* \text{ and } \cdots f_{t,C}^{sos} \leq f_{t+1,C}^{sos} \leq \cdots \leq f_C^*.$$

The foundation of Lasserre relaxation method is to show that these sequences converge to f_C^* , see [13].

We are interested in constructing hierarchies for which, the minimum f_C^* is reached in a finite number of steps. Such hierarchies are called *exact*. We are also interested to compute the minimizers points. For that purpose, we introduce now the truncated Hankel operators, which play a central role in the construction of the minimizer ideal of f on S .

Definition 2.9. For $t \in \mathbb{N}$ and a linear form $\Lambda \in (\mathbb{R}[\mathbf{x}]_{2t})^*$, we define the truncated Hankel operator as the map $M_\Lambda^t : \mathbb{R}[\mathbf{x}]_t \rightarrow (\mathbb{R}[\mathbf{x}]_t)^*$ such that $M_\Lambda^t(p)(q) = \Lambda(pq)$ for $p, q \in \mathbb{R}[\mathbf{x}]_t$. Its matrix in monomial bases of $\mathbb{R}[\mathbf{x}]_t$ and $(\mathbb{R}[\mathbf{x}]_t)^*$ is also called the moment matrix of Λ .

The kernel of the truncated Hankel operator is

$$(2) \quad \ker M_\Lambda^t = \{p \in \mathbb{R}[\mathbf{x}]_t \mid \Lambda(pq) = 0 \forall q \in \mathbb{R}[\mathbf{x}]_t\}.$$

Given $t \in \mathbb{N}$ and $C = \{0\}$ and $\Lambda, \Lambda' \in \mathbb{R}[\mathbf{x}]_{2t}^*$, we easily check the following properties:

- $\forall p \in \mathbb{R}[\mathbf{x}]_t$, $\Lambda(p^2) = 0$ implies $p \in \ker M_\Lambda^t$.
- $\ker M_{\Lambda+\Lambda'}^t = \ker M_\Lambda^t \cap \ker M_{\Lambda'}^t$.

The kernel of truncated Hankel operator is used to compute generators of the minimizer ideal, as we will see.

3. VARIETIES OF CRITICAL POINTS

Before describing how to compute the minimizer points, we analyse the geometry of this minimization problem and the varieties associated to its critical points. In the following, we denote by $\mathbf{y} = (\mathbf{x}, \mathbf{u}, \mathbf{v})$ and $\mathbf{z} = (\mathbf{x}, \mathbf{u}, \mathbf{v}, \mathbf{s})$, the $n + n_1 + n_2$ and $n + n_1 + 2n_2$ variables of these problems. For any ideal $J \subset \mathbb{R}[\mathbf{z}]$, we denote $J^\mathbf{x} = J \cap \mathbb{R}[\mathbf{x}]$. The projection of $\mathbb{C}^n \times \mathbb{C}^{n_1+2n_2}$ (resp. $\mathbb{C}^n \times \mathbb{C}^{n_1+n_2}$) on \mathbb{C}^n is denoted $\pi^\mathbf{x}$.

3.1. The gradient variety. A natural approach to deal with constraints in optimization problems is to introduce Lagrangian multipliers. Replacing the inequalities $g_i^+ \geq 0$ by the equalities $g_i^+ - s_i^2 = 0$ (adding new variables s_i) and introducing new parameters for all the equality constraints yields the following minimization problem:

$$(3) \quad \begin{aligned} \inf_{(\mathbf{x}, \mathbf{u}, \mathbf{v}, \mathbf{s}) \in \mathbb{R}^n \times \mathbb{R}^{n_1+2n_2}} & f(\mathbf{x}) \\ \text{s.t.} & \quad \nabla F(\mathbf{x}, \mathbf{u}, \mathbf{v}, \mathbf{s}) = 0 \end{aligned}$$

where $F(\mathbf{x}, \mathbf{u}, \mathbf{v}, \mathbf{s}) = f(\mathbf{x}) - \sum_{i=1}^{n_1} u_i g_i(\mathbf{x}) - \sum_{j=1}^{n_2} v_j (g_j^+(\mathbf{x}) - s_j^2)$, $\mathbf{u} = (u_1, \dots, u_{n_1})$, $\mathbf{v} = (v_1, \dots, v_{n_2})$ and $\mathbf{s} = (s_1, \dots, s_{n_2})$.

Definition 3.1. *The gradient ideal of $F(\mathbf{z})$ is:*

$$I_{grad} = (\nabla F(\mathbf{z})) = (F_1, \dots, F_n, g_1^0, \dots, g_{n_1}^0, g_1^+ - s_1^2, \dots, g_{n_2}^+ - s_{n_2}^2, v_1 s_1, \dots, v_{n_2} s_{n_2}) \subset \mathbb{R}[\mathbf{z}]$$

where $F_i = \frac{\partial f}{\partial x_i} - \sum_{j=1}^{n_1} u_j \frac{\partial g_j^0}{\partial x_i} - \sum_{j=1}^{n_2} v_j \frac{\partial g_j^+}{\partial x_i}$. The gradient variety is $V_{grad} = \mathcal{V}(I_{grad})$ and we denote $V_{grad}^{\mathbf{x}} = \overline{\pi^{\mathbf{x}}(V_{grad})}$.

Definition 3.2. *For any $F \in \mathbb{R}[\mathbf{z}]$, the values of F at the (resp. real) points of $\mathcal{V}(\nabla F) = V_{grad}$ are called the (resp. real) critical values of F .*

We easily check the following property:

Lemma 3.3. $F|_{V_{grad}} = f|_{V_{grad}}$.

Thus minimizing F on V_{grad} is the same as minimizing f on V_{grad} , that is computing the minimal critical value of F .

3.2. The Karush-Kuhn-Tucker variety. In the case of a constrained problem, one usually introduce the Karush-Kuhn-Tucker (KKT) constraints:

Definition 3.4. *A point \mathbf{x}^* is called a KKT point if there exists $u_1, \dots, u_{n_1}, v_1, \dots, v_{n_2} \in \mathbb{R}$ s.t.*

$$\nabla f(\mathbf{x}^*) - \sum_{i=1}^{n_1} u_i \nabla g_i^0(\mathbf{x}^*) - \sum_{j=0}^{n_2} v_j \nabla g_j^+(\mathbf{x}^*) = 0, \quad g_i^0(\mathbf{x}^*) = 0, \quad v_j g_j^+(\mathbf{x}^*) = 0.$$

The corresponding minimization problem is the following:

$$(4) \quad \begin{aligned} & \inf_{(\mathbf{x}, \mathbf{u}, \mathbf{v}) \in \mathbb{R}^{n+n_1 n_2}} f(\mathbf{x}) \\ & \text{s.t.} \quad F_1 = \dots = F_n = 0 \\ & \quad \quad g_1^0 = \dots = g_{n_1}^0 = 0 \\ & \quad \quad v_1 g_1^+ = \dots = v_{n_2} g_{n_2}^+ = 0 \\ & \quad \quad g_1^+ \geq 0, \dots, g_{n_2}^+ \geq 0 \end{aligned}$$

where $F_i = \frac{\partial f}{\partial x_i} - \sum_{j=1}^{n_1} u_j \frac{\partial g_j^0}{\partial x_i} - \sum_{j=1}^{n_2} v_j \frac{\partial g_j^+}{\partial x_i}$.

This leads to the following definitions:

Definition 3.5. *The Karush-Kuhn-Tucker (KKT) ideal associated to Problem (1) is*

$$(5) \quad I_{KKT} = (F_1, \dots, F_n, g_1^0, \dots, g_{n_1}^0, v_1 g_1^+, \dots, v_{n_2} g_{n_2}^+) \subset \mathbb{R}[\mathbf{y}].$$

The KKT variety is $V_{KKT} = \mathcal{V}(I_{KKT}) \subset \mathbb{C}^n \times \mathbb{C}^{n_1+n_2}$ and the real KKT variety is $V_{KKT}^{\mathbb{R}} = V_{KKT} \cap (\mathbb{R}^n \times \mathbb{R}^{n_1+n_2})$. Its projection on \mathbf{x} is $V_{KKT}^{\mathbf{x}} = \overline{\pi^{\mathbf{x}}(V_{KKT})}$, where $\pi^{\mathbf{x}}$ is the projection of $\mathbb{C}^n \times \mathbb{C}^{n_1+n_2}$ onto \mathbb{C}^n .

The set of KKT points of S is denoted S_{KKT} and a KKT-minimizer of f on S is a point $\mathbf{x}^* \in S_{KKT}$ such that $f(\mathbf{x}^*) = \min_{\mathbf{x} \in S_{KKT}} f(\mathbf{x})$.

Notice that $V_{KKT}^{\mathbf{x}, \mathbb{R}} = \overline{\pi^{\mathbf{x}}(V_{KKT})}^{\mathbb{R}} = \overline{\pi^{\mathbf{x}}(V_{KKT}^{\mathbb{R}})}$, since any linear dependency relation between real vectors can be realized with real coefficients.

The KKT ideal is related to the gradient ideal as follows:

Proposition 3.6. $I_{KKT} = I_{grad} \cap \mathbb{R}[\mathbf{y}]$.

Proof. As $s_i(s_i v_i) + v_i(g_i^+ - s_i^2) = v_i g_i^+ \forall i = 1, \dots, n_2$, we have $I_{KKT} \subset I_{grad} \cap \mathbb{R}[\mathbf{y}]$.

In order to prove the equality, we use the property that if K is a Groebner basis of I_{grad} for an elimination ordering such that $\mathbf{s} \gg \mathbf{x}, \mathbf{u}, \mathbf{v}$ then $K \cap \mathbb{R}[\mathbf{y}]$ is the Groebner basis of $I_{grad} \cap \mathbb{R}[\mathbf{y}]$ (see [3]). Notice that $s_i(s_i v_i) + v_i(g_i^+ - s_i^2) = v_i g_i^+$ ($i = 1, \dots, n_2$) are the only S-polynomials involving the variables s_1, \dots, s_{n_2} which may have a non-trivial reduction. Thus $K \cap \mathbb{R}[\mathbf{y}]$ is also the Groebner basis of $F_1, \dots, F_n, g_1^0, \dots, g_{n_1}^0, v_1 g_1^+, \dots, v_{n_2} g_{n_2}^+$ and we have $(K) \cap \mathbb{R}[\mathbf{y}] = I_{grad} \cap \mathbb{R}[\mathbf{y}] = I_{KKT}$. \square

The KKT points on S are related to the real points of the gradient variety as follows:

Lemma 3.7. $S_{KKT} = \pi^{\mathbf{x}}(V_{grad}^{\mathbb{R}}) = V_{grad}^{\mathbf{x}, \mathbb{R}} \cap \mathcal{S}^+(\mathbf{g})$.

Proof. A real point $\mathbf{y} = (\mathbf{x}, \mathbf{u}, \mathbf{v})$ of $V_{KKT}^{\mathbb{R}}$ lifts to a point $\mathbf{z} = (\mathbf{x}, \mathbf{u}, \mathbf{v}, \mathbf{s})$ in $V_{grad}^{\mathbb{R}}$, if and only if, $g_i^+(\mathbf{x}) \geq 0$ for $i = 1, \dots, n_2$. This implies that $V_{KKT}^{\mathbb{R}} = \pi^{\mathbf{y}}(V_{grad}^{\mathbb{R}}) \cap \mathcal{S}^+(\mathbf{g})$, which gives by projection the equalities $S_{KKT} = \pi^{\mathbf{x}}(V_{grad}^{\mathbb{R}}) \cap \mathcal{S}^+(\mathbf{g}) = \pi^{\mathbf{x}}(V_{grad}^{\mathbb{R}})$ since a point \mathbf{x} of $V_{grad}^{\mathbb{R}}$ satisfies $\mathbf{g}_j^+(\mathbf{x}) \geq 0$ for $j \in [1, n_2]$. \square

This shows that if a minimizer point of f on S is a KKT point, then it is the projection of a real critical point of F .

3.3. The Fritz John variety. A minimizer of f on S is not necessarily a KKT point. More general conditions that are satisfied by minimizers were given by F. John for polynomial non-negativity constraints and further refined for general polynomial constraints [11, 21]. To describe these conditions, we introduce a new variable u_0 and denote by \mathbf{y}' the set of variables $\mathbf{y}' = (\mathbf{x}, u_0, \mathbf{u}, \mathbf{v})$. Let $F_i^{u_0} = u_0 \frac{\partial f}{\partial x_i} - \sum_{j=1}^{n_1} u_j \frac{\partial g_j^0}{\partial x_i} - \sum_{j=1}^{n_2} v_j \frac{\partial g_j^+}{\partial x_i}$.

Definition 3.8. For any $\gamma \subset [1, n_1]$, let

$$(6) \quad I_{FJ}^{\gamma} = (F_1^{u_0}, \dots, F_n^{u_0}, g_1^0, \dots, g_{n_1}^0, v_1 g_1^+, \dots, v_{n_2} g_{n_2}^+, u_i, i \notin \gamma) \subset \mathbb{R}[\mathbf{y}'].$$

For $m \in \mathbb{N}$, the m^{th} Fritz-John (FJ) ideal associated to Problem (1) is

$$(7) \quad I_{FJ}^m = \bigcap_{|\gamma|=m} I_{FJ}^{\gamma}.$$

Let $V_{FJ}^{\gamma} = \mathcal{V}(I_{FJ}^{\gamma}) \subset \mathbb{C}^n \times \mathbb{P}^{n_1+n_2}$. The m^{th} FJ variety is $V_{FJ}^m = \mathcal{V}(I_{FJ}^m) = \bigcup_{|\gamma|=m} V_{FJ}^{\gamma}$, and the real FJ variety is $V_{FJ}^{m, \mathbb{R}} = V_{FJ}^m \cap \mathbb{R}^n \times \mathbb{R}\mathbb{P}^{n_1+n_2}$. Its projection on \mathbf{x} is $V_{FJ}^{m, \mathbf{x}} = \pi^{\mathbf{x}}(V_{FJ}^m) = \overline{\pi^{\mathbf{x}}(V_{FJ}^m)}$. When $m = \max_{\mathbf{x} \in S} \text{rank}([\nabla g_1^0(\mathbf{x}), \dots, \nabla g_{n_1}^0(\mathbf{x})])$, the m^{th} FJ variety is denoted V_{FJ} .

Notice that this definition slightly differs from the classical one [11, 14, 21], which does not provide any information when the gradient vectors $\nabla g_i^0(\mathbf{x}), i = 1 \dots n_1$ are linearly dependent on S .

Proposition 3.9. Any minimizer \mathbf{x}^* of f on S is the projection of a real point of $V_{FJ}^{\mathbb{R}}$.

Proof. The proof is similar to Theorem 4.3.2 of [21]. At a minimizer point \mathbf{x}^* (if it exists) We consider a maximal set of linearly independent gradients $\nabla g_j^0(\mathbf{x}^*)$ for $j \in \gamma$ (with $|\gamma| \leq m$) and apply the same proof as [21][Theorem 4.3.2]. This shows that $\mathbf{x}^* \in V_{FJ}^{\gamma, \mathbb{R}} \subset V_{FJ}^{\mathbb{R}}$. \square

Definition 3.10. We denote by $V_{sing} = V_{FJ} \cap \mathcal{V}(u_0)$ the intersection of V_{FJ} with the hyperplane $u_0 = 0$.

We easily check that the ‘‘affine part’’ of V_{FJ} corresponding to $u_0 \neq 0$ is the variety V_{KKT} . Thus, we have the decomposition

$$V_{FJ} = V_{sing} \cup V_{KKT},$$

Its projection on \mathbb{C}^n decomposes as

$$(8) \quad V_{FJ}^{\mathbf{x}} = V_{sing}^{\mathbf{x}} \cup V_{KKT}^{\mathbf{x}}.$$

Let us describe more precisely the projection $V_{FJ}^{\mathbf{x}}$ onto \mathbb{C}^n . For $\nu = \{j_1, \dots, j_k\} \subset [1, n_2]$, we define

$$\begin{aligned} A_\nu &= [\nabla f, \nabla g_1^0, \dots, \nabla g_{n_1}^0, \nabla g_{j_1}^+, \dots, \nabla g_{j_k}^+] \\ V_\nu &= \{\mathbf{x} \in \mathbb{C}^n \mid g_1^0(\mathbf{x}) = 0, i = 1 \dots n_1, g_j^+(\mathbf{x}) = 0, j \in \nu, \text{rank}(A_\nu) \leq m + |\nu|\}. \end{aligned}$$

Let $\Delta_1^\nu, \dots, \Delta_{l_\nu}^\nu$ be polynomials defining the variety $\{\mathbf{x} \in \mathbb{C}^n \mid \text{rank}(A_\nu) \leq m + |\nu|\}$. If $n > m + |\nu|$, these polynomials can be chosen as linear combinations of $(m + |\nu| + 1)$ -minors of the matrix A_ν , as described in [2, 25]. If $n \leq m + |\nu|$, we take $l_\nu = 0$, $\Delta_i^\nu = 0$. Let Γ_{FJ} be the union of \mathbf{g}^0 and the set of polynomials

$$(9) \quad g_{\nu,i} := \Delta_i^\nu \prod_{j \notin \nu} g_j^+,$$

for $i = 1, \dots, l_\nu, \nu \subset [0, n_2]$.

Lemma 3.11. $V_{FJ}^{\mathbf{x}} = \cup_{\nu \subset [0, n_2]} V_\nu = \mathcal{V}(\Gamma_{FJ})$.

Proof. For any $\mathbf{x} \in \mathbb{C}^n$, let $\nu(\mathbf{x}) = \{j \in [1, n_2] \mid g_j^+(\mathbf{x}) = 0\}$.

Let \mathbf{y}' be a point of V_{FJ} , \mathbf{x} its projection on \mathbb{C}^n and $\nu(\mathbf{x}) = \nu = \{j_1, \dots, j_k\}$. We have $g_j^+(\mathbf{x}) \neq 0$, $v_j = 0$ for $j \notin \nu$ and $\Delta_i^\nu = 0$ for $i = 1, \dots, l_\nu$. This implies that $\text{rank}(A_\nu(\mathbf{x})) \leq m + |\nu|$ and there exists $(u_0, u_1, \dots, u_{n_1}, v_1, \dots, v_{n_2}) \neq 0$ and $\gamma \subset [1, n_1]$ of size $|\gamma| \leq m$ such that

$$u_0 \nabla f + u_1 \nabla g_1^0 + \dots + u_{n_1} \nabla g_{n_1}^0 + v_1 \nabla g_{j_1}^+ + \dots + v_{n_2} \nabla g_{j_k}^+ = 0,$$

with $u_i = 0$, $i \notin \gamma \subset [1, n_1]$. Therefore $\mathbf{x} \in \pi^{\mathbf{x}}(V_{FJ})$, which proves that $\mathcal{V}(\mathbf{g}^0, g_{\nu,i}, \nu \subset [0, n_2], i = 1 \dots l_\nu) \subset \pi^{\mathbf{x}}(V_{FJ})$.

Conversely, if $\mathbf{x} \in \pi^{\mathbf{x}}(V_{FJ})$ then $\mathbf{x} \in V_{\nu(\mathbf{x})} \subset \cup_\nu V_\nu$ which is defined by the polynomials $g_1^0, \dots, g_{n_1}^0$ and $g_{\nu,i} := \Delta_i^\nu \prod_{j \notin \nu} g_j^+$, for $i = 1, \dots, l_\nu, \nu \subset [0, n_2]$. \square

Remark 3.12. The real variety $\pi^{\mathbf{x}}(V_{FJ}^{\mathbb{R}}) = V_{FJ}^{\mathbf{x}} \cap \mathbb{R}^n$ can also be defined by \mathbf{g}^0 and the set Φ_{FJ} of polynomials

$$(10) \quad g_\nu := \Delta^\nu \prod_{j \notin \nu} g_j^+ \text{ where } \Delta^\nu = \det(A_\nu A_\nu^T),$$

for $\nu \subset [0, n_2]$ and $n > m + |\nu|$, as described in [9].

Similarly the projection $V_{sing}^{\mathbf{x}}$ onto \mathbb{C}^n can be described as follows. For $\nu = \{j_1, \dots, j_k\} \subset [1, n_2]$,

$$\begin{aligned} B_\nu &= [\nabla g_1^0, \dots, \nabla g_{n_1}^0, \nabla g_{j_1}^+, \dots, \nabla g_{j_k}^+] \\ W_\nu &= \{\mathbf{x} \in \mathbb{C}^n \mid g_1^0(\mathbf{x}) = 0, i = 1 \dots n_1, g_j^+(\mathbf{x}) = 0, j \in \nu, \text{rank}(B_\nu) \leq m + |\nu| - 1\}. \end{aligned}$$

Let $\Theta_1^\nu, \dots, \Theta_{l_\nu}^\nu$ be polynomials defining the variety $\{\mathbf{x} \in \mathbb{C}^n \mid \text{rank}(B_\nu) \leq m + |\nu| - 1\}$ and let Γ_{sing} be the union of \mathbf{g}^0 and the set of polynomials

$$(11) \quad \sigma_{\nu,i} := \Theta_i^\nu \prod_{j \notin \nu} g_j^+,$$

for $\nu \subset [0, n_2], i = 1 \dots l_\nu$.

We similar arguments, we prove the following

Lemma 3.13. $V_{sing}^{\mathbf{x}} = \cup_{\nu \subset [0, n_2]} W_\nu = \mathcal{V}(\Gamma_{sing})$.

3.4. The minimizer variety. By the decomposition (8) and Proposition 3.9, we know that the minimizer points of f on S are in

$$(12) \quad S_{FJ} = S_{KKT} \cup S_{sing}$$

where $S_{FJ} = \pi^{\mathbf{x}}(V_{FJ}^{\mathbb{R}}) \cap S = \pi^{\mathbf{x}}(V_{FJ}^{\mathbb{R}}) \cap \mathcal{S}^+(\mathbf{g})$, $S_{KKT} = \pi^{\mathbf{x}}(V_{KKT}^{\mathbb{R}}) \cap S = \pi^{\mathbf{x}}(V_{KKT}^{\mathbb{R}}) \cap \mathcal{S}^+(\mathbf{g})$, $S_{sing} = \pi^{\mathbf{x}}(V_{sing}^{\mathbb{R}}) \cap S = \pi^{\mathbf{x}}(V_{sing}^{\mathbb{R}}) \cap \mathcal{S}^+(\mathbf{g})$. Therefore, we can decompose the initial optimization problem (1) into two subproblems:

- (1) find the infimum of f on S_{KKT} ;
- (2) find the infimum of f on S_{sing} ;

and take the least of these two infima. Since the second problem is of the same type as (1) but with the additional constraints $\sigma_{\nu,i} = 0$ described in (11), we analyse only the first subproblem. The approach developed for this first sub-problem is applied recursively to the second subproblem, in order to obtain the solution of Problem (1).

Definition 3.14. We define the KKT-minimizer set and ideal of f on S as:

$$\begin{aligned} S_{min} &= \{\mathbf{x}^* \in S_{KKT} \text{ s.t. } \forall \mathbf{x} \in S_{KKT}, f(\mathbf{x}^*) \leq f(\mathbf{x})\} \\ I_{min} &= \mathcal{I}(S_{min}) \subset \mathbb{R}[\mathbf{x}]. \end{aligned}$$

A point \mathbf{x}^* in S_{min} is called a KKT-minimizer. Notice that $I_{KKT} \subset I_{min}$ and that I_{min} is a real radical ideal.

We have $I_{min} \neq (1)$, if and only if, the KKT-minimum f^* is reached in S_{KKT} .

If $n_1 = n_2 = 0$, I_{min} is the vanishing ideal of the *critical points* \mathbf{x}^* of f (satisfying $\nabla f(\mathbf{x}^*) = 0$) where $f(\mathbf{x}^*)$ reaches its minimal critical value.

Remark 3.15. If we take $f = 0$ in the minimization problem (1), then all the points of S are KKT-minimizers and $I_{min} = \mathcal{I}(S) = \mathfrak{s}^+ \sqrt{\mathbf{g}^0}$. Moreover, $I_{KKT} \cap \mathbb{R}[\mathbf{x}] = (g_1^0, \dots, g_{n_1}^0) = (\mathbf{g}^0)$ since $F_1, \dots, F_n, v_1 g_1^+, \dots, v_{n_2} g_{n_2}^+$ are homogeneous of degree 1 in the variables \mathbf{u}, \mathbf{v} .

4. REPRESENTATION OF POSITIVE POLYNOMIALS

In this section, we analyse the decomposition of polynomials as sum of squares modulo the gradient ideal. Hereafter, J_{grad} is an ideal of $\mathbb{R}[\mathbf{z}]$ such that $\mathcal{V}(J_{grad}) = V_{grad}$ and C is a set of constraints in $\mathbb{R}[\mathbf{x}]$ such that $\mathcal{S}^+(C) = \mathcal{S}^+(\mathbf{g})$.

The first steps consists in decomposing V_{grad} in components on which f has a constant value. We recall here a result, which also appears (with slightly different hypotheses) in [28, Lemma 3.3]³.

Lemma 4.1. Let $f \in \mathbb{R}[\mathbf{x}]$ and let V be an irreducible subvariety contained in $\mathcal{V}^{\mathbb{C}}(\nabla f)$. Then $f(x)$ is constant on V .

Proof. If V is irreducible in the Zariski topology induced from $\mathbb{C}[\mathbf{x}]$, then it is connected in the strong topology on \mathbb{C}^n and even piecewise smoothly path-connected [34]. Let x, y be two arbitrary points of V . There exists a piecewise smooth path $\varphi(t)$ ($0 \leq t \leq 1$) lying inside V such that $x = \varphi(0)$ and $y = \varphi(1)$. Without loss of generality, we can assume

³In its proof, the Mean Value Theorem is applied for a complex valued function, which is not valid. We correct the problem in the proof of Lemma 4.1.

that φ is smooth between x and y in order to prove that $f(x) = f(y)$. By the Mean Value Theorem, it holds that for some $t_1 \in (0, 1)$

$$\operatorname{Re}(f(y) - f(x)) = \operatorname{Re}(f(\varphi(t)))'(t_1) = \operatorname{Re}((\nabla f(\varphi(t_1)) * \varphi'(t_1))) = 0$$

since ∇f vanishes on V . Then $\operatorname{Re}(f(y)) = \operatorname{Re}(f(x))$. We have the same result for the imaginary part: for some $t_2 \in (0, 1)$

$$\operatorname{Im}(f(y)) - \operatorname{Im}(f(x)) = \operatorname{Im}(f(\varphi(t)))'(t_2) = \operatorname{Im}((\nabla f(\varphi(t_2)) * \varphi'(t_2))) = 0$$

since ∇f vanishes on V . Then $\operatorname{Im}(f(y)) = \operatorname{Im}(f(x))$. We conclude that $f(y) = f(x)$ and hence f is constant on V . \square

Lemma 4.2. *The ideal J_{grad} can be decomposed as $J_{grad} = J_0 \cap J_1 \cap \dots \cap J_s$ with $V_i = \mathcal{V}(J_i)$ and $W_i = \overline{\pi^{\mathbf{x}}(V_i)}$ where $\pi^{\mathbf{x}}(V_i)$ is the projection of V_i on \mathbb{C}^n such that*

- $f(V_j) = f_j \in \mathbb{C}$, $f_i \neq f_j$ if $i \neq j$,
- $W_i^{\mathbb{R}} \cap \mathcal{S}^+(C) \neq \emptyset$ for $i = 0, \dots, r$,
- $W_i^{\mathbb{R}} \cap \mathcal{S}^+(C) = \emptyset$ for $i = r+1, \dots, s$,
- $f_0 < \dots < f_r$.

Proof. Consider a minimal primary decomposition of J_{grad} :

$$J_{grad} = Q_0 \cap \dots \cap Q_s,$$

where Q_i is a primary component, and $\mathcal{V}(Q_i)$ is an irreducible variety in $\mathbb{C}^{n+n_1+2n_2}$ included in V_{grad} . By Lemma 4.1, f is constant on $\mathcal{V}(Q_i)$. By Lemma 3.3, it coincides with f on each variety $\mathcal{V}(Q_i)$. We group the primary components Q_i according to the values f_0, \dots, f_s of f on these components, into J_0, \dots, J_s so that $f(\mathcal{V}(J_j)) = f_j$ with $f_i \neq f_j$ if $i \neq j$.

We can number them so that $\overline{\pi^{\mathbf{x}}(V_i)^{\mathbb{R}}} \cap \mathcal{S}^+(C)$ is empty for $i = r+1, \dots, s$ and contains a real point \mathbf{x}_i for $i = 0, \dots, r$. Notice that such a point \mathbf{x}_i is in \mathcal{S} , since it satisfies $g^0(\mathbf{x}_i) = 0 \forall g^0 \in C^0$ and $g^+(\mathbf{x}_i) \geq 0 \forall g^+ \in C^+$. As it is the limit of the projection of points in $\mathcal{V}(J_i)$ on which f is constant, we have $f_i = f(\mathbf{x}_i) \in \mathbb{R}$ for $i = 0, \dots, r$. We can then order J_0, \dots, J_r so that $f_0 < \dots < f_r$. \square

Remark 4.3. *If the minimum of f on S is reached at a KKT-point, then we have $f_0 = \min_{\mathbf{x} \in S} f(\mathbf{x})$.*

Remark 4.4. *If $V_{grad}^{\mathbb{R}} = \emptyset$, then for all $i = 0, \dots, s$, $W_i^{\mathbb{R}} \cap \mathcal{S}^+(C) = \emptyset$ and by convention, we take $r = -1$.*

Lemma 4.5. *There exist $p_0, \dots, p_s \in \mathbb{C}[\mathbf{x}]$ such that*

- $\sum_{i=0}^s p_i = 1 \pmod{J_{grad}}$,
- $p_i \in \bigcap_{j \neq i} J_j$,
- $p_i \in \mathbb{R}[\mathbf{x}]$ for $i = 0, \dots, r$.

Proof. Let $(L_i)_{i=0, \dots, s}$ be the univariate Lagrange interpolation polynomials at the values $f_0, \dots, f_s \in \mathbb{C}$ and let $q_i(\mathbf{x}) = L_i(f(\mathbf{x}))$.

The polynomials q_i are constructed so that

- $q_i(V_j) = 0$ if $j \neq i$,
- $q_i(V_i) = 1$,

where $V_i = \mathcal{V}(J_i)$. As the set $\{f_{r+1}, \dots, f_s\}$ is stable by conjugation and $f_0, \dots, f_r \in \mathbb{R}$, by construction of the Lagrange interpolation polynomials we deduce that $q_0, \dots, q_r \in \mathbb{R}[\mathbf{x}]$.

By Hilbert's Nullstellensatz, there exists $N \in \mathbb{N}$ such that $q_i^N \in \bigcap_{j \neq i} J_j$. As $\sum_{j=0}^s q_j^N = 1$ on V_{grad} and $q_i^N q_j^N = 0 \pmod{\bigcap_i J_i = J_{grad}}$ for $i \neq j$, we deduce that there exists $N' \in \mathbb{N}$ such that

$$\begin{aligned} 0 &= \left(1 - \sum_{j=0}^s q_j^N\right)^{N'} \pmod{J_{grad}} \\ &= 1 - \sum_{j=0}^s (1 - (1 - q_j^N)^{N'}) \pmod{J_{grad}}. \end{aligned}$$

As the polynomial $p_i = 1 - (1 - q_j^N)^{N'} \in \mathbb{C}[\mathbf{x}]$ is divisible by q_j^N , it belongs to $\bigcap_{j \neq i} J_j$. Since $q_j \in \mathbb{R}[\mathbf{x}]$ for $j = 0, \dots, r$, we have $p_j \in \mathbb{R}[\mathbf{x}]$ for $j = 0, \dots, r$, which ends the proof of this lemma. \square

Lemma 4.6. $-1 \in \mathcal{P}^+(C) + (\bigcap_{i>r} J_i^{\mathbf{x}})$.

Proof. As $\bigcup_{i>r} \overline{\pi^{\mathbf{x}}(V_i)}^{\mathbb{R}} \cap \mathcal{S}^+(C) = \mathcal{V}^{\mathbb{R}}(\bigcap_{i>r} J_i \cap \mathbb{R}[\mathbf{x}]) \cap \mathcal{S}^+(C) = \mathcal{V}^{\mathbb{R}}(\bigcap_{i>r} J_i^{\mathbf{x}}) \cap \mathcal{S}^+(C) = \emptyset$, we have $\mathcal{I}(\mathcal{V}^{\mathbb{R}}(\bigcap_{i>r} J_i^{\mathbf{x}}) \cap \mathcal{S}^+(C)) = \mathbb{R}[\mathbf{x}] \ni 1$ and by the Positivstellensatz (Theorem 2.6 (iii)),

$$-1 \in \mathcal{P}^+(C) + \left(\bigcap_{i>r} J_i^{\mathbf{x}}\right).$$

\square

Corollary 4.7. If $S_{min} = \emptyset$, then $-1 \in \mathcal{P}^+(C) + J_{grad}^{\mathbf{x}}$.

Proof. If $S_{min} = \emptyset$, then f has no real KKT critical value on $S(C)$ and $r = -1$. Lemma 4.6 implies that $-1 \in \mathcal{P}^+(C) + (\bigcap_{i=0}^s J_i^{\mathbf{x}}) = \mathcal{P}^+(C) + J_{grad}^{\mathbf{x}}$. \square

In this case, $\forall p \in \mathbb{R}[\mathbf{x}]$, $p = \frac{1}{4}((p+1)^2 - (p-1)^2) \in \mathcal{P}^+(C) + J_{grad}^{\mathbf{x}}$. If C^0 is chosen such that $V(C^0) \subset V_{grad}^{\mathbf{x}}$ then $S_{min} = \emptyset$ if and only if $-1 \in \mathcal{P}(C)$.

We recall another useful result on the representation of positive polynomials (see for instance [5]):

Lemma 4.8. Let $J \subset \mathbb{R}[\mathbf{z}]$ and $V = \mathcal{V}(J)$ such that $f(V) = f^*$ with $f^* \in \mathbb{R}^+$. There exists $t \in \mathbb{N}$, s.t. $\forall \epsilon > 0$, $\exists q \in \mathbb{R}[\mathbf{x}]$ with $\deg(q) \leq t$ and $f + \epsilon = q^2 \pmod{J}$.

Proof. We know that $\frac{f+\epsilon}{f^*+\epsilon} - 1$ vanishes on V . By Hilbert's Nullstellensatz $(\frac{f+\epsilon}{f^*+\epsilon} - 1)^l \in J$ for some $l \in \mathbb{N}$. From the binomial theorem, it follows that

$$\left(1 + \left(\frac{f+\epsilon}{f^*+\epsilon} - 1\right)\right)^{1/2} \equiv \sum_i^{l-1} \binom{1/2}{i} \left(\frac{f+\epsilon}{f^*+\epsilon} - 1\right)^i \stackrel{def}{=} \frac{q}{\sqrt{f^*+\epsilon}} \pmod{J}$$

Then $f + \epsilon = q^2 \pmod{J}$. \square

In particular, if $f^* > 0$ this lemma implies that $f = (f - \frac{1}{2}f^*) + \frac{1}{2}f^* = q^2 \pmod{J}$ for some $q \in \mathbb{R}[\mathbf{x}]$.

Theorem 4.9. Let $C \subset \mathbb{R}[\mathbf{x}]$ be a set of constraints such that $\mathcal{S}^+(C) = \mathcal{S}^+(\mathbf{g})$, let $f \in \mathbb{R}[\mathbf{x}]$, let $f_0 < \dots < f_r$ be the real KKT critical values of f on S and let p_0, \dots, p_r be the associated polynomials defined in Lemma 4.5.

- (1) $f - \sum_{i=0}^r f_i p_i^2 \in \mathcal{P}^+(C) + \sqrt{J_{grad}^{\mathbf{x}}}$.
- (2) If $f \geq 0$ on S_{KKT} , then $f \in \mathcal{P}^+(C) + \sqrt{J_{grad}^{\mathbf{x}}}$.

(3) If $f > 0$ on S_{KKT} , then $f \in \mathcal{P}^+(C) + J_{grad}^{\mathbf{x}}$.

Proof. By Lemma 4.5, we have

$$1 = \left(\sum_{i=0}^s p_i \right)^2 = \sum_{i=0}^s p_i^2 \pmod{J_{grad}}.$$

Thus $f = \sum_{i=0}^s f p_i^2 \pmod{J_{grad}}$.

By Lemma 4.6, $-1 \in \mathcal{P}^+(C) + (\bigcap_{j>r} J_j^{\mathbf{x}})$ so that $f = \frac{1}{4}((f+1)^2 - (f-1)^2) \in \mathcal{P}^+(C) + \bigcap_{j>r} J_j^{\mathbf{x}}$ and

$$(13) \quad \sum_{i>r} f p_i^2 \in \mathcal{P}^+(C) + \bigcap_{j=0}^s J_j^{\mathbf{x}} = \mathcal{P}^+(C) + J_{grad}^{\mathbf{x}}.$$

As the polynomial $(f - f_i) p_i^2$ vanishes on V_{grad} , we deduce that

$$f = \sum_{i=0}^r f_i p_i^2 + \sum_{i=r+1}^s f p_i^2 + \sqrt{J_{grad}^{\mathbf{x}}} = \sum_{i=0}^r f_i p_i^2 + \mathcal{P}^+(C) + \sqrt{J_{grad}^{\mathbf{x}}},$$

which proves the first point.

If $f \geq 0$ on S_{KKT} , then $f_i \geq 0$ for $i = 0, \dots, r$ and $\sum_{i=0}^r f_i p_i^2 \in \mathcal{P}^+(C)$ so that

$$f \in \mathcal{P}^+(C) + \sqrt{J_{grad}^{\mathbf{x}}},$$

which proves the second point.

If $f > 0$ on S_{KKT} by Lemma 4.8, we have $f p_i^2 = q_i^2 \pmod{J_{grad}^{\mathbf{x}}}$ with $q_i \in \mathbb{R}[\mathbf{x}]$, which shows that

$$\sum_{i=0}^r f_i p_i^2 = \sum_{i=0}^r q_i^2 \pmod{J_{grad}^{\mathbf{x}}}$$

Therefore, $\sum_{i=0}^r f_i p_i^2 \in \mathcal{P}^+(C) + J_{grad}^{\mathbf{x}}$ and $f \in \mathcal{P}^+(C) + J_{grad}^{\mathbf{x}}$ by (13), which proves the third point. \square

This theorem involves only polynomials in $\mathbb{R}[\mathbf{x}]$ and the points (2) and (3) generalize results of [5] on the representation of positive polynomials.

Let us give now a refinement of Theorem 4.9 with a control of the degrees of the polynomials involved in the representation of f as an element of $\mathcal{P}^+(C) + J_{grad}^{\mathbf{x}}$.

Theorem 4.10. *Let $C \subset \mathbb{R}[\mathbf{x}]$ be a set of constraints such that $\mathcal{V}(C^0) \subset V_{grad}^{\mathbf{x}}$ and $\mathcal{S}^+(C) = \mathcal{S}^+(\mathbf{g})$. If $f \geq 0$ on S_{KKT} , then there exists t_0 such that $\forall \epsilon > 0$,*

$$f + \epsilon \in \mathcal{P}_{t_0}(C).$$

Proof. Let $J_{grad} = (C^0) \cap I_{grad} \subset \mathbb{R}[\mathbf{z}]$, so that $\mathcal{V}(J_{grad}) = V_{grad}$ since $\mathcal{V}(C^0) \subset V_{grad}^{\mathbf{x}}$. Using the decomposition (13) obtained in the proof of Theorem 4.9, we can choose $t'_0 \in \mathbb{N}$ and $t_0 \geq t'_0 \in \mathbb{N}$ big enough such that $\deg(p_i) \leq t_0/2$ and

$$\sum_{i>r} f p_i^2 \in \mathcal{P}_{t'_0}^+(C) + J_{grad} \cap \mathbb{R}[\mathbf{x}]_{t'_0} \subset \mathcal{P}_{t_0}(C),$$

since $J_{grad}^{\mathbf{x}} = (C^0) \cap I_{grad}^{\mathbf{x}} \subset (C^0)$. Then $\forall \epsilon > 0$,

$$(14) \quad \sum_{i>r} (f + \epsilon) p_i^2 = \sum_{i>r} f p_i^2 + \sum_{i>r} \epsilon p_i^2 \in \mathcal{P}_{t_0}(C).$$

As $\forall \epsilon > 0$, $f + \epsilon > 0$ on S_{KKT} , i.e., $f_i + \epsilon > 0$ for $i = 0, \dots, r$, we deduce from Lemma 4.8 that if t_0 is big enough, we have

$$(15) \quad (f + \epsilon) p_i^2 = q_i^2 \pmod{C_{\langle t_0 \rangle}^0 \cap \mathbb{R}[\mathbf{x}]}$$

with $\deg(q_i) \leq t_0/2$ for $i = 0, \dots, r$.

Since $1 - \sum_{i=0}^s p_i^2 = 0 \pmod{C^0}$, we can choose t_0 big enough so that

$$(16) \quad (f + \epsilon) - \sum_{i=0}^s (f + \epsilon) p_i^2 \in C_{\langle t_0 \rangle}^0 \cap \mathbb{R}[\mathbf{x}].$$

From Equations (14), (15), (16), we deduce that if $t_0 \in \mathbb{N}$ is big enough, $\forall \epsilon > 0$

$$f + \epsilon \in \mathcal{P}_{t_0}(C),$$

which concludes the proof of the theorem. \square

5. FINITE CONVERGENCE

In this section, we show that the sequence of relaxation problems attains its limit in a finite number of steps and that the minimizer ideal can be recovered from an optimal solution of the corresponding relaxation problem. We use the following notation:

- $f^* = \inf_{\mathbf{x} \in S_{KKT}} f(\mathbf{x})$
- $S_{min} = \{\mathbf{x}^* \in S_{KKT} \mid f(\mathbf{x}^*) = f^*\}$

We first show that $S_{min} = \emptyset$ can be detected from an adapted relaxation sequence:

Proposition 5.1. *Let $C = (C^0; C^+)$ be a set of constraints of $\mathbb{R}[\mathbf{x}]$, such that $S_{min} \subset \mathcal{S}(C)$ and $\mathcal{V}(C^0) \subset V_{KKT}^{\mathbf{x}}$ and $C^+ = \mathbf{g}^+$. Then $S_{min} = \emptyset$, if and only if, there $t_0 \in \mathbb{N}$ such that $\forall t \geq t_0$, $\mathcal{L}_t(C) = \emptyset$.*

Proof. Let $J_{grad} = (C^0) \cap I_{grad}$ and let C' be a set of constraints such that $(C'^0) = J_{grad} \cap \mathbb{R}[\mathbf{x}] = J_{grad}^{\mathbf{x}}$ and $C'^+ = \mathbf{g}^+$ be a finite set. By hypothesis, $\mathcal{V}(J_{grad}) = V_{grad}$. We deduce from Corollary 4.7 that if $S_{min} = \emptyset$, then

$$-1 \in \mathcal{P}^+(C') + (C'^0) \subset \mathcal{P}(C) = \cup_{t \in \mathbb{N}} \mathcal{P}_t(C).$$

Thus there exists t_0 such that $-1 \in \mathcal{P}_t(C)$ for $t \geq t_0$, which implies that $\mathcal{L}_t(C) = \emptyset$, since if there exists $\Lambda \in \mathcal{L}_t(C)$, then $\Lambda(1) = 1$ and $\Lambda(-1) \geq 0$.

Conversely, suppose that $S_{min} \neq \emptyset$ contains a point \mathbf{x}^* . As $S_{min} \subset \mathcal{S}(C)$, for all $t \in \mathbb{N}$ the evaluation $\underline{1}_{\mathbf{x}^*}$ at \mathbf{x}^* restricted to $\mathbb{R}[\mathbf{x}]_{2t}$ is an element of $\mathcal{L}_t(C) \neq \emptyset$. \square

This proposition gives a way to check whether $S_{min} = \emptyset$, using the relaxation sequence $\mathcal{L}_t(C)$. We are now going to analyse the case where f has KKT minimizers on S .

From now on, we assume that $S_{min} \neq \emptyset$.

First, we recall a property similar to [15, Claim 4.7]:

Proposition 5.2. *Let $C = (C^0; C^+)$ be a set of constraints of $\mathbb{R}[\mathbf{x}]$. There exists $t_0 \in \mathbb{N}$ such that $\forall t \geq t_0$, $\forall \Lambda \in \mathcal{L}_t(C)$, ${}^{c^+}\sqrt{C^0} \subset (\ker M_{\Lambda}^t)$.*

Proof. Let $C^0 = \{g_1, \dots, g_l\}$ and let q_1, \dots, q_k be generators of $J := {}^{c^+}\sqrt{C^0}$. By the Positivstellensatz, for $j \in 1, \dots, k$, there exist $m_j \in \mathbb{N}^*$ and polynomials $u_r^{(j)} \in \mathbb{R}[\mathbf{x}]$ and $\sigma_j \in \mathcal{P}^+(C)$ such that

$$q_j^{2m_j} + \sigma_j = \sum_{r=1}^l u_r^{(j)} g_r.$$

Let us take $t_0 \in \mathbb{N}$ big enough such that $u_r^{(j)} g_r \in C_{(t_0)}$ and $\sigma_j \in \mathcal{P}_{t_0}^+(C)$. Then for all $t \geq t_0$ and all $\Lambda \in \mathcal{L}_t(C)$, we have $\Lambda(u_r^{(j)} g_r) = 0$, $\Lambda(q_j^{2m_j}) \geq 0$, $\Lambda(\sigma_j) \geq 0$ and $\Lambda(q_j^{2m_j}) + \Lambda(\sigma_j) = 0$, which implies that $\Lambda(q_j^{2m_j}) = 0$ and $q_j \in \ker M_\Lambda^t$. This proves that $(q_1, \dots, q_l) = J \subset (\ker M_\Lambda^t)$. \square

Remark 5.3. *With the same arguments, we can show that for any $t' \in \mathbb{N}$, there exists $t'_0 \geq t'$ such that $\forall t \geq t'_0, \forall \Lambda \in \mathcal{L}_t(C)$,*

$$Q_{\langle t' \rangle} \subset \ker M_\Lambda^{t'},$$

where $Q = \{q_1, \dots, q_k\}$ generates $J = \sqrt[\mathbb{C}]{C^0}$.

The next result shows that in the sequence of optimization problems that we consider, the minimum of f on S_{KKT} is reached from some degree.

Theorem 5.4. *Let C be a set of constraints of $\mathbb{R}[\mathbf{x}]$ such that $S_{min} \subset \mathcal{S}(C) \subset V_{KKT}^{\mathbf{x}, \mathbb{R}}$. There exists $t_1 \geq 0$ such that $\forall t \geq t_1$,*

- (1) $f_{t,C}^\mu = f^*$ is reached for some $\Lambda^* \in \mathcal{L}_t(C)$,
- (2) $\forall \Lambda^* \in \mathcal{L}_t(C)$ with $\Lambda^*(f) = f_{t,C}^\mu = f^*$, we have $p_i \in \ker M_{\Lambda^*}^t, \forall i = 1, \dots, r$,
- (3) if $\mathcal{V}(C^0) \subset V_{KKT}^{\mathbf{x}}$ then $f_{t,C}^{sos} = f_{t,C}^\mu = f^*$.

Proof. By Theorem 4.9(1) applied to $f - f^*$, we can write

$$f - f^* \equiv \sum_{i=1}^r (f_i - f^*) p_i^2 + h + g.$$

with $h \in \mathcal{P}^+(C)$ and $g \in \sqrt{I_{grad}} \cap \mathbb{R}[\mathbf{x}] = \sqrt{I_{KKT}} \cap \mathbb{R}[\mathbf{x}] \subset \sqrt[\mathbb{R}]{I_{KKT}} \cap \mathbb{R}[\mathbf{x}]$ (by Proposition 3.6). Since $\mathcal{S}(C) \subset V_{KKT}^{\mathbf{x}, \mathbb{R}} = \pi^{\mathbf{x}}(V_{KKT}^{\mathbb{R}})$, we have $\sqrt[\mathbb{R}]{I_{KKT}} \cap \mathbb{R}[\mathbf{x}] \subset \mathcal{I}(\mathcal{S}(C)) = \sqrt[\mathbb{C}]{(C^0)}$ by the Positivstellensatz. We deduce that $g \in \sqrt[\mathbb{C}]{(C^0)}$. By proposition 5.2, there exists $t_1 \geq t_0$ such that for all $t \geq t_1$, for all $\Lambda \in \mathcal{L}_t(C)$, $\Lambda(g) = 0$, $\Lambda(h) \geq 0$.

Let us fix $t \geq t_1$ and $\Lambda^* \in \mathcal{L}_t(C)$ such that $\Lambda^*(f) = f_{t,C}^\mu$. Then

$$\Lambda^*(f - f^*) = \sum_{i=1}^r (f_i - f^*) \Lambda^*(p_i^2) + \Lambda^*(h).$$

As $f_i - f^* = f_i - f_0 > 0$, $\Lambda^*(p_i^2) \geq 0$ and $\Lambda^*(h) \geq 0$ ($h \in \mathcal{P}_t^+(C)$), we deduce that $\Lambda^*(f - f^*) = \Lambda^*(f) - f^* \geq 0$.

As $\emptyset \neq S_{min} \subset \mathcal{S}(C)$, we have $\Lambda^*(f) \leq f^*$ (by Remark 2.8), so that $\Lambda^*(f) = f_{t,C}^\mu = f^*$, which proves the first point. Hence for $i = 1, \dots, r$, $\Lambda^*(p_i^2) = 0$ and $p_i \in \ker M_{\Lambda^*}^t$, which proves the second point.

To prove that $f_{t,C}^{sos} = f^*$ when $\mathcal{V}(C^0) \subset V_{KKT}^{\mathbf{x}}$, we apply Theorem 4.10 to $f - f^*$ which is positive on S_{KKT} . Let us take $J_{grad} = (C^0) \cap I_{grad} \subset \mathbb{R}[\mathbf{z}]$. We denote by \tilde{C} the set of constraints such that \tilde{C}^0 is a finite family of generators of $J_{grad} \cap \mathbb{R}[\mathbf{x}]$ and $\tilde{C}^+ = C^+$.

By Theorem 4.10, there exists t_0 such that $\forall \epsilon > 0$,

$$f - f^* + \epsilon \in \mathcal{P}_{t_0}(\tilde{C}).$$

As $(\tilde{C}^0) = (C^0) \cap I_{grad} \subset (C^0)$, we can choose $t_1 \geq t_0$ such that $\tilde{C}_{\langle t_0 \rangle} \subset C_{\langle t_1 \rangle}$ and $\mathcal{P}_{t_0}(\tilde{C}) \subset \mathcal{P}_{t_1}(C)$.

Then $\forall t \geq t_1$, $f - f^* + \epsilon \in \mathcal{P}_t(C)$. Hence by maximality, $\forall \epsilon > 0, f^* - \epsilon \leq f_{t,C}^{sos}$. We deduce that $f^* \leq f_{t,C}^{sos}$, which implies that $f_{t,C}^{sos} = f_{t,C}^\mu = f^*$ and proves the third point. \square

As for the construction of generators of ${}^{c^+}\sqrt{I_{KKT}}$ (Proposition 5.2), we can construct generators of I_{min} from the kernel of a truncated Hankel operator associated to any linear form which minimizes f , using the following propositions:

Proposition 5.5. $I_{min} = (p_1, \dots, p_r) + {}^{c^+}\sqrt{I_{KKT}^x}$.

Proof. First of all, we proof that $I_{min}^z = (p_1, \dots, p_r) + {}^{c^+}\sqrt{I_{grad}} = (p_1, \dots, p_r) + \sqrt[\mathbb{R}]{I_{grad}}$. Using the decomposition of Lemma 4.2 and the polynomials p_i of Lemma 4.5, we have

$$V_{grad}^{\mathbb{R}} = (V_0 \cup V_1 \cup \dots \cup V_s) \cap \mathbb{R}^{n+n_1+2n_2} = V_0^{\mathbb{R}} \cup \dots \cup V_r^{\mathbb{R}},$$

By construction, $\mathcal{I}(V_0^{\mathbb{R}}) = I_{min}^z$, $p_i(V_0^{\mathbb{R}}) = 0$ for $i = 1, \dots, s$ and $p_i \in \mathbb{R}[\mathbf{x}]$ for $i = 0, \dots, r$. This implies that $p_i \in I_{min}^z$ for $i = 1, \dots, r$.

As $V_0^{\mathbb{R}} \subset V_{grad}^{\mathbb{R}}$, we also have ${}^{c^+}\sqrt{I_{grad}} \subset I_{min}^z$.

We have proved so far that $(p_1, \dots, p_r) + {}^{c^+}\sqrt{I_{grad}} \subset I_{min}^z$. In order to prove the reverse inclusion, we denote by q_1, \dots, q_m a family of generators of the ideal I_{min}^z . Take one of these generators q_j ($1 \leq j \leq m$). By construction, $q_j p_0(V_0^{\mathbb{R}}) = 0$ and $q_j p_0(V_i^{\mathbb{R}}) = 0$ for $i = 1, \dots, r$, which implies that $q_j p_0 \in {}^{c^+}\sqrt{I_{grad}}$.

By Lemma 4.5, we have the decomposition

$$q_j \equiv q_j(p_0 + p_1 + \dots + p_s) \pmod{I_{grad}} \subset {}^{c^+}\sqrt{I_{grad}}.$$

Moreover $(p_{r+1} + \dots + p_s) \in \mathbb{R}[\mathbf{z}]$ and vanishes on $V_k^{\mathbb{R}}$ for $k = 0, \dots, r$. Thus $(p_{r+1} + \dots + p_s) \in {}^{c^+}\sqrt{I_{grad}}$ and we deduce that $q_j \in (p_1, \dots, p_r) + {}^{c^+}\sqrt{I_{grad}}$. This proves the other inclusion and the first equality.

As $V_{grad}^{\mathbb{R}} = V_{grad}^{\mathbb{R}} \cap \mathcal{S}^+(C)$ (Remark 3.7), by the Positivstellensatz, ${}^{c^+}\sqrt{I_{grad}} = \sqrt[\mathbb{R}]{I_{grad}}$, which proves the second equality.

By the Positivstellensatz and Remark 3.7, we have

$${}^{c^+}\sqrt{I_{grad}} \cap \mathbb{R}[\mathbf{x}] = \sqrt[\mathbb{R}]{I_{grad}} \cap \mathbb{R}[\mathbf{x}] = \mathcal{I}(\pi^{\mathbf{x}}(V_{grad}^{\mathbb{R}})) = \mathcal{I}(\pi^{\mathbf{x}}(V_{KKT})^{\mathbb{R}} \cap \mathcal{S}^+(C)) = {}^{c^+}\sqrt{I_{KKT}^x}$$

and

$$I_{min} = I_{min}^z \cap \mathbb{R}[\mathbf{x}] = (p_1, \dots, p_r) \cap \mathbb{R}[\mathbf{x}] + {}^{c^+}\sqrt{I_{grad}} \cap \mathbb{R}[\mathbf{x}] = (p_1, \dots, p_r) + {}^{c^+}\sqrt{I_{KKT}^x}$$

which proves the equality. \square

Theorem 5.6. For $C \subset \mathbb{R}[\mathbf{x}]$ with $S_{min} \subset \mathcal{S}(C) \subset V_{KKT}^{\mathbf{x}, \mathbb{R}}$, there exists $t_2 \in \mathbb{N}$ such that $\forall t \geq t_2$, for $\Lambda^* \in \mathcal{L}_t(C)$ with $\Lambda^* = f_{t,C}^{\mu}$, we have $I_{min} \subset (\ker M_{\Lambda^*}^t)$.

Proof. To prove the inclusion we take $t_2 = \max\{t_0, t_1\}$ and we combine Proposition 5.5 with Proposition 5.2 for $C \subset \mathbb{R}[\mathbf{x}]$ and Theorem 5.4. \square

We introduce now the notion of *optimal linear form for f* . Such a linear form allows us to compute I_{min} .

Proposition 5.7. For $\Lambda^* \in \mathcal{L}_t(C)$ and $p \in \mathbb{R}[\mathbf{x}]$, the following assertions are equivalent:

- (i) $\text{rank} M_{\Lambda^*}^t = \max_{\Lambda \in \mathcal{L}_t(C), \Lambda(p) = p_{t,C}^{\mu}} \text{rank} M_{\Lambda}^t$.
- (ii) $\forall \Lambda \in \mathcal{L}_t(C)$ such that $\Lambda(p) = p_{t,C}^{\mu}$, $\ker M_{\Lambda^*}^t \subset \ker M_{\Lambda}^t$.

We say that $\Lambda^* \in \mathcal{L}_t(C)$ is *optimal for p* if it satisfies one of the equivalent conditions (i)-(ii).

A proof of this proposition can be found in [12](Proposition 4.7).

Remark 5.8. A linear form $\Lambda^* \in \mathcal{L}_t(C)$ optimal for p can be computed by solving a Semi-Definite Programming problem by an interior point method [14]. In this case, the solution Λ^* obtained by convex optimization is in the interior of the face of linear forms that minimize f .

The next result, which refines Theorem 5.6, shows that only elements in I_{min} are involved in the kernel of a truncated Hankel operator associated to an optimal linear form for f .

Theorem 5.9. Let $t \in \mathbb{N}$ such that $f \in \mathbb{R}[\mathbf{x}]_{2t}$ and let $C \subset \mathbb{R}[\mathbf{x}]_{2t}$ with $S_{min} \subset \mathcal{S}(C)$. If $\Lambda^* \in \mathcal{L}_t(C)$ is optimal for f and such that $\Lambda^*(f) = f^*$, then $\ker M_{\Lambda^*}^t \subset I_{min}$.

Proof. It is similar to proof of Theorem 4.9 in [12]. \square

The last result of this section shows that an optimal linear form for f yields the generators of the minimizer ideal I_{min} in high enough degree.

Theorem 5.10. Let $\mathbf{g} \subset \mathbb{R}[\mathbf{x}]$ be a set of constraints with $S_{min} \neq \emptyset$. For a set of constraints $C \subset \mathbb{R}[\mathbf{x}]$ with $S_{min} \subset \mathcal{S}(C) \subset V_{KKT}^{\mathbf{x}, \mathbb{R}}$, there exists $t_2 \in \mathbb{N}$ (defined in Theorem 5.6) such that $\forall t \geq t_2$,

- $f_{t,C}^\mu = \min_{\mathbf{x} \in S_{KKT}} f(\mathbf{x})$ is reached for some $\Lambda^* \in \mathcal{L}_t(C)$,
- $\forall \Lambda^* \in \mathcal{L}_t(C)$ optimal for f , we have $\Lambda^*(f) = f^*$ and $(\ker M_{\Lambda^*}^t) = I_{min}$,
- if $\mathcal{V}(C^0) \subset V_{KKT}^{\mathbf{x}, \mathbb{R}}$ then $f_{t,C}^{sos} = f_{t,C}^\mu = f^*$.

Proof. We obtain the result as a consequence of Theorem 5.4, Theorem 5.6 and Theorem 5.9. \square

The same results hold if we replace C by any other finite set defining a real variety such that $S_{min} \subset \mathcal{S}(C) \subset V_{KKT}^{\mathbf{x}, \mathbb{R}}$.

Remark 5.11. We can also replace the initial set of constraints \mathbf{g} by any other set $\tilde{\mathbf{g}}$ defining the same semi-algebraic set $S = \mathcal{S}(\mathbf{g}) = \mathcal{S}(\tilde{\mathbf{g}})$ and consider the KKT variety associated to $\tilde{\mathbf{g}}$.

6. CONSEQUENCES

Let us describe now some consequences of these results in specific cases, which have been previously studied.

6.1. Global optimization. We consider here the case $n_1 = n_2 = 0$. Theorem 4.9 implies the following result (compare with [28]):

Theorem 6.1. Let $f \in \mathbb{R}[\mathbf{x}]$.

- (1) If the real critical values of f are positive, then $f \in \mathcal{Q}^+ + \sqrt{(\frac{\partial f}{\partial x_1}, \dots, \frac{\partial f}{\partial x_n})}$.
- (2) If the real critical values of f are strictly positive, then $f \in \mathcal{Q}^+ + (\frac{\partial f}{\partial x_1}, \dots, \frac{\partial f}{\partial x_n})$.

In particular, if there is no real critical value, then $f \in \mathcal{Q}^+ + (\frac{\partial f}{\partial x_1}, \dots, \frac{\partial f}{\partial x_n})$.

A consequence of Proposition 5.1 and Theorem 5.10 is the following:

Theorem 6.2. Let $f \in \mathbb{R}[\mathbf{x}]$ and $C = \{\frac{\partial f}{\partial x_1}, \dots, \frac{\partial f}{\partial x_n}\}$. Then, there exists $t_0 \in \mathbb{N}$, such that $\forall t \geq t_0$ either $\mathcal{L}_t(C) = \emptyset$ and $S_{min} = \emptyset$ or

- (1) $f_{t,C}^{sos} = f_{t,C}^\mu = f^* = \min_{\mathbf{x} \in \mathbb{R}^n} f(\mathbf{x})$ is reached for some $\Lambda^* \in \mathcal{L}_t(C)$,
- (2) $\forall \Lambda^* \in \mathcal{L}_t(C)$ optimal for f , $\ker M_{\Lambda^*}^t$ generates I_{min} .

The first point of this theorem can also be found in [28].

6.2. General case. A direct consequence of Proposition 5.1 and Theorem 5.10 is the following:

Theorem 6.3. *Let $C \subset \mathbb{R}[\mathbf{x}]$ be a set of constraints such that*

- $(C^0) = I_{KKT} \cap \mathbb{R}[\mathbf{x}]$,
- $C^+ = \mathbf{g}^+$.

Then there exists $t_0 \in \mathbb{N}$ such that $\forall t \geq t_0$, either $\mathcal{L}_t(C) = \emptyset$ and $S_{min} = \emptyset$ or

- $f_{t,C}^{sos} = f_{t,C}^\mu = \min_{\mathbf{x} \in S_{KKT}} f(\mathbf{x})$ is reached for some $\Lambda^* \in \mathcal{L}_t(C)$,
- $\forall \Lambda^* \in \mathcal{L}_t(C)$ optimal for f , we have $\Lambda^*(f) = f^*$ and $(\ker M_{\Lambda^*}^t) = I_{min}$.

The set C^0 is constructed so that $\mathcal{V}(C^0) = V_{KKT}^{\mathbf{x}}$. As we have seen, the weaker condition $S_{min} \subset \mathcal{S}(C) \subset V_{KKT}^{\mathbf{x}}$ is sufficient to have an exact relaxation sequence.

The generators C^0 of $I_{KKT} \cap \mathbb{R}[\mathbf{x}]$ can be computed by elimination techniques (for instance by Groebner basis computation with a product order on monomials [3]).

6.3. Regular case. We consider here a semi-algebraic set S such that its defining constraints intersect properly. For any $\mathbf{x} \in \mathbb{C}^n$, let $\nu(\mathbf{x}) = \{j \in [1, n_2] \mid g_j^+(\mathbf{x}) = 0\}$.

Definition 6.4. *We say that a set of constraints $\mathbf{g} = (g_1^0, \dots, g_{n_1}^0; g_1^+, \dots, g_{n_2}^+)$ is regular if for all points $\mathbf{x} \in \mathcal{S}(\mathbf{g})$ with $\nu(\mathbf{x}) = \{j_1, \dots, j_k\}$, the vectors $\nabla g_1^0(\mathbf{x}), \dots, \nabla g_{n_1}^0(\mathbf{x}), \nabla g_{j_1}^+(\mathbf{x}), \dots, \nabla g_{j_k}^+(\mathbf{x})$ are linearly independent.*

This condition is used for instance in [9]. It implies that $\forall \mathbf{x} \in S$, $|\nu(\mathbf{x})| \leq n - n_1$ and that $B_{\nu(\mathbf{x})}(\mathbf{x})$ is of rank $n_1 + |\nu(\mathbf{x})|$. A stronger condition, called the \mathbb{C} -regularity, corresponds to sets of constraints such that $\forall \mathbf{x} \in \mathbb{C}^n$, $B_{\nu(\mathbf{x})}(\mathbf{x})$ is of rank $n_1 + |\nu(\mathbf{x})|$. This condition is used for instance in [25]. It is satisfied for semi-algebraic sets defined by “generic” constraints when $n_1 \leq n$ as shown in [25].

If \mathbf{g} is regular, then for all points \mathbf{x} in S the rank of $B_{\nu(\mathbf{x})}(\mathbf{x})$ is $n_1 + |\nu(\mathbf{x})|$ and $S_{sing} = \emptyset$. The decomposition (12) implies that $S_{FJ} = S_{KKT}$ and that all minimizer points of f on S are KKT points. If moreover \mathbf{g} is \mathbb{C} -regular, then $V_{FJ}^{\mathbf{x}} = \mathcal{V}(\Gamma_{FJ}) = V_{KKT}^{\mathbf{x}}$.

We deduce from Theorem 5.10 the following result:

Theorem 6.5. *Let $\mathbf{g} \subset \mathbb{R}[\mathbf{x}]$ be a regular set of constraints and let $C \subset \mathbb{R}[\mathbf{x}]$ be the set of constraints such that*

- $C^0 = \Gamma_{FJ}$ defined in (9) (resp. $C^0 = \Phi_{FJ}$ defined in (10)),
- $C^+ = \mathbf{g}^+$.

Suppose that $\min_{\mathbf{x} \in \mathcal{S}(\mathbf{g})} f(\mathbf{x})$ is reached at some point of $\mathcal{S}(\mathbf{g})$. Then, there exists $t_0 \in \mathbb{N}$ such that $\forall t \geq t_0$,

- (1) $f_{t,C}^\mu = f^* = \min_{\mathbf{x} \in \mathcal{S}(\mathbf{g})} f(\mathbf{x})$ is reached for some $\Lambda^* \in \mathcal{L}_t(C)$,
- (2) $\forall \Lambda^* \in \mathcal{L}_t(C)$ optimal for f , $\ker M_{\Lambda^*}^t$ generates $I_{min}^{\mathbf{x}}$,
- (3) If \mathbf{g} is \mathbb{C} -regular and $C^0 = \Gamma_{FJ}$, then $f_{t,C}^{sos} = f_{t,C}^\mu = f^*$.

By Lemma 3.11 and Remark 3.12, C is constructed so that $S_{min} \subset \mathcal{S}(C) = S_{KKT} \subset V_{KKT}^{\mathbf{x}, \mathbb{R}}$.

Points (1) and (3) are proved for $C^0 = \Gamma_{FJ}$ in [25] under the condition that \mathbf{g} is \mathbb{C} -regular. These points can also be found in [9] for $C^0 = \mathbf{g}^0 \cup \Phi_{FJ}$ under the condition that \mathbf{g} regular (but a problem appears in the proof: the vanishing of the polynomials Φ_{FJ} at a point $\mathbf{x} \in \mathbb{C}^n$ does not imply that $\text{rank } A_{\nu(\mathbf{x})}(\mathbf{x}) < n_1 + |\nu(\mathbf{x})|$).

In this case, the relaxation constructed with Γ_{FJ} (or Φ_{FJ}) is exact and can be used to compute the minimizer ideals of f on the semi-algebraic set S .

6.4. Zero dimensional real variety. Let $\mathbf{g} \subset \mathbb{R}[\mathbf{x}]$ be a set of constraints such that $\mathcal{V}^{\mathbb{R}}(\mathbf{g}^0)$ is finite and let $S := \mathcal{S}(\mathbf{g})$. By remark 5.11, we can assume that S is defined by a set of constraints $\tilde{\mathbf{g}}$ such that $(\tilde{\mathbf{g}}^0)$ is radical. Then $\forall \mathbf{x} \in \mathcal{V}(\mathbf{g}^0) = \mathcal{V}(\tilde{\mathbf{g}}^0)$, the Jacobian matrix $\tilde{B}_{\nu(\mathbf{x})}(\mathbf{x})$ associated to $\tilde{\mathbf{g}}^0$ is of rank n . Therefore we have $\mathcal{V}(\mathbf{g}^0) = \mathcal{V}(\tilde{\mathbf{g}}^0) = V_{KKT}^{\mathbf{x}}(\mathbf{g}^0)$ and any point of S is a KKT -point: $S = S_{FJ} = S_{KKT}$. Consequently, we deduce from Theorem 5.10 the following result:

Theorem 6.6. *Let $\mathbf{g} = (\mathbf{g}^0, \mathbf{g}^+) \subset \mathbb{R}[\mathbf{x}]$ be a set of constraints such that $\mathcal{V}^{\mathbb{R}}(\mathbf{g}^0)$ is finite. Then there exists $t_0 \in \mathbb{N}$ such that $\forall t \geq t_0$,*

- (1) $f_{t,\mathbf{g}}^{sos} = f_{t,\mathbf{g}}^{\mu} = f^* = \min_{\mathbf{x} \in \mathcal{S}(\mathbf{g})} f(\mathbf{x})$ is reached for some $\Lambda^* \in \mathcal{L}_t(\mathbf{g})$,
- (2) $\forall \Lambda^* \in \mathcal{L}_t(\mathbf{g})$ optimal for f , $\ker M_{\Lambda^*}^t$ generates I_{min} .

This answers an open question in [18]. The first point was also solved in [27] using dedicated techniques.

6.5. Smooth real variety. We consider a set of constraints $\mathbf{g} = \{g_1^0, \dots, g_{n_1}^0\} \subset \mathbb{R}[\mathbf{x}]$ such that $\mathcal{V}^{\mathbb{R}}(\mathbf{g}^0)$ is equidimensional smooth and $\mathbf{g}^+ = \emptyset$. This means that $S = \mathcal{S}(\mathbf{g}) = \mathcal{V}^{\mathbb{R}}(\mathbf{g}^0)$ is the union of irreducible components of the same dimension d and that for any point $\mathbf{x} \in S$, $B_{\emptyset}(\mathbf{x}) = [\nabla g_1^0(\mathbf{x}), \dots, \nabla g_{n_1}^0(\mathbf{x})]$ is of rank $m = \dim S = n - d$. Therefore, $S_{sing} = \emptyset$. In this case, $\nabla f(\mathbf{x})$ is a linear combination of $\nabla g_1^0(\mathbf{x}), \dots, \nabla g_{n_1}^0(\mathbf{x})$, if and only if, $\text{rank} A_{\emptyset}(\mathbf{x}) \leq r$.

The set Γ_{FJ} defined in (9) (or $C^0 = \mathbf{g}^0 \cup \Phi_{FJ}$ defined in (10)), or the union Δ^{n-d} of \mathbf{g}^0 and the set of $(n - d + 1) \times (n - d + 1)$ minors of the Jacobian matrix of $\{f, g_1^0, \dots, g_{n_1}^0\}$, which contain the first column ∇f define the variety S_{KKT} .

We deduce from Theorem 5.10, the following result:

Theorem 6.7. *Let $\mathbf{g} = \{g_1^0, \dots, g_{n_1}^0\} \subset \mathbb{R}[\mathbf{x}]$ such that $S = \mathcal{V}^{\mathbb{R}}(\mathbf{g})$ is an equidimensional and smooth variety of dimension d .*

Let $C \subset \mathbb{R}[\mathbf{x}]$ be the set of constraints such that $C^0 = \Gamma_{FJ}$ defined in (9) (or $C^0 = \Phi_{FJ}$ defined in (10), $C^0 = \Delta^{n-d}$) Then there exists $t_0 \in \mathbb{N}$ such that $\forall t \geq t_0$, either $\mathcal{L}_t(C) = \emptyset$ and $S_{min} = \emptyset$ or

- (1) $f_{t,C}^{\mu} = f^* = \min_{\mathbf{x} \in S} f(\mathbf{x})$ is reached for some $\Lambda^* \in \mathcal{L}_t(C)$,
- (2) $\forall \Lambda^* \in \mathcal{L}_t(C)$ optimal for f , $\ker M_{\Lambda^*}^t$ generates I_{min} .

6.6. Known minimum. In the case where we know the minimum f^* of f on the basic closed semi-algebraic set S , we take \mathbf{g}' with $\mathbf{g}'^0 = \{\mathbf{g}^0, f - f^*\}$ and $\mathbf{g}'^+ = \mathbf{g}^+$. Let $S = \mathcal{S}(\mathbf{g})$, $S' = \mathcal{S}(\mathbf{g}')$. By construction $S_{min} \subset S'$ and $S' = S'_{KKT}$ and $\mathcal{V}(\mathbf{g}'^0) \subset V_{KKT}^{\mathbf{x}}(\mathbf{g}'^0)$. Theorem 5.10 applied to \mathbf{g}' implies the following result:

Theorem 6.8. *Let $\mathbf{g} = \{g_1^0, \dots, g_{n_1}^0; g_1^+, \dots, g_{n_2}^+\} \subset \mathbb{R}[\mathbf{x}]$. Let f^* be the minimum of f and $C \subset \mathbb{R}[\mathbf{x}]$ the set of constraints such that $C^0 = \{\mathbf{g}^0, f - f^*\}$ and $C^+ = \mathbf{g}^+$. Then there exists $t_0 \in \mathbb{N}$ such that $\forall t \geq t_0$,*

- (1) $f_{t,C}^{sos} = f_{t,C}^{\mu} = f^* = \min_{\mathbf{x} \in \mathcal{S}(C)} f(\mathbf{x})$ is reached for some $\Lambda^* \in \mathcal{L}_t(C)$,
- (2) $\forall \Lambda^* \in \mathcal{L}_t(C)$ optimal for f , $\ker M_{\Lambda^*}^t$ generates I_{min} .

6.7. \mathbf{g}^+ -radical computation. In the case where $f = 0$, by Remark 3.15 all the points of S are KKT points and minimizers of f so that $S_{min} = S = S_{KKT}$. Moreover, $I_{KKT}^{\mathbf{x}} = (g_1^0, \dots, g_{n_1}^0)$ since $F_1, \dots, F_n, v_1 g_1^+, \dots, v_{n_2} g_{n_2}^+$ are homogeneous of degree 1 in the variables $u_1, \dots, u_{n_1}, v_1, \dots, v_{n_2}$. We deduce the following result:

Theorem 6.9. *Let $\mathbf{g} = \{g_1^0, \dots, g_{n_1}^0; g_1^+, \dots, g_{n_2}^+\} \subset \mathbb{R}[\mathbf{x}]$. There exists $t_2 \in \mathbb{N}$ such that $\forall t \geq t_2$, $\forall \Lambda^* \in \mathcal{L}_t(\mathbf{g})$ optimal for 0, we have $(\ker M_{\Lambda^*}^t) = \mathcal{I}(S) = \mathfrak{s}^+/\sqrt{(\mathbf{g}^0)}$.*

This gives a way to compute $\sqrt[C^+]{C^0}$ (see also [20]), which generalizes the approach of [16], [12] or [31] to compute the real radical of an ideal.

7. EXAMPLES

This section contains examples that illustrate different aspects of our method. In the case of a finite number of minimizers for a function f on the semi-algebraic set S defined by the set of constraints \mathbf{g} , the approach we describe leads to the following algorithm:

- (1) Compute $C \subset \mathbb{R}[\mathbf{x}]$ such that C^0 generates $I_{KKT} \cap \mathbb{R}[\mathbf{x}]$ and $C^+ = \mathbf{g}^+$;
- (2) $t := \lceil \frac{1}{2} \max\{\deg(f), \deg(g_i^0), \deg(g_j^+)\} \rceil$;
- (3) Compute $\Lambda^* \in \mathcal{L}_t(C)$ optimal for f (solving a finite dimensional SDP problem by an interior point method);
- (4) Check the convergence certificate for $M_{\Lambda^*}^t$ (by flat extension [10, 19]);
- (5) If it is not satisfied, then $t := t + 1$ and repeat from step (2);
Otherwise compute $K := \ker M_{\Lambda^*}^t$.

Output $f^* = \Lambda^*(f)$ and the generators K of I_{min} .

Example 7.1. We consider the “ill-posed” problem

$$\min x \text{ s.t. } x^3 \geq 0.$$

The ideal I_{KKT} is $I_{KKT} = (1 - 3v_1x^2, v_1x^3) = (1)$. Thus $V_{KKT} = \emptyset$. According to the decomposition (12), $S_{FJ} = S_{sing}$ and we compute the minimum of x on S_{sing} , which is defined by $x^2 = 0$:

$$\min x \text{ s.t. } x^2 = 0.$$

Now according to section 6.4, the relaxation associated to this problem is exact and yields the solution $x = 0$.

Example 7.2. We consider the following problem

$$\begin{aligned} \min \quad & f(x, y, z) = x^2 + y^2 + z^2; \\ \text{s.t.} \quad & \text{rank} \begin{pmatrix} x+z+1 & x+y & y+z \\ x+y & y+z & x+z+1 \end{pmatrix} \leq 1 \end{aligned}$$

or equivalently

$$\begin{aligned} \min \quad & f(x, y, z) = x^2 + y^2 + z^2; \\ \text{s.t.} \quad & (x+z+1)(y+z) - (x+y)^2 = 0; \\ & (x+z+1)^2 - (y+z)(x+y) = 0; \\ & (x+z+1)(x+y) - (y+z)^2 = 0; \end{aligned}$$

This corresponds to computing the closest point on a twisted cubic defined by 2×2 minors. The set of constraints \mathbf{g} is not regular but $\mathcal{S}(\mathbf{g}) = \mathcal{V}^{\mathbb{R}}(\mathbf{g}^0)$ is a smooth real variety.

In the first iteration of the algorithm, the order is 1, the size of the Hankel matrix M_{Λ}^1 is 3, $\min \Lambda(f) = 1$ and there is no duality gap. The flat extension condition is satisfied for M_{Λ}^1 and thus we have found the minimum. The algorithm stops and we obtain $I_{min} = (x, y-1, z)$. The points that minimize f are $\{(x=0, y=1, z=0)\}$.

Example 7.3. We consider the Motzkin polynomial,

$$\min f(x, y) = 1 + x^4y^2 + x^2y^4 - 3x^2y^2$$

which is non negative on \mathbb{R}^2 but not a sum of squares in $\mathbb{R}[x, y]$. We compute its gradient ideal, $I_{grad}(f) = (-6xy^2 + 2xy^4 + 4x^3y^2, -6yx^2 + 2yx^4 + 4y^3x^2)$, which is not zero-dimensional.

In the first iteration of the algorithm, the order is 3, the size of the Hankel matrix M_Λ^3 is 10, $\min \Lambda(f) = -216$. The flat extension condition is not satisfied hence we try with degree 4.

In the second iteration the order is 4, the size of the Hankel matrix M_Λ^4 is 15, $\min \Lambda(f) = 0$, there is no duality gap. The flat extension condition is satisfied for M_Λ^4 and we have found the minimum. The algorithm stops and we obtain $I_{\min} = (x^2 - 1, y^2 - 1)$. The points that minimize f are $\{(x = 1, y = 1), (x = 1, y = -1), (x = -1, y = 1), (x = -1, y = -1)\}$.

For this example Gloptipoly must go until order 9 in order to satisfy the flat extension condition.

Example 7.4. We consider the Robinson polynomial

$$\min f(x, y) = 1 + x^6 - x^4 - x^2 + y^6 - y^4 - y^2 - x^4 y^2 - x^2 y^4 + 3x^2 y^2$$

which is non negative on \mathbb{R}^2 but not a sum of squares in $\mathbb{R}[x, y]$. We compute its gradient ideal,

$$I_{\text{grad}}(f) = (6x^5 - 4x^3 - 2x - 4x^3 y^2 - 2xy^4 + 6xy^2, 6y^5 - 4y^3 - 2y - 4y^3 x^2 - 2yx^4 + 6yx^2)$$

which is not zero-dimensional.

In the first iteration, the order is 3, the size of the Hankel matrix M_Λ^3 is 10, $\min \Lambda(f) = -0.93$. The flat extension condition is not satisfied hence we try with degree 4.

In the second iteration the degree is 4, the size of the Hankel matrix M_Λ^4 is 15, $\min \Lambda(f) = 0$. There is no duality gap. The flat extension condition is satisfied for M_Λ^4 and we have found the minimum.

The algorithm stops and we obtain $I_{\min} = (x^3 - x, y^3 - y, x^2 y^2 - x^2 - y^2 + 1)$. The points that minimize f are $\{(x = 1, y = 1), (x = 1, y = -1), (x = -1, y = 1), (x = -1, y = -1), (x = 1, y = 0), (x = -1, y = 0), (x = 0, y = 1), (x = 0, y = -1)\}$.

For this example, Gloptipoly must go until order 7 in order to satisfy the flat extension condition.

Example 7.5. We consider the homogeneous Motzkin polynomial with a perturbation $\epsilon = 0.005$,

$$\begin{aligned} \min f(x, y, z) &= x^4 y^2 + x^2 y^4 - 3x^2 y^2 z^2 + z^6 + \epsilon(x^2 + y^2 + z^2); \\ \text{s.t } h(x, y, z) &= 1 - x^2 - y^2 - z^2 \geq 0 \end{aligned}$$

This example coming from [18, Example 6.25] is a case where the constraints \mathbf{g} define a compact semi-algebraic set, but the direct relaxation using the associated quadratic module or preordering is not exact.

We add the projection of the KKT ideal and we have the similar problem

$$\begin{aligned} \min x^4 y^2 + x^2 y^4 - 3x^2 y^2 z^2 + z^6 + 0.005(x^2 + y^2 + z^2); \\ \text{s.t } -4zx^4 y - 20zx^2 y^3 + 12x^2 y z^3 - 0.06zy^5 + 12.06yz^5 &= 0; \\ -20zx^3 y^2 - 4zxy^4 + 12xy^2 z^3 - 0.06zx^5 + 12.06xz^5 &= 0; \\ (4x^3 y^2 + 2xy^4 - 6xy^2 z^2 + 0.03x^5)(-x^2 - y^2 - z^2 + 1) &= 0; \\ (2x^4 y + 4x^2 y^3 - 6x^2 y z^2 + 0.03y^5)(-x^2 - y^2 - z^2 + 1) &= 0; \\ (-6x^2 y^2 z + 6.03z^5)(-x^2 - y^2 - z^2 + 1) &= 0; \end{aligned}$$

where the first three equations are the 2×2 minors of the Jacobian matrix of f and h and the last three equations are the gradient ideal of f multiplied by h .

In the first iteration the order is 5, the size of the Hankel matrix M_Λ^5 is 167, $\min \Lambda(f) = 0$, there is no duality gap. The flat extension condition is satisfied for M_Λ^5 and we have found the minimum. The algorithm stops and we obtain $I_{\min} = (x, y, z)$. The point that minimize f is $(0, 0, 0)$.

For this example, the flat extension condition does not hold with Gloptipoly if $\epsilon \leq 0.01$.

Finally with these two last examples we show that even the minimizer ideal I_{min} is not zero-dimensional we can recover it from a solution of the relaxation problem.

Example 7.6. *We consider Motzkin polynomial over the unit ball:*

$$\begin{aligned} \min \quad & f(x, y, z) = x^4y^2 + x^2y^4 - 3x^2y^2z^2 + z^6; \\ \text{s.t.} \quad & h(x, y, z) = 1 - x^2 - y^2 - z^2 \geq 0 \end{aligned}$$

The polynomial f is homogeneous and non negative on \mathbb{R}^3 but not a sum of squares in $\mathbb{R}[x, y, z]$.

We add the projections of KKT ideal and we have the similar problem

$$\begin{aligned} \min \quad & x^4y^2 + x^2y^4 - 3x^2y^2z^2 + z^6; \\ \text{s.t.} \quad & -4xy^5 + 12xy^3z^2 + 4yx^5 - 12x^3yz^2 = 0; \\ & -4zx^4y - 20zx^2y^3 + 12x^2yz^3 + 12yz^5 = 0; \\ & -20zx^3y^2 - 4zxy^4 + 12xy^2z^3 + 12xz^5 = 0; \\ & (4x^3y^2 + 2xy^4 - 6xy^2z^2)(-x^2 - y^2 - z^2 + 1) = 0; \\ & (2x^4y + 4x^2y^3 - 6x^2yz^2)(-x^2 - y^2 - z^2 + 1) = 0; \\ & (-6x^2y^2z + 6z^5)(-x^2 - y^2 - z^2 + 1) = 0; \end{aligned}$$

where the first three equations are the 2×2 minors of the Jacobian matrix of f and h and the last three equations are the gradient ideal of f multiplied by h .

In the first iteration the order is 5, the size of the Hankel matrix M_Λ^5 is 156, $\min \Lambda(f) = 0$, there is no duality gap. We compute the kernel of this matrix: $\ker M_\Lambda^5 = \langle z(y^2 - z^2), x(y^2 - z^2), z(x^2 - z^2), y(x^2 - z^2) \rangle$. It generates the minimizer ideal $I_{min} = (z(y^2 - z^2), x(y^2 - z^2), z(x^2 - z^2), y(x^2 - z^2))$ defining 6 lines: $(y \pm z, x \pm z), (x, z), (y, z)$. Here $\mathcal{V}(I_{min})$ is not included in S .

Example 7.7. *We consider minimization of a linear function on a torus:*

$$\begin{aligned} \min \quad & f(x, y, z) = z \\ \text{s.t.} \quad & 9 - 10x^2 - 10y^2 + 6z^2 + x^4 + 2x^2y^2 + 2x^2z^2 + 2y^2z^2 + y^4 + z^4 = 0 \end{aligned}$$

In the first iteration, the order is 2, the size of the Hankel matrix M_Λ^2 is 10, $\min \Lambda(f) = -1$, there is no duality gap. We compute the kernel of this matrix: $\ker M_\Lambda^2 = \langle x^2 + y^2 - 4, x(z + 1), y(z + 1), z(z + 1), (z + 1) \rangle$ which generates the minimizer ideal $I_{min} = (x^2 + y^2 - 4, z + 1)$, defining a circle which is the intersection of the torus with a tangent plane. Notice that the multiplicity of this intersection has been removed in I_{min} .

REFERENCES

- [1] J. Bochnak, M. Coste, and M.-F. Roy. *Real Algebraic Geometry*. Springer, 1998.
- [2] W. Bruns and U. Vetter. *Determinantal rings*, volume 1327 of *Lecture Notes in Math*. Springer, Berlin, 1988.
- [3] D.A. Cox, J.B. Little, and D.B. O’Shea. *Ideals, Varieties, and Algorithms : An Introduction to Computational Algebraic Geometry and Commutative Algebra (Undergraduate Texts in Mathematics)*. Springer, 2005.
- [4] R.E. Curto and L. Fialkow. Solution of the truncated complex moment problem for flat data. *Memoirs of the American Mathematical Society*, 119(568):1–62, 1996.
- [5] James Demmel, Jiawang Nie, and Victoria Powers. Representations of positive polynomials on non-compact semialgebraic sets via kkt ideals. *Journal of Pure and Applied Algebra*, 209(1):189 – 200, 2007.
- [6] A. Greuet and M. Safey El Din. Deciding reachability of the infimum of a multivariate polynomial. In *Proceedings of the 36th international symposium on Symbolic and algebraic computation, ISSAC ’11*, pages 131–138, New York, NY, USA, 2011. ACM.

- [7] F. Guo, M. Safey El Din, and L. Zhi. Global optimization of polynomials using generalized critical values and sums of squares. In *Proceedings of the 2010 International Symposium on Symbolic and Algebraic Computation*, ISSAC '10, pages 107–114, New York, NY, USA, 2010. ACM.
- [8] H. V. Ha and T.S. Pham. Global optimization of polynomials using the truncated tangency variety. *SIAM Journal on Optimization*, 19(2):941–951, 2008.
- [9] H. V. Ha and T.S. Pham. Representation of positive polynomials and optimization on noncompact semialgebraic sets. *SIAM Journal on Optimization*, 20(6):3082–3103, 2010.
- [10] D. Henrion and J.B. Lasserre. *Positive Polynomials in Control*, chapter Detecting Global Optimality and Extracting Solutions in GloptiPoly., pages 293–310. Lectures Notes in Control and Information Sciences. Springer, 2005.
- [11] F. John. Extremum problems with inequalities as side conditions. In *Studies and Essays, Courant Anniversary Volume*, pages 187–204. Wiley (Interscience), New York, 1948.
- [12] J.-B. Lasserre, M. Laurent, B. Mourrain, P. Rostalski, and P. Trébuchet. Moment matrices, border bases and real radical computation. *Journal of Symbolic Computation*, 2012.
- [13] J.B. Lasserre. Global optimization with polynomials and the problem of moments. *SIAM J. Optim.*, 11:796–817, 2001.
- [14] J.B. Lasserre. *Moments, positive polynomials and their applications*. Imperial College Press, 210.
- [15] J.B. Lasserre, M. Laurent, and P. Rostalski. Semidefinite characterization and computation of real radical ideals. *Foundations of Computational Mathematics*, 8(5):607–647, 2008.
- [16] J.B. Lasserre, M. Laurent, and P. Rostalski. A unified approach for real and complex zeros of zero-dimensional ideals. In M. Putinar and S. Sullivant, editors, *Emerging Applications of Algebraic Geometry*, volume 149, pages 125–156. Springer, 2009.
- [17] M. Laurent. Semidefinite representations for finite varieties. *Math. Progr*, 109:1–26, 2007.
- [18] M. Laurent. *Sums of squares, moment matrices and optimization over polynomials*, volume 149 of *IMA Volumes in Mathematics and its Applications*, pages 157–270. Springer, 2009.
- [19] M. Laurent and B. Mourrain. A generalized flat extension theorem for moment matrices. *Arch. Math. (Basel)*, 93(1):87–98, July 2009.
- [20] Y. Ma, Ch. Wang, and Zhi L. A certificate for semidefinite relaxations in computing positive dimensional real varieties. <http://arxiv.org/abs/1212.4924>, 2013.
- [21] O.L. Mangasarian and S. Fromovitz. The Fritz John necessary optimality conditions in the presence of equality and inequality constraints. *Journal of Mathematical Analysis and Applications*, 17:37–47, 1967.
- [22] M. Marshall. Optimization of polynomial functions. *Canad. Math. Bull.*, 46:575–587, 2003.
- [23] M. Marshall. Representations of non-negative polynomials, degree bounds and applications to optimization. *Can. J. Math.*, 61(1):205–221, 2009.
- [24] Y. Nesterov. Squared functional systems and optimization problems. In H. Frenk, K. Roos, T. Terlaky, and S. Zhang, editors, *High performance optimization*, chapter 17, pages 405–440. Kluwer academic publishers, Dordrecht, The Netherlands, 2000.
- [25] J. Nie. An exact jacobian SDP relaxation for polynomial optimization. *Mathematical Programming*, pages 1–31, 2011.
- [26] J. Nie. Certifying convergence of Lasserre’s hierarchy via flat truncation. *Mathematical Programming*, pages 1–26, 2012.
- [27] J. Nie. Polynomials optimization with real variety. *SIAM Journal On Optimization*, 23(3):1634–1646, 2013.
- [28] J. Nie, J. Demmel, and B. Sturmfels. Minimizing polynomials via sum of squares over gradient ideal. *Math. Program.*, 106(3):587–606, 2006.
- [29] P.A. Parrilo. Semidefinite programming relaxations for semialgebraic problems. *Mathematical Programming Ser. B*, 96(2):293–320, 2003.
- [30] P.A. Parrilo and B. Sturmfels. Minimizing polynomial functions. In *Proceedings of the DIMACS Workshop on Algorithmic and Quantitative Aspects of Real Algebraic Geometry in Mathematics and Computer Science*, pages 83–100. American Mathematical Society, 2003.
- [31] P. Rostalski. *Algebraic moments, real root finding and related topics*. PhD thesis, ETH Zurich, 2009.
- [32] M. Safey El Din. Computing the global optimum of a multivariate polynomial over the reals. In *Proceedings of the twenty-first international symposium on Symbolic and algebraic computation*, ISSAC '08, pages 71–78, New York, NY, USA, 2008. ACM.
- [33] M. Schweighofer. Global optimization of polynomials using gradient tentacles and sums of squares. *SIAM Journal on Optimization*, 17(3):920–942, 2006.
- [34] Shafarevich. *Basic algebraic geometry*. Springer-Verlag, 1974.

[35] N.Z. Shor. Class of global minimum bounds of polynomial functions. *Cybernetics*, 23:731–734, 1987.

MARTA ABRIL BUCERO, BERNARD MOURRAIN: GALAAD, INRIA MÉDITERRANÉE, 06902 SOPHIA ANTIPOLIS