

**Dynamics and spike trains statistics in
conductance-based Integrate-and-Fire neural networks
with chemical and electric synapses**

Rodrigo Cofré, Bruno Cessac

► **To cite this version:**

Rodrigo Cofré, Bruno Cessac. Dynamics and spike trains statistics in conductance-based Integrate-and-Fire neural networks with chemical and electric synapses. AREADNE 2012. Encoding And Decoding of Neural Ensembles, Jun 2012, Santorini, Greece. 2012. <hal-00850100>

HAL Id: hal-00850100

<https://hal.inria.fr/hal-00850100>

Submitted on 2 Aug 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ROLE OF ELECTRIC SYNAPSES IN SPIKE TRAINS STATISTICS OF LINEAR INTEGRATE AND FIRE NEURAL NETWORKS

Rodrigo Cofre^{*}, Bruno Cessac

INRIA Neuromathcomp Team, Sophia Antipolis, France

^{*} cofre@polytech.unice.fr

Communication between neurons involves chemical synapses as well as electric synapses. On theoretical grounds, the role of gap junctions in encoding and shaping collective dynamics as well as spike train statistics is quite less understood than the role of chemical synapses. In previous work [1] the collective spike train statistics in conductance-based Integrate and Fire neural networks was studied rigorously. It was especially shown that this statistics is characterized by a Gibbs distribution whose potential can be explicitly computed. This provides moreover a firm theoretical ground for recent studies attempting to describe experimental rasters in the retina [4] as well as in the parietal cat cortex [2] by Gibbs distributions and maximal entropy principle. The work presented at AREADNE will extend the mathematical analysis of previous work [1] to conductance-based Integrate and Fire neural networks with chemical synapses as well as electric synapses, in the presence of noise. In opposition to previous paper dealing with this subject [3] we do not consider mean field approximations and the analysis is not limited to pulse type chemical synapses.

The core of the analysis is to show how multiple single neurons interact in the presence of gap junctions. In conductance based models coupled with gap junctions, the sub-threshold dynamics is ruled by a coupled set of linear, non-autonomous and non-homogeneous stochastic differential equations of Ornstein-Uhlenbeck type, where conductances depends upon the spike-history of the network. We compute explicitly the evolution operator and show that given the spike-history of the network and the membrane potentials at a given time, the further dynamical evolution can be explicitly written in a closed form. Under a moderate assumption on the evolution operator the system has a unique strong solution. From this we obtain a family of transition probabilities for spike dynamics, corresponding to a Gibbs distribution. This results emphasizes the role of gap junctions on spike statistics with strong effects on space time correlations and memory. This work suggest that electric synapses could have a strong influence in spike train statistics of biological neural systems, especially the retina where gap junctions connections between several cells-type (e.g. amacrine and ganglion cells or amacrine-bipolar) are ubiquitous.

References

1. Cessac B. Statistics of spike trains in conductance-based neural networks: Rigorous results. *J. Mathemat. Neurosci.*, 1(8):1–42, 2011.
2. Marre O, El Boustani S, Fregnac Y, Destexhe A. Prediction of spatiotemporal patterns of neural activity from pairwise correlations. *Phys. Rev. Lett.*, 102(13), April 2009.
3. Ostojic S, Brunel N, Hakim V. Synchronization properties of networks of electrically coupled neurons in the presence of noise and heterogeneities. *J. Comp. Neurosci.*, 26(3):369–392, 2009.
4. Vasquez J-C, Palacios A, Marre O, Berry MJ II, Cessac B. Gibbs distribution analysis of temporal correlation structure on multicell spike trains from retina ganglion cells. *J. Physiol. Paris*, 2011.