

# Role of electric synapses in spike train statistics of integrate and fire neural networks

Rodrigo Cofré, Bruno Cessac

► **To cite this version:**

Rodrigo Cofré, Bruno Cessac. Role of electric synapses in spike train statistics of integrate and fire neural networks. GDR MEA 2012. Encoding And Decoding of Neural Ensembles, Jun 2012, Marseille, France. 2012. <hal-00850109>

**HAL Id: hal-00850109**

**<https://hal.inria.fr/hal-00850109>**

Submitted on 5 Aug 2013

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.


# Dynamics and spike trains statistics in conductance-based Integrate-and-Fire with chemical and electric synapses

Rodrigo Cofre, Bruno Cessac  
NeuroMathComp project team (INRIA, UNSA, ENS, CNRS)



We consider conductance-based Integrate-and-Fire models coupled with gap junctions and chemical synapses, where conductances depend upon the spike-history of the network. We compute explicitly the time evolution operator and show that given the spike-history of the network and the membrane potentials at a given time, the further dynamical evolution can be written in a closed form. Moreover, spike train statistics is described by a Gibbs distribution whose potential can be approximated with an explicit formula, when the noise is weak.

## Spike Trains



Spike state  
 $\omega_k(n) \in \{0, 1\}$

Spike pattern  
 $\omega(n) = (\omega_k(n))_{k=1}^N$

Spike block  
 $\omega_m^n = \{\omega(m) \omega(m+1) \dots \omega(n)\}$

Raster plot  
 $\omega \stackrel{\text{def}}{=} \omega_0^T$

## The equation

We consider a network of  $N$  neurons, where dynamics depend in continuous and discrete dynamical variables. The sub-threshold variation of the membrane potential of neuron  $k$  at time  $t$  is given by:

$$C_k \frac{dV_k}{dt} = -g_{L,k}(V_k - E_L) - \sum_j g_{kj}(t, \omega) (V_k - E_j) + \sum_j \bar{g}_{kj} (V_j - V_k) + I_k(t)$$

Where :

$C_k$  is the capacitance of each neuron  
 $g_{L,k}$  is the leak conductance.

$\bar{g}_{kj}$  is the electrical conductance which is symmetric.

$I_k(t) = i_k^{(ext)}(t) + \sigma_B \xi_k(t)$  is the current term which is composed by:

$i_k^{(ext)}(t)$  the external current, and white noise term  $\xi_k(t)$  whose magnitude is controlled by  $\sigma_B > 0$

## Conductance adaptation

$$g_{kj}(t) = g_{kj}(t_j^{(r)}(\omega)) + G_{kj} \alpha_{kj}(t - t_j^{(r)}(\omega)), \quad t > t_j^{(r)}(\omega)$$

$G_{kj} \geq 0$  is the maximal conductance

$\alpha_{kj}(t) = h(t) e^{-\frac{t}{\tau_{kj}}} H(t)$  is called "alpha function"

## Matrix-Vector representation

$$C \frac{dV}{dt} + [G(t, \omega) - \bar{G}] V = I(t, \omega)$$

$$G(t, \omega) = (g_{L,k} + \sum_{j=1}^N g_{kj}(t, \omega)) \delta_{kk}$$

$\bar{G}$  is the symmetric matrix of electric conductances, with entries  $\bar{g}_{kj}$

$$I(t, \omega) = I^{cs}(t, \omega) + I^{ext}(t) + I^B(t)$$

## The linear SDE

$$\frac{dV}{dt} = \underbrace{C^{-1}(\bar{G} - G(t, \omega)) V}_{\Phi(t, \omega)} + \underbrace{C^{-1} I^{cs}(t, \omega)}_{f(t, \omega)} + C^{-1} I^{ext}(t) + C^{-1} I^B(t)$$

## The flow and the solution

If we define recursively the following matrices:

$$M_0(t_0, t, \omega) = \mathcal{I}_N$$

$$M_k(t_0, t, \omega) = \mathcal{I}_N + \int_{t_0}^t \Phi(s, \omega) M_{k-1}(s, t, \omega) ds$$

Under very general conditions, this limits exist and is called "flow"

$$\Gamma(t_0, t, \omega) \stackrel{\text{def}}{=} \lim_{k \rightarrow \infty} M_k(t_0, t, \omega)$$

Considering both chemical and electric synapses, there exist a unique strong solution to the stochastic differential equation given by

$$V(t_0, t, \omega) = \Gamma(t_0, t, \omega) v + \int_{t_0}^t \Gamma(s, t, \omega) f(s, \omega) ds + \frac{\sigma_B}{c} \int_{t_0}^t \Gamma(s, t, \omega) dW(s)$$

This are examples when the flow takes an exponential form

(i)  $\bar{G}$  is diagonal;

(ii)  $\bar{G} = 0$ ;

(iii)  $G(t, \omega) = G(t) = \kappa(t) \mathcal{I}_N$  where  $\kappa(t)$  is a real function.

## Spike Trains Statistics

The membrane potential can be decomposed in their deterministic and stochastic part:

$$V(t, \omega) = V^{(d)}(t, \omega) + V^{(noise)}(t, \omega)$$

The main result establishes that spike trains are distributed according to a Gibbs distribution whose potential can be obtained under Gaussian approximation of  $V^{(noise)}(t, \omega)$

$$\hat{\theta}_k(t, \omega) = \theta - V_k^{(d)}(t, \omega) \quad \mathcal{J}_{k(n, \omega)} = \begin{cases} ]-\infty, \hat{\theta}_k(n-1, \omega)_+ & \text{if } \omega_k(n) = 0; \\ [\hat{\theta}_k(n-1, \omega)_-, +\infty[ & \text{if } \omega_k(n) = 1; \end{cases}$$

$$\mathcal{J}(n, \omega) = \prod_{k=1}^N \mathcal{J}_k(n, \omega) \quad dv = \prod_{i=1}^N dv_i$$

$$\mathbb{P}[\omega(n) | \omega_{-\infty}^{n-1}] = \int_{\mathcal{J}(n, \omega)} \frac{e^{-V^T Q^{-1} (n-1, \omega) V}}{(2\pi)^{\frac{N}{2}} |\mathcal{Q}(n-1, \omega)|^{\frac{1}{2}}} dV$$

In the case without electric synapses:

$$\mathbb{P}[\omega(n) | \omega_{-\infty}^{n-1}] = \prod_{k=1}^N \mathbb{P}[\omega_k(n) | \omega_{-\infty}^{n-1}]$$

## Gibbs distribution

The transition probabilities define a stochastic process in the set of rasters, where the probability of having a spike pattern depends on an infinite past. Such process are called "Chains with complete connections". Under suitable conditions define a unique probability distribution called Gibbs distribution.

$$\phi(n, \omega) = \log \mathbb{P}[\omega(n) | \omega_{-\infty}^{n-1}] \quad \text{is called Gibbs potential}$$

$$\mathbb{P}[\omega_m^n | \omega_{-\infty}^{m-1}] = e^{\sum_{i=m}^n \phi(i, \omega)} \quad \text{is the conditional probability in terms of the Gibbs potential}$$

## Conclusions:

Including electric synapses spike statistics becomes **indecomposable**.

$$\mathbb{P}[\omega(n) | \omega_{-\infty}^{n-1}] \neq \prod_{k=1}^N \mathbb{P}[\omega_k(n) | \omega_{-\infty}^{n-1}]$$

Correlations are due to stimulus and **dynamics**.

Electric synapses have an effect in the **memory** of the system.

The **Gibbs potential** is largely **more complex** than the Ising Model used in retina spike train analysis.

## Bibliography :

- [1] Bruno Cessac. Statistics of spike trains in conductance-based neural networks: Rigorous results. *The Journal of Mathematical Neuroscience*, 1(8):1-42, 2011
- [2] Rodrigo Cofre, Bruno Cessac. Dynamics and spike trains statistics in conductance-based Integrate-and-Fire neural networks with chemical and electric synapses. Preprint

**Acknowledgments:** This work was supported by the INRIA, ERC-NERVI number 227747, KEOPS ANR-CONICYT and from the European Union Seventh Framework Programme (FP7/2007- 2013) under grant agreement no. 269921 (BrainScale5). R Cofre is funded by the French ministry of Research and University of Nice (EDSTIC).