

## Mathematical modeling of soil strains growth at batch culture

Miled El Hajji, Lionel Ranjard, Pierre-Alain Maron, Virginie Nowak, Jérôme Harmand, Alain Rapaport

► **To cite this version:**

Miled El Hajji, Lionel Ranjard, Pierre-Alain Maron, Virginie Nowak, Jérôme Harmand, et al.. Mathematical modeling of soil strains growth at batch culture. 4ème Colloque de l'Association Francophone d'Ecologie Microbienne (AFEM), Aug 2009, Lyon, France. 1 p. hal-00858560

**HAL Id: hal-00858560**

**<https://hal.inria.fr/hal-00858560>**

Submitted on 6 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.





# MATHEMATICAL MODELING OF SOIL STRAINS GROWTH AT BATCH CULTURE

AFEM'09

<sup>a,b</sup>M. El Hajji, <sup>a</sup>L. Ranjard, <sup>a</sup>P-A. Maron, <sup>a</sup>V. Nowak, <sup>b,c</sup>J. Harmand and <sup>b</sup>A. Rapaport

<sup>a</sup>UMR-MSE, INRA Dijon, <sup>b</sup>INRA-INRIA MERE Research Team, UMR-ASB, INRA Montpellier, <sup>c</sup>LBE-INRA Narbonne

## Abstract

A mathematical model, validated on experimental data aiming at describing and predicting microbial growth on an essential limited substrate in batch cultures is proposed. This model takes into account viable cell growth, substrate consumption, cell mortality, nonviable cell accumulation in the culture medium and partial dead cell recycling into substrate. A detailed analysis is carried out. Least squares method is used to identify model parameters.

## Material & Methods

The substrate solution was prepared by dissolving glucose (as the only carbon source) in the medium culture (Bergersen medium) at the required concentration. The pH was adjusted to 6.8 due to the high biodegradation efficiency of the strains at this value. Both the medium and the equipment were sterilized before inoculation. Then, the strains previously isolated from soils, were inoculated into the culture medium. The temperature was controlled at 28C and the culture was automatically and continuously shaken. The experiments were carried out in triplicate. The growth of microorganisms on different glucose concentrations was monitored by manual optical density measurements in a spectrometer at 600 nm, where a linear relationship between optical density and concentration of viable and nonviable cells was assumed.

## Modeling and identification of the specific growth rate

Assuming there is initially no dead cell, and that the natural mortality is negligible during the exponential phase, we calculated the regression coefficient in the exponential phase for each glucose concentration. The growth rate of all strains were then identified as functions of glucose concentrations. The best-fit results based on the Haldane equation

$$\mu(s) = \frac{\mu_{max}s}{k_s + s + \frac{s^2}{k_i}}$$

were obtained using the least squares method and are presented hereafter where  $\mu_{max}$  is the maximum growth rate,  $k_s$  is the saturation constant and  $k_i$  is the inhibition coefficient.




Figure 2: Specific growth rates

## Mathematical model

Let  $s, x$  and  $x_d$  denote the substrate, the viable cells and the nonviable cells concentrations, respectively. The model can be written as :

$$\begin{cases} \dot{s} = -\frac{\mu(s)}{k}x + \lambda(1-\delta)mx, \\ \dot{x} = (\mu(s) - m)x, \\ \dot{x}_d = \delta mx, \end{cases} \quad \text{such that } \frac{1}{k} > \lambda(1-\delta) \quad (1)$$

$$s(0) = s_0 > 0, \quad x(0) = x_0 > 0 \quad \text{and} \quad x_d(0) = 0.$$

- The solution of system (1) has positive, bounded components and thus is defined for every positive time  $t$ .
- $\{(s, x, x_d) \in \mathbb{R}_+^3 / ks + x + \frac{1-k\lambda(1-\delta)}{\delta}x_d = ks_0 + x_0\}$  is positively invariant under (1).
- Equilibrium points of system (1) have the following form  $A^* = (s^*, 0, x_d^*)$  such that  $ks^* + \frac{1-k\lambda(1-\delta)}{\delta}x_d^* = ks_0 + x_0$ .  
There exists two values  $s_1^*$  and  $s_2^*$  such  $0 \leq s_1^* < s_2^*$  satisfying  $\mu(s_1^*) = \mu(s_2^*) = m$ .
- If  $s^* < s_1^*$  or  $s_2^* < s^*$ , then the equilibrium point  $A^*$  is semi-stable.
- If  $s_1^* < s^* < s_2^*$ , the equilibrium point  $A^*$  is unstable.
- The equilibrium points set  $\Gamma = \{(s_1^*, 0, x_d^*); x_d^* \geq 0\} \cup \{(s_2^*, 0, x_d^*); x_d^* \geq 0\}$  separate semi-stable equilibrium points set and unstable equilibrium points set one.

## Parameter identification

Available measurements were the substrate concentration and the optical density, modeled as a linear combination of viable and nonviable cells concentrations




$$y = \begin{pmatrix} s \\ \gamma_1 x + \gamma_2 x_d \end{pmatrix}.$$

The least squares method was used to calculate model parameters by minimizing the following criterion :

$$J = \frac{\sigma_1}{2} \sum_{i=1}^n (s(t_i) - s_i)^2 + \frac{\sigma_2}{2} \sum_{i=1}^n (\gamma_1 x(t_i) + \gamma_2 x_d(t) - z_i)^2$$

where  $t_i$  is the time variable,  $s_i$  is the substrate concentration and  $z_i$  is the optical density.  $s(t_i), x(t_i)$  and  $x_d(t_i)$  are the substrate, the viable and the nonviable cells concentrations simulated using the model at time  $t_i, i = 1, \dots, n$  while  $\sigma_1$  and  $\sigma_2$  are weighting coefficients.

## Results



## Conclusions and Perspectives

We have developed and validated simple models in revisiting the way the optical density is modeled. Perspectives include co-cultures modeling in nonhomogeneous media.

## References

- [1] S.J. PIRT (1975) Principles of microbe and cell cultivation. *Blackwell Scientific Publications*, London.

## Acknowledgements

We are grateful to the Laboratoire Microbiologie of Soil and of Environnement (INRA Dijon) for its technical and human support.