



Proteolytic activity of dairy lactic acid bacteria and probiotics as determinant of growth and in vitro angiotensin-converting enzyme inhibitory activity in fermented milk

Osaana N. Donkor, Anders Henriksson, Todor Vasiljevic, Nagendra P. Shah

► To cite this version:

Osaana N. Donkor, Anders Henriksson, Todor Vasiljevic, Nagendra P. Shah. Proteolytic activity of dairy lactic acid bacteria and probiotics as determinant of growth and in vitro angiotensin-converting enzyme inhibitory activity in fermented milk. *Le Lait*, 2007, 87 (1), pp.21-38. hal-00895625

HAL Id: hal-00895625

<https://hal.science/hal-00895625>

Submitted on 11 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Proteolytic activity of dairy lactic acid bacteria and probiotics as determinant of growth and in vitro angiotensin-converting enzyme inhibitory activity in fermented milk

Osaana N. DONKOR^a, Anders HENRIKSSON^b, Todor VASILJEVIC^a,
Nagendra P. SHAH^{a*}

^a School of Molecular Sciences, Victoria University, Werribee Campus, P.O. Box 14428, Melbourne, Vic 8001, Australia

^b DSM Food Specialties Australia Pty Ltd, P.O. Box 83, Moorebank, NSW 1875, Australia

Received 5 June 2006 – Accepted 29 November 2006

Abstract – Two strains each of *Lactobacillus acidophilus* (L10 and La 4962), *Bifidobacterium* spp. (*B. lactis* B94 and *B. longum* BI 536), and *Lactobacillus casei* (L26 and Lc 279), and one strain each of *Streptococcus thermophilus* (St 1342) and *Lactobacillus delbrueckii* ssp. *bulgaricus* (Lb 1466) were assessed for growth characteristics, proteolytic activity and release of in vitro angiotensin-converting enzyme inhibitory peptides in reconstituted skim milk. Single cultures grew well with exception of *Lactobacillus delbrueckii* ssp. *bulgaricus*. Despite slow growth, this culture produced substantial amount of lactic acid, second to *S. thermophilus*. All strains exhibited proteolytic activities with intra- and extracellular specific peptidases including X-prolyl-dipeptidyl aminopeptidase. The latter cleaved proline-containing sequences, which possibly enhanced liberation of various peptides and likely resulted in improved cell growth. The extent of proteolysis varied among strains and appeared to be time dependant. All the cultures released peptides with in vitro ACE-inhibitory activity during growth with *B. longum* BI 536 and *L. acidophilus* L10 having IC₅₀ values of 0.196 and 0.151 mg·mL⁻¹, respectively.

dairy culture / probiotics / proteolytic activity / growth / angiotensin-converting enzyme inhibition (ACE-I)

摘要 – 乳酸菌和益生菌在乳基培养基和发酵乳制品中的生长特性及产生的生物活性肽研究。本文研究了两株嗜酸乳杆菌 (L10, La 4962), 两株双歧杆菌 (*B. lactis* B94, *B. longum* BI 536), 两株干酪乳杆菌 (L26, Lc 279), 一株嗜热链球菌 (St 1342) 和一株德氏乳杆菌保加利亚亚种 (Lb 1466) 在还原脱脂乳中的生长特性、蛋白水解酶的活性以及具有抑制血管紧张素转移酶 (ACE) 活性的生物活性肽的释放。除了德氏乳杆菌保加利亚亚种外, 所有的菌株在乳基培养基中的生长性能良好, 尽管这株菌在乳基培养基中的生长速度较慢, 但却能够产生大量的乳酸, 其产乳酸的量仅次于嗜热链球菌。所有的菌株均能分泌出具有蛋白水解活性的胞内和胞外肽酶, 如 X-脯氨酰-二肽酰基-氨基肽酶。该酶能够选择性地从肽的 N 端水解含有脯氨酸的肽链, 释放出小分子的肽, 而这些小分子肽进一步促进了细胞的生长。各菌株的蛋白水解程度不同, 并且每菌株的蛋白水解活性随着培养时间的增加而提高。所有菌株在发酵乳中均产生能够抑制血管紧张素转移酶活性的生物活性肽。其中长双歧杆菌 BI 536 和嗜酸乳杆菌 L10 产生的这种生物活性肽的活性最高, 其 IC₅₀ 值分别为 0.196 和 0.151 mg·mL⁻¹。

乳基培养基 / 益生菌 / 蛋白水解活力 / 生长特性 / 血管紧张素转移酶抑制作用

* Corresponding author (通讯作者): Nagendra.Shah@vu.edu.au

Résumé – L’activité protéolytique des bactéries lactiques et probiotiques laitiers détermine la croissance et l’activité inhibitrice de l’enzyme de conversion de l’angiotensine (ACE) dans le lait fermenté. Deux souches de *Lactobacillus acidophilus* (L10 et La 4962), de *Bifidobacterium* spp. (*B. lactis* B94 et *B. longum* BI 536), et de *Lactobacillus casei* (L26 et Lc 279), et une souche de *Streptococcus thermophilus* (St 1342) et de *Lactobacillus delbrueckii* ssp. *bulgaricus* (Lb 1466) ont été étudiées pour leurs caractéristiques de croissance, activité protéolytique et libération de peptides inhibant l’ACE in vitro dans du lait écrémé reconstitué. Les cultures pures croissaient bien à l’exception de *Lactobacillus delbrueckii* ssp. *bulgaricus*. Malgré une croissance lente, cette culture produisait des quantités substantielles d’acide lactique, venant en seconde position après *S. thermophilus*. Toutes les souches présentaient des activités protéolytiques avec des peptidases intra- et extracellulaires dont une X-prolyl-dipeptidyl aminopeptidase. Cette enzyme coupait les séquences contenant une proline, ce qui améliorerait potentiellement la libération de peptides variés et, en conséquence, probablement la croissance cellulaire. L’importance de la protéolyse était variable selon les souches, mais apparaissait dépendante du temps. Toutes les cultures libéraient des peptides ayant une activité inhibitrice de l’ACE in vitro au cours de la croissance, avec pour *B. longum* et *L. acidophilus* L10 des valeurs d’IC₅₀ de 0,196 et de 0,151 mg·mL⁻¹ respectivement.

bactérie lactique / probiotique / activité protéolytique / croissance / activité inhibitrice de l’ACE

1. INTRODUCTION

Incorporation of probiotic organisms such as *Lactobacillus acidophilus*, *Bifidobacterium* spp., and *L. casei* in fermented dairy products provides a potential to improve health status of consumers. A growing public awareness of diet related health issues and mounting evidence regarding health benefits of probiotics have increased consumers demand for foods containing probiotic organisms. Probiotics are defined by the Food and Agricultural Organization of the United Nations (FAO) and World Health Organization (WHO) as “live microorganisms, which when administered in adequate amounts confer a health benefit on the host” [14]. A number of health benefits of probiotic organisms has been suggested and described in numerous reviews [25, 37].

However, certain criteria during production of probiotic foods need to be satisfied to ensure maximization of therapeutic properties of probiotics. These include the incorporation of sufficient numbers of probiotic microorganisms into the product, maintenance of viable populations during shelf life of the food and subsequent survival of these organisms through the gastrointestinal tract (GIT) [32, 51]. Fermented dairy foods present ideal delivery systems for probiotics to the human GIT, since they

may provide a favourable environment, which promotes growth and enhances viability of these organisms [25]. On the other hand, several factors such as low pH, presence of hydrogen peroxide and dissolved oxygen as well as the state of culturing and storage conditions may affect the survival of probiotic bacteria in fermented dairy products [36, 38, 44].

Lactic acid bacteria (LAB) including probiotic organisms are fastidious in nature, requiring numerous essential growth factors. Milk, although a rich growth medium, contains low concentration of free amino acids and peptides to efficiently support growth of LAB [39, 48]. In response to this limitation, LAB have developed a complex system of proteinases and peptidases, which enable them to utilise casein as an additional source of organic nitrogen [41]. On the other hand, Klaver et al. [20] reported that *Bifidobacterium* strains were not as proteolytic as other LAB. This may explain why *Bifidobacterium* spp. grows slowly in milk and may require supplementation of peptides and amino acids from external sources [11].

The proteolytic activities of LAB including yoghurt starter bacteria and probiotic organisms have been studied extensively and proteolytic enzymes have been isolated and characterised [22, 39, 50]. Such an interest has been derived from the importance of

proteolytic system of LAB in cheese ripening and rapid growth in milk during fermentation as well as improved survival during storage [12]. This ability of dairy LAB and probiotics has become even more important upon realising that a range of bioactive peptides may be liberated due to microbial action. Biologically active peptides are generated during milk fermentation by proteolytic enzymes produced by various LAB such as *L. helveticus*, *L. lactis* subsp. *cremoris* FT4 and *L. delbrueckii* ssp. *bulgaricus* SS1 [18, 27]. These biologically active peptides include hypotensive peptides which inhibit angiotensin I-converting enzyme (ACE), opioid agonist and antagonist peptides, and mineral binding, immunomodulatory, antibacterial, and antithrombotic peptides [30, 35]. Angiotensin I-converting enzyme regulates blood pressure via formation of vasopressor angiotensin II from angiotensin I [40]. Inhibition of ACE mainly results in an overall hypotensive effect. High proteolytic activity is expected to promote good cell growth and ACE-inhibitory activity in fermented milk.

As stated, the proteolytic activity of dairy cultures is very important governing factor playing a major role in various cellular and physiological processes. Therefore, the aims of our study were to assess the proteolytic activity of selected dairy lactic acid bacteria and probiotic organisms as determinant of growth and in vitro ACE inhibitory activity in fermented milk.

2. MATERIALS AND METHODS

2.1. Bacterial cultures

L. acidophilus LAFTI® L10, *B. lactis* LAFTI® B94 and *L. casei* LAFTI® L26 were obtained from DSM Food Specialties (Moorebank, NSW, Australia) and have been reported to have probiotic properties [8]. *S. thermophilus* St 1342, *L. delbrueckii* subsp. *bulgaricus* Lb 1466, *L. acidophilus* La 4962, *B. longum* B1 536 and *L. casei* Lc 279 were supplied by the Victoria University Culture Collection (Werribee, Australia). Each strain was propagated in de Mann Rogosa Sharpe (MRS) broth (Oxoid,

West Heidelberg, Australia) at 37 °C with the exception of *L. delbrueckii* ssp. *bulgaricus* Lb 1466 which was propagated at 42 °C. For propagation of *Bifidobacterium*, sterile MRS broth was supplemented with 0.05% L-cysteine-hydrochloride to provide anaerobic condition and stimulate their growth [29]. After three successive transfers of 20 h incubation each, the activated organisms were used for the preparation of the pre-inocula for further experiments. The pre-inocula were prepared by transferring 1% (v/v) of activated culture to 10 mL aliquots of reconstituted skim milk (RSM) supplemented with 2% glucose and 1% yeast extract.

2.2. Proteolytic systems of LAB

2.2.1. Preparation of intracellular and cell wall extracts

Individual cultures of *L. acidophilus* L10, *Bifidobacterium* B94, *L. casei* L26, *S. thermophilus* St 1342, *L. delbrueckii* ssp. *bulgaricus* Lb 1466, *L. acidophilus* La 4962, *Bifidobacterium* B1 536 and *L. casei* Lc 279 were propagated three successive times in RSM and finally in MRS broth according to the methods previously reported [39, 50]. The latter was performed to prevent a carryover of milk proteins and their interference with the assay. Thus after two subculturing in MRS medium for 18 h at 37 °C, cultures were cultivated separately in 100 mL batches of MRS broth at 42 °C. The cells were then harvested by centrifugation at 4000× g at 4 °C for 30 min at the end of the logarithmic growth phase at approximately 20 h which was determined by measuring the absorbance at 600 nm. The supernatant was designated as the cell-free extracellular enzymatic extract (EE). The cell pellet obtained was washed twice with 0.9% (w/v) NaCl solution and resuspended in 10% volume of original growth medium with 0.05 M Tris-HCl buffer, pH 8.5. The resulting cell dispersion was sonicated for 5 min at 30 s intervals at 4 °C. The supernatant obtained after centrifugation at 4000× g for 30 min at 4 °C was designated as the intracellular enzymatic extract (IE).

2.2.2. Determination of protein concentration

The protein content of the EE and IE extracts was estimated using the method of Bradford [3]. A 0.1-mL aliquot of the enzyme solution and 3-mL aliquot of the Bradford reagent (Sigma) were vortexed gently to mix thoroughly and the samples were incubated at room temperature for 30 min after which the absorbance was measured at 595 nm. Bovine serum albumin (Sigma) was used as a standard. The protein concentration of the samples was determined by comparing the net absorbance values obtained at 595 nm against the standard curve.

2.2.3. Enzyme assay

Endopeptidase and tripeptidase activities of the EE and IE extracts were detected by thin-layer chromatography (TLC) according to the method of Shihata and Shah [39] and Tan and Konings [45] with some modifications. Each reaction mixture contained 60 μL of a 2 $\text{mmol}\cdot\text{L}^{-1}$ substrate (Gly-Ala-Tyr, Gly-Leu-Phe and Bradykinin) in 20 $\text{mmol}\cdot\text{L}^{-1}$ Tris-HCl pH 7.0 and 20 μL of extract. The reaction mixture was incubated for 60 min at 37 °C and was stopped by adding 10 μL of 30% acetic acid and cooling to 4 °C. Ten microlitres of the mixture were spotted onto a precoated silica gel 60 plate (50 \times 100 \times 0.25) mm (Alltech Associates Pty. Ltd., Baulkham Hills, Australia). A 4:1:1 (v/v/v) mixture of *n*-butanol:acetic acid:water was used as the mobile phase. As a control, 2 $\text{mmol}\cdot\text{L}^{-1}$ of each standard peptide was also spotted onto the plate. Silica gels were stained by spraying with 0.1% (w/v) ninhydrin in 99% ethanol. Peptides and amino acids became visible after incubating the plate in an oven for 5 min at 80 °C.

The proteolytic strains of probiotic and yoghurt bacteria were assessed for selected peptidolytic activities using Pro-Ile, Leu-Tyr, Leu-Gly, Ala-Met, and Ala-His as well as Gly-Ala-Tyr, Gly-Leu-Phe and bradykinin as substrates according to the method of Shihata and Shah [39] and Wohlrab and Bockelmann [50]. The reaction mixture contained 10 μL of enzyme solution, 424 μL of 50 $\text{mmol}\cdot\text{L}^{-1}$ Tris-HCl pH 7.5, 50 μL of

22 $\text{mmol}\cdot\text{L}^{-1}$ substrate, 25 μL of peroxidase (5 $\text{mg}\cdot\text{mL}^{-1}$ in 0.8 $\text{mol}\cdot\text{L}^{-1}$ $(\text{NH}_4)_2\text{SO}_4$), 25 μL of L-amino acid oxidase (2 $\text{mg}\cdot\text{mL}^{-1}$ in distilled water) and 25 μL of *o*-dianisidine (11.5 $\text{mmol}\cdot\text{L}^{-1}$). The reaction mixture was incubated at 50 °C for 20 min and was stopped by the addition of 50 μL of dithiothreitol (120 $\text{mmol}\cdot\text{L}^{-1}$). Oxidation of *o*-dianisidine coupled to substrate hydrolysis resulted in an increase in brown colour which was measured at 436 nm. Enzyme activity was calculated using a molar absorbance coefficient of 8100 $\text{mol}^{-1}\cdot\text{cm}^{-1}$ and defined as the amount of enzyme required to oxidise 1 μmol of *o*-dianisidine per min under assay conditions [50]. The specific activity was expressed as units per milligram of protein.

2.2.4. Assessment of X-prolyl-dipeptidyl aminopeptidase activity

X-prolyl-dipeptidyl aminopeptidase (PepX) activity of dairy cultures is a very important characteristic due to high proline content in milk protein. This activity was assayed as described previously [29] with glycyl-prolyl *p*-nitroanilide (Sigma) (Gly-Pro-pNA) as the substrate with some modifications. The incubation mixture contained 50 μL of 6.4 $\text{mmol}\cdot\text{L}^{-1}$ of substrate, 2.85 mL of 50 $\text{mmol}\cdot\text{L}^{-1}$ Tris-HCl buffer pH 7.0, and 100 μL of cell-free intracellular extract (IE) in 50 $\text{mmol}\cdot\text{L}^{-1}$ Tris-HCl buffer pH 7.0. The mixture was incubated at 37 °C for 20 min and the reaction was stopped by adding 500 μL of 30% acetic acid. The extent of hydrolysis was measured with the Cary IE UV/visible spectrophotometer (Varian Australia Pty. Ltd., Melbourne, Australia) at 410 nm. The same experiment was also performed with cell-free extracellular (EE) enzymatic extracts. One unit of enzyme activity was defined as the amount of enzyme required to release 1 μmol of *p*-nitroanilide per min under assay conditions. The enzyme activity reported is expressed as specific PepX activity defined as units of enzyme activity per milligram of proteins. A unit of activity is derived from the enzyme content required to liberate 1 μmol of *p*-nitroanilide per min under assay conditions.

2.3. Preparation of fermented milk

Individually activated cultures of *L. acidophilus* (L10 and La 4962), *Bifidobacterium* spp. (*B. lactis* B94 and *B. longum* B1 536), *L. casei* (L26 and Lc 279), *S. thermophilus* (St 1342) and *L. delbrueckii* ssp. *bulgaricus* (Lb 1466) in the form of pre-inoculate (Sect. 2.1) were used for milk fermentations. Each organism was transferred into sterile RSM (12%) to obtain approximately 10^8 colony forming units (CFU) per mL. The inoculated milk was incubated at 37 °C for 20 h, and this step was repeated twice to prevent any carry overs (i.e. yeast extract) that would likely interfere with the assays [23]. Batch fermentations were further carried out with each culture in 12% (w/w) RSM. Eight batches of 250 mL aliquots of RSM were aseptically prepared with 1% (v/v) of each of the cultures and incubated at 42 °C for 24 h. The fermentation experiments were performed in triplicate. A control consisted of uninoculated RSM. During fermentation, aliquots of each batch were taken at 0, 3, 6, 9, 12 and 24 h to monitor cell growth, organic acid production and pH changes. Proteolysis was monitored at 0, 6, 12 and 24 h.

2.4. Cell growth

The methods of Leclerc et al. [23] and Ravula and Shah [32] were used to assess the cell growth. One gram of each batch, aseptically sampled at 0, 3, 6, 9, and 24 h, was 10-fold serially diluted (10^3 to 10^7) in 0.15% sterile peptone (Oxoid) and water diluent. Enumeration of the bacteria was performed on MRS agar (Amyl media, Dandenong, Australia) using the pour plate technique as described previously [10, 47]. Anaerobic jars and gas generating kits (Anaerobic system BR 38, Oxoid Ltd., Hampshire, England) were used for creating anaerobic conditions. Plates in duplicate were incubated for 48 h at 37 °C for *L. acidophilus*, *L. casei*, and *Bifidobacterium* spp., 48 h at 42 °C for *L. delbrueckii* ssp. *bulgaricus* and aerobically for 48 h at 37 °C for *S. thermophilus*. All enumeration techniques followed protocols reported previously [10, 47]. Plates with 25–250 colonies were counted and recorded as CFU·mL⁻¹ of

the fermented milk. The pH changes of batches were monitored before and during fermentation at 0, 3, 6, 9, 12 and 24 h using a pH meter (HANNA Instruments 8417, Singapore).

2.5. Organic acids

Organic acid contents were measured following the method of Shah and Ravula [38] with some modifications. Three millilitre aliquots of fermented milk were mixed with 50 µL of 15.5 mol·L⁻¹ nitric acid and then diluted with 1.0 mL of 0.01 mol·L⁻¹ H₂SO₄ solution. The resulting mixture was centrifuged for 30 min at 14 000× g using an Eppendorf centrifuge (Model 5415C, Crown Scientific, Melbourne, Australia) to precipitate the protein. The supernatant was filtered through a 0.20-µm membrane filter (FP point, Schleicher & Schuell, Dassel, Germany) into an HPLC vial for the determination of organic acids. The separation of organic acids was achieved using a Varian HPLC (Varian Analytical Instruments, CA, USA) fitted with an Aminex HPX - 87H, 300×7.8 mm ion exchange column (Biorad Life Science Group, Hercules, CA, USA) and a guard column maintained at 65 °C. Using 0.01 mol·L⁻¹ H₂SO₄ as the mobile phase, the flow rate was maintained at 0.6 mL·min⁻¹. A UV/visible detector was used at 220 nm. A 25-µL injection volume was used for both samples and standards with the retention time of 12.15 (L(+)-lactic acid) and 14.40 (glacial acetic acid) min. Quantification of acetic and lactic acids was performed as described previously [13].

2.6. Proteolytic activity in fermented milk

The degree of proteolysis during fermentation of milk was determined by measuring the release of free NH₃ groups following the *o*-phthaldialdehyde (OPA) method [6, 12]. An aliquot of 2.5 mL from each medium was mixed with 5 mL of 0.75% trichloroacetic acid (TCA) and the mixture was filtered using Advantec 231 filter (M.F.S. Inc., Dublin, CA, USA). The filtrate (150 µL) was added to 3 mL of OPA reagent and after 2 min at room temperature

(20 °C), absorbance of the solution was measured by a spectrophotometer (LKB NOVASPEC II Pharmacia, LKB Biochrom, England) at 340 nm. The proteolytic activity of these bacterial cultures was expressed as the absorbance of free amino groups measured at 340 nm. A relative degree of proteolysis was determined as the difference between proteolytic activity in fermented milk to that of unfermented milk. All the analyses were carried out in triplicate.

2.7. In vitro inhibition of angiotensin I-converting enzyme (ACE inhibitory activity)

ACE inhibitory activity was measured according to Cushman and Cheung [9] with some modifications as described previously [13]. Briefly, 20 g of fermented milk were mixed with 5 mL of 0.75% TCA and the mixture was centrifuged at 4000× *g* for 30 min at 4 °C. The pH of the supernatant was adjusted to 8.3 and 20 µL of the filtrate was added to an aliquot of 200 µL hippuryl-histidyl-leucine (Sigma) dissolved in sodium borate buffer of pH 8.5. Aliquot of 60 µL of the buffer was added to the mixture to maintain the pH of the reaction solution in the range between 8.1 and 8.3 [49]. The reaction was initiated by adding 20 µL of ACE solution followed by incubation at 37 °C for 30 min. The reaction was stopped by adding 250 µL of 1 mol·L⁻¹ HCl and 1.7 mL of ethyl acetate. The ethyl acetate layer (1.4 mL) was evaporated to dryness and the residue containing hippuric acid was dissolved in 1 mL of deionised water. The experimental procedure was repeated with unfermented RSM as a control prepared following the procedures as described above (appropriate MRS transfer into sterile RSM supplemented with yeast extract, followed by subsequent transfers and dilutions in RSM). The absorbance of the solution was measured spectrophotometrically using Cary IE UV/visible spectrophotometer (Varian) at 228 nm against water as a blank. The extent of inhibition was calculated as follows:

$$\text{ACE-inhibitor activity (\%)} = 1 - \left[\frac{C - D}{A - B} \right] \times 100$$

where *A* = absorbance in the presence of ACE and the buffer only;

B = absorbance in the presence of buffer only;

C = absorbance in the presence of ACE, ACE-inhibitory component and the buffer;

D = absorbance in the presence of ACE-inhibitory component and the buffer.

Additionally ACE inhibition was expressed as an IC₅₀, defined as the amount of sample required to inhibit 50% of the original ACE activity. The IC₅₀ was determined using graphical extrapolation by plotting ACE inhibition as a function of different sample concentrations with known protein contents. All the analyses were carried out in triplicate.

2.8. Statistical analysis

The experiment was independently replicated three times in a completely randomized design. All results obtained were analysed as a split plot in time design with strains and time as main factors, using the general linear model (GLM) procedure of the SAS System [34]. The univariate ANOVA test was validated by fulfilling Huynh-Feldt (H-F) condition [24]. Where appropriate, one-way ANOVA and correlational analyses were employed and the multicomparison of means was assessed by Tukey's test. The statistical level of significance was preset at 0.05 (*P* < 0.05).

3. RESULTS AND DISCUSSION

3.1. Assessment of extracellular and intracellular peptidase activity

Table I shows the activities of extracellular and intracellular peptidases towards various peptides detected by TLC. All dairy cultures tested appeared to have produced enzymes capable of hydrolysing large biologically active peptide, bradykinin, at both the extracellular and intracellular levels. The enzymes involved in the hydrolysis may include endopeptidases PepE and PepO and aminopeptidase PepP [5]. PepE

Table I. Aminopeptidase activities of single cultures: *S. thermophilus* St 1342, *L. helveticus* subsp. *bulgaricus* Lb 1466, *L. acidophilus* (L10 and La 4962), *B. lactis* B94, *B. longum* BI 536, *L. casei* (L26 and Lc 279).

Substrates	Aminopeptidase activity	
	EE	IE
Tripeptides		
Gly-Ala-Tyr	+	+
Gly-Leu-Phe	–	+
Oligopeptide		
Bradykinin	+	+

Hydrolysis of peptides was analysed by TLC; + = hydrolysis; – = no hydrolysis; EE = cell wall extract; IE = intracellular extract.

has been identified to hydrolyse the N-terminal and C-terminal blocked substrate N-benzoyl-Phe-Val-Arg-pNA [15]. PepE is capable of hydrolysing bradykinin (Arg-Pro-Pro-Gly-Phe-Ser-Pro-Phe-Arg) at the Gly4-Phe5 bond [5]. Similarly, PepO hydrolysis oligopeptides ranging in length from five to thirty-five including bradykinin [43]. Furthermore, PepP liberates the N-terminal amino acid from peptides with general specificity for Xaa-Pro-Pro-(Yaa)n sequences with high activity reported for peptides ranging from three to nine residues (i.e. bradykinin) but did not hydrolyse dipeptides [26]. Identified tripeptidase, PepT, has been reported to show no activity for any di-, tetra-, or larger oligopeptides and therefore exhibit strict specificity for tripeptides [5]. Consequently, the tripeptide substrates Gly-Ala-Tyr and Gly-Leu-Phe tested might have been hydrolysed by PepT. In addition to PepT, aminopeptidases PepC and PepP have also been reported to hydrolyse a variety of tripeptides with uncharged or basic residues in the amino acid terminal position [5]. However, with the substrate ending with a C-terminal of phenylalanine, Gly-Leu-Phe, the hydrolysis only occurred intracellularly, suggesting the location of these aminopeptidases. This was consistent with the findings of Shihata and Shah [39] and Tan and Konings [45]. The intracellular aminopeptidases are capa-

ble of cleaving N-terminal amino acids from a wide range of peptides differing both in size and composition to release amino acids and peptides [21].

Specific tripeptidase and endopeptidase activities of yoghurt culture and selected strains of probiotic organisms assessed towards Gly-Ala-Tyr, Gly-Leu-Phe and oligopeptide bradykinin as substrates is shown in Table II. As opposed to other strains, the hydrolysis of all oligopeptides by *L. acidophilus* La 4962 was significantly ($P < 0.05$) higher at the IE level. This suggested that *L. acidophilus* La 4962 may have produced substantial amounts of PepT and PepE than the other strains which caused the hydrolysis of Gly-Ala-Tyr, Gly-Leu-Phe and oligopeptide bradykinin. Eventhough, all organisms showed specific tripeptidase activity mainly at the IE level, the extent of which was greatly ($P < 0.05$) strain specific confirmed previous reports (Tab. II) [21, 45, 46]. The low ($P < 0.05$) EE level of enzyme activity shows that tripeptidase and endopeptidase activities may solely take place intracellularly. The observed extracellular enzyme activity may be due to cell lysis and release of intracellular enzymes into the medium [2, 7]. The EE values may further indicate variations in the extent of lysis which may be high with some cultures and likely lead to erroneous conclusions [2].

X-prolyl-dipeptidyl aminopeptidase, PepX, activity of cell-free extract of individual cultures of bacteria is presented in Figure 1. This enzyme is of great importance for the selection of dairy cultures due to high proline content in caseins. All the studied strains exhibited PepX activity towards glycyl-prolyl *p*-nitroanilide (Gly-Pro-pNA) as substrate at both IE and EE cell-free extracts. The IE level of X-prolyl-dipeptidyl aminopeptidase activity was significantly ($P < 0.05$) higher than that at the EE level. This shows that proline-containing peptides were mainly hydrolysed intracellularly by the enzyme (Fig. 1). Kunji et al. [21] reported the presence of PepX in all species of LAB. Pan et al. [29] similarly identified X-prolyl-dipeptidyl aminopeptidase activity from IE cell-free extract of *L. helveticus* in the hydrolysis of skimmed

Table II. Tripeptidase specific activity of yoghurt culture and selected strains of probiotic organisms after 20 h of single culture fermentation in MRS at 42 °C.

Culture	Tripeptidase specific activity, U·mg ⁻¹ protein		
	Substrates		
	Gly-Ala-Tyr	Bradykinin	Gly-Leu-Phe
<i>S. thermophilus</i> St 1342			
EE	370.79 ± 1.14 ^a	349.32 ± 0.38 ^b	359.97 ± 0.72 ^c
IE	555.09 ± 0.40 ^a	534.32 ± 0.70 ^b	551.30 ± 0.91 ^c
<i>L. delbrueckii</i> ssp. <i>bulgaricus</i> Lb 1466			
EE	441.06 ± 6.91 ^a	500.88 ± 0.96 ^b	449.36 ± 0.46 ^a
IE	770.45 ± 0.42 ^a	671.99 ± 1.57 ^b	700.76 ± 1.52 ^c
<i>L. acidophilus</i> L10			
EE	218.20 ± 2.30 ^{Aa}	246.80 ± 0.51 ^{Ab}	252.11 ± 0.51 ^{Ab}
IE	557.47 ± 0.46 ^{Aa}	521.10 ± 1.18 ^{Ab}	485.40 ± 1.66 ^{Ac}
<i>L. acidophilus</i> La 4962			
EE	704.52 ± 32.73 ^{aB}	611.14 ± 7.76 ^{Bab}	675.21 ± 0.29 ^{Bc}
IE	2844.43 ± 3.45 ^{aB}	2921.26 ± 2.84 ^{Bb}	3021.52 ± 1.13 ^{Bc}
<i>B. lactis</i> B94			
EE	671.96 ± 3.63 ^{Aa}	180.72 ± 1.46 ^{Ab}	209.78 ± 2.81 ^{Ac}
IE	1516.73 ± 9.44 ^{Aa}	1360.36 ± 3.12 ^{Ab}	1263.60 ± 1.80 ^{Ac}
<i>B. longum</i> Bl 536			
EE	628.39 ± 2.57 ^{Ba}	656.43 ± 2.08 ^{Bb}	583.77 ± 2.19 ^{Bc}
IE	913.84 ± 0.96 ^{Ba}	741.68 ± 0.34 ^{Bb}	724.69 ± 0.91 ^{Bc}
<i>L. casei</i> L26			
EE	289.78 ± 2.74 ^{Aa}	414.14 ± 4.20 ^{Ab}	726.36 ± 2.11 ^{Ac}
IE	899.92 ± 1.05 ^{Aa}	907.47 ± 9.60 ^{Aab}	927.82 ± 0.23 ^{Ab}
<i>L. casei</i> Lc 279			
EE	470.76 ± 29.51 ^{Ba}	573.84 ± 23.27 ^{Ba}	467.94 ± 49.42 ^{Ba}
IE	646.27 ± 0.80 ^{Ba}	644.55 ± 0.32 ^{Ba}	652.03 ± 0.64 ^{Bb}

Results presented as a mean of three observations. Significant when $P < 0.05$; tripeptidase activity expressed as specific activity defined as units of enzyme activity per milligram of protein in crude cellular extract.

EE = extracellular cell wall extract; IE = intracellular extract.

^{abc} Means in the same row with different small letter superscripts are significantly different;

^{AB} means in the same column for particular strains with different capital letter superscripts are significantly different.




Figure 1. X-prolyl-dipeptidyl aminopeptidase activity of cell-free intracellular (IE) and extracellular (EE) enzymatic extracts of individual cultures of bacteria (*S. thermophilus* St 1342, *L. delbrueckii* subsp. *bulgaricus* Lb 1466, *L. acidophilus* L10, *B. lactis* B94 and *L. casei* L26, *L. acidophilus* La 4962, *B. longum* BI 536 and *L. casei* Lc 279) using Gly-Pro-pNA as the substrate in 50 mmol·L⁻¹ Tris-HCl buffer pH 7.0 at 37 °C. Aminopeptidase specific activity is defined as units (U) of enzyme activity per mg of protein in crude cellular extract. One unit of enzyme activity was defined as the amount of enzyme required to release 1 µmol of *p*-nitroanilide per min under assay conditions (Error bars represent a pooled standard error of the mean).

milk proteins. Fernandez-Espla et al. [16] detected the presence of various proline-specific peptidases in the cell-free extract of *L. casei* ssp. *casei* IFPL 731 and further reported that the enzyme usually cleaves N-terminal X-Pro dipeptides from tri- and oligopeptides. The aminopeptidase activities of microorganisms included in our study may contribute to the production of free amino acids in a fermenting medium as growth factors and peptides, which may have ACE-inhibitory activity [29].

Aminopeptidase activities of all examined strains towards dipeptide substrates are shown in Table III. The strains studied exhibited on average 71% dipeptidase activity at the IE level showing preference towards N-terminal hydrophobic/uncharged residues resulting in hydrolysis of free amino acids for growth. Leu-Gly, Leu-Tyr, Ala-Met and Ala-His were probably hydrolysed by either one or both aminopeptidases (PepC and PepN) of the studied microorganisms. These enzymes, which have been identified in LAB, were reported to exhibit broad specificity for residues that are basic,

acidic, hydrophobic/uncharged and aromatic substrates [5]. Wohlrab and Bockelmann [50] also observed that all dipeptidases were active towards numerous dipeptides. PepC and PepN, on the other hand, have been shown to be inactive towards Pro-Xaa dipeptides [5]. Since PepX has specificity for removal of proline-containing dipeptides (X-Pro ↓ Y...) from the N-termini of peptides [5], it may be assumed that PepX was likely not active towards Pro-Ile. Therefore the probable presence of aminopeptidase PepR in the studied strains might have hydrolysed the tested substrate Pro-Ile from the hydrophobic uncharged C-terminal-Ile. In general, all the organisms showed appreciable aminopeptidase activity at the IE level towards Ala-Met, Leu-Tyr, Leu-Gly, Ala-His, and Pro-Ile, in accordance with the findings of Shihata and Shah [39]. Contrary to our findings, Tan et al. [46] observed very low or no intracellular dipeptidase activity towards Pro-containing dipeptides. Extracellular dipeptidase activity observed in our study was significantly ($P < 0.05$) lower for all substrates than that at the IE level. *Bifidobacterium* spp. showed

Table III. Dipeptidase specific activities of single strains of selected probiotic organisms and yoghurt culture after 20 h of individual fermentations in MRS at 42 °C.

Culture	Dipeptidase specific activity, U·mg ⁻¹ protein				
	Substrates				
	Leu-Gly	Leu-Tyr	Pro-Ile	Ala-Met	Ala-His
<i>S. thermophilus</i>					
St 1342					
EE	90.69 ± 10.78 ^a	82.20 ± 22.34 ^a	71.58 ± 0.30 ^a	92.81 ± 14.56 ^a	93.72 ± 17.36 ^a
IE	271.35 ± 0.99 ^a	329.20 ± 4.36 ^b	317.26 ± 0.64 ^b	283.51 ± 2.01 ^a	283.05 ± 1.89 ^a
<i>L. delbrueckii</i> ssp. <i>bulgaricus</i>					
Lb 1466					
EE	101.94 ± 15.08 ^a	84.68 ± 4.17 ^a	89.75 ± 30.13 ^a	72.76 ± 33.07 ^a	76.63 ± 27.36 ^a
IE	265.66 ± 4.62 ^a	282.77 ± 1.31 ^{ab}	281.17 ± 2.48 ^a	239.77 ± 2.99 ^{bc}	335.36 ± 3.53 ^{bcd}
<i>L. acidophilus</i>					
L10					
EE	81.36 ± 5.52 ^{Aa}	105.17 ± 16.29 ^{Aa}	131.99 ± 1.38 ^{Aa}	86.79 ± 20.34 ^{Aa}	114.81 ± 15.37 ^{Aa}
IE	323.73 ± 2.99 ^{Aa}	234.27 ± 2.99 ^{Ab}	247.54 ± 5.39 ^{Ab}	318.95 ± 2.04 ^{Aa}	292.01 ± 1.27 ^{Abc}
<i>L. acidophilus</i>					
La 4962					
EE	241.78 ± 35.50 ^{Ba}	297.55 ± 5.04 ^{Ba}	270.50 ± 43.51 ^{Ba}	277.85 ± 25.38 ^{Ba}	207.38 ± 28.73 ^{Ba}
IE	1127.62 ± 4.69 ^{Ba}	1097.02 ± 2.84 ^{Ba}	1028.01 ± 6.41 ^{Bb}	1081.39 ± 17.91 ^{Ba}	917.33 ± 5.33 ^{Bbc}
<i>B. lactis</i>					
B94					
EE	114.99 ± 17.53 ^{Aa}	56.25 ± 4.23 ^{Aa}	122.18 ± 27.87 ^{Aa}	107.80 ± 32.97 ^{Aa}	86.56 ± 18.95 ^{Aa}
IE	407.80 ± 4.13 ^{Aa}	469.81 ± 4.01 ^{Ab}	578.82 ± 6.89 ^{Abc}	518.18 ± 3.59 ^{Abcd}	385.31 ± 2.36 ^{Aa}
<i>B. longum</i>					
BI 536					
EE	103.41 ± 32.51 ^{Aa}	60.19 ± 20.80 ^{Aa}	129.68 ± 28.28 ^{Aa}	129.35 ± 25.96 ^{Aa}	66.17 ± 22.87 ^{Aa}
IE	340.20 ± 4.47 ^{Ba}	362.34 ± 5.40 ^{Ba}	320.63 ± 2.89 ^{Bab}	355.82 ± 5.08 ^{Ba}	311.36 ± 12.19 ^{Bab}
<i>L. casei</i>					
L26					
EE	124.17 ± 3.53 ^{Aa}	64.73 ± 6.38 ^{Aa}	87.76 ± 15.02 ^{Aa}	101.14 ± 5.18 ^{Aa}	67.53 ± 30.21 ^{Aa}
IE	417.04 ± 14.67 ^{Aa}	277.57 ± 1.87 ^{Ab}	280.31 ± 0.82 ^{Ab}	307.98 ± 1.19 ^{Ab}	313.01 ± 8.77 ^{Ab}
<i>L. casei</i>					
Lc 279					
EE	63.50 ± 33.45 ^{Ba}	106.71 ± 18.19 ^{Ba}	92.63 ± 19.45 ^{Aa}	91.32 ± 2.47 ^{Aa}	78.23 ± 0.87 ^{Aa}
IE	243.10 ± 0.61 ^{Ba}	290.32 ± 1.74 ^{Ab}	292.04 ± 0.98 ^{Ab}	288.61 ± 2.45 ^{Ab}	286.15 ± 2.33 ^{Ab}

Results presented as a mean of three observations. Significant when $P < 0.05$; dipeptidase activity expressed as specific activity which is defined as units of enzyme activity per milligram of protein in crude cellular extract.

EE = extracellular cell wall extract; IE = intracellular extract.

^{abcd} Means in the same row with different small letter superscripts are significantly different;

^{AB} means in the same column for particular strains with different capital letter superscripts are significantly different.




Figure 2. Proteolytic activity by single strains of bacteria (*S. thermophilus* St 1342, *L. delbrueckii* subsp. *bulgaricus* Lb 1466, *L. acidophilus* L10, *B. lactis* B94 and *L. casei* L26, *L. acidophilus* La 4962, *B. longum* Bl 536 and *L. casei* Lc 279) during 24 h fermentation in RSM at 42 °C (Error bars represent a pooled standard error of the mean SEM = 0.03 absorbance units).

similarities in their intracellular dipeptidase activities, however, *B. lactis* B94 exhibited higher Pro-Ile and Ala-Met hydrolysis than that of *B. longum* Bl 536 (Tab. III). This may explain the similarity in their growth pattern (Fig. 3). Several studies [21, 45, 46] have reported dipeptidase activity solely at the IE level. It is interesting to note that there is little information available in the literature on aminopeptidase activities of *Bifidobacterium* spp. Therefore this study is important in this area of enzyme activity and further work in understanding their proteolytic activity and growth in milk is underway.

3.2. Proteolytic activity

After analysing segments of proteolytic system in the defined medium, we assessed the proteolytic activities of selected LAB in RSM as presented in Figure 2. During fermentation, milk proteins were hydrolysed by LAB proteinases and peptidases resulting in an enhanced amount of free amino groups and peptides (Fig. 2). Juillard et al. [19] reported that the level of free amino acids and peptides in milk is low therefore LAB depend on a proteolytic system that allows for an efficient degradation of milk

proteins. The presence of aminopeptidases is important for the release of amino acids for growth by microorganisms through the hydrolysis of peptides in the growth medium. Christensen and Steele [4] demonstrated that the loss of selected aminopeptidase (PepC, PepN, and PepX) activities resulted in significant impairment of growth rate in milk.

The extent of proteolysis varied among strains and appeared to be time dependant. As depicted in Figure 2, the amount of liberated amino groups and peptides increased only slightly during fermentation from 0 to 12 h for some strains (*L. acidophilus* L10, *L. acidophilus* La 4962, *B. lactis* B94, *B. longum* Bl 536, *L. casei* L26, and *L. casei* Lc 279) but increased significantly ($P < 0.05$) for all strains from 12 to 24 h (Fig. 2). These findings were consistent with those reported by Nielsen et al. [28]. In contrast, however, Leclerc et al. [23] reported a linear increase in the amount of free amino groups until end of fermentation in milk fermented with *L. helveticus* strains. The proteolytic pattern certainly had a strong effect on bacterial growth due to correlation between these parameters, which ranged from 0.76 for L26 to 0.90 for St 1342. This dependence could possibly




Figure 3. The change of cell concentration of selected dairy LAB and probiotic organisms (*S. thermophilus* St 1342, *L. delbrueckii* subsp. *bulgaricus* Lb 1466, *L. acidophilus* L10, *B. lactis* B94 and *L. casei* L26, *L. acidophilus* La 4962, *B. longum* B1 536 and *L. casei* Lc 279) cultivated in reconstituted skim milk (RSM) for 24 h at 42 °C (Error bars represent a pooled standard error of the mean, SEM = 0.11 CFU·mL⁻¹).

explain the general slow cell growth up to 6 h of fermentation at 42 °C [12]. Although *L. delbrueckii* ssp. *bulgaricus* Lb 1466 showed appreciable peptidase activity, it experienced poor growth. This indicated that this organism might require some other growth factors in addition to free amino acids and peptides. *Lactobacillus casei* L26 showed the highest proteolytic activity followed in order by *L. delbrueckii* ssp. *bulgaricus* Lb 1466, *S. thermophilus* St 1342 and *L. acidophilus* (La 4962 and L10), *Bifidobacterium* and *L. casei* (Lc 279) (Fig. 2) with the activity apparently strain specific ($P < 0.0001$). Similar to our findings, those of Shihata and Shah [39], Fuglsang et al. [17] also showed that the amount of free amino groups formed in the medium during fermentation was strain dependent. Thus, the differences in the amounts of amino groups released during fermentation of milk observed for the microorganisms, could probably relate to the different proteinases of the strains. Shihata and Shah [39] reported proteolytic activity of *S. thermophilus*, *L. delbrueckii* ssp. *bulgaricus*,

and *L. acidophilus* to be much greater than that of *Bifidobacterium* spp. The results in our study also showed similar pattern of proteolytic activity for *S. thermophilus* St 1342, *L. delbrueckii* ssp. *bulgaricus* Lb 1466, *L. acidophilus* strains (La 4962 and L10) and *L. casei* L26 being higher than that of *Bifidobacterium* spp. at 24 h of fermentation.

3.3. Cell growth and organic acids production

The change of cell concentration of selected dairy LAB and probiotic organisms cultivated individually in RSM for 24 h at 42 °C is shown in Figure 3. Lactic acid bacteria are nutritionally fastidious organisms which require more free amino acids or peptides than present in milk. Thus proteolytic activity is important requirement for achieving a minimum level of 10^6 to 10^7 CFU·mL⁻¹ of active probiotics in a product to observe positive health effect [36, 44]. In general, examined probiotic cultures achieved the desired therapeutic




Figure 4. pH changes during fermentation of single strains in RSM by *S. thermophilus* St 1342, *L. delbrueckii* subsp. *bulgaricus* Lb 1466, *L. acidophilus* L10, *B. lactis* B94 and *L. casei* L26, *L. acidophilus* La 4962, *B. longum* B1 536 and *L. casei* Lc 279 (Error bars present a pooled standard error of the mean).

level (10^8 CFU·mL⁻¹) during their growth in RSM after 24 h (Fig. 3). Although the cultures showed a consistent increase in cell concentration until 9 h, the required pH of 4.5 was not reached as shown in Figure 4. Sodini et al. [42] also reported that individual probiotic cultures grew well in fermented milk and did not produce organic acids as fast as with mixed starter cultures. *Lactobacillus acidophilus* La 4962 and *L. casei* Lc 279 experienced a superior growth ($P < 0.05$) as compared to other probiotic organisms. On the other hand, *B. lactis* B94 and *B. longum* B1 536 showed slow growth pattern and no significant ($P > 0.05$) difference in the cell counts was observed (Fig. 3). *S. thermophilus* St 1342 achieved a significantly ($P < 0.05$) higher cell concentration in comparison to *L. delbrueckii* ssp. *bulgaricus* Lb 1466; the slow growth of the latter resulted in low viability (10^7 CFU·mL⁻¹) at the end of the fermentation period of 20 h.

Eventhough *L. delbrueckii* ssp. *bulgaricus* Lb 1466 grew slowly in RSM, it produced substantial amount of lactic acid,

second to *S. thermophilus* St 1342 (Fig. 5), and the highest ($P < 0.05$) decline in pH at the end of fermentation (Fig. 4). As previously observed by Donkor et al. [13], the yoghurt culture (*S. thermophilus* and *L. delbrueckii* ssp. *bulgaricus*) produced substantially more lactic acid than any other strain in the study. However, the study also showed that probiotic organisms produced some lactic acid even though not as high as in the case of yoghurt culture, with *L. acidophilus* L10 producing the highest and *L. casei* L26 the lowest concentration (Fig. 5).

In general, the production of acetic acid as presented in Figure 6 was fairly uniform in each batch of fermented milk up to 12 h. All cultures produced a substantial ($P < 0.05$) amount of acetic acid except *L. acidophilus* L10 (Fig. 6). *Lactobacillus casei* Lc 279 on the other hand, produced significantly ($P < 0.05$) higher acetic acid than other probiotics. The upsurge in cell growth for all batches from 9 h to 24 h (Fig. 3) resulted in significant ($P < 0.05$) increases in the concentration of organic acids and decline in pH (Fig. 4) but the concentration of




Figure 5. Production of lactic acid during fermentation of single strains in RSM for 24 h at 42 °C by *S. thermophilus* St 1342, *L. delbrueckii* subsp. *bulgaricus* Lb 1466, *L. acidophilus* L10, *B. lactis* B94 and *L. casei* L26, *L. acidophilus* La 4962, *B. longum* Bl 536 and *L. casei* Lc 279 (Error bars represent a pooled standard error of the mean SEM = 0.01 g·L⁻¹).




Figure 6. Production of acetic acid in fermented milk by *S. thermophilus* St 1342, *L. delbrueckii* subsp. *bulgaricus* Lb 1466, *L. acidophilus* L10, *B. lactis* B94 and *L. casei* L26, *L. acidophilus* La 4962, *B. longum* Bl 536 and *L. casei* Lc 279 during 24 h fermentation of RSM at 42 °C (Error bars represent a pooled standard error of the mean SEM = 0.01 g·L⁻¹).

Table IV. Angiotensin I-converting enzyme (ACE) inhibitory of fermented milk extracts obtained after 24 h of individual culture fermentations in RSM at 42 °C.

In vitro ACE-inhibitor activity		
Culture	24 h inhibition, %	IC ₅₀ *, mg·mL ⁻¹
<i>S. thermophilus</i> St 1342	43.65 ± 4.67	0.225 ± 0.033
<i>L. delbrueckii</i> ssp. <i>bulgaricus</i> Lb 1466	51.10 ± 2.30	0.187 ± 0.012
<i>L. acidophilus</i> L10	62.55 ± 0.24 ^A	0.151 ± 0.001 ^A
<i>L. acidophilus</i> La 4962	53.58 ± 1.73 ^B	0.164 ± 0.007 ^B
<i>B. lactis</i> B94	51.64 ± 3.63 ^A	0.175 ± 0.016 ^A
<i>B. longum</i> B1 536	63.73 ± 1.71 ^B	0.196 ± 0.007 ^A
<i>L. paracasei</i> L26	54.02 ± 1.71 ^A	0.196 ± 0.008 ^A
<i>L. casei</i> Lc 279	41.15 ± 4.13 ^B	0.344 ± 0.053 ^B

* The activity is expressed as the concentration of peptides in mg·mL⁻¹ required to inhibit 50% of the original ACE activity (IC₅₀).

^{AB} Means in the same column for particular strains with different capital letter superscripts are significantly different.

organic acids attained did not affect the cell growth as noted in other studies [1, 36, 42]. The organisms appeared to have maintained appreciable cell counts during the 24 h fermentation in RSM possibly due to efficient proteolytic systems [19].

3.4. ACE-inhibitory activity

In our study, eight strains of LAB were cultured individually in RSM to produce fermented milk with in vitro ACE-inhibitory activity shown in Table IV. The results indicate that the production of ACE inhibitors was not confined to a single species or strains of bacteria but all the strains tested produced peptides, which showed in vitro ACE-inhibitory activity. However, un-inoculated RSM did not show ACE-I activity (data not shown). Furthermore, these selected microorganisms showed X-prolyl-dipeptidyl aminopeptidase activity, which was able to cleave proline-containing sequences (Fig. 1). This may mean that specific peptides could be produced by cleaving N-terminal of X-Pro dipeptides from tri- and oligopeptides which may show ACE-I activity [21]. The activity of the fermented milk obtained for the selected LAB was also compared by determining the protein concentration needed to inhibit

50% of the original ACE activity (IC₅₀) (Tab. IV). There was a significant ($P < 0.05$) difference in IC₅₀ values between the bacterial strains, indicating possible differences in the quality of ACE-inhibitory peptides produced by the organisms. *B. longum* B1 536 and *L. acidophilus* L10 produced the highest activity at 63.7% and 62.6% with IC₅₀ values of 0.196 and 0.151 mg·mL⁻¹, respectively. Similar results were reported by Pihlanto-Leppä et al. [31] using different lactic acid starters and digestive enzymes to obtain inhibition of 35 to 86% and IC₅₀ values between 0.080 to 0.314 mg·mL⁻¹. Several milk protein-derived peptides produced by the enzymatic hydrolysis [33] or by the fermentation of milk with different strains of organisms [18, 23] inhibited the activity of ACE.

Although the proteolytic activity of *B. longum* B1 536 was not as high as *L. casei* L26, the peptides released by *B. longum* B1 536 had higher ACE-inhibitory activity than those of *L. casei* L26 (Tab. IV). Fuglsang et al. [17] found that most LAB produced ACE inhibitors in varying amounts during milk fermentation which varied with strains. Their study was in line with our findings which showed a similar trend using selected strains of bacteria and

further reported that only *L. helveticus* produced such substances in amounts large enough to cause a direct effect on ACE in vivo out of several strains used. The time-dependent release of various peptides observed in our study might have important consequences on the extent of in vitro ACE inhibitory activity in fermented milk, which deserves further elaboration.

4. CONCLUSION

Selected cultures of *S. thermophilus* St 1342, *L. delbrueckii* ssp. *bulgaricus* Lb 1466, *L. acidophilus* L10, *B. lactis* B94, *L. casei* L26, *L. acidophilus* La 4962, *B. longum* B1 536 and *L. casei* Lc 279 were capable of growing in milk likely due to degradation of milk proteins. Thus, this study has shown that for active and rapid growth in milk, dairy cultures may rely on their proteolytic system. The cultures possessed proteolytic enzymes – proteinases, peptidases and aminopeptidases at extra-cellular and intracellular levels which showed specific activities toward certain substrates. All strains exhibited X-prolyl-dipeptidyl aminopeptidase activity cleaving proline-containing sequences. The proteolytic systems including X-prolyl-dipeptidyl aminopeptidase activity of the bacteria strains in RSM resulted in the release of free amino groups and peptides which substantially improved growth and maintained viability in the medium. The proteolytic activity did not appear as the governing factor of growth for *Lactobacillus delbrueckii* ssp. *bulgaricus*, which might require additional growth factors. All selected strains produced a range of bioactive peptides with varying degree of ACE-inhibition. *B. longum* B1 536 and *L. acidophilus* L10 produced the highest activity IC₅₀ values of 0.196 and 0.151 mg·mL⁻¹, respectively. ACE-I was strain and apparently time dependent.

Acknowledgments: This research was funded by the Australian Research Council (ARC) – Linkage grant. The financial contribution by DSM Food Specialties (Moorebank, NSW, Australia) is also gratefully acknowledged.

REFERENCES

- [1] Akalin A.S., Fenderya S., Akbulut N., Viability and activity of bifidobacteria in yoghurt containing fructooligosaccharids during refrigerated storage, *Int. J. Food Sci.* 39 (2004) 613–621.
- [2] Blanc B., Laloi P., Atlan D., Gilbert C., Portalier R., Two cell-wall-associated aminopeptidases from *Lactobacillus helveticus* and the purification and characterization of APII from strain ITGL1, *J. Gen. Microbiol.* 139 (1993) 1441–1448.
- [3] Bradford M.M., A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding, *Anal. Biochem.* 72 (1976) 248–254.
- [4] Christensen J.E., Steele J.L., Impaired growth rates in milk of *Lactobacillus helveticus* peptidase mutants can be overcome by use of amino acid supplements, *J. Bacteriol.* 185 (2003) 3297–3306.
- [5] Christensen J.E., Dudley E.G., Pederson J.A., Steele J.L., Peptidases and amino acid catabolism in lactic acid bacteria, *Antonie Leeuwenhoek* 76 (1999) 217–246.
- [6] Church F.C., Swaisgood H.E., Porter D.H., Catignani G.L., Spectrophotometric assay using o-phthaldialdehyde for determination of proteolysis in milk and isolated milk proteins, *J. Dairy Sci.* 66 (1983) 1219–1227.
- [7] Coolbear T., Pillidge C.J., Crow V.L., The diversity of potential cheese ripening characteristics of lactic acid starter bacteria: resistance to cell lysis and levels and cellular distribution of proteinase activities, *Int. Dairy J.* 4 (1994) 697–721.
- [8] Crittenden R., Bird A.R., Gopal P., Henriksson A., Lee Y.K., Playne M.J., Probiotic research in Australia, New Zealand and the Asia-Pacific Region, *Current Pharmaceut. Design* 11 (2005) 37–53.
- [9] Cushman D.W., Cheung H.S., Spectrophotometric assay and properties of the angiotensin-converting enzyme of rabbit lung, *Biochem. Pharmacol.* 20 (1971) 1637–1648.
- [10] Dave R.I., Shah N.P., Evaluation of media for selective enumeration of *Streptococcus thermophilus*, *Lactobacillus delbrueckii* ssp. *bulgaricus*, *Lactobacillus acidophilus*, and bifidobacteria, *J. Dairy Sci.* 79 (1996) 1529–1536.
- [11] Dave R.I., Shah N.P., Ingredient supplementation effects on viability of probiotic bacteria in yoghurt, *J. Dairy Sci.* 81 (1998) 2804–2816.

- [12] Donkor O.N., Henriksson A., Vasiljevic T., Shah N.P., Probiotic strains as starter cultures improve angiotensin-converting enzyme inhibitory activity in soy yoghurt, *J. Food Sci.* 70 (2005) M375–M381.
- [13] Donkor O.N., Henriksson A., Vasiljevic T., Shah N.P., Effect of acidification on the activity of probiotics in yoghurt during cold storage, *Int. Dairy J.* 16 (2006) 1180–1189.
- [14] FAO/WHO, Guidelines for the evaluation of probiotics in food. Report of a joint FAO/WHO working group on drafting guidelines for the evaluation of probiotics in food, 2002, London ON, Canada.
- [15] Fenster K.M., Parkin K.L., Steele J.L., Characterization of a thiol-dependent endopeptidase from *Lactobacillus helveticus* CNRZ32, *J. Bacteriol.* (1997) 2529–2533.
- [16] Fernandez-Espla M.D., Martin-Hernandez M.C., Fox P.F., Purification and characterization of a prolidase from *Lactobacillus casei* subsp. *casei* IFPL 731, *Appl. Environ. Microbiol.* 63 (1997) 314–316.
- [17] Fuglsang A., Rattray F.P., Nilsson D., Nyborg N.C.B., Lactic acid bacteria: inhibition of angiotensin converting enzymes *in vitro* and *in vivo*, *Antonie Leeuwenhoek* 83 (2003) 27–34.
- [18] Gobbetti M., Ferranti P., Smacchi E., Goffredi F., Addeo F., Production of angiotensin-I-converting enzyme-inhibitory peptides in fermented milk started by *Lactobacillus delbrueckii* subsp. *bulgaricus* SS1 and *Lactobacillus lactis* subsp. *cremoris* FT4, *Appl. Environ. Microbiol.* 66 (2002) 3898–3904.
- [19] Juillard V., Le Bars D., Kunji E.R.S., Konings W.N., Gripon J.C., Richard J., Oligopeptides are the main source of nitrogen for *Lactococcus lactis* during growth in milk, *Appl. Environ. Microbiol.* 61 (1995) 3024–3030.
- [20] Klaver F.A.M., Kingma F., Weerkamp A.H., Growth and survival of bifidobacteria in milk, *Neth. Milk Dairy J.* 47 (1993) 151–164.
- [21] Kunji E.R.S., Mierau I., Hagting A., Poolman B., Konings W.N., The proteolytic systems of lactic acid bacteria, *Antonie Leeuwenhoek* 70 (1996) 187–221.
- [22] Law J., Haandrikman A., Proteolytic enzymes of lactic acid bacteria, *Int. Dairy J.* 7 (1997) 1–11.
- [23] Leclerc P.L., Gauthier S.F., Bachelard H., Santure M., Roy D., Antihypertensive activity of casein-enriched milk fermented by *Lactobacillus helveticus*, *Int. Dairy J.* 12 (2002) 995–1004.
- [24] Littell R.C., Henry P.R., Ammerman C.B., Statistical analysis of repeated measures data using SAS procedures, *J. Anim. Sci.* 76 (1998) 1216–1231.
- [25] Lourens-Hattingh A., Viljoen C.B., Yoghurt as probiotic carrier food, *Int. Dairy J.* 11 (2001) 1–17.
- [26] Mars I., Monnet V., An aminopeptidase P from *Lactococcus lactis* with original specificity, *Biochem. Biophys. Acta* 1243 (1995) 209–215.
- [27] Nakamura Y., Yamamoto N., Sakai K., Okubo A., Yamazaki S., Takano T., Purification and characterization of angiotensin I-converting enzyme inhibitors from sour milk, *J. Dairy Sci.* 78 (1995) 777–783.
- [28] Nielsen P.M., Petersen D., Dambmann C., Improved method for determining food protein degree of hydrolysis, *J. Food Sci.* 66 (2001) 642–646.
- [29] Pan D., Luo Y., Tanokura M., Antihypertensive peptides from skimmed milk hydrolysate digested by cell-free extract of *Lactobacillus helveticus* JCM1004, *Food Chem.* 91 (2005) 123–129.
- [30] Pihlanto A., Korhonen H., Bioactive peptides and proteins, *Adv. Food Nutr. Res.* 47 (2003) 176–275.
- [31] Pihlanto-Leppä A., Rokka T., Korhonen H., Angiotensin I converting enzyme inhibitory peptides derived from bovine milk proteins, *Int. Dairy J.* 8 (1998) 325–333.
- [32] Ravula R.R., Shah N.P., Selection enumeration of *Lactobacillus casei* from yoghurts and fermented milk drinks, *Biotechnol. Techn.* 12 (1998) 819–822.
- [33] Roy M.K., Watanabe Y., Tamai Y., Yeast protease B-digested skimmed milk inhibits angiotensin-I-converting-enzyme activity, *Biotechnol. Appl. Biochem.* 31 (2000) 95–100.
- [34] SAS, SAS/STAT Software: Changes and enhancements through release 6.11, 1996, SAS Inst. Inc., Cary, NC.
- [35] Shah N.P., Effects of milk-derived bioactives: an overview, *Br. J. Nutr.* 84 (2000) S3–S10.
- [36] Shah N.P., Probiotic bacteria: Selective enumeration and survival in dairy foods, *J. Dairy Sci.* 83 (2000) 894–907.
- [37] Shah N.P., Some beneficial effects of probiotic bacteria, *Biosci. Microflora* 19 (2000) 99–106.
- [38] Shah N.P., Ravula R.R., Influence of water activity on fermentation organic acids production and viability of yoghurt and probiotic

- bacteria, *Australian J. Dairy Technol.* 55 (2000) 127–131.
- [39] Shihata A., Shah N.P., Proteolytic profile of yoghurt and probiotic bacteria, *Int. Dairy J.* 10 (2000) 401–408.
- [40] Skeggs L.T., Kahn J.R., Shumway N.P., The preparation and function of the hypertension-converting enzyme, *J. Exp. Medicine* 103 (1956) 295–299.
- [41] Smid E.J., Poolmann B., Konings W.N., Casein utilization by lactococci, *Appl. Environ. Microbiol.* 57 (1991) 2447–2452.
- [42] Sodini I., Lucas A., Oliviera M.N., Remeuf F., Corrieu G., Effect of milk base and starter culture on acidification, texture, and probiotic cell counts in fermented milk processing, *J. Dairy Sci.* 85 (2002) 2479–2488.
- [43] Stepaniak L., Fox P.F., Characterization of principal intracellular endopeptidase from *Lactococcus lactis* subsp. *lactis* MG1363, *Int. Dairy J.* 5 (1995) 699–713.
- [44] Talwalkar A., Kailasapathy K., A review of oxygen toxicity in probiotic yogurts: Influence on the survival of probiotic bacteria and protective techniques, *Rev. Food Sci. Food Safety* 3 (2004) 117–124.
- [45] Tan P.S.T., Konings W.N., Purification and characterization of aminopeptidase from *Lactococcus lactis* subsp. *cremoris* Wg2, *Appl. Environ. Microbiol.* 56 (1990) 526–532.
- [46] Tan P.S.T., Sasaki M., Bosman B.W., Iwasaki T., Purification and characterization of a dipeptidase from *Lactobacillus helveticus* SBT 2171, *Appl. Environ. Microbiol.* 61 (1995) 3430–3435.
- [47] Tharmaraj N., Shah N.P., Selective enumeration of *Lactobacillus delbrueckii* ssp. *bulgaricus*, *Streptococcus thermophilus*, *Lactobacillus acidophilus*, bifidobacteria, *Lactobacillus casei*, *Lactobacillus rhamnosus* and *Propionibacteria*, *J. Dairy Sci.* 86 (2003) 2288–2296.
- [48] Vasiljevic T., Shah N.P., Jelen P., Growth characteristics of *Lactobacillus delbrueckii* subsp. *bulgaricus* ATCC 11842 as affected by different neutralizers, *Aust. J. Dairy Technol.* 60 (2005) 3–9.
- [49] Vermeirssen V., Camp J.V., Verstraete W., Optimisation and validation of an angiotensin-converting enzyme inhibition assay for the screening of bioactive peptides, *J. Biochem. Biophys. Methods* 51 (2002) 75–87.
- [50] Wohlrab Y., Bockelmann W., Purification and characterization of a dipeptidase from *Lactobacillus delbrueckii* subsp. *bulgaricus*, *Int. Dairy J.* 2 (1992) 345–361.
- [51] Ziemer C.J., Gibson G.R., An overview of probiotics, prebiotics and synbiotics in the functional food concept: perspectives and future strategies, *Int. Dairy J.* 8 (1998) 473–479.