



Stochastic neural field equations: A rigorous footing

Olivier Faugeras, James Inglis

► **To cite this version:**

Olivier Faugeras, James Inglis. Stochastic neural field equations: A rigorous footing. Journal of Mathematical Biology, Springer Verlag (Germany), 2014, pp.40. hal-00907555v2

HAL Id: hal-00907555

<https://hal.inria.fr/hal-00907555v2>

Submitted on 16 Dec 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Stochastic neural field equations: A rigorous footing*

O. Faugeras [†] and J. Inglis [‡]

December 16, 2014

Abstract

We here consider a stochastic version of the classical neural field equation that is currently actively studied in the mathematical neuroscience community. Our goal is to present a well-known rigorous probabilistic framework in which to study these equations in a way that is accessible to practitioners currently working in the area, and thus to bridge some of the cultural/scientific gaps between probability theory and mathematical biology. In this way, the paper is intended to act as a reference that collects together relevant rigorous results about notions of solutions and well-posedness, which although may be straightforward to experts from SPDEs, are largely unknown in the neuroscientific community, and difficult to find in a very large body of literature. Moreover, in the course of our study we provide some new specific conditions on the parameters appearing in the equation (in particular on the neural field kernel) that guarantee the existence of a solution.

1 Introduction

Neural field equations have been widely used to study spatiotemporal dynamics of cortical regions. Arising as continuous spatial limits of discrete models, they provide a step towards an understanding of the relationship between the macroscopic spatially structured activity of densely populated regions of the brain, and the underlying microscopic neural circuitry. The discrete models themselves describe the activity of a large number of individual neurons with no spatial dimensions. Such neural mass models have been proposed by Lopes da Silva and colleagues [38, 39] to account for oscillatory phenomena observed in the brain, and were later put on a

*This work was partially supported by the European Union Seventh Framework Programme (FP7/2007- 2013) under grant agreement no. 269921 (BrainScaleS), no. 318723 (MathemacS), and by the ERC advanced grant NerVi no. 227747.

[†]NeuroMathComp Team, INRIA Sophia-Antipolis. *Email: olivier.faugeras@inria.fr*

[‡]NeuroMathComp/TOSCA Team, INRIA Sophia-Antipolis. *Email: james.inglis@inria.fr*

stronger mathematical footing in the study of epileptic-like seizures in [24]. When taking the spatial limit of such discrete models, one typically arrives at a nonlinear integro-differential equation, in which the integral term can be seen as a nonlocal interaction term describing the spatial distribution of synapses in a cortical region. Neural field models build on the original work of Wilson and Cowan [43, 44] and Amari [1], and are known to exhibit a rich variety of phenomena including stationary states, traveling wave fronts, pulses and spiral waves. For a comprehensive review of neural field equations, including a description of their derivation, we refer to [6].

More recently several authors have become interested in stochastic versions of neural field equations (see for example [4, 5, 8, 10, 27]), in order to (amongst other things) model the effects of fluctuations on wave front propagation. In particular, in [8] a multiplicative stochastic term is added to the neural field equation, resulting in a stochastic nonlinear integro-differential equation of the form

$$dY(t, x) = \left[-Y(t, x) + \int_{\mathbb{R}} w(x, y)G(Y(t, y))dy \right] dt + \sigma(Y(t, x))dW(t, x), \quad (1.1)$$

for $x \in \mathbb{R}, t \geq 0$, and some functions G (referred to as the nonlinear gain function), σ (the diffusion coefficient) and w (the neural field kernel, sometimes also called the connectivity function). Here $(W(t, x))_{x \in \mathbb{R}, t \geq 0}$ is a stochastic process (notionally a ‘‘Gaussian random noise’’) that depends on both space and time, and which may possess some spatial correlation.

Of course the first step towards understanding (1.1) rigorously is defining what we mean by a solution. This is in fact not completely trivial and is somewhat glossed-over in the neuroscientific literature. The main point is that any solution must involve an object of the form

$$\text{‘‘} \int \sigma(Y(t, x))dW(t, x)\text{’’} \quad (1.2)$$

which must be precisely defined. Of course, in the case where there is no spatial dimension, the theory of such stochastic integrals is widely disseminated, but for integrals with respect to space-time white noise (for example) it is far less well-known. It is for this reason that we believe it to be extremely worthwhile making a detailed review of how to give sense to these objects, and moreover to solutions to (1.1) when they exist, in a way that is accessible to practitioners. Although such results are quite well-known in probability theory, the body of literature is very large and generalistic, posing a daunting prospect for a mathematical neuroscientist looking to apply a specific result. The fact that the equation fits into well-studied frameworks also opens up opportunities to apply existing abstract results (for example large deviation principles – see Remark 2.3).

There are in fact two distinct approaches to defining and interpreting the quantity (1.2), both of which allow one to build up a theory of stochastic *partial* differential equations (SPDEs). Although (1.1) does not strictly classify as a SPDE (since there

is no derivative with respect to the spatial variable), both approaches provide a rigorous underlying theory upon which to base a study of such equations.

The first approach generalizes the theory of stochastic processes in order to give sense to solutions of SPDEs as random processes that take their values in a Hilbert space of functions (as presented by Da Prato and Zabczyk in [35] and more recently by Prévôt and Röckner in [36]). With this approach, the quantity (1.2) is interpreted as a *Hilbert space-valued* integral i.e. “ $\int \mathbf{B}(Y(t))dW(t)$ ”, where $(Y(t))_{t \geq 0}$ and $(W(t))_{t \geq 0}$ take their values in a Hilbert space of functions, and $\mathbf{B}(Y(t))$ is an operator between Hilbert spaces (depending on σ). The second approach is that of J. B. Walsh (as described in [42]), which, in contrast, takes as its starting point a PDE with a random and highly irregular “white-noise” term. This approach develops integration theory with respect to a class of random measures, so that (1.2) can be interpreted as a random field in both t and x .

In the theory of SPDEs, there are advantages and disadvantages of taking both approaches. This is also the case with regards to the stochastic neural field equation (1.1), as described in the conclusion below (Section 5), and it is for this reason that we here review both approaches. Taking the functional approach of Da Prato and Zabczyk is perhaps more straightforward for those with knowledge of stochastic processes, and the existing general results can be applied more directly in order to obtain, for example, existence and uniqueness. This was the path taken in [28] where the emphasis was on large deviations, though in a much less general setup than we consider here (see Remark 2.3). However, it can certainly be argued that solutions constructed in this way may be “non-physical”, since the functional theory tends to ignore any spatial regularity properties (solutions are typically L^2 -valued in the spatial direction). We argue that the approach of Walsh is more suited to looking for “physical” solutions that are at least continuous in the spatial dimension. A comparison of the two approaches in a general setting is presented in [15] or [25, 26], and in our setting in Section 4 below. Our main conclusion is that in typical cases of interest for practitioners, the approaches are equivalent (see Example 4.2), but one or the other may be more suited to a particular need.

To reiterate, the main aim of this article is to present a review of an existing theory, which is accessible to readers unfamiliar with stochastic partial differential equations, that puts the study of stochastic neural field equations on a rigorous mathematical footing. As a by product we will be able to give general conditions on the functions G , σ and w that, as far as we know, do not appear anywhere else in the literature and guarantee the existence of a solution to (1.1) in some sense. Moreover, these conditions are weak enough to be satisfied for all typical choices of functions made by practitioners (see Sections 2.6, 2.7 and 2.8). By collecting all these results in a single place, we hope this will provide a reference for practitioners in future works.

The layout of the article is as follows. We first present in Section 2 the necessary material in order to consider the stochastic neural field equation (1.1) as an evolution equation in a Hilbert space. This involves introducing the notion of a

Q -Wiener process taking values in a Hilbert space and stochastic integration with respect to Q -Wiener processes. A general existence result from [35] is then applied in Section 2.5 to yield a unique solution to (1.1) interpreted as a Hilbert space valued process. The second part of the paper switches track, and describes Walsh’s theory of stochastic integration (Section 3.1), with a view of giving sense to a solution to (1.1) as a random field in both time and space. To avoid dealing with distribution-valued solutions, we in fact consider a Gaussian noise that is smoothed in the spatial direction (Section 3.2), and show that, under some weak conditions, the neural field equation driven by such a smoothed noise has a unique solution in the sense of Walsh that is continuous in both time and space (Section 3.3). We finish with a comparison of the two approaches in Section 4, and summarize our findings in a conclusion (Section 5).

Notation: Throughout the article $(\Omega, \mathcal{F}, \mathbb{P})$ will be a probability space, and $L^2(\Omega, \mathcal{F}, \mathbb{P})$ will be the space of square-integrable random variables on $(\Omega, \mathcal{F}, \mathbb{P})$. We will use the standard notation $\mathcal{B}(\mathcal{T})$ to denote the Borel σ -algebra on \mathcal{T} for any topological space \mathcal{T} . The Lebesgue space of p -integrable (with respect to the Lebesgue measure) functions over \mathbb{R}^N for $N \in \mathbb{N} = \{1, 2, \dots\}$ will be denoted by $L^p(\mathbb{R}^N)$, $p \geq 1$, as usual, while $L^p(\mathbb{R}^N, \rho)$, $p \geq 1$, will be the Lebesgue space weighted by a measurable function $\rho : \mathbb{R}^N \rightarrow \mathbb{R}^+$.

2 Stochastic neural field equations as evolution equations in Hilbert spaces

As stated in the introduction, the goal of this section is to provide the theory and conditions needed to interpret the solution to (1.1) as a process $(Y(t))_{t \geq 0}$ that takes its values in a Hilbert space of functions i.e. for each $t \geq 0$, $Y(t)$ is a function of the spatial variable x . This is in order to try and cast the problem into the well-known theoretical framework of stochastic evolution equations in Hilbert spaces, as detailed in [35]. In particular we will look for solutions to

$$dY(t) = (-Y(t) + \mathbf{F}(Y(t))) dt + \mathbf{B}(Y(t))dW(t), \quad t \geq 0, \quad (2.1)$$

such that $Y(t) \in L^2(\mathbb{R}^N, \rho)$ for some measurable $\rho : \mathbb{R}^N \rightarrow \mathbb{R}^+$ (to be determined), where \mathbf{F} is now an operator on $L^2(\mathbb{R}^N, \rho)$ given by

$$\mathbf{F}(Y(t))(x) = \int_{\mathbb{R}^N} w(x, y)G(Y(t, y))dy, \quad x \in \mathbb{R}^N.$$

Here $w : \mathbb{R}^N \times \mathbb{R}^N \rightarrow \mathbb{R}$ is the neural field kernel, and $G : \mathbb{R} \rightarrow \mathbb{R}$ is the nonlinear gain function. Note that we have made a slight generalization here in comparison with (1.1) in that we in fact work on \mathbb{R}^N , rather than \mathbb{R} . The term $\mathbf{B}(Y(t))dW(t)$ represents a stochastic differential term that must be made sense of as a differential

in the Hilbert space $L^2(\mathbb{R}^N, \rho)$. This is done with the help of Sections 2.1 and 2.2 below.

Notation: In this section we will also need the following basic notions from functional analysis. Let U and H be two separable Hilbert spaces. We will write $L_0(U, H)$ to denote the space of all bounded linear operators from U to H with the usual norm¹ (with the shorthand $L_0(H)$ when $U = H$), and $L_2(U, H)$ for the space of all Hilbert-Schmidt operators from U to H , i.e. those bounded linear operators $B : U \rightarrow H$ such that

$$\sum_{k \geq 1} \|B(e_k)\|_H^2 < \infty,$$

for some (and hence all) complete orthonormal systems $\{e_k\}_{k \geq 1}$ of U . Finally, a bounded linear operator $Q : U \rightarrow U$ will be said to be trace-class if $\text{Tr}(Q) := \sum_{k \geq 1} \langle Q(e_k), e_k \rangle_U < \infty$, again for some (and hence all) complete orthonormal systems $\{e_k\}_{k \geq 1}$ of U .

2.1 Hilbert space valued Q -Wiener processes

The purpose of this section is to provide a basic understanding of how we can generalize the idea of an \mathbb{R}^d -valued Wiener process to one that takes its values in an infinite dimensional Hilbert space, which for convenience we fix to be $U = L^2(\mathbb{R}^N)$ (this is simply for the sake of being concrete).

In the finite dimensional case, it is well-known that \mathbb{R}^d -valued Wiener processes are characterized by their $d \times d$ covariance matrices, which are symmetric and non-negative. The basic idea is that in the infinite dimensional setup the covariance matrices are replaced by covariance *operators*, which are linear, non-negative, symmetric and bounded.

Indeed, let $Q : U \rightarrow U$ be a non-negative, symmetric bounded linear operator on U . To avoid introducing extra embeddings, we also suppose $\text{Tr}(Q) < \infty$. Then, completely analogously to the finite dimensional case, there exists a sequence of non-negative real numbers $(\lambda_k)_{k \geq 1}$ which are eigenvalues of Q , associated with a sequence of eigenfunctions $\{e_k\}_{k \geq 1}$ (i.e. $Qe_k = \lambda_k e_k$) that form a complete orthonormal basis for U . Moreover, since $\text{Tr}(Q) < \infty$, it holds that

$$\sum_{k=1}^{\infty} \lambda_k < \infty.$$

By a Q -Wiener process $W = (W(t))_{t \geq 0}$ on U we will simply mean that $W(t)$ can be expanded as

$$W(t) = \sum_{k=1}^{\infty} \sqrt{\lambda_k} \beta_k(t) e_k, \tag{2.2}$$

¹The norm of $B \in L_0(U, H)$ is classically defined as $\sup_{x \neq 0} \frac{\|Bx\|_H}{\|x\|_U}$.

where $(\beta_k(t))_{t \geq 0}$, $k = 1, 2, \dots$ are mutually independent standard real-valued Brownian motions. We note that $W(t)$ exists as a U -valued square-integrable random variable i.e. $W(t) \in L^2(\Omega, \mathcal{F}, \mathbb{P})$.

Equation (2.2) shows the role played by Q : the eigenvectors e_k are functions that determine “where” the noise “lives” in U , while the eigenvalues λ_k determine its dimensionality and relative strength. As an example of a covariance operator², let us compute the covariance operator of W . An easy computation based on (2.2) and the elementary properties of the standard real-valued Brownian motion shows that

$$\mathbb{E}[\langle W(s), g \rangle_U \langle W(t), h \rangle_U] = (s \wedge t) \langle Qg, h \rangle_U \quad \forall g, h \in U. \quad (2.3)$$

It turns out that W is white in both space and time. The whiteness in time is apparent from the above expression. The whiteness in space is shown explicitly in Section 2.7.

2.2 Stochastic integration with respect to Q -Wiener processes

The second point is that we would like to be able to define stochastic integration with respect to these Hilbert space valued Wiener processes. In particular we must determine for which integrands this can be done (exactly as in [35]).

As above, let $U = L^2(\mathbb{R}^N)$, $Q : U \rightarrow U$ a non-negative, symmetric bounded linear operator on U such that $\text{Tr}(Q) < \infty$, and $W = (W(t))_{t \geq 0}$ be a Q -Wiener process on U (given by (2.2)).

Unfortunately, in order to define stochastic integrals with respect to W , we need a couple of technical definitions from functional analysis. This is simply in order to control the convergence of the infinite series that appear in the construction, as we will see in the example below. Indeed, let $Q^{\frac{1}{2}}(U)$ be the subspace of U , which is a Hilbert space under the inner product

$$\langle u, v \rangle_{Q^{\frac{1}{2}}(U)} := \langle Q^{-\frac{1}{2}}u, Q^{-\frac{1}{2}}v \rangle_U, \quad u, v \in Q^{\frac{1}{2}}(U).$$

$Q^{\frac{1}{2}}(U)$ is in fact simply the space generated by the orthonormal basis $\{\sqrt{\lambda_k}e_k\}$ whenever $\{e_k\}$ is the orthonormal basis for U consisting of eigenfunctions of Q . Moreover, let $H = L^2(\mathbb{R}^N, \rho)$ for some measurable $\rho : \mathbb{R}^N \rightarrow \mathbb{R}^+$ (again this is just for the sake of concreteness – one could instead take any separable Hilbert space). It turns out that the space $L_2(Q^{\frac{1}{2}}(U), H)$ of all Hilbert-Schmidt operators from $Q^{\frac{1}{2}}(U)$ into H plays an important role in the theory of stochastic integration with respect to W , and for this reason we detail the following simple but illuminating example.

²The covariance operator $C : U \rightarrow U$ of W is defined as $\mathbb{E}[\langle W(s), g \rangle_U \langle W(t), h \rangle_U] = \langle Cg, h \rangle_U$ for all $g, h \in U$.

Example 2.1. Let $B : U \rightarrow H$ be a bounded linear operator from U to H i.e. $B \in L_0(U, H)$. Then, by definition,

$$\begin{aligned} \|B\|_{L_2(Q^{\frac{1}{2}}(U), H)}^2 &= \sum_{k=1}^{\infty} \|B(Q^{\frac{1}{2}}(e_k))\|_H^2 \\ &\leq \|B\|_{L_0(U, H)}^2 \sum_{k=1}^{\infty} \|Q^{\frac{1}{2}}(e_k)\|_U^2 = \|B\|_{L_0(U, H)}^2 \sum_{k=1}^{\infty} \langle Q^{\frac{1}{2}}(e_k), Q^{\frac{1}{2}}(e_k) \rangle_U \\ &= \|B\|_{L_0(U, H)}^2 \sum_{k=1}^{\infty} \langle Q(e_k), e_k \rangle_U = \|B\|_{L_0(U, H)}^2 \text{Tr}(Q) < \infty, \end{aligned}$$

since $\text{Tr}(Q) < \infty$, where $\{e_k\}_{k \geq 1}$ is again a complete orthonormal system for U . In other words $B \in L_0(U, H) \Rightarrow B \in L_2(Q^{\frac{1}{2}}(U), H)$.

The main point of the section is the following. According to the construction detailed in Chapter 4 of [35], we have that for a (random) process $(\Phi(t))_{t \geq 0}$ the integral

$$\int_0^t \Phi(s) dW(s) \tag{2.4}$$

has a sense as an element of H when $\Phi(s) \in L_2(Q^{\frac{1}{2}}(U), H)$, $\Phi(s)$ is knowable³ at time s , and if

$$\mathbb{P} \left(\int_0^t \|\Phi(s)\|_{L_2(Q^{\frac{1}{2}}(U), H)}^2 ds < \infty \right) = 1.$$

Now in view of Example 2.1, the take-away message is simply that the stochastic integral (2.4) has a sense in H if $\Phi(s) : U \rightarrow H$ is a bounded linear operator i.e. is in $L_0(U, H)$ for all $s \in [0, t]$, and that the norm of $\Phi(s)$ is bounded on $[0, t]$. In fact this is the only knowledge that will be needed below.

2.3 The stochastic neural field equation: interpretation in language of Hilbert space valued processes

With the previous two sections in place, we can now return to (2.1) and interpret it (and in particular the noise term) in a rigorous way. Indeed, as above, let W be an $L^2(\mathbb{R}^N)$ -valued Q -Wiener process, with Q a non-negative, symmetric bounded linear operator on $L^2(\mathbb{R}^N)$ such that $\text{Tr}(Q) < \infty$ (trace-class). The rigorous interpretation of (2.1) as an equation for a process $(Y(t))_{t \geq 0}$ taking its values in the Hilbert space $L^2(\mathbb{R}^N, \rho)$ is then

$$dY(t) = (-Y(t) + \mathbf{F}(Y(t))) dt + \mathbf{B}(Y(t)) dW(t), \quad Y(0) = Y_0 \in L^2(\mathbb{R}^N, \rho) \tag{2.5}$$

³Technically this means that $\Phi(s)$ is measurable with respect the σ -algebra generated by all left-continuous processes that are known at time s when $(W(u))_{u \leq s}$ is known (these process are said to be adapted to the filtration generated by W).

where \mathbf{B} is a map from $L^2(\mathbb{R}^N, \rho)$ into the space of bounded linear operators $L_0(L^2(\mathbb{R}^N), L^2(\mathbb{R}^N, \rho))$. Note that if \mathbf{B} is such a map, then the integrated noise term of this equation has a sense thanks to Section 2.2.

We in fact work with a general map \mathbf{B} satisfying a Lipschitz condition (see below), but we keep in mind the following example which provides the link with the diffusion coefficient σ in (1.1):

$$\mathbf{B}(h)(u)(x) = \sigma(h(x)) \int_{\mathbb{R}^N} \varphi(x-y)u(y)dy, \quad x \in \mathbb{R}^N, \quad (2.6)$$

for $h \in L^2(\mathbb{R}^N, \rho)$ and $u \in L^2(\mathbb{R}^N)$, where σ and φ are some functions that must be chosen to ensure the conditions stated below are satisfied. We detail potential choices of σ and φ (and their significance from a modeling point of view – in particular how φ controls the spatial correlation) in Section 2.7 below.

To summarize, we are here concerned with the solvability of (2.5) in $L^2(\mathbb{R}^N, \rho)$ (for some measurable $\rho : \mathbb{R}^N \rightarrow \mathbb{R}^+$ to be determined), where

$$\mathbf{F}(h)(x) = \int_{\mathbb{R}^N} w(x,y)G(h(y))dy, \quad x \in \mathbb{R}^N, \quad h \in L^2(\mathbb{R}^N, \rho), \quad (2.7)$$

and $\mathbf{B} : L^2(\mathbb{R}^N, \rho) \rightarrow L_0(L^2(\mathbb{R}^N), L^2(\mathbb{R}^N, \rho))$. To this end, we make the following two Lipschitz assumptions on \mathbf{B} and the nonlinear gain function G :

- $\mathbf{B} : H \rightarrow L_0(U, H)$ is such that

$$\|\mathbf{B}(g) - \mathbf{B}(h)\|_{L_0(U,H)} \leq C_\sigma \|g - h\|_U, \quad g, h \in L^2(\mathbb{R}^N, \rho),$$

where $U = L^2(\mathbb{R}^N)$ and $H = L^2(\mathbb{R}^N, \rho)$ for notational simplicity;

- $G : \mathbb{R} \rightarrow \mathbb{R}$ is bounded and globally Lipschitz i.e such that there exists a constant C_G with $\sup_{a \in \mathbb{R}} |G(a)| \leq C_G$ and

$$|G(a) - G(b)| \leq C_G |a - b|, \quad \forall a, b \in \mathbb{R}.$$

Typically the nonlinear gain function G is taken to be a sigmoid function, for example $G(a) = (1 + e^{-a})^{-1}$, $a \in \mathbb{R}$, which certainly satisfies this assumption.

2.4 Discussion of conditions on the neural field kernel w and

ρ

Of particular interest to us are the conditions on the neural field kernel w which will allow us to prove existence and uniqueness of a solution to (2.5) by quoting a standard result from [35].

In [28, footnote 1] it is suggested that the condition

$$\int_{\mathbb{R}^N} \int_{\mathbb{R}^N} |w(x,y)|^2 dx dy < \infty \quad (\mathbf{C1})$$

together with symmetry of w is enough to ensure that there exists a unique $L^2(\mathbb{R}^N)$ -valued solution to (2.5). However, the problem is that it does not follow from (C1) that the operator \mathbf{F} is stable on the space $L^2(\mathbb{R}^N)$. For instance, suppose that in fact $G \equiv 1$ (so that G is trivially globally Lipschitz). Then for $h \in L^2(\mathbb{R}^N)$ (and assuming $w \geq 0$) we have that

$$\|\mathbf{F}(h)\|_{L^2(\mathbb{R}^N)}^2 = \int_{\mathbb{R}^N} \|w(x, \cdot)\|_{L^1(\mathbb{R}^N)}^2 dx. \quad (2.8)$$

The point is that we can choose positive w such that (C1) holds, while (2.8) is not finite. For example in the case $N = 1$ we could take $w(x, y) = (1 + |x|)^{-1}(1 + |y|)^{-1}$ for $x, y \in \mathbb{R}$. In such a case the equation (2.5) is ill-posed: if $Y(t) \in L^2(\mathbb{R})$ then $F(t, Y(t))$ is not guaranteed to be in $L^2(\mathbb{R})$, which in turn implies that $Y(t) \notin L^2(\mathbb{R})$!

With this in mind we argue two points. Firstly, if we want a solution in $L^2(\mathbb{R}^N)$, we must make the additional strong assumption that

$$\forall x \in \mathbb{R}^N \quad (y \mapsto w(x, y)) \in L^1(\mathbb{R}^N), \quad \text{and} \quad \|w(x, \cdot)\|_{L^1(\mathbb{R}^N)} \in L^2(\mathbb{R}^N). \quad (\mathbf{C2})$$

Indeed, below we will show that (C1) together with (C2) are enough to yield the existence of a unique $L^2(\mathbb{R}^N)$ -valued solution to (2.5).

On the other hand, if we don't want to make the strong assumptions that (C1) and (C2) hold, then we have to work instead in a weighted space $L^2(\mathbb{R}^N, \rho)$, in order to ensure that \mathbf{F} is stable. In this case, we will see that if

$$\exists \rho_w \in L^1(\mathbb{R}^N) \quad \text{s.t.} \quad \int_{\mathbb{R}^N} |w(x, y)| \rho_w(x) dx \leq \Lambda_w \rho_w(y) \quad \forall y \in \mathbb{R}^N, \quad (\mathbf{C1}')$$

for some $\Lambda_w > 0$, and

$$\forall x \in \mathbb{R}^N \quad (y \mapsto w(x, y)) \in L^1(\mathbb{R}^N), \quad \text{and} \quad \sup_{x \in \mathbb{R}^N} \|w(x, \cdot)\|_{L^1(\mathbb{R}^N)} \leq C_w \quad (\mathbf{C2}')$$

for some constant C_w , then we can prove the existence of a unique $L^2(\mathbb{R}^N, \rho_w)$ -valued solution to (2.5).

Condition (C1') is in fact a non-trivial eigenvalue problem, and it is not straightforward to see whether it is satisfied for a given function w . However, we chose to state the theorem below in a general way, and then below provide some important examples of when it can be applied.

We will discuss these abstract conditions from a modeling point of view below. However, we first present the existence and uniqueness result.

2.5 Existence and uniqueness

Theorem 2.2. *Suppose that the neural field kernel w either*

- (i) *satisfies conditions (C1) and (C2); or*

(ii) satisfies conditions **(C1')** and **(C2')**.

If (i) holds set $\rho_w \equiv 1$, while if (ii) holds let ρ_w be the function appearing in condition **(C1')**.

Then, whenever Y_0 is an $L^2(\mathbb{R}^N, \rho_w)$ -valued random variable with finite p -moments for all $p \geq 2$, the neural field equation (2.5) has a unique solution taking values in the space $L^2(\mathbb{R}^N, \rho_w)$. To be precise, there exists a unique $L^2(\mathbb{R}^N, \rho_w)$ -valued process $(Y(t))_{t \geq 0}$ such that for all $T > 0$

$$\mathbb{P} \left(\int_0^T \|Y(s)\|_{L^2(\mathbb{R}^N, \rho_w)}^2 ds < \infty \right) = 1,$$

and

$$Y(t) = e^{-t}Y_0 + \int_0^t e^{-(t-s)} \mathbf{F}(Y(s)) ds + \int_0^t e^{-(t-s)} \mathbf{B}(Y(s)) dW(s), \quad \mathbb{P} - a.s.$$

Moreover, $(Y(t))_{t \geq 0}$ has a continuous modification, and satisfies the bounds

$$\sup_{t \in [0, T]} \mathbb{E} \left[\|Y(t)\|_{L^2(\mathbb{R}^N, \rho_w)}^p \right] \leq C_T^{(p)} \left(1 + \mathbb{E} \left[\|Y_0\|_{L^2(\mathbb{R}^N, \rho_w)}^p \right] \right), \quad T > 0, \quad (2.9)$$

for all $p \geq 2$, while for $p > 2$,

$$\mathbb{E} \left[\sup_{t \in [0, T]} \|Y(t)\|_{L^2(\mathbb{R}^N, \rho_w)}^p \right] \leq C_T^{(p)} \left(1 + \mathbb{E} \left[\|Y_0\|_{L^2(\mathbb{R}^N, \rho_w)}^p \right] \right) \quad T > 0. \quad (2.10)$$

Proof. We simply check the hypotheses of [35, Theorem 7.4] (a standard reference in the theory) in both cases (i) and (ii). This involves showing that (a) $\mathbf{F} : L^2(\mathbb{R}^N, \rho_w) \rightarrow L^2(\mathbb{R}^N, \rho_w)$; (b) the operator $\mathbf{B}(h) \in L_2(Q^{\frac{1}{2}}(U), H)$, for all $h \in H$ (recalling that $U = L^2(\mathbb{R}^N)$ and $H = L^2(\mathbb{R}^N, \rho)$); and (c) \mathbf{F} and \mathbf{B} are globally Lipschitz.

(a): We check that the function $\mathbf{F} : L^2(\mathbb{R}^N, \rho_w) \rightarrow L^2(\mathbb{R}^N, \rho_w)$. In case (i) this holds since $\rho_w \equiv 1$ and for any $h \in L^2(\mathbb{R}^N)$

$$\begin{aligned} \|\mathbf{F}(h)\|_{L^2(\mathbb{R}^N)}^2 &= \int_{\mathbb{R}^N} \left| \int_{\mathbb{R}^N} w(x, y) G(h(y)) dy \right|^2 dx \\ &\leq C_G^2 \int_{\mathbb{R}^N} \|w(x, \cdot)\|_{L^1(\mathbb{R}^N)}^2 dx < \infty, \end{aligned}$$

by assumption **(C2)**. Similarly in case (ii) for any $h \in L^2(\mathbb{R}^N, \rho_w)$

$$\begin{aligned} \|\mathbf{F}(h)\|_{L^2(\mathbb{R}^N, \rho_w)}^2 &= \int_{\mathbb{R}^N} \left| \int_{\mathbb{R}^N} w(x, y) G(h(y)) dy \right|^2 \rho_w(x) dx \\ &\leq C_G^2 \sup_{x \in \mathbb{R}^N} \|w(x, \cdot)\|_{L^1(\mathbb{R}^N)}^2 \|\rho_w\|_{L^1(\mathbb{R}^N)} < \infty. \end{aligned}$$

Hence in either case \mathbf{F} in fact maps $L^2(\mathbb{R}^N, \rho_w)$ into a metric ball in $L^2(\mathbb{R}^N, \rho_w)$.

(b): To show (b) in both cases, we know by Example 2.1 that for $h \in H$, $\mathbf{B}(h) \in L_2(Q^{\frac{1}{2}}(U), H)$ whenever $\mathbf{B}(h) \in L_0(U, H)$, which is true by assumption.

(c): To show (c), we first want $\mathbf{F} : L^2(\mathbb{R}^N, \rho_w) \rightarrow L^2(\mathbb{R}^N, \rho_w)$ to be globally Lipschitz. To this end, for any $g, h \in L^2(\mathbb{R}^N, \rho_w)$, we see that in either case

$$\begin{aligned} \|\mathbf{F}(g) - \mathbf{F}(h)\|_{L^2(\mathbb{R}^N, \rho_w)}^2 &= \int_{\mathbb{R}^N} |\mathbf{F}(g) - \mathbf{F}(h)|^2(x) \rho_w(x) dx \\ &\leq \int_{\mathbb{R}^N} \left(\int_{\mathbb{R}^N} |w(x, y)| |G(g(y)) - G(h(y))| dy \right)^2 \rho_w(x) dx \\ &\leq C_G^2 \int_{\mathbb{R}^N} \left(\int_{\mathbb{R}^N} |w(x, y)| |g(y) - h(y)| dy \right)^2 \rho_w(x) dx, \end{aligned}$$

where we have used the Lipschitz property of G . Now in case (i) it clearly follows from the Cauchy-Schwartz inequality that

$$\|\mathbf{F}(g) - \mathbf{F}(h)\|_{L^2(\mathbb{R}^N)}^2 \leq C_G^2 \left(\int_{\mathbb{R}^N} \int_{\mathbb{R}^N} |w(x, y)|^2 dx dy \right) \|g - h\|_{L^2(\mathbb{R}^N)},$$

so that by condition (C1), \mathbf{F} is indeed Lipschitz.

In case (ii), by Cauchy-Schwartz and the specific property of ρ_w given by (C1'), we see that

$$\begin{aligned} \|\mathbf{F}(g) - \mathbf{F}(h)\|_{L^2(\mathbb{R}^N, \rho_w)}^2 &\leq C_G^2 \sup_{x \in \mathbb{R}^N} \|w(x, \cdot)\|_{L^1(\mathbb{R}^N)} \int_{\mathbb{R}^N} |g(y) - h(y)|^2 \left(\int_{\mathbb{R}^N} |w(x, y)| \rho_w(x) dx \right) dy \\ &\leq C_G^2 \Lambda_w \sup_{x \in \mathbb{R}^N} \|w(x, \cdot)\|_{L^1(\mathbb{R}^N)} \|g - h\|_{L^2(\mathbb{R}^N, \rho_w)}^2, \end{aligned}$$

so that again \mathbf{F} is Lipschitz. Since we have assumed that $\mathbf{B} : H \rightarrow L_0(U, H)$ is Lipschitz, we are done. \square

Remark 2.3 (Large Deviation Principle). *The main focus of [28] was a large deviation principle for the stochastic neural field equation (2.5) with small noise, but in a less general situation than we consider here. In particular, the authors only considered the neural field equation driven by a simple additive noise, white in both space and time.*

We would therefore like to remark that in our more general case, and under much weaker conditions than those imposed in [28] (our conditions are for example satisfied for a connectivity function w that is homogeneous, as we will see in Example 2 below), an LDP result for the solution identified by the above theorem still holds and can be quoted from the literature. Indeed, such a result is presented in [33,

Theorem 7.1]. The main conditions required for the application of this result have essentially already been checked above (global Lipschitz properties of \mathbf{F} and \mathbf{B}), and it thus remains to check conditions (E.1) – (E.4) as they appear in [33]. In fact these are trivialities, since the strongly continuous contraction semigroup $S(t)$ is generated by the identity in our case.

2.6 Discussion of conditions on w and ρ in practice

Our knowledge about the kinds of neural field kernels that are found in the brains of mammals is still quite limited. Since visual perception is the most active area of research, it should not come as a surprise that it is in cortical regions involved in visual perception that this knowledge is the most extensive, and in particular in the primary visual area called V1 in humans. In models of this region it is usually assumed that w is the sum of two parts: a local part w_{loc} corresponding to local neuronal connections, and a non-local part w_{lr} corresponding to longer range connections. As suggested in [29, 30], w_{loc} is well approximated by a Gaussian function (or a difference of of such functions, see below):

$$w_{loc}(x, y) = K \exp(-|x - y|^2 / 2\beta_{loc}^2) \quad x, y \in \mathbb{R}^N, \quad K > 0, \quad (2.11)$$

where β_{loc} is the extent of the local connectivity. Hence w_{loc} is isotropic and homogeneous. In fact for practitioners, a very common assumption on w is that it is homogeneous and in $L^1(\mathbb{R}^N)$, which thus concentrates on modeling the local interactions ([9, 8, 10, 22, 27, 31]). However, when w is homogeneous it is clear that neither (C1) nor (C2) of the above theorem are satisfied, and so we instead must try to show that (C1') is satisfied ((C2') trivially holds), and look for solutions in a weighted L^2 space. This is done in the second example below.

Long range connectivity is best described by assuming $N = 2$. It is built upon the existence of maps of orientation sensitivity in which the preferred visual orientation at each point x is represented by a function $\theta(x) \in [0, \pi)$. This function is smooth except at countably many points called the pinwheels where it is undefined⁴. Depending on the species, the long range connections feature an anisotropy, meaning that they tend to align themselves with the preferred orientation at x . One way to take this into account is to introduce the function $A(\chi, x) = \exp[-((1 - \chi)^2 x_1^2 + x_2^2) / 2\beta_{lr}^2]$, where $x = (x_1, x_2)$, $\chi \in [0, 1)$, and β_{lr} is the extent of the long range connectivity. When $\chi = 0$ there is no isotropy (as for the macaque monkey for example) and when $\chi \in (0, 1)$ there is some anisotropy (as for the tree shrew, for example). Let R_α represent the rotation by angle α around the origin. The long range neural field kernel is then defined by [2, 3]

$$w_{lr}(x, y) = \varepsilon_{lr} A(\chi, R_{-2\theta(x)}(x - y)) \cdot G_{\beta_\theta}(\theta(x) - \theta(y)),$$

⁴This would be for an infinite size cortex. The cortex is in effect of finite size but the spatial extents of w_{loc} and w_{lr} are very small with respect to this size and hence the model in which the cortex is \mathbb{R}^2 is acceptable.

where $\varepsilon_{lr} \ll 1$ and G_{β_θ} is the one-dimensional Gaussian density with 0 mean and variance β_θ^2 . Note that w_{lr} is not homogeneous, even in the case $\chi = 0$, because $\theta(x) - \theta(y)$ is not a function of $x - y$. It is easy to verify that $w_{lr} \in L^2(\mathbb{R}^2)$.

Combining the local and non-local parts, one then writes for the neural field kernel of the primary visual area:

$$w_{pva}(x, y) = w_{loc}(x - y) + w_{lr}(x, y). \quad (2.12)$$

In view of our results, in the case where $w = w_{pva}$, since the first part is homogeneous while the second is non-homogeneous but is in $L^2(\mathbb{R}^2)$, we need a combination of the results above. Indeed, the homogeneous part dictates to work in $L^2(\mathbb{R}^2, \rho_{w_{loc}})$ ($\rho_{w_{loc}} \in L^1(\mathbb{R}^2)$). The second kernel dictates to work in $L^2(\mathbb{R}^2)$. But $L^2(\mathbb{R}^2) \in L^2(\mathbb{R}^2, \rho_{w_{loc}})$, because, as shown in Example 2 below $\rho_{w_{loc}}$ can be chosen to be bounded, and hence there is no problem.

Another commonly used type of (homogeneous) neural field kernel, when modeling excitatory and inhibitory populations of neurons is the so-called ‘‘Mexican hat’’ kernel defined by

$$w_{mh}(x, y) = K_1 \exp(-|x - y|^2/2\beta_1^2) - K_2 \exp(-|x - y|^2/2\beta_2^2), \quad x, y \in \mathbb{R}^N, \quad (2.13)$$

for some $K_1, K_2 > 0$. If $\beta_2 > \beta_1$ and $K_1 > K_2$ for example, this is locally excitatory and remotely inhibitory.

It is also important to mention the role of ρ_w from a modeling perspective. The first point is that in the case where w is homogeneous, it is very natural to look for solutions that live in $L^2(\mathbb{R}^N, \rho)$ for some $\rho \in L^1(\mathbb{R}^N)$, rather than in $L^2(\mathbb{R}^N)$. This is because in the deterministic case (see [18]), solutions of interest are of the form of traveling waves, which are constant at ∞ , and thus are not integrable.

Moreover, we emphasize that in Theorem 2.2 and the examples in the next section we identify a single $\rho_w \in L^1(\mathbb{R}^N)$ so that the standard existence result of [35] can be directly applied through Theorem 2.2. We do not claim that this is the only weight ρ for which the solution can be shown to exist in $L^2(\mathbb{R}^N, \rho)$ (see also Example 2 below).

Remark 2.4. *If we replace the spatial coordinate space \mathbb{R}^N by a bounded domain $\mathcal{D} \subset \mathbb{R}^N$, so that the neural field equation (2.5) describes the activity of a neuron found at position $x \in \mathcal{D}$ then checking the conditions as done Theorem 2.2 becomes rather trivial (under appropriate boundary conditions). Indeed, by doing this one can see that there exists a unique $L^2(\mathcal{D})$ -valued solution to (2.5) under the condition (C2’) only (with \mathbb{R}^N replaced by \mathcal{D}). Although working in a bounded domain seems more physical (since any physical section of cortex is clearly bounded), the unbounded case is still often used, see [8] or the review [6], and is mathematically more interesting. The problem in passing to the unbounded case stems from the fact that the nonlocal term in (2.5) naturally ‘lives’ in the space of bounded functions, while according to the theory the noise naturally lives in an L^2 space. These are not compatible when the underlying space is unbounded.*

2.7 Discussion of the noise term in (2.5)

It is important to understand the properties of the noise term in the neural field equation (2.5) which we now know has a solution in some sense. As mentioned above, one particular form of the noise operator \mathbf{B} that is of special importance from a modeling point of view is given by (2.6) i.e.

$$\mathbf{B}(h)(u)(x) = \sigma(h(x)) \int_{\mathbb{R}^N} \varphi(x-y)u(y)dy, \quad x \in \mathbb{R}^N, \quad (2.14)$$

for $h \in L^2(\mathbb{R}^N, \rho)$ and $u \in L^2(\mathbb{R}^N)$, and some functions σ , and φ . This is because such noise terms are spatially correlated depending on φ (as we will see below) and make the link with the original equation (1.1) considered in [8], where spatial correlations are important.

An obvious question is then for which choices of σ and φ can we apply the above results? In particular we need to check that $\mathbf{B}(h)$ is a bounded linear operator from $L^2(\mathbb{R}^N)$ to $L^2(\mathbb{R}^N, \rho)$ for all $h \in L^2(\mathbb{R}^N, \rho)$, and that \mathbf{B} is Lipschitz (assuming as usual that $\rho \in L^1(\mathbb{R}^N)$).

To this end, suppose $\varphi \in L^2(\mathbb{R}^N)$ and that there exists a constant C_σ such that

$$|\sigma(a) - \sigma(b)| \leq C_\sigma |a - b|, \quad \text{and} \quad |\sigma(a)| \leq C_\sigma(1 + |a|), \quad \forall a, b \in \mathbb{R}. \quad (2.15)$$

In other words $\sigma : \mathbb{R} \rightarrow \mathbb{R}$ is assumed to be Lipschitz and of linear growth. Then for any $h \in L^2(\mathbb{R}^N, \rho)$ and $u \in L^2(\mathbb{R}^N)$,

$$\begin{aligned} \|\mathbf{B}(h)(u)\|_{L^2(\mathbb{R}^N, \rho)}^2 &= \int_{\mathbb{R}^N} \sigma^2(h(x)) \left(\int_{\mathbb{R}^N} \varphi(x-y)u(y)dy \right)^2 \rho(x)dx \\ &\leq 2\|u\|_{L^2(\mathbb{R}^N)}^2 \|\varphi\|_{L^2(\mathbb{R}^N)}^2 (\|\rho\|_{L^1(\mathbb{R}^N)} + \|h\|_{L^2(\mathbb{R}^N, \rho)}^2). \end{aligned}$$

Thus $\mathbf{B}(h)$ is indeed a bounded linear operator from $L^2(\mathbb{R}^N)$ to $L^2(\mathbb{R}^N, \rho)$. Moreover, a similar calculation yields the Lipschitz property of \mathbf{B} , so that the above results can be applied. In particular our results hold when $\sigma(a) = \lambda a$, for some $\lambda \in \mathbb{R}$. This is important because it is this choice of σ that is used for the simulations carried out in [8, Section 2.3].

To see the spatial correlation in the noise term in (2.5) when \mathbf{B} has the form (2.14) with $\varphi \in L^2(\mathbb{R}^N)$, consider the case $\sigma \equiv 1$ (so that the noise is purely additive). Then

$$\int_0^t \mathbf{B}(Y(t))dW(t) = \int_0^t B dW(t) =: X(t), \quad t \geq 0,$$

where

$$B(u)(x) = \int_{\mathbb{R}^N} \varphi(x-y)u(y)dy, \quad x \in \mathbb{R}^N, \quad u \in L^2(\mathbb{R}^N),$$

and $X(t)$ is a well-defined $L^2(\mathbb{R}^N, \rho)$ -valued process since B is bounded from $L^2(\mathbb{R}^N)$ into $L^2(\mathbb{R}^N, \rho)$ (see Section 2.2). Moreover, by Theorem 5.2⁵ of [35], $(X(t))_{t \geq 0}$ is Gaussian with mean zero and

$$\text{Cov}(X(t)X(s)) = s \wedge t BQB^*, \quad s, t \geq 0,$$

where $B^* : L^2(\mathbb{R}^N, \rho) \rightarrow L^2(\mathbb{R}^N)$ is the adjoint of B . In other words, for all $g, h \in L^2(\mathbb{R}^N, \rho)$, $s, t \geq 0$, we have, by definition of the covariance operator, that

$$\mathbb{E} \left[\langle g, X(s) \rangle_{L^2(\mathbb{R}^N, \rho)} \langle h, X(t) \rangle_{L^2(\mathbb{R}^N, \rho)} \right] = s \wedge t \langle BQB^*g, h \rangle_{L^2(\mathbb{R}^N, \rho)}.$$

That is, for any $g, h \in L^2(\mathbb{R}^N, \rho)$

$$\begin{aligned} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \mathbb{E} [X(s, x)X(t, y)] g(x)h(y)\rho(x)\rho(y) dx dy &= s \wedge t \langle QB^*h, B^*g \rangle_{L^2(\mathbb{R}^N)} \\ &= s \wedge t \int_{\mathbb{R}^N} QB^*g(z)B^*h(z) dz \\ &= s \wedge t \int_{\mathbb{R}^N} Q^{1/2}B^*g(z)Q^{1/2}B^*h(z) dz. \end{aligned} \quad (2.16)$$

Now, by definition, for $u \in L^2(\mathbb{R}^N)$ and $f \in L^2(\mathbb{R}^N, \rho)$

$$\begin{aligned} \int_{\mathbb{R}^N} u(y)B^*(f)(y) dy &= \int_{\mathbb{R}^N} B(u)(x)f(x)\rho(x) dx \\ &= \int_{\mathbb{R}^N} u(y) \int_{\mathbb{R}^N} \varphi(x-y)f(x)\rho(x) dx dy \end{aligned}$$

so that $B^*(f)(y) = \int_{\mathbb{R}^N} \varphi(x-y)f(x)\rho(x) dx$. Using this in (2.16), we see that

$$\begin{aligned} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \mathbb{E} [X(s, x)X(t, y)] g(x)h(y)\rho(x)\rho(y) dx dy \\ = s \wedge t \int_{\mathbb{R}^N} \left(\int_{\mathbb{R}^N} Q^{1/2}\varphi(x-z)g(x)\rho(x) dx \right) \left(\int_{\mathbb{R}^N} Q^{1/2}\varphi(y-z)h(y)\rho(y) dy \right) dz, \end{aligned}$$

for all $g, h \in L^2(\mathbb{R}^N, \rho)$, since Q is a linear operator and is self-adjoint. We can then conclude that

$$\mathbb{E} [X(s, x)X(t, y)] = s \wedge t \int_{\mathbb{R}^N} Q^{1/2}\varphi(x-z)Q^{1/2}\varphi(y-z) dz = (s \wedge t)c(x-y), \quad (2.17)$$

where $c(x) = Q^{1/2}\varphi * Q^{1/2}\tilde{\varphi}(x)$ and $\tilde{\varphi}(x) = \varphi(-x)$. Hence $(X(t))_{t \geq 0}$ is white in time but stationary and colored in space with covariance function $(s \wedge t)c(x)$. We remark that the manipulations above are certainly not new (they are for example used in [12]), but they illustrate nicely the spatial correlation property of the noise we consider.

We conclude that (2.14) is exactly the rigorous interpretation of the noise described in [8], when interpreting a solution to the stochastic neural field equation as a process taking values in $L^2(\mathbb{R}^N, \rho_w)$.

⁵This can also be obtained by applying the operator B to the representation (2.2) of W .

Remark 2.5. Note that in the case where B is the identity, $X(t) = W(t)$. We can, at least formally, carry out the above computation with $\varphi = \delta_0$ and find that

$$\mathbb{E} [W(s, x)W(t, y)] = (s \wedge t)Q\delta_0(x - y), \quad (2.18)$$

which yields for any $g, h \in L^2(\mathbb{R}^N)$

$$\begin{aligned} & \mathbb{E} [\langle W(s), g \rangle_{L^2(\mathbb{R}^N)} \langle W(t), h \rangle_{L^2(\mathbb{R}^N)}] \\ &= \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \mathbb{E} [W(s, x)W(t, y)] g(x)h(y) dx dy = (s \wedge t) \langle Qg, h \rangle_{L^2(\mathbb{R}^N)}, \end{aligned}$$

which is equation (2.3). Equation (2.18) is the reason why we stated in Section 2.1 that W was a white noise in space and time.

2.8 Examples

As mentioned we now present two important cases where the conditions (C1') and (C2') are satisfied. For convenience, in both cases we in fact show that (C1') is satisfied for some $\rho_w \in L^1(\mathbb{R}^N)$ that is also bounded.

Example 1: $|w|$ defines a compact integral operator. Suppose that

- given $\varepsilon > 0$, there exists $\delta > 0$ and $R > 0$ such that for all $\theta \in \mathbb{R}^N$ with $|\theta| < \delta$

(i) for almost all $x \in \mathbb{R}^N$,

$$\int_{\mathbb{R}^N \setminus B(0, R)} |w(x, y)| dy < \varepsilon, \quad \int_{\mathbb{R}^N} |w(x, y + \theta) - w(x, y)| dy < \varepsilon,$$

(ii) for almost all $y \in \mathbb{R}^N$,

$$\int_{\mathbb{R}^N \setminus B(0, R)} |w(x, y)| dx < \varepsilon, \quad \int_{\mathbb{R}^N} |w(x + \theta, y) - w(x, y)| dx < \varepsilon,$$

where $B(0, R)$ denotes the ball of radius R in \mathbb{R}^N centered at the origin;

- There exists a bounded subset $\Omega \subset \mathbb{R}^N$ of positive measure such that

$$\inf_{y \in \Omega} \int_{\Omega} |w(x, y)| dx > 0, \quad \text{or} \quad \inf_{x \in \Omega} \int_{\Omega} |w(x, y)| dy > 0;$$

- w satisfies (C2') and moreover

$$\forall y \in \mathbb{R}^N \quad (x \mapsto w(x, y)) \in L^1(\mathbb{R}^N), \quad \text{and} \quad \sup_{y \in \mathbb{R}^N} \|w(\cdot, y)\|_{L^1(\mathbb{R}^N)} < \infty.$$

We claim that these assumptions are sufficient for **(C1')** so that we can apply Theorem 2.2 in this case. Indeed, let \mathbb{X} be the Banach space of functions in $L^1(\mathbb{R}^N) \cap L^\infty(\mathbb{R}^N)$ equipped with the norm $\|\cdot\|_{\mathbb{X}} = \max\{\|\cdot\|_{L^1(\mathbb{R}^N)}, \|\cdot\|_{L^\infty(\mathbb{R}^N)}\}$. Thanks to the last point above, we can well-define the map $J : \mathbb{X} \rightarrow \mathbb{X}$ by

$$Jh(y) = \int_{\mathbb{R}^N} |w(x, y)|h(x)dx, \quad h \in \mathbb{X}.$$

Moreover, it follows from [19, Corollary 5.1] that the first condition we have here imposed on w is in fact necessary and sufficient for both the operators $J : L^1(\mathbb{R}^N) \rightarrow L^1(\mathbb{R}^N)$ and $J : L^\infty(\mathbb{R}^N) \rightarrow L^\infty(\mathbb{R}^N)$ to be compact. We therefore clearly also have that the condition is necessary and sufficient for the operator $J : \mathbb{X} \rightarrow \mathbb{X}$ to be compact.

Note now that the space \mathbb{K} of positive functions in \mathbb{X} is a cone in \mathbb{X} such that $J(\mathbb{K}) \subset \mathbb{K}$, and that the cone is *reproducing* (i.e. $\mathbb{X} = \{f - g : f, g \in \mathbb{K}\}$). If we can show that $r(J)$ is strictly positive, we can thus finally apply the Krein-Rutman Theorem (see for example [17, Theorem 1.1]) to see that $r(J)$ is an eigenvalue with corresponding non-zero eigenvector $\rho \in \mathbb{K}$.

To show that $r(J) > 0$, suppose first of all that there exists a bounded $\Omega \subset \mathbb{R}^N$ of positive measure such that $\inf_{y \in \Omega} \int_{\Omega} |w(x, y)|dx > 0$. Define $h = 1$ on Ω , 0 elsewhere, so that $\|h\|_{\mathbb{X}} = \max\{1, |\Omega|\}$. Then, trivially,

$$\|Jh\|_{\mathbb{X}} \geq \sup_{y \in \mathbb{R}^N} \int_{\Omega} |w(x, y)|dx \geq \inf_{y \in \Omega} \int_{\Omega} |w(x, y)|dx =: m > 0,$$

by assumption. Replacing h by $\tilde{h} = h / \max\{1, |\Omega|\}$ yields $\|\tilde{h}\|_{\mathbb{X}} = 1$ and

$$\|J\tilde{h}\|_{\mathbb{X}} \geq m / \max\{1, |\Omega|\}.$$

Thus $\|J\| \geq m / \max\{1, |\Omega|\}$. Similarly

$$\begin{aligned} \|J^2h\|_{\mathbb{X}} &\geq \sup_{y \in \mathbb{R}^N} \int_{\mathbb{R}^N} |w(x_1, y)| \left(\int_{\Omega} |w(x_2, x_1)|dx_2 \right) dx_1 \\ &\geq \int_{\mathbb{R}^N} |w(x_1, y)| \left(\int_{\Omega} |w(x_2, x_1)|dx_2 \right) dx_1, \quad \forall y \in \mathbb{R}^N \\ &\geq \inf_{x_1 \in \Omega} \left(\int_{\Omega} |w(x_2, x_1)|dx_2 \right) \int_{\Omega} |w(x_1, y)|dx_1, \quad \forall y \in \mathbb{R}^N. \end{aligned}$$

Therefore

$$\|J^2h\|_{\mathbb{X}} \geq m^2,$$

so that $\|J^2\| \geq m^2 / \max\{1, |\Omega|\}$. In fact we have $\|J^k\| \geq m^k / \max\{1, |\Omega|\}$ for all $k \geq 1$, so that, by the spectral radius formula, $r(J) \geq m > 0$. The case where $\inf_{x \in \Omega} \int_{\Omega} |w(x, y)|dy > 0$ holds instead is proved similarly, by instead taking

$h = 1/|\Omega|$ on Ω (0 elsewhere) and working with the $L^1(\mathbb{R}^N)$ norm of Jh in place of the $L^\infty(\mathbb{R}^N)$ norm.

We have thus found a non-negative, non-zero function $\rho = \rho_w \in L^1(\mathbb{R}^N) \cap L^\infty(\mathbb{R}^N)$ such that

$$\int_{\mathbb{R}^N} |w(x, y)| \rho_w(x) dx = r(J) \rho_w(y), \quad \forall y \in \mathbb{R}^N,$$

so that **(C1')** is satisfied.

Example 2: Homogeneous case. Suppose that

- w is homogeneous i.e $w(x, y) = w(x - y)$ for all $x, y \in \mathbb{R}^N$;
- $w \in L^1(\mathbb{R}^N)$ and is continuous;
- $\int_{\mathbb{R}^N} |x|^{2N} |w(x)| dx < \infty$.

These conditions are satisfied for many typical choices of the neural field kernel in the literature (e.g. the ‘‘Mexican hat’’ kernel [7, 20, 31, 41] and (2.13) above). However, it is clear that we are not in the case of the previous example, since for any $R > 0$

$$\sup_{x \in \mathbb{R}^N} \int_{\mathbb{R}^N \setminus B(0, R)} |w(x - y)| dy = \|w\|_{L^1(\mathbb{R}^N)},$$

which is not uniformly small. We thus again show that **(C1')** is satisfied in this case so that (since **(C2')** is trivially satisfied) Theorem 2.2 yields the existence of a unique $L^2(\mathbb{R}^N, \rho_w)$ -valued solution to (2.5).

In order to do this, we use the Fourier transform. Let $v = |w|$, so that v is continuous and in $L^1(\mathbb{R}^N)$. Let $\mathfrak{F}v$ be the Fourier transform of v i.e.

$$\mathfrak{F}v(\xi) := \int_{\mathbb{R}^N} e^{-2\pi i x \cdot \xi} v(x) dx, \quad \xi \in \mathbb{R}^N.$$

Therefore $\mathfrak{F}v$ is continuous and bounded by

$$\sup_{\xi \in \mathbb{R}^N} |\mathfrak{F}v(\xi)| \leq \|v\|_{L^1(\mathbb{R}^N)} = \|w\|_{L^1(\mathbb{R}^N)}.$$

Now let $\Lambda_w = \|w\|_{L^1(\mathbb{R}^N)} + 1$, and $z(x) := e^{-|x|^2/2}$, $x \in \mathbb{R}^N$, so that z is in the Schwartz space of smooth rapidly decreasing functions, which we denote by $\mathcal{S}(\mathbb{R}^N)$. Then define

$$\hat{\rho}(\xi) := \frac{\mathfrak{F}z(\xi)}{\Lambda_w - \mathfrak{F}v(\xi)}. \quad (2.19)$$

We note that the denominator is continuous and strictly bounded away from 0 (indeed by construction $\Lambda_w - \mathfrak{F}v(\xi) \geq 1$ for all $\xi \in \mathbb{R}^N$). Thus $\hat{\rho}$ is continuous,

bounded and in $L^1(\mathbb{R}^N)$ (since $\mathfrak{F}z \in \mathcal{S}(\mathbb{R}^N)$ by the standard stability result for the Fourier transform on $\mathcal{S}(\mathbb{R}^N)$).

We now claim that $\mathfrak{F}^{-1}\hat{\rho}(x) \in L^1(\mathbb{R}^N)$, where the map \mathfrak{F}^{-1} is defined by

$$\mathfrak{F}^{-1}g(x) := \int_{\mathbb{R}^N} e^{2\pi i x \cdot \xi} g(\xi) d\xi, \quad g \in L^1(\mathbb{R}^N).$$

Indeed, we note that for any $k \in \{1, \dots, N\}$,

$$\partial_k^{2N} \mathfrak{F}v(\xi) = (-2\pi i)^{2N} \int_{\mathbb{R}^N} e^{-2\pi i x \cdot \xi} x_k^{2N} v(x) dx,$$

which is well-defined and bounded thanks to our assumption on the integrability of $x \mapsto |x|^{2N}|w(x)|$. Since $\mathfrak{F}z$ is rapidly decreasing, we can thus see that the function $\hat{\rho}(\xi)$ is $2N$ times differentiable with respect to every component and $\partial_k^{2N} \hat{\rho}(\xi)$ is absolutely integrable for every $k \in \{1, \dots, N\}$. Finally, since $\mathfrak{F}^{-1}(\partial_k^{2N} \hat{\rho})(x) = (2\pi i)^{2N} x_k^{2N} \mathfrak{F}^{-1}\hat{\rho}(x)$ for each $k \in \{1, \dots, N\}$, we have that

$$|\mathfrak{F}^{-1}\hat{\rho}(x)| \leq \frac{\sum_{k=1}^N |\mathfrak{F}^{-1}(\partial_k^{2N} \hat{\rho})(x)|}{(2\pi)^{2N} \sum_{k=1}^N x_k^{2N}} \leq \frac{N^{N-1} \sum_{k=1}^N \|\partial_k^{2N} \hat{\rho}\|_{L^1(\mathbb{R}^N)}}{(2\pi)^{2N} |x|^{2N}},$$

for all $x \in \mathbb{R}^N$. Thus there exists a constant K such that $|\mathfrak{F}^{-1}\hat{\rho}(x)| \leq K/|x|^{2N}$. Moreover, since we also have the trivial bound

$$|\mathfrak{F}^{-1}\hat{\rho}(x)| \leq \|\hat{\rho}\|_{L^1(\mathbb{R}^N)},$$

for all $x \in \mathbb{R}^N$, it follows that $|\mathfrak{F}^{-1}\hat{\rho}(x)| \leq K/(1+|x|^{2N})$, by adjusting the constant K . Since this is integrable over \mathbb{R}^N , the claim is proved.

Now, by the classical Fourier Inversion Theorem (which is applicable since $\hat{\rho}$ and $\mathfrak{F}^{-1}\hat{\rho}$ are both in $L^1(\mathbb{R}^N)$), we thus have that

$$\mathfrak{F}(\mathfrak{F}^{-1}\hat{\rho})(\xi) = \hat{\rho}(\xi),$$

for all $\xi \in \mathbb{R}^N$.

By setting $\rho(x) = \mathfrak{F}^{-1}\hat{\rho}(x)$, we see that

$$\Lambda_w \mathfrak{F}\rho(\xi) - \mathfrak{F}\rho(\xi) \mathfrak{F}v(\xi) := \mathfrak{F}z(\xi).$$

We may finally again apply the inverse Fourier transform \mathfrak{F}^{-1} to both sides, so that by the Inversion Theorem again (along with the standard convolution formula) it holds that

$$\Lambda_w \rho(y) - \int_{\mathbb{R}^N} v(x-y) \rho(x) dx = e^{-\frac{|y|^2}{2}}, \quad y \in \mathbb{R}^N.$$

It then follows that

$$\int_{\mathbb{R}^N} |w(x-y)| \rho(x) dx \leq \Lambda_w \rho(y), \quad y \in \mathbb{R}^N,$$

as claimed.

Moreover, equation (2.19) shows that $\hat{\rho}(\xi)$ is in Schwartz space, hence so is ρ , implying that it is bounded. Note that equation (2.19) provides a way of explicitly computing one possible function ρ_w appearing in condition (C1') in the cases where the neural field kernel is homogeneous (for example given by (2.11) and (2.13)). That particular function can be varied for example by changing the function z and/or the constant Λ_w .

3 Stochastic neural fields as Gaussian random fields

In this section we take an alternative approach, and try to give sense to a solution to the stochastic neural field equation (1.1) as a random field, using Walsh's theory of integration.

This approach generally takes as its starting point a deterministic PDE, and then attempts include a term which is random in both space and time. With this in mind, consider first the well studied deterministic neural field equation

$$\partial_t Y(t, x) = -Y(t, x) + \int_{\mathbb{R}^N} w(x, y)G(Y(t, y))dy, \quad x \in \mathbb{R}^N, \quad t \geq 0. \quad (3.1)$$

Under some conditions on the neural field kernel w (boundedness, condition (C2') above and L^1 -Lipschitz continuity), this equation has a unique solution $(t, x) \mapsto Y(t, x)$ that is bounded and continuous in x and continuously differentiable in t , whenever $x \mapsto Y(0, x)$ is bounded and continuous ([34]).

The idea then is to directly add a noise term to this equation, and try and give sense to all the necessary objects in order to be able to define what we mean by a solution. Indeed, consider the following stochastic version of (3.1),

$$\partial_t Y(t, x) = -Y(t, x) + \int_{\mathbb{R}^N} w(x, y)G(Y(t, y))dy + \sigma(Y(t, x))\dot{W}(t, x) \quad (3.2)$$

where \dot{W} is a "space-time white noise". Informally we may think of the object $\dot{W}(t, x)$ as the random *distribution* which, when integrated against a test function $h \in L^2(\mathbb{R}^+ \times \mathbb{R}^N)$

$$\dot{W}(h) := \int_0^\infty \int_{\mathbb{R}^N} h(t, x)\dot{W}(t, x)dt dx, \quad h \in L^2(\mathbb{R}^+ \times \mathbb{R}^N),$$

yields a zero-mean Gaussian random field $(\dot{W}(h))_{h \in L^2(\mathbb{R}^+ \times \mathbb{R}^N)}$ with covariance

$$\mathbb{E} \left[\dot{W}(g)\dot{W}(h) \right] = \int_0^\infty \int_{\mathbb{R}^N} g(t, x)h(t, x)dx dt, \quad g, h \in L^2(\mathbb{R}^+ \times \mathbb{R}^N).$$

The point is that with this interpretation of space-time white noise, since equation (3.2) specifies no regularity in the spatial direction (the map $x \mapsto Y(t, x)$ is

simply assumed to be Lebesgue measurable so that the integral makes sense), it is clear that any solution will be *distribution*-valued in the spatial direction, which is rather unsatisfactory. Indeed, consider the extremely simple linear case when $G \equiv 0$ and $\sigma \equiv 1$, so that (3.2) reads

$$\partial_t Y(t, x) = -Y(t, x) + \dot{W}(t, x). \quad (3.3)$$

Formally, the solution to this equation is given by

$$Y(t, x) = e^{-t}Y(0, x) + \int_0^t e^{-(t-s)}\dot{W}(s, x)ds, \quad t \geq 0, x \in \mathbb{R}^N,$$

and since the integral is only over time it is clear (at least formally) that $x \mapsto Y(t, x)$ is a distribution for all $t \geq 0$. This differs significantly from the usual SPDE situation, when one would typically have an equation such as (3.3) where a second order differential operator in space is applied to the first term on the right-hand side (leading to the much studied *stochastic heat equation*). In such a case, the semigroup generated by the second order differential operator can be enough to smooth the space-time white noise in the spatial direction, leading to solutions that are continuous in both space and time (at least when the spatial dimension is 1 – see for example [32, Chapter 3] or [42, Chapter 3]).

Of course one can develop a theory of distribution-valued processes (as is done in [42, Chapter 4]) to interpret solutions of (3.2) in the obvious way: one says that the random field $(Y(t, x))_{t \geq 0, x \in \mathbb{R}^N}$ is a (weak) solution to (3.2) if for all $\phi \in C_0^\infty(\mathbb{R}^N)$ it holds that

$$\begin{aligned} \int_{\mathbb{R}^N} \phi(x)Y(t, x)dx &= e^{-t} \int_{\mathbb{R}^N} \phi(x)Y(0, x)dx \\ &+ \int_0^t \int_{\mathbb{R}^N} e^{-(t-s)}\phi(x) \int_{\mathbb{R}^N} w(x, y)G(Y(s, y))dydxds \\ &+ \int_0^t \int_{\mathbb{R}^N} e^{-(t-s)}\phi(x)\sigma(Y(s, x))\dot{W}(s, x)dxds, \end{aligned}$$

for all $t \geq 0$. Here all the integrals can be well-defined, which makes sense intuitively if we think of $\dot{W}(t, x)$ as a distribution. In fact it is more common to write $\int_0^t \int_{\mathbb{R}^N} e^{-(t-s)}\phi(x)W(dsdx)$ for the stochastic integral term, once it has been rigorously defined.

However, we argue that it is not worth developing this theory here, since distribution valued solutions are of little interest physically. It is for this reason that we instead look for other types of random noise to add to the deterministic equation (3.1) which in particular will be correlated in space that will produce solutions that are real-valued random fields, and are at least Hölder continuous in both space and time. In the theory of SPDEs, when the spatial dimension is 2 or more, the problem of an equation driven by space-time white noise having no real-valued solution is a well-known and much studied one (again see for example [32, Chapter 3] or [42,

Chapter 3] for a discussion of this). To get around the problem, a common approach ([14, 21, 37]) is to consider random noises that are smoother than white noise, namely a Gaussian noise that is white in time but has a smooth spatial covariance. Such random noise is known as either spatially colored or spatially homogeneous white-noise. One can then formulate conditions on the covariance function to ensure that real-valued Hölder continuous solutions to the specific SPDE exist.

It should also be mentioned, as remarked in [14], that in trying to model physical situations, there is some evidence that white-noise smoothed in the spatial direction is more natural, since spatial correlations are typically of a much larger order of magnitude than time correlations.

In the stochastic neural field case, since we have no second order differential operator, our solution will only ever be as smooth as the noise itself. We therefore look to add a noise term to (3.1) that is at least Hölder continuous in the spatial direction instead of pure white noise, and then proceed to look for solutions to the resulting equation in the sense of Walsh.

The section is structured as follows. First we briefly introduce Walsh's theory of stochastic integration, for which the classical reference is [42]. This theory will be needed to well-define the stochastic integral in our definition of a solution to the neural field equation. We then introduce the spatially smoothed space-time white noise that we will consider, before finally applying the theory to analyze solutions of the neural field equation driven by this spatially smoothed noise under certain conditions.

3.1 Walsh's stochastic integral

We will not go into the details of the construction of Walsh's stochastic integral, since a very nice description is given by D. Khoshnevisan in [13] (see also [42]). Instead we present the bare essentials needed in the following sections.

The elementary object of study is the centered Gaussian random field⁶

$$\dot{W} := (\dot{W}(A))_{A \in \mathcal{B}(\mathbb{R}^+ \times \mathbb{R}^N)}$$

indexed by $A \in \mathcal{B}(\mathbb{R}^+ \times \mathbb{R}^N)$ (where $\mathbb{R}^+ := [0, \infty)$) with covariance function

$$\mathbb{E} \left[\dot{W}(A) \dot{W}(B) \right] = |A \cap B|, \quad A, B, \in \mathcal{B}(\mathbb{R}^+ \times \mathbb{R}^N), \quad (3.4)$$

where $|A \cap B|$ denotes the Lebesgue measure of $A \cap B$. We say that \dot{W} is a *white noise* on $\mathbb{R}^+ \times \mathbb{R}^N$. We then define the *white noise process* $W := (W_t(A))_{t \geq 0, A \in \mathcal{B}(\mathbb{R}^N)}$ by

$$W_t(A) := \dot{W}([0, t] \times A), \quad t \geq 0. \quad (3.5)$$

⁶Recall that a collection of random variables $X = \{X(\theta)\}_{\theta \in \Theta}$ indexed by a set Θ is a Gaussian random field on Θ if $(X(\theta_1), \dots, X(\theta_k))$ is a k -dimensional Gaussian random vector for every $\theta_1, \dots, \theta_k \in \Theta$. It is characterized by its mean and covariance functions.

Now define the norm

$$\|f\|_W^2 := \mathbb{E} \left[\int_0^T \int_{\mathbb{R}^N} |f(t, x)|^2 dt dx \right], \quad (3.6)$$

for any (random) function f that is knowable⁷ at time t given $(W_s(A))_{s \leq t, A \in \mathcal{B}(\mathbb{R}^N)}$. Then let \mathfrak{P}_W be the set of all such functions f for which $\|f\|_W < \infty$. The point is that this space forms the set of integrands that can be integrated against the white noise process according to Walsh's theory.

Indeed, we have then following theorem ([42, Theorem 2.5]).

Theorem 3.1. *For all $f \in \mathfrak{P}_W$, $t \in [0, T]$ and $A \in \mathcal{B}(\mathbb{R}^N)$,*

$$\int_0^t \int_A f(s, x) W(ds dx)$$

can be well-defined in $L^2(\Omega, \mathcal{F}, \mathbb{P})$. Moreover for all $t \in (0, T]$ and $A, B \in \mathcal{B}(\mathbb{R}^N)$, $\mathbb{E}[\int_0^t \int_A f(s, x) W(ds dx)] = 0$ and

$$\mathbb{E} \left[\int_0^t \int_A f(s, x) W(ds dx) \int_0^t \int_B f(s, x) W(ds dx) \right] = \mathbb{E} \left[\int_0^t \int_{A \cap B} f^2(s, x) dx dt \right].$$

The following inequality will also be fundamental:

Theorem 3.2 (Burkholder's inequality). *For all $p \geq 2$ there exists a constant c_p (with $c_2 = 1$) such that for all $f \in \mathfrak{P}_W$, $t \in (0, T]$ and $A \in \mathcal{B}(\mathbb{R}^N)$,*

$$\mathbb{E} \left[\left| \int_0^t \int_A f(s, x) W(ds dx) \right|^p \right] \leq c_p \mathbb{E} \left[\left(\int_0^T \int_{\mathbb{R}^N} |f(t, x)|^2 dt dx \right)^{\frac{p}{2}} \right].$$

3.2 Spatially smoothed space-time white noise

Let $W = (W_t(A))_{t \geq 0, A \in \mathcal{B}(\mathbb{R}^N)}$ be a white-noise process as defined in the previous section. For $\varphi \in L^2(\mathbb{R}^N)$, we can well-define the (Gaussian) random field $(W^\varphi(t, x))_{t \geq 0, x \in \mathbb{R}^N}$ for any $T > 0$ by

$$W^\varphi(t, x) := \int_0^t \int_{\mathbb{R}^N} \varphi(x - y) W(ds dy). \quad (3.7)$$

To see this one just needs to check that $\varphi(x - \cdot) \in \mathfrak{P}_W$ for every x , where \mathfrak{P}_W is as above. The function $\varphi(x - \cdot)$ is clearly completely determined by W for each x

⁷Precisely we consider functions f such that $(t, x, \omega) \mapsto f(t, x, \omega)$ is measurable with respect to the σ -algebra generated by linear combinations of functions of the form $X(\omega) \mathbf{1}_{(a, b]}(t) \mathbf{1}_A(x)$, where $a, b \in \mathbb{R}^+$, $A \in \mathcal{B}(\mathbb{R}^N)$, and $X : \Omega \rightarrow \mathbb{R}$ is bounded and measurable with respect to the σ -algebra generated by $(W_s(A))_{s \leq a, A \in \mathcal{B}(\mathbb{R}^N)}$.

(since it is non-random) and for every $T > 0$

$$\begin{aligned}\|\varphi(x - \cdot)\|_W^2 &= \mathbb{E} \left[\int_0^T \int_{\mathbb{R}^N} |\varphi(x - z)|^2 dt dz \right] \\ &= T \|\varphi\|_{L^2(\mathbb{R}^N)}^2 < \infty,\end{aligned}$$

so that the integral in (3.7) is indeed well-defined in the sense of the above construction. Moreover, by Theorem 3.1 the random field $(W^\varphi(t, x))_{t \geq 0, x \in \mathbb{R}^N}$ has spatial covariance

$$\begin{aligned}\mathbb{E}[W^\varphi(t, x)W^\varphi(t, y)] &= \mathbb{E} \left[\int_0^t \int_{\mathbb{R}^N} \varphi(x - z) W(ds dz) \int_0^t \int_{\mathbb{R}^N} \varphi(y - z) W(ds dz) \right] \\ &= \int_0^t \int_{\mathbb{R}^N} \varphi(x - z)\varphi(y - z) dz ds = t\varphi \star \tilde{\varphi}(x - y),\end{aligned}$$

where \star denotes the convolution operator as usual, and $\tilde{\varphi}(x) = \varphi(-x)$. Thus the random field $(W^\varphi(t, x))_{t \geq 0, x \in \mathbb{R}^N}$ is spatially correlated.

The regularity in time of this process is the same as that of a Brownian path:

Lemma 3.3. *For any $x \in \mathbb{R}^N$, the path $t \mapsto W^\varphi(t, x)$ has an η -Hölder continuous modification for any $\eta \in (0, 1/2)$.*

Proof. For $x \in \mathbb{R}^N$, $s, t \geq 0$ with $s \leq t$ and any $p \geq 2$ we have by Burkholder's inequality (Theorem 3.2 above) that

$$\mathbb{E}[|W^\varphi(t, x) - W^\varphi(s, x)|^p] \leq c_p \|\varphi\|_{L^2(\mathbb{R}^N)}^2 (t - s)^{\frac{p}{2}}.$$

The result follows from the standard Kolmogorov continuity theorem (see for example Theorem 4.3 of [13, Chapter 1]). \square

More importantly, if we impose some (very weak) regularity on φ then W^φ inherits some spatial regularity:

Lemma 3.4. *Suppose that there exists a constant C_φ such that*

$$\|\varphi - \tau_z(\varphi)\|_{L^2(\mathbb{R}^N)} \leq C_\varphi |z|^\alpha, \quad \forall z \in \mathbb{R}^N, \quad (3.8)$$

for some $\alpha \in (0, 1]$, where τ_z indicates the shift by z operator (so that $\tau_z(\varphi)(y) := \varphi(y + z)$ for all $y, z \in \mathbb{R}^N$). Then for all $t \geq 0$, the map $x \mapsto W^\varphi(t, x)$ has an η -Hölder continuous modification, for any $\eta \in (0, \alpha)$.

Proof. For $x, \tilde{x} \in \mathbb{R}^N$, $t \geq 0$, and any $p \geq 2$ we have (again by Burkholder's inequality) that

$$\begin{aligned}\mathbb{E}[|W^\varphi(t, x) - W^\varphi(t, \tilde{x})|^p] &\leq t^{\frac{p}{2}} c_p \left(\int_{\mathbb{R}^N} |\varphi(x - y) - \varphi(\tilde{x} - y)|^2 dy \right)^{\frac{p}{2}} \\ &= t^{\frac{p}{2}} c_p \left(\int_{\mathbb{R}^N} |\varphi(y) - \varphi(y + \tilde{x} - x)|^2 dy \right)^{\frac{p}{2}} \\ &\leq t^{\frac{p}{2}} c_p C_\varphi^p |x - \tilde{x}|^{p\alpha}.\end{aligned}$$

The result follows by Kolmogorov's continuity theorem. \square

Remark 3.5. The condition (3.8) with $\alpha = 1$ is true if and only if the function φ is in the Sobolev space $W^{1,2}(\mathbb{R}^N)$ ([11, Proposition 9.3]).

When $\alpha < 1$ the set of functions $\varphi \in L^2(\mathbb{R}^N)$ which satisfy (3.8) defines a Banach space denoted by $N^{\alpha,2}(\mathbb{R}^N)$ which is known as the Nikolskii space. This space is closely related to the more familiar fractional Sobolev space $W^{\alpha,2}(\mathbb{R}^N)$ though they are not identical. We refer to [40] for a detailed study of such spaces and their relationships. An example of when (3.8) holds with $\alpha = 1/2$ is found by taking φ to be an indicator function. It is in this way we see that (3.8) is a rather weak condition.

3.3 The stochastic neural field equation driven by spatially smoothed space-time white noise

We now have everything in place to define and study the solution to the stochastic neural field equation driven by a spatially smoothed space-time white noise. Indeed, consider the equation

$$\partial_t Y(t, x) = -Y(t, x) + \int_{\mathbb{R}^N} w(x, y) G(Y(t, y)) dy + \sigma(Y(t, x)) \frac{\partial}{\partial t} W^\varphi(t, x), \quad (3.9)$$

with initial condition $Y(0, x) = Y_0(x)$ for $x \in \mathbb{R}^N$ and $t \geq 0$, where $(W^\varphi(t, x))_{t \geq 0, x \in \mathbb{R}^N}$ is the spatially smoothed space-time white noise defined by (3.7) for some $\varphi \in L^2(\mathbb{R}^N)$. As above, we will impose Lipschitz assumptions on σ and G , by supposing that

- $\sigma : \mathbb{R} \rightarrow \mathbb{R}$ is globally Lipschitz (exactly as in (2.15)) i.e. there exists a constant C_σ such that

$$|\sigma(a) - \sigma(b)| \leq C_\sigma |a - b|, \quad \text{and} \quad |\sigma(a)| \leq C_\sigma (1 + |a|), \quad \forall a, b \in \mathbb{R};$$

- $G : \mathbb{R} \rightarrow \mathbb{R}$ is bounded and globally Lipschitz (exactly as above) i.e such that there exists a constant C_G with $\sup_{a \in \mathbb{R}} |G(a)| \leq C_G$ and

$$|G(a) - G(b)| \leq C_G |a - b|, \quad \forall a, b \in \mathbb{R}.$$

Although the above equation is not well-defined ($\frac{\partial}{\partial t} W^\varphi(t, x)$ does not exist), we will interpret a solution to (3.9) in the following way.

Definition 3.6. By a solution to (3.9) we will mean a real-valued random field $(Y(t, x))_{t \geq 0, x \in \mathbb{R}^N}$ such that

$$\begin{aligned} Y(t, x) = & e^{-t} Y_0(x) + \int_0^t e^{-(t-s)} \int_{\mathbb{R}^N} w(x, y) G(Y(s, y)) dy ds \\ & + \int_0^t \int_{\mathbb{R}^N} e^{-(t-s)} \sigma(Y(s, x)) \varphi(x - y) W(ds dy), \end{aligned} \quad (3.10)$$

almost surely for all $t \geq 0$ and $x \in \mathbb{R}^N$, where the stochastic integral term is understood in the sense described in Section 3.1.

Once again we are interested in the conditions on the neural field kernel w that allow us to prove the existence of solutions in this new sense. Recall that in Section 2 we either required conditions **(C1)** and **(C2)** or **(C1')** and **(C2')** to be satisfied. The difficulty was to keep everything well-behaved in the Hilbert space $L^2(\mathbb{R}^N)$ (or $L^2(\mathbb{R}^N, \rho)$). However, when looking for solutions in the sense of random fields $(Y(t, x))_{t \geq 0, x \in \mathbb{R}^N}$ such that (3.10) is satisfied, such restrictions are no longer needed, principally because we no longer have to concern ourselves with the behavior in space at infinity. Indeed, in this section we simply work with the condition **(C2')** i.e. that

$$\forall x \in \mathbb{R}^N \ (y \mapsto w(x, y)) \in L^1(\mathbb{R}^N), \quad \text{and} \quad \sup_{x \in \mathbb{R}^N} \|w(x, \cdot)\|_{L^1(\mathbb{R}^N)} \leq C_w,$$

for some constant C_w . Using the standard technique of a Picard iteration scheme (closely following [42, Theorem 3.2]) and the simple properties of the Walsh stochastic integral stated in Section 3.1, we can prove the following:

Theorem 3.7. *Suppose that the map $x \mapsto Y_0(x)$ is Borel measurable almost surely, and that*

$$\sup_{x \in \mathbb{R}^N} \mathbb{E} [|Y_0(x)|^2] < \infty.$$

*Suppose moreover that the neural field kernel w satisfies condition **(C2')**. Then there exists an almost surely unique predictable random field $(Y(t, x))_{t \geq 0, x \in \mathbb{R}^N}$ which is a solution to (3.9) in the sense of Definition 3.6 such that*

$$\sup_{t \in [0, T], x \in \mathbb{R}^N} \mathbb{E} [|Y(t, x)|^2] < \infty, \tag{3.11}$$

for any $T > 0$.

Proof. The proof proceeds in a classical way, but where we are careful to interpret all stochastic integrals as described in Section 3.1, and so we provide the details.

Uniqueness: Suppose that $(Y(t, x))_{t \geq 0, x \in \mathbb{R}^N}$ and $(Z(t, x))_{t \geq 0, x \in \mathbb{R}^N}$ are both solutions to (3.9) in the sense of Definition 3.6. Let $D(t, x) = Y(t, x) - Z(t, x)$ for $x \in \mathbb{R}^N$ and $t \geq 0$. Then we have

$$\begin{aligned} D(t, x) &= \int_0^t e^{-(t-s)} \int_{\mathbb{R}^N} w(x, y) [G(Y(s, y)) - G(Z(s, y))] dy ds \\ &\quad + \int_0^t \int_{\mathbb{R}^N} e^{-(t-s)} [\sigma(Y(s, x)) - \sigma(Z(s, x))] \varphi(x - y) W(ds dy). \end{aligned}$$

Therefore

$$\begin{aligned}
\mathbb{E} [|D(t, x)|^2] &\leq 2\mathbb{E} \left[\left(\int_0^t e^{-(t-s)} \int_{\mathbb{R}^N} |w(x, y)| |G(Y(s, y)) - G(Z(s, y))| dy ds \right)^2 \right] \\
&\quad + 2\mathbb{E} \left[\left(\int_0^t \int_{\mathbb{R}^N} e^{-(t-s)} [\sigma(Y(s, x)) - \sigma(Z(s, x))] \varphi(x - y) W(ds dy) \right)^2 \right] \\
&\leq 2t \int_0^t e^{-2(t-s)} \mathbb{E} \left[\left(\int_{\mathbb{R}^N} |w(x, y)| |G(Y(s, y)) - G(Z(s, y))| dy \right)^2 \right] ds \\
&\quad + 2 \int_0^t \int_{\mathbb{R}^N} e^{-2(t-s)} \mathbb{E} [|\sigma(Y(s, x)) - \sigma(Z(s, x))|^2] |\varphi(x - y)|^2 ds dy,
\end{aligned}$$

where we have used Cauchy-Schwarz and Burkholder's inequality (Theorem 3.2) with $p = 2$. Thus, using the Lipschitz property of σ and G ,

$$\begin{aligned}
\mathbb{E} [|D(t, x)|^2] &\leq 2tC_G^2 \int_0^t e^{-2(t-s)} \mathbb{E} \left[\left(\int_{\mathbb{R}^N} |w(x, y)| |D(s, y)| dy \right)^2 \right] ds \\
&\quad + 2C_\sigma^2 \|\varphi\|_{L^2(\mathbb{R}^N)}^2 \int_0^t e^{-2(t-s)} \mathbb{E} [|D(s, x)|^2] ds.
\end{aligned}$$

By the Cauchy-Schwarz inequality once again

$$\begin{aligned}
\mathbb{E} [|D(t, x)|^2] &\leq 2tC_G^2 \|w(x, \cdot)\|_{L^1(\mathbb{R}^N)} \int_0^t e^{-2(t-s)} \int_{\mathbb{R}^N} |w(x, y)| \mathbb{E} [|D(s, y)|^2] dy ds \\
&\quad + 2C_\sigma^2 \|\varphi\|_{L^2(\mathbb{R}^N)}^2 \int_0^t e^{-2(t-s)} \mathbb{E} [|D(s, x)|^2] ds.
\end{aligned}$$

Let $H(s) := \sup_{x \in \mathbb{R}^N} \mathbb{E} [|D(s, x)|^2]$, which is finite since we are assuming Y and Z satisfy (3.11). Writing $K = 2 \max\{C_\sigma^2, C_G^2\}$, we have

$$\begin{aligned}
\mathbb{E} [|D(t, x)|^2] &\leq K \left[tC_w^2 + \|\varphi\|_{L^2(\mathbb{R}^N)}^2 \right] \int_0^t e^{-2(t-s)} H(s) ds \\
&\Rightarrow H(t) \leq K \left[tC_w^2 + \|\varphi\|_{L^2(\mathbb{R}^N)}^2 \right] \int_0^t H(s) ds.
\end{aligned}$$

An application of Gronwall's lemma then yields $\sup_{s \leq t} H(s) = 0$ for all $t \geq 0$. Hence $Y(t, x) = Z(t, x)$ almost surely for all $t \geq 0, x \in \mathbb{R}^N$.

Existence: Let $Y_0(t, x) = Y_0(x)$. Then define iteratively for $n \in \mathbb{N}_0, t \geq 0, x \in \mathbb{R}^N$,

$$\begin{aligned}
Y_{n+1}(t, x) &:= e^{-t} Y_0(x) + \int_0^t e^{-(t-s)} \int_{\mathbb{R}^N} w(x, y) G(Y_n(s, y)) dy ds \\
&\quad + \int_0^t \int_{\mathbb{R}^N} e^{-(t-s)} \sigma(Y_n(s, x)) \varphi(x - y) W(ds dy). \tag{3.12}
\end{aligned}$$

We first check that the stochastic integral is well-defined, under the assumption that

$$\sup_{t \in [0, T], x \in \mathbb{R}^N} \mathbb{E}(|Y_n(t, x)|^2) < \infty, \quad (3.13)$$

for any $T > 0$, which we know is true for $n = 0$ by assumption, and we show by induction is also true for each integer $n \geq 1$ below. To this end for any $T > 0$

$$\begin{aligned} & \mathbb{E} \left[\int_0^T \int_{\mathbb{R}^N} e^{-2(t-s)} \sigma^2(Y_n(s, x)) \varphi^2(x - y) ds dy \right] \\ & \leq 2C_\sigma^2 \|\varphi\|_{L^2(\mathbb{R}^N)}^2 \int_0^T (1 + \mathbb{E}[|Y_n(s, x)|^2]) ds \\ & \leq 2C_\sigma^2 \|\varphi\|_{L^2(\mathbb{R}^N)}^2 T \left[1 + \sup_{t \in [0, T], x \in \mathbb{R}^N} \mathbb{E}[|Y_n(t, x)|^2] \right] < \infty. \end{aligned}$$

This shows that the integrand in the stochastic integral is in the space \mathfrak{B}_W (for all $T > 0$), which in turn implies that the stochastic integral in the sense of Walsh is indeed well-defined (by Theorem 3.1).

Now define $D_n(t, x) := Y_{n+1}(t, x) - Y_n(t, x)$ for $n \in \mathbb{N}_0$, $t \geq 0$ and $x \in \mathbb{R}^N$. Then exactly as in the uniqueness calculation we have

$$\begin{aligned} \mathbb{E}[|D_n(t, x)|^2] & \leq 2tC_G^2 C_w \int_0^t e^{-2(t-s)} \int_{\mathbb{R}^N} |w(x, y)| \mathbb{E}[|D_{n-1}(s, y)|^2] dy ds \\ & \quad + 2C_\sigma^2 \|\varphi\|_{L^2(\mathbb{R}^N)}^2 \int_0^t \mathbb{E}[|D_{n-1}(s, x)|^2] e^{-2(t-s)} ds. \end{aligned}$$

This implies that by setting $H_n(s) = \sup_{x \in \mathbb{R}^N} \mathbb{E}[|D_n(s, x)|^2]$,

$$H_n(t) \leq K^n \left[tC_w^2 + \|\varphi\|_{L^2(\mathbb{R}^N)}^2 \right]^n \int_0^t \int_0^{t_1} \dots \int_0^{t_{n-1}} H_0(t_n) dt_n \dots dt_1, \quad (3.14)$$

for all $n \in \mathbb{N}_0$ and $t \geq 0$. Now, similarly, we can find a constant C_t such that

$$\mathbb{E}[|D_0(s, x)|^2] \leq C_t \left(1 + \sup_{x \in \mathbb{R}^N} \mathbb{E}[|Y_0(x)|^2] \right),$$

for any $x \in \mathbb{R}^N$ and $s \in [0, t]$, so that for $s \in [0, t]$,

$$H_0(s) = \sup_{x \in \mathbb{R}^N} \mathbb{E}[|D_0(s, x)|^2] \leq C_t \left(1 + \sup_{x \in \mathbb{R}^N} \mathbb{E}[|Y_0(x)|^2] \right),$$

Using this in (3.14) we see that,

$$H_n(t) \leq C_t K^n \left[tC_w^2 + \|\varphi\|_{L^2(\mathbb{R}^N)}^2 \right]^n \left(1 + \sup_{x \in \mathbb{R}^N} \mathbb{E}[|Y_0(x)|^2] \right) \frac{t^n}{n!},$$

for all $t \geq 0$. This is sufficient to see that (3.13) holds uniformly in n . By completeness, for each $t \geq 0$ and $x \in \mathbb{R}^N$ there exists $Y(t, x) \in L^2(\Omega, \mathcal{F}, \mathbb{P})$ such that $Y(t, x)$ is the limit in $L^2(\Omega, \mathcal{F}, \mathbb{P})$ of the sequence of square-integrable random variables $(Y_n(t, x))_{n \geq 1}$. Moreover, the convergence is uniform on $[0, T] \times \mathbb{R}^N$, i.e.

$$\sup_{t \in [0, T], x \in \mathbb{R}^N} \mathbb{E} |Y_n(t, x) - Y(t, x)|^2 \rightarrow 0.$$

From this we can see that (3.11) is satisfied for the random field $(Y(t, x))_{t \geq 0, x \in \mathbb{R}^N}$. It remains to show that $(Y(t, x))_{t \geq 0, x \in \mathbb{R}^N}$ satisfies (3.10) almost surely. By the above uniform convergence, we have that

$$\mathbb{E} \left[\left| \int_0^t \int_{\mathbb{R}^N} e^{-(t-s)} [\sigma(Y_n(s, x)) - \sigma(Y(s, x))] \varphi(x - y) W(dsdy) \right|^2 \right] \rightarrow 0,$$

and

$$\mathbb{E} \left[\left| \int_0^t e^{-(t-s)} \int_{\mathbb{R}^N} w(x, y) [G(Y_n(s, y)) - G(Y(s, y))] dsdy \right|^2 \right] \rightarrow 0,$$

uniformly for all $t \geq 0$ and $x \in \mathbb{R}^N$. Thus taking the limit as $n \rightarrow \infty$ in (3.12) (in the $L^2(\Omega, \mathcal{F}, \mathbb{P})$ sense) proves that $(Y(t, x))_{t \geq 0, x \in \mathbb{R}^N}$ does indeed satisfy (3.10) almost surely. \square

In a very similar way, one can also prove that the solution remains L^p -bounded whenever the initial condition is L^p -bounded for any $p > 2$. Moreover, this also allows us to conclude that the solution has time continuous paths for all $x \in \mathbb{R}^N$.

Theorem 3.8. *Suppose that we are in the situation of Theorem 3.7, but in addition we have that $\sup_{x \in \mathbb{R}^N} \mathbb{E} [|Y_0(x)|^p] < \infty$ for some $p > 2$. Then the solution $(Y(t, x))_{t \geq 0, x \in \mathbb{R}^N}$ to (3.9) in the sense of Definition 3.6 is L^p -bounded on $[0, T] \times \mathbb{R}^N$ for any T i.e.*

$$\sup_{t \in [0, T], x \in \mathbb{R}^N} \mathbb{E} [|Y(t, x)|^p] < \infty,$$

and the map $t \mapsto Y(t, x)$ has a continuous version for all $x \in \mathbb{R}^N$. If the initial condition has finite p -moments for all $p > 2$, then $t \mapsto Y(t, x)$ has an η -Hölder continuous version, for any $\eta \in (0, 1/2)$ and any $x \in \mathbb{R}^N$.

Proof. The proof of the first part of this result uses similar techniques as in the proof of Theorem 3.7 in order to bound $\mathbb{E} [|Y(t, x)|^p]$ uniformly in $t \in [0, T]$ and $x \in \mathbb{R}^N$. In particular, we use the form of $Y(t, x)$ given by (3.10), Burkholder's inequality (see Theorem 3.2), Hölder's inequality and Gronwall's lemma, as well as the conditions imposed on w , σ , G and φ .

For the time continuity, we again use similar techniques to achieve the bound

$$\mathbb{E} [|Y(t, x) - Y(s, x)|^p] \leq C_T^{(p)} \left(1 + \sup_{r \in [0, T], y \in \mathbb{R}^N} \mathbb{E} [|Y(r, y)|^p] \right) (t - s)^{\frac{p}{2}},$$

for all $s, t \in [0, T]$ with $s \leq t$ and $x \in \mathbb{R}^N$, for some constant $C_T^{(p)}$. The results then follow from Kolmogorov's continuity theorem once again. \square

Spatial regularity of solution

As mentioned in the introduction to this section, the spatial regularity of the solution $(Y(t, x))_{t \geq 0, x \in \mathbb{R}^N}$ to (3.9) is of interest. In particular we would like to find conditions under which it is at least continuous in space. As we saw in Lemma 3.4, under the weak condition on φ given by (3.8), we have that the spatially smoothed space-time white noise is continuous in space. We here show that under this assumption together with a Hölder continuity type condition on the neural field kernel w , the solution $(Y(t, x))_{t \geq 0, x \in \mathbb{R}^N}$ inherits the spatial regularity of the the driving noise.

It is worth mentioning that the neural field equation fits into the class of degenerate diffusion SPDEs (indeed there is no diffusion term), and that regularity theory for such equations is an area that is currently very active (see for example [23] and references therein). However, in our case we are not concerned with any kind of sharp regularity results (in contrast to those found in [16] for the stochastic wave equation), and simply want to assert that for most typical choices of neural field kernels w made by practitioners, the random field solution to the neural field equation is at least regular in space. The results of the section are simple applications of standard techniques to prove continuity in space of random field solutions to SPDEs, as is done for example in [42, Corollary 3.4].

The condition we introduce on w is the following:

$$\exists K_w \geq 0 \text{ s.t. } \|w(x, \cdot) - w(\tilde{x}, \cdot)\|_{L^1(\mathbb{R}^N)} \leq L_w |x - \tilde{x}|^\alpha, \quad \forall x, \tilde{x} \in \mathbb{R}^N, \quad (\mathbf{C3}')$$

for some $\alpha \in (0, 1]$.

Remark 3.9. *This condition is certainly satisfied for all typical choices of neural field kernel w . In particular, any smooth rapidly decaying function will satisfy (C3').*

Theorem 3.10 (Regularity). *Suppose that we are in the situation of Theorem 3.7 and*

$$\sup_{x \in \mathbb{R}^N} \mathbb{E}[|Y_0(x)|^p] < \infty$$

for all $p \geq 2$. Suppose moreover that there exists $\alpha \in (0, 1]$ such that

- w satisfies (C3');
- φ satisfies (3.8) i.e.

$$\|\varphi - \tau_z(\varphi)\|_{L^2(\mathbb{R}^N)} \leq C_\varphi |z|^\alpha, \quad \forall z \in \mathbb{R}^N,$$

where τ_z indicates the shift by $z \in \mathbb{R}^N$ operator;

- $x \mapsto Y_0(x)$ is α -Hölder continuous.

Then $(Y(t, x))_{t \geq 0, x \in \mathbb{R}^N}$ has a modification such that $(t, x) \mapsto Y(t, x)$ is (η_1, η_2) -Hölder continuous, for any $\eta_1 \in (0, 1/2)$ and $\eta_2 \in (0, \alpha)$.

Proof. Let $(Y(t, x))_{t \geq 0, x \in \mathbb{R}^N}$ be the mild solution to (3.9), which exists and is unique by Theorem 3.7. The stated regularity in time is given in Theorem 3.8. It thus remains to prove the regularity in space.

Let $t \geq 0$, $x \in \mathbb{R}^N$. Then by (3.10)

$$Y(t, x) = e^{-t}Y_0(x) + I_1(t, x) + I_2(t, x), \quad (3.15)$$

for all $t \geq 0$ and $x \in \mathbb{R}^N$, where $I_1(t, x) = \int_0^t e^{-(t-s)} \int_{\mathbb{R}^N} w(x, y)G(Y(s, y))dyds$ and $I_2(t, x) = \int_0^t \int_{\mathbb{R}^N} e^{-(t-s)} \sigma(Y(s, x))\varphi(x - y)W(dsdy)$.

Now let $p \geq 2$. The aim is to estimate $\mathbb{E}[|Y(t, x) - Y(t, \tilde{x})|^p]$ for $x, \tilde{x} \in \mathbb{R}^N$ and then to use Kolmogorov's theorem to get the stated spatial regularity. To this end, we have that

$$\begin{aligned} & \mathbb{E}[|I_1(t, x) - I_1(t, \tilde{x})|^p] \\ & \leq \mathbb{E}\left[\left(\int_0^t \int_{\mathbb{R}^N} |w(x, y) - w(\tilde{x}, y)| |G(Y(s, y))| dyds\right)^p\right] \\ & \leq C_G^p t^p \|w(x, \cdot) - w(\tilde{x}, \cdot)\|_{L^1(\mathbb{R}^N)}^p \\ & \leq C_G^p t^p K_w^p |x - \tilde{x}|^{p\alpha}, \end{aligned} \quad (3.16)$$

where we have used (C3'). Moreover, by Hölder's and Burkholder's inequalities once again, we see that

$$\begin{aligned} & \mathbb{E}[|I_2(t, x) - I_2(t, \tilde{x})|^p] \\ & \leq 2^{p-1} \mathbb{E}\left[\left|\int_0^t \int_{\mathbb{R}^N} e^{-(t-s)} [\sigma(Y(s, x)) - \sigma(Y(s, \tilde{x}))] \varphi(x - y)W(dyds)\right|^p\right] \\ & \quad + 2^{p-1} \mathbb{E}\left[\left|\int_0^t \int_{\mathbb{R}^N} e^{-(t-s)} \sigma(Y(s, \tilde{x})) [\varphi(x - y) - \varphi(\tilde{x} - y)]W(dyds)\right|^p\right] \\ & \leq 2^{p-1} c_p \mathbb{E}\left[\left(\int_0^t \int_{\mathbb{R}^N} |\sigma(Y(s, x)) - \sigma(Y(s, \tilde{x}))|^2 |\varphi(x - y)|^2 dyds\right)^{\frac{p}{2}}\right] \\ & \quad + 2^{p-1} c_p \mathbb{E}\left[\left(\int_0^t \int_{\mathbb{R}^N} |\sigma(Y(s, \tilde{x}))|^2 |\varphi(x - y) - \varphi(\tilde{x} - y)|^2 dyds\right)^{\frac{p}{2}}\right], \end{aligned}$$

for all $x, \tilde{x} \in \mathbb{R}^N$ and $p \geq 2$. Thus

$$\begin{aligned} & \mathbb{E}[|I_2(t, x) - I_2(t, \tilde{x})|^p] \\ & \leq 2^{p-1} c_p C_\sigma^p t^{\frac{p}{2}-1} \|\varphi\|_{L^2(\mathbb{R}^N)}^p \int_0^t \mathbb{E}[|Y(s, x) - Y(s, \tilde{x})|^p] ds \\ & \quad + 2^{2(p-1)} c_p C_\sigma^p t^{\frac{p}{2}} \|\varphi - \tau_{\tilde{x}-x}(\varphi)\|_{L^2(\mathbb{R}^N)}^p \left(1 + \sup_{s \in [0, T], y \in \mathbb{R}^N} \mathbb{E}[|Y(s, y)|^p]\right), \end{aligned} \quad (3.17)$$

where we note that the right-hand side is finite thanks to Theorem 3.8. Returning to (3.15) and using estimates (3.16) and (3.17) we see that there exists a constant $C_T^{(p)}$ (depending on $T, p, C_G, K_w, C_\sigma, C_\varphi, \|\varphi\|_{L^2(\mathbb{R}^N)}$ as well as $\sup_{s \in [0, T], y \in \mathbb{R}^N} \mathbb{E}[|Y(s, y)|^p]$), such that

$$\begin{aligned} & \mathbb{E}[|Y(t, x) - Y(t, \tilde{x})|^p] \\ & \leq C_T^{(p)} \left[\mathbb{E}[|Y_0(x) - Y_0(\tilde{x})|^p] + |x - \tilde{x}|^{p\alpha} + \int_0^t \mathbb{E}[|Y(s, x) - Y(s, \tilde{x})|^p] ds \right] \\ & \leq C_T^{(p)} \left[|x - \tilde{x}|^{p\alpha} + \int_0^t \mathbb{E}[|Y(s, x) - Y(s, \tilde{x})|^p] ds \right], \end{aligned}$$

where the last line follows from our assumptions on Y_0 and by adjusting the constant $C_T^{(p)}$. This bound holds for all $t \geq 0, x, \tilde{x} \in \mathbb{R}^N$ and $p \geq 2$. The proof is then completed using Gronwall's inequality, and Kolmogorov's continuity theorem once again. \square

4 Comparison of the two approaches

The purpose of this section is to compare the two different approaches taken in Sections 2 and 3 above to give sense to the stochastic neural field equation. Such a comparison of the two approaches in a general setting has existed for a long time in the probability literature (see for example [25, 26], or more recently [15]), but we provide a proof of the main result (Theorem 4.1) in the Appendix for completeness.

Our starting point is the random field solution, given by Theorem 3.7. Suppose that the conditions of Theorem 3.7 are satisfied (i.e. $\varphi \in L^2(\mathbb{R}^N)$, $\sigma : \mathbb{R} \rightarrow \mathbb{R}$ Lipschitz, $G : \mathbb{R} \rightarrow \mathbb{R}$ Lipschitz and bounded, w satisfies **(C2')** and the given assumptions on the initial condition). Then, by that result, there exists a unique random field $(Y(t, x))_{t \geq 0, x \in \mathbb{R}^N}$ such that

$$\begin{aligned} Y(t, x) &= e^{-t} Y_0(x) + \int_0^t e^{-(t-s)} \int_{\mathbb{R}^N} w(x, y) G(Y(s, y)) dy ds \\ &\quad + \int_0^t \int_{\mathbb{R}^N} e^{-(t-s)} \sigma(Y(s, x)) \varphi(x - y) W(ds dy) \end{aligned} \quad (4.1)$$

where

$$\sup_{t \in [0, T], x \in \mathbb{R}^N} \mathbb{E}[|Y(t, x)|^2] < \infty, \quad (4.2)$$

for all $T > 0$, and we say that $(Y(t, x))_{t \geq 0, x \in \mathbb{R}^N}$ is the random field solution to the stochastic neural field equation.

It turns out that this random field solution is equivalent to the Hilbert space valued solution constructed in Section 2, in the following sense.

Theorem 4.1. *Suppose the conditions of Theorem 3.7 and Theorem 3.8 are satisfied. Moreover suppose that condition (C1') is satisfied for some $\rho_w \in L^1(\mathbb{R}^N)$. Then the random field $(Y(t, x))_{t \geq 0}$ satisfying (4.1) and (4.2) is such that $(Y(t))_{t \geq 0} := (Y(t, \cdot))_{t \geq 0}$ is the unique $L^2(\mathbb{R}^N, \rho_w)$ -valued solution to the stochastic evolution equation*

$$dY(t) = [-Y(t) + \mathbf{F}(Y(t))]dt + \mathbf{B}(Y(t))dW(t), \quad t \in [0, T], \quad (4.3)$$

constructed in Theorem 2.2, where $\mathbf{B} : H \rightarrow L_0(U, H)$ (with $U = L^2(\mathbb{R}^N)$ and $H = L^2(\mathbb{R}^N, \rho_w)$) is given by (2.14) i.e.

$$\mathbf{B}(h)(u)(x) := \sigma(h(x)) \int_{\mathbb{R}^N} \varphi(x - y)u(y)dy, \quad h \in H, u \in U.$$

Example 4.2. *We finish this section with an example illustrating the above result, and the applicability of the two approaches. Indeed, we make the same choices for the neural field kernel w and noise term as in [8], by taking*

$$w(x, y) = \frac{1}{2\beta} e^{-\frac{|x-y|}{\beta}}, \quad x, y \in \mathbb{R}^N, \quad \sigma(a) = \lambda a, \quad a \in \mathbb{R},$$

where β and λ are constants. As noted in Section 2.6, β determines the range of the local synaptic connections. Then, first of all, it is clear that condition (C2') is satisfied (indeed $\|w(x - \cdot)\|_{L^1(\mathbb{R}^N)}$ is constant) and σ is Lipschitz and of linear growth, so that (assuming the initial condition has finite moments), Theorem 3.7 and Theorem 3.8 can be applied to yield a unique random field solution $(Y(t, x))_{t \geq 0}$ to the stochastic neural field equation. Moreover, by Example 2 in Section 2.8, we also see that (C1') is satisfied. Thus Theorem 2.2 can also be applied to construct a Hilbert space valued solution to the stochastic neural field equation (equation (4.3)). By Theorem 4.1, the solutions are equivalent.

5 Conclusion

We have here explored two rigorous frameworks in which stochastic neural field equations can be studied in a mathematically precise fashion. Both these frameworks are useful in the mathematical neuroscience literature: the approach of using the theory of Hilbert space valued processes is adopted in [28], while the random field framework is more natural for Bressloff, Ermentrout and their associates in [8, 10, 27].

It turns out that the constructions are equivalent (see Section 4), when all the conditions are satisfied (which we emphasize is certainly the case for all usual modeling choices of the neural field kernel w and noise terms made in the literature – see Sections 2.6, 2.7 and Example 4.2). However, there are still some advantages and disadvantages for taking one approach over the other, depending on the purpose. For example, an advantage of the construction of a solution as a stochastic process taking values in a Hilbert space carried out in Section 2, is that it allows one

to consider more general diffusion coefficients. Moreover, it is easy to apply results from a large body of literature taking this approach (for example LDP results – see Remark 2.3). A disadvantage is that we have to be careful to impose conditions which control the behavior of the solution in space at infinity and guarantee the integrability of the solution. In particular we require that the connectivity function w either satisfies the strong conditions (C1) and (C2), or the weaker but harder to check conditions (C1') and (C2').

On the other hand, the advantage of the random field approach developed in Section 3 is that one no longer needs to control what happens at infinity. We therefore require fewer conditions on the connectivity function w to ensure the existence of a solution ((C2') is sufficient – see Theorem 3.7). Moreover, with this approach, it is easier to write down conditions that guarantee the existence of a solution that is continuous in both space and time (as opposed to the Hilbert space approach, where spatial regularity is somewhat hidden). However, in order to avoid non-physical distribution valued solutions, we had to impose *a priori* some extra spatial regularity on the noise (see Section 3.2).

Appendix

of Theorem 4.1. The proof of the result involves some technical definition chasing, and in fact is contained in [15], though rather implicitly, but see also [25, 26]. It is for this reason that we carry out the proof explicitly in our situation, by closely following [15, Proposition 4.10]. The most important point is to relate the stochastic integrals that appear in the two different formulations of a solution. To this end, define

$$\mathcal{I}(t, x) := \int_0^t \int_{\mathbb{R}^N} e^{-(t-s)} \sigma(Y(s, x)) \varphi(x - y) W(ds dy), \quad x \in \mathbb{R}^N, t \geq 0,$$

to be the Walsh integral that appears in the random field solution (4.1). Our aim is to show that

$$\mathcal{I}(t, \cdot) = \int_0^t e^{-(t-s)} \mathbf{B}(Y(s)) dW(s), \quad (5.1)$$

where the integral on the right-hand side is the H -valued stochastic integral which appears in the solution to (4.3).

Step 1: Adapting Proposition 2.6 of [15] very slightly, we have that the Walsh integral $\mathcal{I}(t, x)$ can be written as the integral with respect to the *cylindrical* Wiener process $\mathcal{W} = \{\mathcal{W}_t(u) : t \geq 0, u \in U\}$ with covariance \mathbf{Id}_U .⁸ Precisely, we have

$$\mathcal{I}(t, x) = \int_0^t g_s^{t,x} d\mathcal{W}_s,$$

⁸This is a family of random variables such that for each $u \in U$, $(\mathcal{W}_t(u))_{t \geq 0}$ is a Brownian motion with variance $t \|u\|_U^2$, and for all $s, t \geq 0$, $u_1, u_2 \in U$, $\mathbb{E}[\mathcal{W}_t(u_1) \mathcal{W}_s(u_2)] = (s \wedge t) \langle u_1, u_2 \rangle_U$. See for example [15] Section 2.1

for all $t \geq 0, x \in \mathbb{R}^N$, where $g_s^{t,x}(y) := e^{-(t-s)}\sigma(Y(s,x))\varphi(x-y)$, $y \in \mathbb{R}^N$, which is in $L^2(\Omega \times [0, T]; U)$ for any $T > 0$ thanks to (4.2). By definition, the integral with respect to the cylindrical Wiener process \mathcal{W} is given by

$$\int_0^t g_s^{t,x} d\mathcal{W}_s = \sum_{k=1}^{\infty} \int_0^t \langle g_s^{t,x}, e_k \rangle_U d\beta_k(s),$$

where $\{e_k\}_{k=1}^{\infty}$ is a complete orthonormal basis for U , and $(\beta_k(t))_{t \geq 0} := (\mathcal{W}_t(e_k))_{t \geq 0}$ are independent real-valued Brownian motions. This series is convergent in $L^2(\Omega)$.

Step 2: Fix arbitrary $T > 0$. As in Section 3.5 of [15], we can consider the process $\{W(t), t \in [0, T]\}$ defined by

$$W(t) = \sum_{k=1}^{\infty} \beta_k(t) J(e_k) \quad (5.2)$$

where $J : U \rightarrow U$ is a Hilbert-Schmidt operator. $W(t)$ takes its values in U , where it is a $Q(= JJ^*)$ -Wiener process with $\text{Tr}(Q) < \infty$ (Proposition 3.6 of [15]). We define $J(u) := \sum_k \sqrt{\lambda_k} \langle u, e_k \rangle_U e_k$ for a sequence of positive real numbers $(\lambda_k)_{k \geq 1}$ such that $\sum_k \lambda_k < \infty$.

Now define

$$\Phi_s^{t,x}(u) = \langle g_s^{t,x}, u \rangle_U,$$

which takes values in \mathbb{R} . Proposition 3.10 of [15] tells us that the process $\{\Phi_s^{t,x}, s \in [0, T]\}$ defines a predictable process with values in $L_2(U, \mathbb{R})$ and

$$\int_0^t \Phi_s^{t,x} dW(s) = \int_0^t g_s^{t,x} d\mathcal{W}_s, \quad (5.3)$$

where the integral on the left is defined according to Section (2.2), with values in \mathbb{R} .

Step 3: We now note that the original Walsh integral $\mathcal{I}(\cdot, \cdot) \in L^2(\Omega \times [0, T]; H)$. Indeed, because of Burkholder's inequality with $p = 2$,

$$\begin{aligned} \|\mathcal{I}\|_{L^2(\Omega \times [0, T]; H)}^2 &= \mathbb{E} \left[\int_0^T \|\mathcal{I}(t, \cdot)\|_H^2 dt \right] = \int_0^T \int_{\mathbb{R}^N} \mathbb{E} [|\mathcal{I}(t, x)|^2] \rho_w(x) dx dt \\ &\leq \|\varphi\|_{L^2(\mathbb{R}^N)}^2 \int_0^T \int_0^t \int_{\mathbb{R}^N} e^{-2(t-s)} \mathbb{E} [\sigma^2(Y(s, x))] ds \rho_w(x) dx dt, \end{aligned}$$

which is finite, again thanks to (4.2). Hence $\mathcal{I}(t, \cdot)$ takes values in H , and we can therefore write

$$\mathcal{I}(t, \cdot) = \sum_{j=1}^{\infty} \langle \mathcal{I}(t, \cdot), f_j \rangle_H f_j = \sum_{j=1}^{\infty} \left\langle \int_0^t \Phi_s^{t, \cdot} dW(s), f_j \right\rangle_H f_j,$$

by (5.3), where $\{f_j\}_{j=1}^\infty$ is a complete orthonormal basis in H . Moreover, by using (5.2)

$$\begin{aligned}\mathcal{I}(t, \cdot) &= \sum_{j=1}^\infty \left(\int_{\mathbb{R}^N} \left(\int_0^t \Phi_s^{t,x} dW(s) \right) f_j(x) \rho_w(x) dx \right) f_j \\ &= \sum_{j=1}^\infty \left(\int_{\mathbb{R}^N} \left(\sum_{k=1}^\infty \int_0^t \Phi_s^{t,x} (\sqrt{\lambda_k} e_k) d\beta_k(s) \right) f_j(x) \rho_w(x) dx \right) f_j.\end{aligned}\quad (5.4)$$

Finally, consider the H -valued stochastic integral

$$\int_0^t e^{-(t-s)} \mathbf{B}(Y(s)) dW(s),$$

where $\mathbf{B} : H \rightarrow L_0(U, H)$ is given above. Then similarly

$$\begin{aligned}\int_0^t e^{-(t-s)} \mathbf{B}(Y(s)) dW(s) &= \sum_{j=1}^\infty \left\langle \int_0^t e^{-(t-s)} \mathbf{B}(Y(s)) dW(s), f_j \right\rangle_H f_j \\ &= \sum_{j=1}^\infty \left\langle \sum_{k=1}^\infty \int_0^t e^{-(t-s)} \sqrt{\lambda_k} \mathbf{B}(Y(s))(e_k) d\beta_k(s), f_j \right\rangle_H f_j \\ &= \sum_{j=1}^\infty \left(\int_{\mathbb{R}^N} \left(\sum_{k=1}^\infty \int_0^t e^{-(t-s)} \sqrt{\lambda_k} \mathbf{B}(Y(s))(e_k)(x) d\beta_k(s) \right) f_j(x) \rho_w(x) dx \right) f_j.\end{aligned}$$

Here, by definition, for $x \in \mathbb{R}^N$, $0 \leq s \leq t$,

$$\begin{aligned}e^{-(t-s)} \sqrt{\lambda_k} \mathbf{B}(Y(s))(e_k)(x) &= \int_{\mathbb{R}^N} e^{-(t-s)} \sigma(Y(s, x)) \varphi(x - y) \sqrt{\lambda_k} e_k(y) dy \\ &= e^{-(t-s)} \sigma(Y(s, x)) \langle \varphi(x - \cdot), \sqrt{\lambda_k} e_k \rangle_U = \Phi_s^{t,x}(\sqrt{\lambda_k} e_k),\end{aligned}$$

which proves (5.1) by comparison with (5.4).

Step 4: To conclude it suffices to note that the pathwise integrals in (4.1) and the H -valued solution to (4.3) coincide as elements of H . Indeed, it is clear that, by definition of \mathbf{F} ,

$$\int_0^t e^{-(t-s)} \int_{\mathbb{R}^N} w(\cdot, y) G(Y(s, y)) dy ds = \int_0^t e^{-(t-s)} \mathbf{F}(Y(s)) ds,$$

where the later in an element of H . □

Acknowledgements

The authors are grateful to James Maclaurin for suggesting the use of the Fourier transform in Example 2 on page 18, to Etienne Tanré for discussions, and to the referees for their useful suggestions and references.

References

- [1] Amari, S.I.: Dynamics of pattern formation in lateral-inhibition type neural fields. *Biological Cybernetics* **27**(2), 77–87 (1977)
- [2] Baker, T., Cowan, J.: Spontaneous pattern formation and pinning in the primary visual cortex. *Journal of Physiology-Paris* **103**(1-2), 52–68 (2009)
- [3] Bressloff, P.: Spatially periodic modulation of cortical patterns by long-range horizontal connections. *Physica D: Nonlinear Phenomena* **185**(3-4), 131–157 (2003)
- [4] Bressloff, P.: Stochastic neural field theory and the system-size expansion. *SIAM J. Appl. Math* **70**, 1488–1521 (2009)
- [5] Bressloff, P.: Metastable states and quasicycles in a stochastic wilson-cowan model of neuronal population dynamics. *Physical Review E* **82**(5), 051,903 (2010)
- [6] Bressloff, P.: Spatiotemporal dynamics of continuum neural fields. *Journal of Physics A: Mathematical and Theoretical* **45** (2012)
- [7] Bressloff, P., Cowan, J., Golubitsky, M., Thomas, P., Wiener, M.: Geometric visual hallucinations, Euclidean symmetry and the functional architecture of striate cortex. *Phil. Trans. R. Soc. Lond. B* **306**(1407), 299–330 (2001)
- [8] Bressloff, P., Webber, M.: Front propagation in stochastic neural fields. *SIAM Journal on Applied Dynamical Systems* (2012)
- [9] Bressloff, P.C., Folias, S.E.: Front bifurcations in an excitatory neural network. *SIAM Journal on Applied Mathematics* **65**(1), 131–151 (2004)
- [10] Bressloff, P.C., Wilkerson, J.: Traveling pulses in a stochastic neural field model of direction selectivity. *Frontiers in computational neuroscience* **6** (2012)
- [11] Brezis, H.: *Functional analysis, Sobolev spaces and Partial Differential Equations*. Springer (2010)
- [12] Brzeźniak, Z., Peszat, S.: Space-time continuous solutions to SPDE’s driven by a homogeneous Wiener process. *Studia Math.* **137**(3), 261–299 (1999)
- [13] Dalang, R., Khoshnevisan, D., Mueller, C., Nualart, D., Xiao, Y.: *A Minicourse on Stochastic Partial Differential Equations, Lecture Notes in Mathematics*, vol. 1962. Springer-Verlag, Berlin (2009). Held at the University of Utah, Salt Lake City, UT, May 8–19, 2006, Edited by Khoshnevisan and Firas Rassoul-Agha
- [14] Dalang, R.C., Frangos, N.E.: The stochastic wave equation in two spatial dimensions. *Ann. Probab.* **26**(1), 187–212 (1998)

- [15] Dalang, R.C., Quer-Sardanyons, L.: Stochastic integrals for spde's: a comparison. *Expo. Math.* **29**(1), 67–109 (2011)
- [16] Dalang, R.C., Sanz-Solé, M.: Hölder-Sobolev regularity of the solution to the stochastic wave equation in dimension three. *Mem. Amer. Math. Soc.* **199**(931), vi+70 (2009)
- [17] Du, Y.: Order Structure and Topological Methods in Nonlinear Partial Differential Equations. Vol. 1, *Series in Partial Differential Equations and Applications*, vol. 2. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ (2006)
- [18] Ermentrout, G., McLeod, J.: Existence and uniqueness of travelling waves for a neural network. In: Royal Society(Edinburgh), *Proceedings*, vol. 123, pp. 461–478 (1993)
- [19] Eveson, S.P.: Compactness criteria for integral operators in L^∞ and L^1 spaces. *Proc. Amer. Math. Soc.* **123**(12), 3709–3716 (1995)
- [20] Faye, G., Chossat, P., Faugeras, O.: Analysis of a hyperbolic geometric model for visual texture perception. *The Journal of Mathematical Neuroscience* **1**(4) (2011)
- [21] Ferrante, M., Sanz-Solé, M.: SPDEs with coloured noise: analytic and stochastic approaches. *ESAIM Probab. Stat.* **10**, 380–405 (electronic) (2006)
- [22] Folias, S.E., Bressloff, P.C.: Breathing pulses in an excitatory neural network. *SIAM Journal on Applied Dynamical Systems* **3**(3), 378–407 (2004)
- [23] Hofmanová, M.: Degenerate parabolic stochastic partial differential equations. *Stochastic Process. Appl.* **123**(12), 4294–4336 (2013)
- [24] Jansen, B.H., Rit, V.G.: Electroencephalogram and visual evoked potential generation in a mathematical model of coupled cortical columns. *Biological Cybernetics* **73**, 357–366 (1995)
- [25] Jetschke, G.: Different approaches to stochastic parabolic differential equations. *Proceedings of the 10th Winter School on Abstract Analysis* pp. 161–169 (1982)
- [26] Jetschke, G.: On the equivalence of different approaches to stochastic partial differential equations. *Mathematische Nachrichten* **128**(1), 315–329 (1986)
- [27] Kilpatrick, Z.P., Ermentrout, B.: Wandering bumps in stochastic neural fields. *SIAM Journal on Applied Dynamical Systems* **12**(1), 61–94 (2013)
- [28] Kuehn, C., Riedler, M.G.: Large deviations for nonlocal stochastic neural fields. *Journal of Mathematical Neuroscience* **4**(1) (2014)

- [29] Lund, J.S., Angelucci, A., Bressloff, P.C.: Anatomical substrates for functional columns in macaque monkey primary visual cortex. *Cerebral Cortex* **12**, 15–24 (2003)
- [30] Mariño, J., Schummers, J., Lyon, D., Schwabe, L., Beck, O., Wiesing, P., Obermayer, K., Sur, M.: Invariant computations in local cortical networks with balanced excitation and inhibition. *Nature neuroscience* **8**(2), 194–201 (2005)
- [31] Owen, M., Laing, C., Coombes, S.: Bumps and rings in a two-dimensional neural field: splitting and rotational instabilities. *New Journal of Physics* **9**(10), 378–401 (2007)
- [32] Pardoux, E.: *Stochastic Partial Differential Equations. Lectures given in Fudan University, Shanghai* (2007)
- [33] Peszat, S.: Large deviation principle for stochastic evolution equations. *Probab. Theory Related Fields* **98**(1), 113–136 (1994)
- [34] Potthast, R., Beim Graben, P.: Existence and properties of solutions for neural field equations. *Mathematical Methods in the Applied Sciences* **33**(8), 935–949 (2010)
- [35] Prato, G.D., Zabczyk, J.: *Stochastic Equations in Infinite Dimensions*. Cambridge University Press (1992)
- [36] Prévôt, C., Röckner, M.: *A Concise Course on Stochastic Partial Differential Equations, Lecture Notes in Mathematics*, vol. 1905. Springer, Berlin (2007)
- [37] Sanz-Solé, M., Sarrà, M.: Hölder continuity for the stochastic heat equation with spatially correlated noise. In: *Seminar on Stochastic Analysis, Random Fields and Applications, III (Ascona, 1999)*, *Progr. Probab.*, vol. 52, pp. 259–268. Birkhäuser, Basel (2002)
- [38] Lopes da Silva, F., Hoeks, A., Zetterberg, L.: Model of brain rhythmic activity. *Kybernetik* **15**, 27–37 (1974)
- [39] Lopes da Silva, F., van Rotterdam, A., Barts, P., van Heusden, E., Burr, W.: Model of neuronal populations. the basic mechanism of rhythmicity. M.A. Corner, D.F. Swaab (eds) *Progress in brain research*, Elsevier, Amsterdam **45**, 281–308 (1976)
- [40] Simon, J.: Sobolev, Besov and Nikol’ skiĭ fractional spaces: imbeddings and comparisons for vector valued spaces on an interval. *Ann. Mat. Pura Appl.* (4) **157**, 117–148 (1990)

- [41] Veltz, R., Faugeras, O.: Local/global analysis of the stationary solutions of some neural field equations. *SIAM Journal on Applied Dynamical Systems* **9**(3), 954–998 (2010)
- [42] Walsh, J.B.: An introduction to stochastic partial differential equations. In: École d’été de probabilités de Saint-Flour, XIV—1984, *Lecture Notes in Math.*, vol. 1180, pp. 265–439. Springer, Berlin (1986)
- [43] Wilson, H., Cowan, J.: Excitatory and inhibitory interactions in localized populations of model neurons. *Biophys. J.* **12**, 1–24 (1972)
- [44] Wilson, H., Cowan, J.: A mathematical theory of the functional dynamics of cortical and thalamic nervous tissue. *Biological Cybernetics* **13**(2), 55–80 (1973)