

The probability of non-confluent systems

Alejandro Díaz-Caro, Gilles Dowek

► **To cite this version:**

Alejandro Díaz-Caro, Gilles Dowek. The probability of non-confluent systems. Mauricio Ayala Rincón and Eduardo Bonelli and Ian Mackie. DCM - 9th International Workshop on Developments in Computational Models - 2013, Aug 2013, Buenos Aires, Argentina. 144, pp.1-15, 2014, Electronic Proceedings in Theoretical Computer Science. <<http://dcm-workshop.org.uk/2013/>>. <10.4204/EPTCS.144.1>. <hal-00919546>

HAL Id: hal-00919546

<https://hal.inria.fr/hal-00919546>

Submitted on 17 Dec 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The probability of non-confluent systems

Alejandro Díaz-Caro

Université Paris Ouest
200 avenue de la République
92001 Nanterre, France
alejandro@diaz-caro.info

INRIA
23 avenue d'Italie, CS 81321
75214 Paris Cedex 13, France

Gilles Dowek

INRIA
23 avenue d'Italie, CS 81321
75214 Paris Cedex 13, France
gilles.dowek@inria.fr

We show how to provide a structure of probability space to the set of execution traces on a non-confluent abstract rewrite system, by defining a variant of a Lebesgue measure on the space of traces. Then, we show how to use this probability space to transform a non-deterministic calculus into a probabilistic one. We use as example λ_+ , a recently introduced calculus defined through type isomorphisms.

1 Introduction

Many probabilistic calculi has been developed in the pasts years, e.g. [1, 9, 11, 14, 20]. In particular, the algebraic versions of λ -calculus [5, 24] are extensions to λ -calculus where a linear combination of terms, e.g. $\alpha.r + \beta.s$, is also a term. One way to interpret such a linear combination is that it represents a term which is the term r with probability α , or the term s with probability β . However, endowing such a calculus with a non-restrictive type system is a challenge [3, 4].


A simpler framework is that of non deterministic calculi which can be seen as algebraic calculi without scalars. They have been studied, for instance in [8, 10, 12, 13, 15–18, 21], however moving back from non-determinism to probabilities is not trivial. In this paper we propose, instead of changing these models, to define a probability measure on reductions in non-deterministic systems. In fact, as we shall see, such a probability measure can be defined on any abstract non-deterministic transition systems, or non-confluent abstract rewrite systems (ARS) (cf. [23, Chapter 1]). Our goal is to show that explicit probabilities are not needed in the syntax, and that the simpler non-deterministic calculi are as powerful as the more complicated probabilistic calculi.

Consider for example the following non-confluent ARS

$$\mathbf{a} \rightarrow \mathbf{b} , \quad \mathbf{a} \rightarrow \mathbf{c} , \quad \mathbf{c} \rightarrow \mathbf{d} , \quad \mathbf{c} \rightarrow \mathbf{e} ,$$

we want to associate a probability to events such as

$$\mathbf{a} \rightarrow^* \mathbf{b} , \quad \mathbf{a} \rightarrow^* \mathbf{c} , \quad \mathbf{a} \rightarrow^* \mathbf{d} , \quad \mathbf{a} \rightarrow^* \mathbf{e} .$$



In this example, assuming equiprobability, we have $P(\mathbf{a} \rightarrow^* \mathbf{b}) = \frac{1}{2}$, $P(\mathbf{a} \rightarrow^* \mathbf{c}) = \frac{1}{2}$, $P(\mathbf{a} \rightarrow^* \mathbf{d}) = \frac{1}{4}$, $P(\mathbf{a} \rightarrow^* \mathbf{e}) = \frac{1}{4}$. Notice that these events are not disjoint and that their sum is larger than 1. In particular, $\mathbf{a} \rightarrow^* \mathbf{d}$ implies $\mathbf{a} \rightarrow^* \mathbf{c}$. Defining the elements of the set Ω of elementary events is not completely straightforward, in particular because we want to make it general enough to also consider infinite cases. For example, in the following system

$$\mathbf{a}_i \rightarrow \mathbf{a}_{i+1}, \quad \mathbf{a}_i \rightarrow \mathbf{a}'_{i+1} ,$$

we naturally would like that $P(\mathbf{a}_0 \rightarrow^* \mathbf{a}_n) = \frac{1}{2^n}$.

Besides defining the elements of the set Ω , we need to define a notion of a measurable subset of Ω and endow such a subset with a probability distribution verifying the Kolmogorov axioms.

Our idea is to follow Lebesgue: define first the probability of rectangles, or boxes, then the probability of any set and finally measurable sets as those that verify Lebesgue's property. Thus besides defining the set Ω , we need to define a subset of $\mathcal{P}(\Omega)$ of boxes.

The first intuition would be to take paths as elements of the set Ω , for instance assigning the probability $\frac{1}{2}$ to the paths $\mathbf{a} \rightarrow \mathbf{b}$, $\frac{1}{4}$ to $\mathbf{a} \rightarrow \mathbf{c} \rightarrow \mathbf{d}$ and $\frac{1}{4}$ to $\mathbf{a} \rightarrow \mathbf{c} \rightarrow \mathbf{e}$. In fact it seems more convenient to extend such paths to strategies prescribing one reduct for each non normal object. Boxes are then defined as sets of strategies agreeing on a finite domain. We show in this paper that this is sufficient to define a probability space on strategies, consistent with the intuitive probability of events of the form $\mathbf{a} \rightarrow^* \mathbf{b}$.

Our study is generic enough to be applicable to several settings, such as automata, or any other kind of transition systems. We use the nomenclature of abstract rewrite systems, but that of states and transitions could be used as well. Finally, we apply this construction to λ_+ [15, 16].

Plan of the paper. Section 2 introduces the basic concepts of strategies and boxes, it defines the Lebesgue measures. Section 3 proves that the space of strategies forms a probability space. Finally, in section 4 we show how to modify the calculus λ_+ into a probabilistic calculus λ_+^p . Also, we provide an encoding of an algebraic λ -calculus into λ_+^p and, to some extent, the inverse translation.

2 Preliminaries

Let Λ be a set of objects and \rightarrow a function from $\Lambda \times \Lambda$ to \mathbb{N} such that for all \mathbf{a} the set $\{\mathbf{b} \mid \rightarrow(\mathbf{a}, \mathbf{b}) \neq 0\}$ is finite. We write $\mathbf{a} \rightarrow \mathbf{b}$ if $\rightarrow(\mathbf{a}, \mathbf{b}) \neq 0$. We allow a term to be written to the same symbol more than once, so its probability increases, e.g. if $\rightarrow(\mathbf{a}, \mathbf{b}) = 2$ and $\rightarrow(\mathbf{a}, \mathbf{c}) = 1$, then the probability of getting \mathbf{b} will be the double than the probability of getting \mathbf{c} . Think for example in a non-deterministic choice between two objects, which happen to be equal, then there would be two ways to get such an object by doing the choice. For a given object $\mathbf{a} \in \Lambda$, we denote by $\rho(\mathbf{a})$ its degree, that is, the number of objects to which it can be rewritten to in one step. Definition 2.1 formalises this.

Definition 2.1 (Degree of an object). $\rho : \Lambda \rightarrow \mathbb{N}$ is a function defined by $\rho(\mathbf{a}) = \sum_{\mathbf{b}} \rightarrow(\mathbf{a}, \mathbf{b})$.

An object is normal if its degree is 0. We denote by $\Lambda^+ = \{\mathbf{a} \mid \mathbf{a} \in \Lambda \text{ and } \rho(\mathbf{a}) \geq 1\}$ to the set of non-normal objects, that is, objects that can be rewritten to other objects.

A strategy prescribing one reduct for each non-normal object is defined as a function from Λ^+ to Λ (cf. [23, Def. 4.9.1]).

Definition 2.2 (Strategy). A strategy is a total function $f : \Lambda^+ \rightarrow \Lambda$ such that $f(\mathbf{a}) = \mathbf{b}$ implies $\mathbf{a} \rightarrow \mathbf{b}$. For instance, if $\mathbf{a} \rightarrow \mathbf{b}$ and $\mathbf{a} \rightarrow \mathbf{b}'$, there are two functions, f and f' assigning different results to \mathbf{a} . We denote by Ω the set of all such functions.

A box is a set of strategies agreeing on a finite domain.

Definition 2.3 (Box). A box $B \subseteq \Omega$ is a set of the form $\{f \mid f(\mathbf{a}_1) = \mathbf{a}'_1, \dots, f(\mathbf{a}_n) = \mathbf{a}'_n\}$ for some objects $\mathbf{a}_i, \mathbf{a}'_i$. We write $\mathcal{B}(\Omega)$ the subset of $\mathcal{P}(\Omega)$ containing all the boxes.

Example 2.4. Continuing with the example given at the introduction, $\Lambda^+ = \{\mathbf{a}, \mathbf{c}\}$. Let $f_1(\mathbf{a}) = \mathbf{b}, f_1(\mathbf{c}) = \mathbf{d}$ and $f_2(\mathbf{a}) = \mathbf{b}, f_2(\mathbf{c}) = \mathbf{e}$ be two of the four strategies of Ω . Then the box $\{f \mid f(\mathbf{a}) = \mathbf{b}, f(\mathbf{c}) = \mathbf{d}\} =$

$\{f_1\}$, and the box $\{f \mid f(\mathbf{a}) = \mathbf{b}\}$ is $\{f_1, f_2\}$.

$$\left\{ \begin{array}{l} f_1 = \begin{array}{c} \mathbf{a} \\ \swarrow \quad \searrow \\ \mathbf{b} \quad \mathbf{c} \\ \quad \searrow \\ \mathbf{d} \end{array} \quad ; \quad f_2 = \begin{array}{c} \mathbf{a} \\ \swarrow \quad \searrow \\ \mathbf{b} \quad \mathbf{c} \\ \quad \searrow \\ \mathbf{e} \end{array} \end{array} \right\} = \text{Box} \begin{array}{c} \mathbf{a} \\ \swarrow \quad \searrow \\ \mathbf{b} \quad \mathbf{c} \end{array}$$

A probability distribution can be defined in term of boxes, and then be extended to arbitrary sets of strategies.

Definition 2.5 (Probability function). Let $p : \mathcal{B}(\Omega) \rightarrow [0, 1]$ be a total function defined over boxes as follows. If $B = \{f \mid f(\mathbf{a}_1) = \mathbf{a}'_1, \dots, f(\mathbf{a}_n) = \mathbf{a}'_n\}$, then

$$p(B) = \prod_{i=1}^n \frac{\rho(\mathbf{a}_i, \mathbf{a}'_i)}{\rho(\mathbf{a}_i)} .$$


By convention, if no condition is given in B (i.e. $B = \Omega$), we have $n = 0$, and we consider the product of zero elements to be 1, the neutral element of the product.

Then we define the probability measure $P : \mathcal{P}(\Omega) \rightarrow [0, 1]$ for arbitrary sets of strategies as follows

$$P(S) = \begin{cases} 0 & \text{if } S = \emptyset \\ \inf \{ \sum_{B \in \mathcal{C}} p(B) \mid \mathcal{C} \text{ is a countable family of boxes s.t. } S \subseteq \bigcup_{B \in \mathcal{C}} B \} & \text{in other case} \end{cases}$$

Example 2.6. Consider the ARS $\mathbf{a} \rightarrow \mathbf{b}$ with multiplicity 2 and $\mathbf{a} \rightarrow \mathbf{c}$ with multiplicity 1.

Let B be the box $B = \{f \mid f(\mathbf{a}) = \mathbf{b}\}$. Then we have $p(B) = \frac{\rho(\mathbf{a}, \mathbf{b})}{\rho(\mathbf{a})} = \frac{2}{3}$. Intuitively, $P(B)$ is the same as $p(B)$ (this will be later formalised in Lemma 3.10), because B is the minimum cover of B , that is, $\{B\}$ is the minimum family of boxes such that B is in its union. Hence $P(B) = \frac{2}{3}$.



Example 2.7. We continue with the same running example depicted in the introduction. Let $f_1(\mathbf{a}) = \mathbf{b}$, $f_1(\mathbf{c}) = \mathbf{d}$ and $f_3(\mathbf{a}) = \mathbf{c}$, $f_3(\mathbf{c}) = \mathbf{e}$ be two strategies. Then the set $S = \{f_1, f_3\}$ is minimally covered by the boxes $B_1 = \{f_1\} = \{f \mid f(\mathbf{a}) = \mathbf{b}, f(\mathbf{c}) = \mathbf{d}\}$ and $B_2 = \{f_3\} = \{f \mid f(\mathbf{a}) = \mathbf{c}, f(\mathbf{c}) = \mathbf{e}\}$. So we have $P(S) = p(B_1) + p(B_2) = \frac{1}{2 \times 2} + \frac{1}{2 \times 2} = \frac{1}{2}$.

$$S = \left\{ \begin{array}{l} f_1 = \begin{array}{c} \mathbf{a} \\ \swarrow \quad \searrow \\ \mathbf{b} \quad \mathbf{c} \\ \quad \searrow \\ \mathbf{d} \end{array} \quad ; \quad f_3 = \begin{array}{c} \mathbf{a} \\ \searrow \\ \mathbf{c} \\ \searrow \\ \mathbf{e} \end{array} \end{array} \right\}$$

Now we can define the Lebesgue measure in terms of the given probability measure.

Definition 2.8 (Measurable). Let A be an element of $\mathcal{P}(\Omega)$, we write A^\sim for the complement of A , that is $\Omega \setminus A$. The set A is Lebesgue measurable if $\forall S \in \mathcal{P}(\Omega)$, we have $P(S) = P(S \cap A) + P(S \cap A^\sim)$.

We define $\mathcal{A} = \{A \mid A \text{ is measurable}\}$.

3 A probability space of strategies

The aim of this section is to prove that (Ω, \mathcal{A}, P) is a probability space. That is, the sample space Ω (the set of all possible strategies), the set of events \mathcal{A} , which is the set of the Lebesgue measurable sets of

strategies, and the probability measure P , form a probability space. Our proof follows [7]. We proceed by proving that this triplet satisfies the Kolmogorov axioms, that is the probability of any event is between 0 and 1, the probability of Ω is 1, and the probability of any countable sequence of pairwise disjoint (that is incompatible) events, is the sum of their probabilities. In order to do so, we need first to prove several properties.

Lemma 3.1 establishes several known properties of Lebesgue measurable sets.

Lemma 3.1.

1. Let $A \in \mathcal{A}$ and $S \in \mathcal{P}(\Omega)$. If $A \cap S = \emptyset$, then $P(A \cup S) = P(A) + P(S)$.
2. Let $A_1, A_2 \in \mathcal{A}$. If $A_1 \subseteq A_2$, then $P(A_1) \leq P(A_2)$.
3. \emptyset , the empty set, is Lebesgue measurable.
4. A is Lebesgue measurable if and only if A^\sim is Lebesgue measurable.
5. If A_1, A_2 are Lebesgue measurable, then $A_1 \cup A_2$ is Lebesgue measurable.

Proof.

1. $P(A \cup S) = P((A \cup S) \cap A) + P((A \cup S) \cap A^\sim) = P(A) + P(S)$.
2. First notice that by definition, $P(S) \geq 0$ for any $S \in \mathcal{P}(\Omega)$. Hence, $P(A_2) = P(A_2 \cap A_1) + P(A_2 \cap A_1^\sim) = P(A_1) + P(A_2 \cap A_1^\sim) \geq P(A_1)$.
3. Notice that $P(\emptyset) = 0$. Then, $\forall S \in \mathcal{P}(\Omega)$, $P(S) = P(S \cap \emptyset) + P(S \cap \Omega)$, so \emptyset is Lebesgue measurable.
4. Let A be Lebesgue measurable, then $\forall S \subseteq \Omega$, $P(S) = P(S \cap A) + P(S \cap A^\sim) = P(S \cap A^\sim) + P(S \cap A^{\sim\sim})$, so A^\sim is Lebesgue measurable.
5. Let A_1, A_2 be Lebesgue measurable, then $\forall S \subseteq \Omega$, we have

$$P(S) = P(S \cap A_1) + P(S \cap A_1^\sim) \quad (1) \quad \text{and} \quad P(S) = P(S \cap A_2) + P(S \cap A_2^\sim) \quad (2)$$

$$\text{From set theory} \quad S \cap (A_1 \cup A_2) = S \cap (A_1 \cup (A_1^\sim \cap A_2)) = (S \cap A_1) \cup (S \cap A_1^\sim \cap A_2) \quad (3)$$

Using $S \cap A_1^\sim$ for S in (2) gives

$$P(S \cap A_1^\sim) = P(S \cap A_1^\sim \cap A_2) + P(S \cap A_1^\sim \cap A_2^\sim) = P(S \cap A_1^\sim \cap A_2) + P(S \cap (A_1 \cup A_2)^\sim) \quad (4)$$

From (3), using items 1 and 2, we have $P(S \cap (A_1 \cup A_2)) = P(S \cap A_1) + P(S \cap A_1^\sim \cap A_2)$. Adding $P(S \cap (A_1 \cup A_2)^\sim)$ to both sides gives $P(S \cap (A_1 \cup A_2)) + P(S \cap (A_1 \cup A_2)^\sim) = P(S \cap A_1) + P(S \cap A_1^\sim \cap A_2) + P(S \cap (A_1 \cup A_2)^\sim)$. Using (4) and (1) we obtain $P(S \cap (A_1 \cup A_2)) + P(S \cap (A_1 \cup A_2)^\sim) = P(S \cap A_1) + P(S \cap A_1^\sim) = P(S)$. \square

The concept of algebra (Definition 3.2) gives a closure property of subsets. As a corollary of the Lemma 3.1 we can show that the set \mathcal{A} of Lebesgue measurable sets form an algebra (Corollary 3.3).

Definition 3.2 (Algebra). Let X be a set. We say that a set $\mathbb{A} \in \mathcal{P}(X)$ is an algebra over X if for all $A, B \in \mathbb{A}$, $A \cup B$, A^\sim and X itself are also in \mathbb{A} .

Corollary 3.3. \mathcal{A} is an algebra over Ω .

Proof. $\mathcal{A} \in \mathcal{P}(\Omega)$. Let $A, B \in \mathcal{A}$, then by Lemma 3.1(5), $A \cup B \in \mathcal{A}$. By Lemma 3.1(4), $A^\sim \in \mathcal{A}$. Finally, by Lemma 3.1(3) and (4), $\Omega \in \mathcal{A}$. \square

Moreover, we can show that \mathcal{A} is a σ -algebra, that is an algebra, completed to include countably infinite operations. Definition 3.4 formalises it.

Definition 3.4 (σ -algebra). Let X be a set. We say that a set $\Sigma \in \mathcal{P}(X)$ is a σ -algebra over X if it is an algebra and it is closed under countable unions, that is, if A_1, A_2, A_3, \dots are in Σ , then so is $\bigcup A_i$.

Theorem 3.7 states that the set \mathcal{A} of Lebesgue measurable sets is a σ -algebra. We need to prove two properties of Lebesgue measurable sets first (Lemmas 3.5 and 3.6).

Lemma 3.5. *Let $S \subseteq \Omega$ and $A_1, \dots, A_n \in \mathcal{A}$ be a disjoint family. Then*

$$\mathbb{P} \left(S \cap \left(\bigcup_{i=1}^n A_i \right) \right) = \sum_{i=1}^n \mathbb{P}(S \cap A_i) .$$

Proof. We proceed by induction on n . If $n = 1$ it is trivial. Assume it is true for $n - 1$. Notice that

$$S \cap \left(\bigcup_{i=1}^n A_i \right) \cap A_n = S \cap A_n \quad (5) \quad \text{and} \quad S \cap \left(\bigcup_{i=1}^n A_i \right) \cap A_n^c = S \cap \left(\bigcup_{i=1}^{n-1} A_i \right) \quad (6)$$

Equation (5) is clear, and (6) follows since $(\bigcup_{i=1}^n A_i) \cap A_n^c = \bigcup_{i=1}^{n-1} (A_i \cap A_n^c) = (\bigcup_{i=1}^{n-1} (A_i \cap A_n^c)) \cup (A_n \cap A_n^c)$.

Thus, since A_n is measurable, we have that

$$\mathbb{P} \left(S \cap \left(\bigcup_{i=1}^n A_i \right) \right) = \mathbb{P} \left(S \cap \left(\bigcup_{i=1}^n A_i \right) \cap A_n \right) + \mathbb{P} \left(S \cap \left(\bigcup_{i=1}^n A_i \right) \cap A_n^c \right)$$

and from (5) and (6) this is equal to $\mathbb{P}(S \cap A_n) + \mathbb{P}(S \cap (\bigcup_{i=1}^{n-1} A_i))$, which by the induction hypothesis is equal to $\sum_{i=1}^n \mathbb{P}(S \cap A_i)$. \square

Lemma 3.6. *Let $S_1, S_2, \dots \subseteq \Omega$. Then*

$$\mathbb{P} \left(\bigcup_{i=1}^{\infty} S_i \right) \leq \sum_{i=1}^{\infty} \mathbb{P}(S_i) .$$

Proof. If $\mathbb{P}(S_i) = \infty$ for some i , then we are finished. Therefore, assume $\mathbb{P}(S_i) < \infty$ for each $i \in \mathbb{N}$.

Without loss of generality, assume $S_i \neq \emptyset$, for all i . Indeed, since $\mathbb{P}(\emptyset) = 0$, an empty set would not add anything to any side of the equation. For a given $\varepsilon > 0$ and i , there is a sequence $\{B_{ij} \mid i = 1, \dots, j = 1, \dots\}$ of boxes such that $S_i \subseteq \bigcup_{j=1}^{\infty} B_{ij}$ and $\sum_{j=1}^{\infty} \mathbb{P}(B_{ij}) < \mathbb{P}(S_i) + 2^{-i}\varepsilon$, by the definition of \mathbb{P} . Now, $\#\{B_{ij} \mid i, j\} \leq \aleph_0$ and $\bigcup_{i=1}^{\infty} S_i \subseteq \bigcup_{i=1}^{\infty} \bigcup_{j=1}^{\infty} B_{ij}$. Therefore, using the definition of \mathbb{P} ,

$$\mathbb{P} \left(\bigcup_{i=1}^{\infty} S_i \right) \leq \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \mathbb{P}(B_{ij}) \leq \sum_{i=1}^{\infty} \mathbb{P}(S_i) + \varepsilon \sum_{i=1}^{\infty} \frac{1}{2^i} = \sum_{i=1}^{\infty} \mathbb{P}(S_i) + \varepsilon$$

Since this is true for each ε , the lemma holds. \square

Using these properties, we can prove that \mathcal{A} is a σ -algebra (Theorem 3.7).

Theorem 3.7. *\mathcal{A} is a σ -algebra over Ω .*

Proof. By Corollary 3.3, \mathcal{A} is an algebra. We only have to prove that \mathcal{A} is closed under any countable unions. That is, if $B_1, B_2, \dots \in \mathcal{A}$, then $\bigcup_{i=1}^{\infty} B_i \in \mathcal{A}$. Since \mathcal{A} is an algebra (Corollary 3.3), there is a disjoint family $A_1, A_2, \dots \in \mathcal{A}$ such that $A = \bigcup_{i=1}^{\infty} B_i = \bigcup_{i=1}^{\infty} A_i$. For example, we can take $A_1 = B_1$, $A_2 = B_2 \setminus B_1$, $A_3 = B_3 \setminus (B_1 \cup B_2)$, \dots . Let $C_n = \bigcup_{i=1}^n A_i$, so $C_n \in \mathcal{A}$ again using that \mathcal{A} is an algebra. Also notice that $A^{\sim} \subseteq C_n^{\sim}$ because $C_n \subseteq A$.

Since C_n is measurable, take any $S \subseteq \Omega$ and, using Lemma 3.1(2), we can calculate $P(S) = P(S \cap C_n) + P(S \cap C_n^{\sim}) \geq P(S \cap C_n) + P(S \cap A^{\sim})$. Since $P(S \cap C_n) = P(S \cap (\bigcup_{i=1}^n A_i))$, using Lemma 3.5, we obtain $P(S) \geq \sum_{i=1}^n P(S \cap A_i) + P(S \cap A^{\sim})$ and, since the left-hand side is independent of n , $P(S) \geq \sum_{i=1}^{\infty} P(S \cap A_i) + P(S \cap A^{\sim})$. Thus, by Lemma 3.6, $P(S) \geq P(S \cap (\bigcup_{i=1}^{\infty} A_i)) + P(S \cap A^{\sim}) = P(S \cap A) + P(S \cap A^{\sim})$.

For the converse inequality, notice that $S = (S \cap A) \cup (S \cap A^{\sim})$, so using Lemma 3.6 we have $P(S) = P((S \cap A) \cup (S \cap A^{\sim})) \leq P(S \cap A) + P(S \cap A^{\sim})$. Hence, $A \in \mathcal{A}$. \square

As intuited in Example 2.6, the probability of a box B is $p(B)$. Lemma 3.10 formalises it. Before proving this lemma, we need two auxiliary ones (Lemmas 3.8 and 3.9). For short, we use the notation $B \cap \mathbf{a} = \mathbf{b}$ for $B \cap \{f \mid f(\mathbf{a}) = \mathbf{b}\}$.

Lemma 3.8. *Let $N \subseteq \mathbb{N}$ and for all $i \in N$, let $B, B_i \subseteq \Omega$ be boxes s.t. $B \subseteq \bigcup_{i \in N} B_i$ and $p(B) > \sum_{i \in N} p(B_i)$. Then for every object \mathbf{a} , there exists an object \mathbf{b} such that, $p(B \cap \mathbf{a} = \mathbf{b}) > \sum_{i \in N} p(B_i \cap \mathbf{a} = \mathbf{b})$.*

Proof. Let $\mathbf{a} \rightarrow \mathbf{b}_i$, with $i = 1, \dots, n$. Hence notice that $p(B) = \sum_{j=1}^n p(B \cap \mathbf{a} = \mathbf{b}_j)$, and this happens for any B . Then, from $p(B) > \sum_{i \in N} p(B_i)$, we have $\sum_{j=1}^n p(B \cap \mathbf{a} = \mathbf{b}_j) > \sum_{i \in N} \sum_{j=1}^n p(B_i \cap \mathbf{a} = \mathbf{b}_j) = \sum_{j=1}^n \sum_{i \in N} p(B_i \cap \mathbf{a} = \mathbf{b}_j)$. Therefore, there must be at least one h such that $p(B \cap \mathbf{a} = \mathbf{b}_h) > \sum_{i \in N} p(B_i \cap \mathbf{a} = \mathbf{b}_h)$. \square

Lemma 3.9. *Let $N \subseteq \mathbb{N}$ and for all $i \in N$, let $B, B_i \subseteq \Omega$ be boxes s.t. $B \subseteq \bigcup_{i \in N} B_i$ and $p(B) > \sum_{i \in N} p(B_i)$. Then for all family $\{\mathbf{a}_j\}$ of objects, there exists a family $\{\mathbf{b}_j\}$ such that, for every k , $p(B \cap \mathbf{a}_1 = \mathbf{b}_1 \cap \dots \cap \mathbf{a}_k = \mathbf{b}_k) > \sum_{i \in N} p(B_i \cap \mathbf{a}_1 = \mathbf{b}_1 \cap \dots \cap \mathbf{a}_k = \mathbf{b}_k)$.*

Proof. We proceed by induction on k . For $k = 1$, use Lemma 3.8. By the induction hypothesis, we have $p(B \cap \mathbf{a}_1 = \mathbf{b}_1 \cap \dots \cap \mathbf{a}_{k-1} = \mathbf{b}_{k-1}) > \sum_{i \in N} p(B_i \cap \mathbf{a}_1 = \mathbf{b}_1 \cap \dots \cap \mathbf{a}_{k-1} = \mathbf{b}_{k-1})$. We conclude by Lemma 3.8. \square

Lemma 3.10. *Let $B \subseteq \Omega$ be a box, then $P(B) = p(B)$.*

Proof. Let $B = \{f \mid f(\mathbf{a}_1) = \mathbf{a}'_1, \dots, f(\mathbf{a}_n) = \mathbf{a}'_n\}$. Since $B \subseteq B$, by definition of P , we have $P(B) \leq p(B)$. We must prove $p(B) \leq P(B) = \inf\{\sum_{i \in N} p(B_i) \mid B \subseteq \bigcup_{i \in N} B_i\}$. In other words, we must prove that $B \subseteq \bigcup_{i \in N} B_i$ implies $p(B) \leq \sum_{i \in N} p(B_i)$. We proceed by induction on n .

- If $n = 0$, $p(B) = 1$. Notice that, without restrictions in B , $B = \Omega$. We prove this case by contradiction. Let $p(F) > \sum_{i \in N} p(B_i)$. Then by Lemma 3.9, there exists g such that for all k ,

$$p(\mathbf{a}_1 = g(\mathbf{a}_1) \cap \dots \cap \mathbf{a}_k = g(\mathbf{a}_k)) > \sum_{i \in N} p(B_i \cap \mathbf{a}_1 = g(\mathbf{a}_1) \cap \dots \cap \mathbf{a}_k = g(\mathbf{a}_k)) \quad (7)$$

Since $g \in \Omega \subseteq \bigcup_{i \in N} B_i$, there exists j such that $g \in B_j$. Let B_j be defined with constraints on objects $\mathbf{a}_1, \dots, \mathbf{a}_q$. Let $k = q$ and from equation (7),

$$p(\mathbf{a}_1 = g(\mathbf{a}_1) \cap \dots \cap \mathbf{a}_q = g(\mathbf{a}_q)) > \sum_{i \in N} p(B_i \cap \mathbf{a}_1 = g(\mathbf{a}_1) \cap \dots \cap \mathbf{a}_q = g(\mathbf{a}_q)) \quad (8)$$

We know that $\mathbf{p}(\mathbf{a}_1 = g(\mathbf{a}_1) \cap \dots \cap \mathbf{a}_q = g(\mathbf{a}_q)) = \prod_{h=1}^q \frac{\rightarrow(\mathbf{a}_h, g(\mathbf{a}_h))}{\rho(\mathbf{a}_h)}$, and since $g \in B_j$, we know that this is also equal to $\mathbf{p}(B_j \cap \mathbf{a}_1 = g(\mathbf{a}_1) \cap \dots \cap \mathbf{a}_q = g(\mathbf{a}_q))$. Hence equation (8) leads to a contradiction.

- Consider the case $n - 1$. Let $B' = \{f \mid \exists g \in B \text{ s.t. } \forall \mathbf{a} \neq \mathbf{a}_n, f(\mathbf{a}) = g(\mathbf{a})\}$. Then if $B' \subseteq \bigcup_{i \in N'} B'_i$ we have $\mathbf{p}(B') \leq \sum_{i \in N'} \mathbf{p}(B'_i)$. Notice that either $B'_i = B_i$ or B_i has a constraint on \mathbf{a}_n and so $\frac{\rightarrow(\mathbf{a}_n, g(\mathbf{a}_n))}{\rho(\mathbf{a}_n)} \mathbf{p}(B'_i) = \mathbf{p}(B_i)$. In any case, $\frac{\rightarrow(\mathbf{a}_n, g(\mathbf{a}_n))}{\rho(\mathbf{a}_n)} \mathbf{p}(B'_i) \leq \mathbf{p}(B_i)$. Then $\mathbf{p}(B) = \frac{\rightarrow(\mathbf{a}_n, g(\mathbf{a}_n))}{\rho(\mathbf{a}_n)} \mathbf{p}(B') \leq \sum_{i \in N'} \frac{\rightarrow(\mathbf{a}_n, g(\mathbf{a}_n))}{\rho(\mathbf{a}_n)} \mathbf{p}(B'_i) \leq \sum_{i \in N'} \mathbf{p}(B_i)$. \square

Theorem 3.11 (Space of strategies). $(\Omega, \mathcal{A}, \mathbf{P})$ is a probability space.

Proof. We prove it satisfies the Kolmogorov axioms.

1st axiom: $\forall A \in \mathcal{A}, 0 \leq \mathbf{P}(A) \leq 1$.

Since \mathbf{P} is defined as an inf of sums of p , and p is always positive, so \mathbf{P} cannot be negative. By the second Kolmogorov axiom $\mathbf{P}(\Omega) = 1$. Notice that A is measurable and $A \subseteq \Omega$, so $1 = \mathbf{P}(\Omega) = \mathbf{P}(\Omega \cap A) + \mathbf{P}(\Omega \setminus A) = \mathbf{P}(A) + \mathbf{P}(\Omega \setminus A)$, hence $1 - \mathbf{P}(\Omega \setminus A) = \mathbf{P}(A)$. Since \mathbf{P} is not negative, $\mathbf{P}(A) \leq 1$.

2nd axiom: $\mathbf{P}(\Omega) = 1$.

Notice that Ω is the box including all the functions. Hence, there is no condition on the functions and so $n = 0$. Then $\mathbf{p}(\Omega) = 1$. By Lemma 3.10, $\mathbf{P}(\Omega) = \mathbf{p}(\Omega) = 1$.

3rd axiom: Any countable sequence of pairwise disjoint (i.e. incompatible) events $A_1, A_2, \dots \in \mathcal{A}$, satisfies $\mathbf{P}(A_1 \cup A_2 \dots) = \sum_{i=1}^{\infty} \mathbf{P}(A_i)$.


Let $\emptyset \neq I \subsetneq \mathbb{N}$. Since the sets A_i are in \mathcal{A} , consider $n \in \mathbb{N} \setminus I$ and we have

$$\mathbf{P}\left(\bigcup_{i \in \mathbb{N} \setminus I} A_i\right) = \mathbf{P}\left(\left(\bigcup_{i \in \mathbb{N} \setminus I} A_i\right) \cap A_n\right) + \mathbf{P}\left(\left(\bigcup_{i \in \mathbb{N} \setminus I} A_i\right) \cap A_n^{\sim}\right)$$

Notice that $\left(\bigcup_{i \in \mathbb{N} \setminus I} A_i\right) \cap A_n = A_n$ and since the A_i 's are pairwise disjoint $\left(\bigcup_{i \in \mathbb{N} \setminus I} A_i\right) \cap A_n^{\sim} = \bigcup_{i \in \mathbb{N} \setminus (I \cup \{n\})} A_i$. Therefore, considering that this is valid for any I and $n \notin I$, we have

$$\mathbf{P}\left(\bigcup_{i=1}^{\infty} A_i\right) = \mathbf{P}(A_1) + \mathbf{P}\left(\bigcup_{i=2}^{\infty} A_i\right) = \mathbf{P}(A_1) + \mathbf{P}(A_2) + \mathbf{P}\left(\bigcup_{i=3}^{\infty} A_i\right) = \dots = \sum_{i=1}^{\infty} \mathbf{P}(A_i). \quad \square$$

Example 3.12. Consider the non-strongly-normalising non-confluent rewrite system described in the introduction $\mathbf{a}_i \rightarrow \mathbf{a}_{i+1}$, $\mathbf{a}_i \rightarrow \mathbf{a}'_{i+1}$, where each reduction is equiprobable and each symbol is different from each other. It can be depicted as follows.



The probability that this rewrite system stops after exactly n steps, starting from term \mathbf{a}_0 is $\mathbf{P}(B)$, with $B = \{f \mid f(\mathbf{a}_0) = \mathbf{a}_1, \dots, f(\mathbf{a}_{n-2}) = \mathbf{a}_{n-1} \text{ and } f(\mathbf{a}_{n-1}) = \mathbf{a}'_n\}$, and since B is a box, by Lemma 3.10 it is the same to $\mathbf{P}(B) = \mathbf{p}(B) = \frac{1}{\rho(\mathbf{a}_0) \dots \rho(\mathbf{a}_{n-1})} = \frac{1}{2^n}$.

The probability of stopping at the step n or before, starting at any point before a_{n-1} , is just the probability of the box $\{f \mid f(\mathbf{a}_{n-1}) = \mathbf{a}'_n\}$, which is $\frac{1}{2}$.

The probability of stopping at the step n or m , starting at any point before \mathbf{a}_{n-1} and \mathbf{a}_{m-1} is the probability of the union of two boxes, however they are not independent events (its intersection is not empty). Hence let $B_1 = \{f \mid \mathbf{a}_{n-1} = \mathbf{a}'_n\}$ and $B_2 = \{f \mid \mathbf{a}_{m-1} = \mathbf{a}'_m\}$. The probability $P(B_1 \cup B_2) = P((B_1 \setminus B_2) \cup B_2) = P(B_1 \setminus B_2) \cup P(B_2) = P(\{f \mid \mathbf{a}_{n-1} = \mathbf{a}'_n, \mathbf{a}_{m-1} = \mathbf{a}'_m\}) + P(B_2) = \frac{1}{4} + \frac{1}{2} = \frac{3}{4}$.

Finally, the probability of not stopping at all, is the probability of the set $S = \{f \mid f(\mathbf{a}_i) = \mathbf{a}_{i+1} \text{ for } i \in \mathbb{N}\}$, which is not a box, since there is an infinite number of conditions. It is easy to check that we need an infinite number of boxes to cover such a set, however we can chose boxes as small as we want (that is, with a big number of conditions), which makes the infimum of their sums to be 0, and so the probability of not stopping is, as expected, 0.

In other words, $P(S) \leq \{f \mid f(\mathbf{a}_i) = \mathbf{a}_{i+1}, i \in [0, n]\} = \frac{1}{2^n}$, for any n . Hence when n tends to ∞ , $P(S)$ tends to 0.

4 Transforming a non-deterministic into a probabilistic calculus

4.1 The calculus λ_+

In [15, 16] we have introduced a non-deterministic calculus called λ_+ , which is a simplification of an earlier probabilistic calculus by keeping non-determinism but removing explicit probabilities. Now we can transform this calculus into a probabilistic one.

The full calculus is depicted in Table 1. Typing judgements are of the form $\mathbf{r} : A$. A term \mathbf{r} is typable if there exists a type A such that $\mathbf{r} : A$. Following [19, 22], we use a presentation of typed lambda-calculus without contexts and where each variable occurrence is labelled by its type, such as $\lambda x^A.x^A$ or $\lambda x^A.y^B$. We sometimes omit the labels when they are clear from the context and write, for example, $\lambda x^A.x$ for $\lambda x^A.x^A$. We use different letters for different variables and the type system forbids terms such as $\lambda x^A.x^B$ when A and B are different, by imposing preconditions to when the typing rules apply. Let $S = \{x_1^{A_1}, \dots, x_n^{A_n}\}$ be a set of declarations, we write S^f when this set is functional, that is when $x_i = x_j$ implies $A_i = A_j$. For example $\{x^A, y^{A \Rightarrow B}\}^f$, but not $\{x^A, x^{A \Rightarrow B}\}^f$. Typing rules have the following structure:

$$\frac{\text{[Preconditions]} \quad \text{Hypotheses}}{\text{Derived judgement}} \quad (\text{Rule name})$$

The α -conversion and the sets $FV(\mathbf{r})$ of free variables of \mathbf{r} and $FV(A)$ of free variables of A are defined as usual in the λ -calculus (cf. [6, §2.1]). For example $FV(x^A y^B) = \{x^A, y^B\}$. We say that a term \mathbf{r} is closed whenever $FV(\mathbf{r}) = \emptyset$. If $FV(\mathbf{r}) = \{x_1^{A_1}, \dots, x_n^{A_n}\}$, we write $\Gamma(\mathbf{r}) = \{A_1, \dots, A_n\}$. $FV(\{A_1, \dots, A_n\})$ is defined by $\bigcup_{i=1}^n FV(A_i)$. Given two terms \mathbf{r} and \mathbf{s} we denote by $\mathbf{r}[\mathbf{s}/x]$ the term obtained by simultaneously substituting the term \mathbf{s} for all the free occurrences of x in \mathbf{r} , subject to the usual proviso about renaming bound variables in \mathbf{r} to avoid capture of the free variables of \mathbf{s} . Analogously $A[B/X]$ denotes the substitution of the type B for all the free occurrences of X in A , and $\mathbf{r}[B/X]$ the substitution in \mathbf{r} . For example, $(x^A)[B/Y] = x^{A[B/Y]}$, $(\lambda x^A.\mathbf{r})[B/X] = \lambda x^{A[B/X]}. \mathbf{r}[B/X]$ and $(\pi_A(\mathbf{r}))[B/X] = \pi_{A[B/X]}(\mathbf{r}[B/X])$. Simultaneous substitutions are defined in the same way. Finally, terms and types are considered up to α -conversion.

Each term of the language has a main type associated, which can be obtained from the type annotations, and other types induced by the type equivalences.

The operational semantics of λ_+ is also given in Table 1, where there are two distinct relations between terms: a symmetric relation \rightleftharpoons and a reduction relation \hookrightarrow . We write \rightleftharpoons^* and \hookrightarrow^* for the transitive and reflexive closures of \rightleftharpoons and \hookrightarrow respectively. In particular, notice that \rightleftharpoons^* is an equivalence relation. We just write \rightarrow when we do not want to make the distinction between these relations. We write $n.\mathbf{r}$ in λ_+ as a shorthand for $\underbrace{\mathbf{r} + \dots + \mathbf{r}}_{n \text{ times}}$.

This calculus has a non-deterministic projector. Indeed, the rule “If $\mathbf{r} : A$, then $\pi_A(\mathbf{r} + \mathbf{s}) \hookrightarrow \mathbf{r}$ ” is not-deterministic because the symbol $+$ is commutative, so if $\mathbf{s} : A$, this rule can produce either \mathbf{r} or \mathbf{s} non-deterministically. In any case, both reducts are valid proofs of A , and so the proof system is consistent. Refer to [15] for details.

Grammar of types and terms		
$A, B, C, \dots ::= X \mid A \Rightarrow B \mid A \wedge B \mid \forall X.A \ .$		
$\mathbf{r}, \mathbf{s}, \mathbf{t} ::= x^A \mid \lambda x^A.\mathbf{r} \mid \mathbf{r}\mathbf{s} \mid \mathbf{r} + \mathbf{s} \mid \pi_A(\mathbf{r}) \mid \Lambda X.\mathbf{r} \mid \mathbf{r}\{A\} \ .$		
Equivalence between types		
$A \wedge B \equiv B \wedge A \ , \quad (A \wedge B) \wedge C \equiv A \wedge (B \wedge C) \ , \quad A \Rightarrow (B \wedge C) \equiv (A \Rightarrow B) \wedge (A \Rightarrow C) \ .$		
Rewriting system		
<i>Symmetric relation:</i>		
$\mathbf{r} + \mathbf{s} \rightleftharpoons \mathbf{s} + \mathbf{r} \ ,$	$(\mathbf{r} + \mathbf{s})\mathbf{t} \rightleftharpoons \mathbf{r}\mathbf{t} + \mathbf{s}\mathbf{t} \ ,$	If $\mathbf{r} : A \Rightarrow (B \wedge C)$, then
$(\mathbf{r} + \mathbf{s}) + \mathbf{t} \rightleftharpoons \mathbf{r} + (\mathbf{s} + \mathbf{t}) \ ,$	$\lambda x^A.(\mathbf{r} + \mathbf{s}) \rightleftharpoons \lambda x^A.\mathbf{r} + \lambda x^A.\mathbf{s} \ ,$	$\pi_{A \Rightarrow B}(\mathbf{r})\mathbf{s} \rightleftharpoons \pi_B(\mathbf{r}\mathbf{s}) \ .$
<i>Reductions:</i>		
$(\lambda x^A.\mathbf{r})\mathbf{s} \hookrightarrow \mathbf{r}[\mathbf{s}/x] \ ,$	$(\Lambda X.\mathbf{r})\{A\} \hookrightarrow \mathbf{r}[A/X] \ ,$	If $\mathbf{r} : A$, then $\pi_A(\mathbf{r} + \mathbf{s}) \hookrightarrow \mathbf{r} \ .$
Typing system		
$\frac{[A \equiv B] \mathbf{r} : A}{\mathbf{r} : B} (\equiv)$	$\frac{}{x^A : A} (ax)$	$\frac{[(FV(\mathbf{r}) \cup \{x^A\})^f] \mathbf{r} : B}{\lambda x^A.\mathbf{r} : A \Rightarrow B} (\Rightarrow_i)$
$\frac{[FV(\mathbf{r} + \mathbf{s})^f] \mathbf{r} : A \quad \mathbf{s} : B}{\mathbf{r} + \mathbf{s} : A \wedge B} (\wedge_i)$	$\frac{\mathbf{r} : A \wedge B}{\pi_A(\mathbf{r}) : A} (\wedge_e)$	$\frac{[X \notin FV(\Gamma(\mathbf{r}))] \mathbf{r} : A}{\Lambda X.\mathbf{r} : \forall X.A} (\forall_i)$
		$\frac{\mathbf{r} : \forall X.A}{\mathbf{r}\{B\} : A[B/X]} (\forall_e)$

Table 1: The non-deterministic calculus λ_+

4.2 From non-determinism to probabilities (or from λ_+ to λ_+^p)

Consider the following example (cf. [16, Example 5]). Two possible reduction paths can be fired from $(\Lambda X.(\pi_A(x^A + y^X)))\{A\}$: Reducing first the projection, $(\Lambda X.x^A)\{A\} \hookrightarrow x^A$, or reducing first the beta $\pi_A(x^A + y^A) \hookrightarrow x^A$. The former path is deterministic and will always reduce to x^A , on the contrary, the latter can non-deterministically chose between x^A and y^A . However, in both cases a proof of A is obtained.

Hence, the non-determinism is present not only due to the projector, but also by a combination of not defining a reduction strategy and the polymorphism, which can turn a deterministic projection into a non-deterministic one. We want to associate a probability to the second case, that is, to the non-deterministic projector (the π reduction). With this aim, we consider the following ARS, called λ_+^\downarrow . The closed normal terms of λ_+ are objects of λ_+^\downarrow . If $\mathbf{r}_1, \dots, \mathbf{r}_n$ are objects, then it is also an object. The function \rightarrow is

given by the relations \Leftrightarrow and \hookrightarrow . In particular, if $\mathbf{r} : A$, then $\pi_A(\mathbf{r} + \mathbf{r}) \rightarrow \mathbf{r}$, with multiplicity 2, i.e. $\rightarrow (\pi_A(\mathbf{r} + \mathbf{r}), \mathbf{r}) = 2$.

Theorem 4.1. Let $(\Omega, \mathcal{A}, \mathbb{P})$ be a probability space over λ_+^\downarrow . Let $B_{\mathbf{r}_i} = \{f \mid f(\pi_A(\sum_{j=1}^n m_j \cdot \mathbf{r}_j)) = \mathbf{r}_i\}$ be a box. Then $\mathbb{P}(B_{\mathbf{r}_i}) = \frac{m_i}{\sum_{j=1}^n m_j}$.

Proof. Notice that $\rho(\pi_A(\sum_{i=1}^n m_i \cdot \mathbf{r}_i)) = \sum_{\mathbf{r}} \rightarrow (\pi_A(\sum_{i=1}^n m_i \cdot \mathbf{r}_i), \mathbf{r}) = \#[\overbrace{\mathbf{r}_1, \dots, \mathbf{r}_1}^{m_1 \text{ times}}, \dots, \overbrace{\mathbf{r}_n, \dots, \mathbf{r}_n}^{m_n \text{ times}}] = \sum_{j=1}^n m_j$. And $\rightarrow (\pi_A(\sum_{i=1}^n m_i \cdot \mathbf{r}_i), \mathbf{r}_i) = m_i$. Hence, $\mathbb{P}(B_{\mathbf{r}_i}) = \mathbb{p}(B_{\mathbf{r}_i}) = \frac{m_i}{\sum_{j=1}^n m_j}$. \square

Definition 4.2 (The probabilistic calculus λ_+^p). Let λ_+^p be the language of Table 1, with the following modification:

Replace rule “If $\mathbf{r} : A$, then $\pi_A(\mathbf{r} + \mathbf{s}) \hookrightarrow \mathbf{r}$ ” by

“For $i = 1, \dots, n$, let $\mathbf{r}_i : A$ and $\mathbf{s} \not\vdash A$, be closed normal terms. Then

$$\pi_A\left(\sum_{i=1}^n m_i \cdot \mathbf{r}_i + \mathbf{s}\right) \hookrightarrow \mathbf{r}_i \quad \text{with probability } \frac{m_i}{\sum_{j=1}^n m_j} \text{”} .$$

Remark 4.3. Notice that by Theorem 4.1 the probabilistic reduction is well defined.

4.3 The calculus Alg_F^p

The calculus Alg_F^p is inspired from [5, 24]. We restrict the algebraic calculus to only have probabilistic superpositions, and we type it with a simple extension of System F (cf. [2, Def. 5.1]). The grammar of terms ensures that the linear combinations of terms are probability distributions, however the type system allows typing pseudo-terms, that is, terms that are not probability distributions. A term in this language, is a term produced by the grammar of terms, and typed. The full calculus is depicted in Table 2.

4.4 From Alg_F^p to λ_+^p

We give a translation from the probabilistic calculus Alg_F^p , including scalars, to the probabilistic calculus λ_+^p .

$$\begin{aligned} \llbracket x^A \rrbracket &= x^A & \llbracket \mathbf{r}\mathbf{s} \rrbracket &= \llbracket \mathbf{r} \rrbracket \llbracket \mathbf{s} \rrbracket & \llbracket \mathbf{r}\{A\} \rrbracket &= \llbracket \mathbf{r} \rrbracket \{A\} \\ \llbracket \lambda x^A \cdot \mathbf{r} \rrbracket &= \lambda x^A \cdot \llbracket \mathbf{r} \rrbracket & \llbracket \Lambda X \cdot \mathbf{r} \rrbracket &= \Lambda X \cdot \llbracket \mathbf{r} \rrbracket & \llbracket \sum_{i=1}^n \frac{n_i}{d_i} \cdot \mathbf{r}_i \rrbracket &= \pi_A(\sum_{i=1}^n m_i \cdot \llbracket \mathbf{r}_i \rrbracket) \\ & & & & \text{where } \mathbf{r}_i : A, d_i \in \mathbb{N}^*, m_i = n_i \left(\prod_{\substack{k=1 \\ k \neq i}}^n d_k \right), \text{ for } i = 1, \dots, n. \end{aligned}$$

Example 4.4. Let $\mathbf{r} : A$, $\mathbf{t} : A$ and $\mathbf{s} : A$. $\llbracket \frac{3}{4} \cdot \mathbf{r} + \frac{1}{8} \cdot \mathbf{t} + \frac{1}{8} \cdot \mathbf{s} \rrbracket = \pi_A(192 \cdot \llbracket \mathbf{r} \rrbracket + 32 \cdot \llbracket \mathbf{t} \rrbracket + 32 \cdot \llbracket \mathbf{s} \rrbracket)$. By Theorem 4.1, this last term reduces to $\llbracket \mathbf{r} \rrbracket$ with probability $\frac{192}{192+32+32} = \frac{3}{4}$, to $\llbracket \mathbf{t} \rrbracket$ with probability $\frac{32}{192+32+32} = \frac{1}{8}$, and to $\llbracket \mathbf{s} \rrbracket$ with probability $\frac{32}{192+32+32} = \frac{1}{8}$.

Lemma 4.5.

1. $\llbracket \mathbf{r} \rrbracket [A/X] = \llbracket \mathbf{r}[A/X] \rrbracket$.
2. $\llbracket \mathbf{r} \rrbracket [\llbracket \mathbf{s} \rrbracket / x] = \llbracket \mathbf{r}[\mathbf{s}/x] \rrbracket$.

Proof.

1. We proceed by induction on \mathbf{r} .

- Let $\mathbf{r} = x^B$. $\llbracket x^B \rrbracket [A/X] = x^B [A/X] = x^{B[A/X]} = \llbracket x^{B[A/X]} \rrbracket = \llbracket x^B [A/X] \rrbracket$.

Grammar of types		
$A, B, C, \dots ::= X \mid A \Rightarrow B \mid \forall X.A$.		
Grammar of pseudo-terms		
$\mathbf{r}, \mathbf{s}, \mathbf{t} ::= x^A \mid \lambda x^A.\mathbf{r} \mid \mathbf{rs} \mid \Lambda X.\mathbf{r} \mid \mathbf{r}\{A\} \mid p.\mathbf{r} \mid \mathbf{r} + \mathbf{s}$		
Grammar of terms		
$\mathbf{r}, \mathbf{s}, \mathbf{t} ::= x^A \mid \lambda x^A.\mathbf{r} \mid \mathbf{rs} \mid \Lambda X.\mathbf{r} \mid \mathbf{r}\{A\} \mid \sum_{i=1}^n p_i.\mathbf{r}_i$ with $\begin{cases} n > 0, \\ p_i \in \mathbb{Q}(0, 1] \text{ and} \\ \sum_{i=1}^n p_i = 1 \end{cases}$		
Rewriting system		
<i>Symmetric relation:</i>		
$\mathbf{r} + \mathbf{s} \rightleftharpoons \mathbf{s} + \mathbf{r}$,	$(\mathbf{r} + \mathbf{s})\mathbf{t} \rightleftharpoons \mathbf{rt} + \mathbf{st}$,	$1.\mathbf{r} \rightleftharpoons \mathbf{r}$.
$(\mathbf{r} + \mathbf{s}) + \mathbf{t} \rightleftharpoons \mathbf{r} + (\mathbf{s} + \mathbf{t})$,	$\lambda x^A.(\mathbf{r} + \mathbf{s}) \rightleftharpoons \lambda x^A.\mathbf{r} + \lambda x^A.\mathbf{s}$,	
<i>Reductions:</i>		
Beta	Elementary	Factorisation
$(\lambda x^A.\mathbf{r})\mathbf{s} \hookrightarrow \mathbf{r}[s/x]$,	$p.q.\mathbf{r} \hookrightarrow pq.\mathbf{r}$,	$p.\mathbf{r} + q.\mathbf{r} \hookrightarrow (p+q).\mathbf{r}$.
$(\Lambda X.\mathbf{r})\{A\} \hookrightarrow \mathbf{r}[A/X]$,	$p.(\mathbf{r} + \mathbf{s}) \hookrightarrow p.\mathbf{r} + p.\mathbf{s}$,	
Typing system		
$\frac{}{x^A : A} \text{ (ax)}$	$\frac{\mathbf{r} : B}{\lambda x^A.\mathbf{r} : A \Rightarrow B} \text{ (}\Rightarrow\text{i)}$	$\frac{[FV(\mathbf{rs})]^f \mathbf{r} : A \Rightarrow B \quad \mathbf{s} : A}{\mathbf{rs} : B} \text{ (}\Rightarrow\text{e)}$
$\frac{[FV(\mathbf{r+s})]^f \mathbf{r} : A \quad \mathbf{s} : A}{\mathbf{r} + \mathbf{s} : A} \text{ (+i)}$	$\frac{\mathbf{r} : A}{p.\mathbf{r} : A} \text{ (p}_i\text{)}$	$\frac{[X \notin FV(\Gamma(\mathbf{r}))] \mathbf{r} : A}{\Lambda X.\mathbf{r} : \forall X.A} \text{ (}\forall\text{i)}$
		$\frac{\mathbf{r} : \forall X.A}{\mathbf{r}\{B\} : A[B/X]} \text{ (}\forall\text{e)}$

Table 2: The algebraic calculus Alg_F^p .

- Let $\mathbf{r} = \lambda x^B.\mathbf{t}$. $\llbracket \lambda x^B.\mathbf{t} \rrbracket[A/X] = \lambda x^B.\llbracket \mathbf{t} \rrbracket[A/X] = \lambda x^{B[A/X]}\llbracket \mathbf{t} \rrbracket[A/X] \stackrel{IH}{=} \lambda x^{B[A/X]}\llbracket \mathbf{t} \rrbracket[A/X] = \llbracket \lambda x^{B[A/X]}\mathbf{t} \rrbracket[A/X] = \llbracket (\lambda x^B.\mathbf{t}) \rrbracket[A/X]$.
- Let $\mathbf{r} = \mathbf{t}_1\mathbf{t}_2$. $\llbracket \mathbf{t}_1\mathbf{t}_2 \rrbracket[A/X] = \llbracket \mathbf{t}_1 \rrbracket[A/X]\llbracket \mathbf{t}_2 \rrbracket[A/X] \stackrel{IH}{=} \llbracket \mathbf{t}_1 \rrbracket[A/X]\llbracket \mathbf{t}_2 \rrbracket[A/X] = \llbracket \mathbf{t}_1 \rrbracket[A/X]\mathbf{t}_2[A/X] = \llbracket (\mathbf{t}_1\mathbf{t}_2) \rrbracket[A/X]$.
- Let $\mathbf{r} = \Lambda Y.\mathbf{t}$, with $Y \notin FV(A)$. $\llbracket \Lambda Y.\mathbf{t} \rrbracket[A/X] = \Lambda Y.\llbracket \mathbf{t} \rrbracket[A/X] \stackrel{IH}{=} \Lambda Y.\llbracket \mathbf{t} \rrbracket[A/X] = \llbracket \Lambda Y.\mathbf{t} \rrbracket[A/X] = \llbracket (\Lambda Y.\mathbf{t}) \rrbracket[A/X]$.
- Let $\mathbf{r} = \mathbf{t}\{B\}$. $\llbracket \mathbf{t}\{B\} \rrbracket[A/X] = \llbracket \mathbf{t} \rrbracket\{B\}[A/X] = \llbracket \mathbf{t} \rrbracket[A/X]\{B[A/X]\} \stackrel{IH}{=} \llbracket \mathbf{t} \rrbracket[A/X]\{B[A/X]\} = \llbracket \mathbf{t}\{B\} \rrbracket[A/X]$.
- Let $\mathbf{r} = \sum_{i=1}^n \frac{n_i}{d_i}.\mathbf{r}_i$. $\llbracket \sum_{i=1}^n \frac{n_i}{d_i}.\mathbf{r}_i \rrbracket[A/X] = \pi_A(\sum_{i=1}^n m_i.\llbracket \mathbf{r}_i \rrbracket[A/X]) = \pi_A(\sum_{i=1}^n m_i.\llbracket \mathbf{r}_i \rrbracket[A/X]) \stackrel{IH}{=} \pi_A(\sum_{i=1}^n m_i.\llbracket \mathbf{r}_i \rrbracket[A/X]) = \llbracket \sum_{i=1}^n \frac{n_i}{d_i}.\mathbf{r}_i \rrbracket[A/X] = \llbracket (\sum_{i=1}^n \frac{n_i}{d_i}.\mathbf{r}_i) \rrbracket[A/X]$.

2. We proceed by induction on \mathbf{r} .

- Let $\mathbf{r} = x^A$. $\llbracket x^A \rrbracket[\llbracket \mathbf{s} \rrbracket/x] = x^A[\llbracket \mathbf{s} \rrbracket/x] = \llbracket \mathbf{s} \rrbracket = \llbracket x^A[s/x] \rrbracket$.
- Let $\mathbf{r} = y^A$. $\llbracket y^A \rrbracket[\llbracket \mathbf{s} \rrbracket/x] = y^A[\llbracket \mathbf{s} \rrbracket/x] = y^A = \llbracket y^A \rrbracket = \llbracket y^A[s/x] \rrbracket$.
- Let $\mathbf{r} = \lambda y^B.\mathbf{t}$. $\llbracket \lambda y^B.\mathbf{t} \rrbracket[\llbracket \mathbf{s} \rrbracket/x] = \lambda y^B.\llbracket \mathbf{t} \rrbracket[\llbracket \mathbf{s} \rrbracket/x] \stackrel{IH}{=} \lambda y^B.\llbracket \mathbf{t} \rrbracket[\llbracket \mathbf{s} \rrbracket/x] = \llbracket \lambda y^B.\mathbf{t} \rrbracket[\llbracket \mathbf{s} \rrbracket/x] = \llbracket (\lambda y^B.\mathbf{t})[s/x] \rrbracket$.

- Let $\mathbf{r} = \mathbf{t}_1 \mathbf{t}_2$. $\llbracket \mathbf{t}_1 \mathbf{t}_2 \rrbracket \llbracket \llbracket \mathbf{s} \rrbracket / x \rrbracket = \llbracket \mathbf{t}_1 \rrbracket \llbracket \llbracket \mathbf{s} \rrbracket / x \rrbracket \llbracket \mathbf{t}_2 \rrbracket \llbracket \llbracket \mathbf{s} \rrbracket / x \rrbracket \stackrel{IH}{=} \llbracket \mathbf{t}_1 \llbracket \mathbf{s} / x \rrbracket \rrbracket \llbracket \mathbf{t}_2 \llbracket \mathbf{s} / x \rrbracket \rrbracket = \llbracket \mathbf{t}_1 \llbracket \mathbf{s} / x \rrbracket \mathbf{t}_2 \llbracket \mathbf{s} / x \rrbracket \rrbracket = \llbracket (\mathbf{t}_1 \mathbf{t}_2) \llbracket \mathbf{s} / x \rrbracket \rrbracket$.
- Let $\mathbf{r} = \Lambda X. \mathbf{t}$. $\llbracket \Lambda X. \mathbf{t} \rrbracket \llbracket \llbracket \mathbf{s} \rrbracket / x \rrbracket = \Lambda X. \llbracket \llbracket \mathbf{t} \rrbracket \llbracket \llbracket \mathbf{s} \rrbracket / x \rrbracket \stackrel{IH}{=} \Lambda X. \llbracket \llbracket \mathbf{t} \rrbracket \llbracket \mathbf{s} / x \rrbracket \rrbracket = \llbracket \Lambda X. \mathbf{t} \llbracket \mathbf{s} / x \rrbracket \rrbracket = \llbracket (\Lambda X. \mathbf{t}) \llbracket \mathbf{s} / x \rrbracket \rrbracket$.
- Let $\mathbf{r} = \mathbf{t}\{B\}$. Let $FV(\mathbf{s}) = \vec{X}$ and \vec{Y} be a set of free variables such that $\mathbf{s}[\vec{Y}/\vec{X}][\vec{X}/\vec{Y}] = \mathbf{s}$. Then, $\llbracket \mathbf{t}\{B\} \rrbracket \llbracket \llbracket \mathbf{s} \rrbracket / x \rrbracket = \llbracket \llbracket \mathbf{t} \rrbracket \{B\} \llbracket \llbracket \mathbf{s} \rrbracket / x \rrbracket \rrbracket = \llbracket \llbracket \mathbf{t} \rrbracket \llbracket \llbracket \mathbf{s}[\vec{Y}/\vec{X}]/x \rrbracket \rrbracket \{B\}[\vec{X}/\vec{Y}] \stackrel{IH}{=} \llbracket \llbracket \mathbf{t} \llbracket \mathbf{s}[\vec{Y}/\vec{X}]/x \rrbracket \rrbracket \{B\}[\vec{X}/\vec{Y}]$
 $= \llbracket \llbracket \mathbf{t} \llbracket \mathbf{s}[\vec{Y}/\vec{X}]/x \rrbracket \{B\} \rrbracket [\vec{X}/\vec{Y}] \stackrel{item 1}{=} \llbracket \llbracket \mathbf{t} \llbracket \mathbf{s}[\vec{Y}/\vec{X}]/x \rrbracket \{B\} \rrbracket [\vec{X}/\vec{Y}] = \llbracket (\mathbf{t}\{B\}) \llbracket \mathbf{s} / x \rrbracket \rrbracket$.
- Let $\mathbf{r} = \sum_{i=1}^n \frac{n_i}{d_i} \cdot \mathbf{r}_i$. $\llbracket \sum_{i=1}^n \frac{n_i}{d_i} \cdot \mathbf{r}_i \rrbracket \llbracket A/X \rrbracket = \pi_A(\sum_{i=1}^n m_i \cdot \llbracket \mathbf{r}_i \rrbracket) \llbracket A/X \rrbracket = \pi_A(\sum_{i=1}^n m_i \cdot \llbracket \mathbf{r}_i \rrbracket \llbracket A/X \rrbracket) \stackrel{IH}{=} \pi_A(\sum_{i=1}^n m_i \llbracket \mathbf{r}_i \llbracket A/X \rrbracket \rrbracket) = \llbracket \sum_{i=1}^n \frac{n_i}{d_i} \cdot \mathbf{r}_i \llbracket A/X \rrbracket \rrbracket = \llbracket (\sum_{i=1}^n \frac{n_i}{d_i} \cdot \mathbf{r}_i) \llbracket A/X \rrbracket \rrbracket$. \square

Theorem 4.6. If $\mathbf{r} \rightarrow^* \sum_{i=1}^n p_i \cdot \mathbf{t}_i$, with \mathbf{t}_i in Alg_F^p , with $\sum_{i=1}^n p_i = 1$ and $\llbracket \mathbf{t}_i \rrbracket \rightarrow^* \mathbf{s}_i$, then $\llbracket \mathbf{r} \rrbracket \rightarrow^* \mathbf{s}$ with probability $p_i (\sum_{j=1}^n p_j)^{-1}$ in λ_+^p .

Proof. Let $\mathbf{r} : A$ in Alg_F^p . For $i = 1, \dots, n$, assume $p_i = \frac{n_i}{d_i}$ with $n_i, d_i \in \mathbb{N}^*$. We proceed by a case analysis on the last reduction step to reach $\sum_{i=1}^n p_i \cdot \mathbf{t}_i$.

- If $\mathbf{r} = \sum_{i=1}^n p_i \cdot \mathbf{t}_i$, then $\pi_A(\sum_{i=1}^n (\prod_{k=1, k \neq i}^n d_k n_i) \cdot \llbracket \mathbf{t}_i \rrbracket) \rightarrow^* \pi_A(\sum_{i=1}^n (\prod_{k=1, k \neq i}^n d_k n_i) \cdot \mathbf{s}'_i)$ By Theorem 4.1, this term reduces in one step to \mathbf{s}'_i with probability

$$\frac{\prod_{k=1, k \neq i}^n d_k n_i}{\sum_{i=1}^n \left(\prod_{k=1, k \neq i}^n d_k n_i \right)} = \left(\frac{\frac{n_i}{d_i}}{\sum_{i=1}^n \frac{n_i}{d_i}} \right) \cdot \left(\frac{\prod_{k=1}^n d_k}{\prod_{k=1}^n d_k} \right) = p_i \left(\sum_{j=1}^n p_j \right)^{-1}.$$

- Consider $1.\mathbf{r} \rightleftharpoons \mathbf{r}$, with $\mathbf{r} = \sum_{i=1}^n p_i \cdot \mathbf{t}_i$. We have, $\llbracket 1.\mathbf{r} \rrbracket = \pi_A(1, \llbracket \mathbf{r} \rrbracket) \rightarrow^* \pi_A(1, \mathbf{s})$, which reduces with probability one to \mathbf{s} . Notice that \mathbf{s} is a reduct of $\llbracket \sum_{i=1}^n p_i \cdot \mathbf{t}_i \rrbracket = \pi_A(\sum_{i=1}^n m_i \cdot \llbracket \mathbf{t}_i \rrbracket)$. We conclude with Theorem 4.1.
- Consider $(\sum_{i=m+1}^n p_i \cdot \mathbf{t}_i) + (\sum_{i=1}^m p_i \cdot \mathbf{t}_i) \rightleftharpoons \sum_{i=1}^n p_i \cdot \mathbf{t}_i$, with $1 \leq m < n$. Since $\mathbf{r} : A$, then each $\mathbf{t}_i : A$. We have,

$$\llbracket \sum_{i=m+1}^n p_i \cdot \mathbf{t}_i + \sum_{i=1}^m p_i \cdot \mathbf{t}_i \rrbracket = \pi_A \left(\sum_{i=m+1}^n m_i \cdot \llbracket \mathbf{t}_i \rrbracket + \sum_{i=1}^m m_i \cdot \llbracket \mathbf{t}_i \rrbracket \right) \rightleftharpoons \pi_A \left(\sum_{i=1}^n m_i \cdot \llbracket \mathbf{t}_i \rrbracket \right).$$

where $m_i = \prod_{k=1, k \neq i}^n d_k n_i$. We conclude with Theorem 4.1.

- Consider $\lambda x^A. (\mathbf{r} + \mathbf{s}) \rightleftharpoons \lambda x^A. \mathbf{r} + \lambda x^A. \mathbf{s}$. We have $\llbracket \lambda x^A. (\mathbf{r} + \mathbf{s}) \rrbracket = \lambda x^A. (\llbracket \mathbf{r} + \mathbf{s} \rrbracket) = \lambda x^A. \pi_A(\llbracket \mathbf{r} \rrbracket + \llbracket \mathbf{s} \rrbracket) \rightarrow^* \lambda x^A. \pi_A(\mathbf{r}' + \mathbf{s}')$ By Theorem 4.1, $\lambda x^A. \pi_A(\mathbf{r}' + \mathbf{s}')$ reduces to $\lambda x^A. \mathbf{r}'$ (which is a reduct of $\llbracket \lambda x^A. \mathbf{r} \rrbracket = \lambda x^A. \llbracket \mathbf{r} \rrbracket$), with probability $\frac{1}{2}$, and to $\lambda x^A. \mathbf{s}'$ (which is a reduct of $\llbracket \lambda x^A. \mathbf{s} \rrbracket = \lambda x^A. \llbracket \mathbf{s} \rrbracket$), with probability $\frac{1}{2}$.
- Consider $(\lambda x^A. \mathbf{r}) \mathbf{s} \hookrightarrow \mathbf{r}[\mathbf{s}/x]$, with $\mathbf{r}[\mathbf{s}/x] = \sum_{i=1}^n p_i \cdot \mathbf{t}_i$. Then $\llbracket (\lambda x^A. \mathbf{r}) \mathbf{s} \rrbracket = (\lambda x^A. \llbracket \mathbf{r} \rrbracket) \llbracket \mathbf{s} \rrbracket \hookrightarrow \llbracket \mathbf{r} \rrbracket \llbracket \llbracket \mathbf{s} \rrbracket / x \rrbracket$ which, by Lemma 4.5(2), is equal to $\llbracket \mathbf{r}[\mathbf{s}/x] \rrbracket = \llbracket \sum_{i=1}^n p_i \cdot \mathbf{t}_i \rrbracket$ and this, by definition is equal to $\pi_A \left(\sum_{i=1}^n (\prod_{k=1, k \neq i}^n d_k n_i) \cdot \llbracket \mathbf{t}_i \rrbracket \right)$. We conclude with Theorem 4.1.
- Consider $(\Lambda X. \mathbf{r}) \{A\} \hookrightarrow \mathbf{r}[A/X]$, with $\mathbf{r}[A/X] = \sum_{i=1}^n p_i \cdot \mathbf{t}_i$. Then, $\llbracket \Lambda X. \mathbf{r} \{A\} \rrbracket = \Lambda X. \llbracket \mathbf{r} \{A\} \rrbracket \hookrightarrow \llbracket \mathbf{r} \{A\} \rrbracket [A/X]$, which by Lemma 4.5(1) is equal to $\llbracket \mathbf{r}[A/X] \rrbracket = \llbracket \sum_{i=1}^n p_i \cdot \mathbf{t}_i \rrbracket = \pi_A \left(\sum_{i=1}^n (\prod_{k=1, k \neq i}^n d_k n_i) \cdot \llbracket \mathbf{t}_i \rrbracket \right)$. We conclude with Theorem 4.1.

- Consider $p.q.r \hookrightarrow pq.r$. Let $p.q.r : A$. Since $p.q.r = \sum_{i=1}^n p_i.t_i$ with $\sum_{i=1}^n p_i = 1$, we have $n = 1$ and $pq = p_1 = 1$. Also, since $p.q.r$ is a term, $p = q = 1$. So, we have $\llbracket 1.1.r \rrbracket = \pi_A(1.\llbracket 1.r \rrbracket) = \pi_A(1.\pi_A(1.\llbracket r \rrbracket)) \rightarrow^* \pi_A(1.\pi_A(1.s))$. Notice that this term reduces with probability 1 to $\pi_A(1.s)$, which is a reduct of $\pi_A(1.\llbracket r \rrbracket) = \llbracket 1.r \rrbracket$.
- Consider $p.(r_1 + r_2) \hookrightarrow p.r_1 + p.r_2$. Since $p.r_1 + p.r_2 = \sum_{i=1}^n p_i.t_i$, with $\sum_{i=1}^n p_i = 1$, we have $n = 2$ and $p = \frac{1}{2}$, however in such case $\frac{1}{2}.(r_1 + r_2)$ is a pseudo-term, not a term.
- Consider $p.r + q.r \hookrightarrow (p + q).r$. Since $(p + q).r = \sum_{i=1}^n p_i.t_i$, with $\sum_{i=1}^n p_i = 1$, we have $n = 1$ and $p + q = 1$. Let $p = \frac{m}{d}$, then $q = \frac{d-m}{d}$. So, $\llbracket p.r + q.r \rrbracket = \pi_A(dm.\llbracket r \rrbracket + (d(d-m)).\llbracket r \rrbracket) \hookrightarrow \pi_A(d^2.\llbracket r \rrbracket)$, which reduces with probability 1 to s , where $\llbracket r \rrbracket \rightarrow^* s$.
- Contextual rules are straightforward. □

4.5 Back from λ_+^p to Alg_F^p

The inverse translation is given by

$$\begin{aligned} \langle x^A \rangle &= x^A & \langle \mathbf{rs} \rangle &= \langle \mathbf{r} \rangle \langle \mathbf{s} \rangle & \langle \mathbf{r}\{A\} \rangle &= \langle \mathbf{r} \rangle \{A\} \\ \langle \lambda x^A.r \rangle &= \lambda x^A.\langle \mathbf{r} \rangle & \langle \Lambda X.r \rangle &= \Lambda X.\langle \mathbf{r} \rangle & \langle \mathbf{r} + \mathbf{s} \rangle &= \langle \mathbf{r} \rangle + \langle \mathbf{s} \rangle \\ \text{If } \pi_A(\mathbf{t}) \hookrightarrow s_i \text{ with probability } p_i, \text{ for } i = 1, \dots, n, & & \langle \pi_A(\mathbf{t}) \rangle &= \sum_{i=1}^n p_i.\langle s_i \rangle \end{aligned}$$

Remark 4.7. This translation does not admit translating a term of the form $\pi_A(\mathbf{t})$ in normal form. Moreover, let Π be the rule “ $\pi_{A \Rightarrow B}(\mathbf{r})\mathbf{s} \rightleftharpoons \pi_B(\mathbf{rs})$ with $\mathbf{r} : A \Rightarrow (B \wedge C)$ ”, then the translation keep reductions, except for the one using rule Π , as expressed in Theorem 4.9.

Lemma 4.8.

1. $\langle \mathbf{r} \rangle [A/X] = \langle \mathbf{r}[A/X] \rangle$
2. $\langle \mathbf{r} \rangle [\langle \mathbf{s} \rangle / x] = \langle \mathbf{r}[s/x] \rangle$

Proof. Both items follow by induction on \mathbf{r} . Cases x^B , $\lambda x^B.t$, $\mathbf{t}_1\mathbf{t}_2$, $\Lambda Y.t$ and $\mathbf{t}\{B\}$ are analogous to those in proof of Lemma 4.5. Hence we only need to verify the case $\pi_B(\mathbf{t})$, when $\mathbf{r} \hookrightarrow \mathbf{r}_i$ with probability p_i , for $i = 1, \dots, n$.

1. $\langle \pi_B(\mathbf{t}) \rangle [A/X] = \langle \sum_{i=1}^n p_i.\langle \mathbf{r}_i \rangle \rangle [A/X] = \sum_{i=1}^n p_i.\langle \mathbf{r}_i \rangle [A/X]$, which by the induction hypothesis, is equal to $\sum_{i=1}^n p_i.\langle \mathbf{r}_i[A/X] \rangle = \langle \pi_{B[A/X]}(\mathbf{t}[A/X]) \rangle = \langle (\pi_B(\mathbf{t})) [A/X] \rangle$.
2. $\langle \pi_B(\mathbf{t}) \rangle [s/x] = \langle \sum_{i=1}^n p_i.\langle \mathbf{r}_i \rangle \rangle [s/x] = \sum_{i=1}^n p_i.\langle \mathbf{r}_i \rangle [s/x]$, which by the induction hypothesis, is equal to $\sum_{i=1}^n p_i.\langle \mathbf{r}_i[s/x] \rangle = \langle \pi_B(\mathbf{t}[s/x]) \rangle = \langle (\pi_B(\mathbf{t})) [s/x] \rangle$. □

Theorem 4.9. Let $\mathbf{r}, \mathbf{s}, s_i$ in λ_+^p .

- If $\mathbf{r} \rightleftharpoons \mathbf{s}$, then $\langle \mathbf{r} \rangle \rightleftharpoons \langle \mathbf{s} \rangle$.
- If $\mathbf{r} \hookrightarrow \mathbf{s}$, with probability 1, then $\langle \mathbf{r} \rangle \hookrightarrow \langle \mathbf{s} \rangle$, except if the reduction is done by rule Π .
- If $\mathbf{r} \hookrightarrow s_i$ with probability p_i , for $i = 1, \dots, n$, then $\langle \mathbf{r} \rangle = \sum_{i=1}^n p_i.\langle s_i \rangle$.

Proof. Case by case analysis.

- Consider $\mathbf{r} + \mathbf{s} \rightleftharpoons \mathbf{s} + \mathbf{r}$. Notice that $\langle \mathbf{r} + \mathbf{s} \rangle = \langle \mathbf{r} \rangle + \langle \mathbf{s} \rangle \rightleftharpoons \langle \mathbf{s} \rangle + \langle \mathbf{r} \rangle = \langle \mathbf{s} + \mathbf{r} \rangle$.
- Consider $(\mathbf{r} + \mathbf{s}) + \mathbf{t} \rightleftharpoons \mathbf{r} + (\mathbf{s} + \mathbf{t})$. Notice that $\langle (\mathbf{r} + \mathbf{s}) + \mathbf{t} \rangle = \langle (\mathbf{r} + \mathbf{s}) \rangle + \langle \mathbf{t} \rangle \rightleftharpoons \langle \mathbf{r} \rangle + \langle (\mathbf{s} + \mathbf{t}) \rangle = \langle \mathbf{r} + (\mathbf{s} + \mathbf{t}) \rangle$.

- Consider $(\mathbf{r} + \mathbf{s})\mathbf{t} \rightleftharpoons \mathbf{r}\mathbf{t} + \mathbf{s}\mathbf{t}$. Notice that $\langle (\mathbf{r} + \mathbf{s})\mathbf{t} \rangle = (\langle \mathbf{r} \rangle + \langle \mathbf{s} \rangle)\langle \mathbf{t} \rangle \rightleftharpoons \langle \mathbf{r} \rangle\langle \mathbf{t} \rangle + \langle \mathbf{s} \rangle\langle \mathbf{t} \rangle = \langle \mathbf{r}\mathbf{t} + \mathbf{s}\mathbf{t} \rangle$.
- Consider $\lambda x^A.(\mathbf{r} + \mathbf{s}) \rightleftharpoons \lambda x^A.\mathbf{r} + \lambda x^A.\mathbf{s}$. Notice that $\langle \lambda x^A.(\mathbf{r} + \mathbf{s}) \rangle = \lambda x^A.(\langle \mathbf{r} \rangle + \langle \mathbf{s} \rangle) \rightleftharpoons \lambda x^A.\langle \mathbf{r} \rangle + \lambda x^A.\langle \mathbf{s} \rangle = \langle \lambda x^A.\mathbf{r} + \lambda x^A.\mathbf{s} \rangle$.
- Consider $(\lambda x^A.\mathbf{r})\mathbf{s} \hookrightarrow \mathbf{r}[\mathbf{s}/x]$. Notice that $\langle (\lambda x^A.\mathbf{r})\mathbf{s} \rangle = (\lambda x^A.\langle \mathbf{r} \rangle)\langle \mathbf{s} \rangle \hookrightarrow \langle \mathbf{r} \rangle[\langle \mathbf{s} \rangle/x]$, and this, by Lemma 4.8(2), is equal to $\langle \mathbf{r}[\mathbf{s}/x] \rangle$.
- Consider $(\Lambda X.\mathbf{r})\{A\} \hookrightarrow \mathbf{r}[A/X]$. Notice that $\langle (\Lambda X.\mathbf{r})\{A\} \rangle = \Lambda X.\langle \mathbf{r} \rangle[A/X] \hookrightarrow \langle \mathbf{r} \rangle[A/X]$, and this, by Lemma 4.8(1), is equal to $\langle \mathbf{r}[A/X] \rangle$.
- Consider $\pi_A(\sum_{i=1}^n m_i.\mathbf{r}_i + \mathbf{s}) \hookrightarrow \mathbf{r}_i$ with probability $\frac{m_i}{\sum_{j=1}^n m_j}$, where $\mathbf{r}_i : A$ and $\mathbf{s} \not\vdash A$ are closed normal terms. Notice that, by definition, $\langle \pi_A(\sum_{i=1}^n m_i.\mathbf{r}_i + \mathbf{s}) \rangle = \sum_{i=1}^n \frac{m_i}{\sum_{j=1}^n m_j}.\langle \mathbf{r}_i \rangle$. \square

5 Conclusion

In this paper we have defined a probability space on the execution traces of non-confluent abstract rewrite systems. We define a sample space on strategies deciding the rewrite to apply at each state (cf. Definition 2.2).

Our main motivation has been to be able to use this probability space in non-deterministic calculi, hence being able to encode a probability superposition of the kind $\alpha.\mathbf{t} + \beta.\mathbf{r}$, with $\alpha + \beta = 1$, as a term having probability α of rewriting to \mathbf{t} and probability β of rewriting to \mathbf{r} . As an example, we provided such an encoding from an algebraic calculus into a non-deterministic calculus.

Acknowledgements We would like to thank Laurent Duvern et for enlightening discussions. This work has been funded by the ANR-10-JCJC-0208 CausaQ grant.

References

- [1] Suzana Andova (1999): *Process Algebra with Probabilistic Choice*. In Joost-Pieter Katoen, editor: *Formal Methods for Real-Time and Probabilistic Systems, Lecture Notes in Computer Science* 1601, pp. 111–129, doi:10.1007/3-540-48778-6_7.
- [2] Pablo Arrighi & Alejandro D iaz-Caro (2012): *A System F Accounting for Scalars*. *Logical Methods in Computer Science* 8(1):11, doi:10.2168/LMCS-8(1:11).
- [3] Pablo Arrighi, Alejandro D iaz-Caro & Beno t Valiron (2012): *A Type System for the Vectorial Aspects of the Linear-Algebraic Lambda-Calculus*. In Elham Kashefi, Jean Krivine & Femke van Raamsdonk, editors: *Developments in Computational Models, Electronic Proceedings in Theoretical Computer Science* 88, pp. 1–15, doi:10.4204/EPTCS.88.1.
- [4] Pablo Arrighi, Alejandro D iaz-Caro & Beno t Valiron (2013): *The Vectorial Lambda-Calculus*. Available at <http://www.arxiv.org/abs/1308.1138>. (Under review).
- [5] Pablo Arrighi & Gilles Dowek (2008): *Linear-algebraic λ -calculus: higher-order, encodings, and confluence*. In Andrei Voronkov, editor: *Rewriting Techniques and Applications, Lecture Notes in Computer Science* 5117, pp. 17–31, doi:10.1007/978-3-540-70590-1_2. Available at <http://www.arxiv.org/abs/quant-ph/0612199>.
- [6] Henk Barendregt (1984): *The Lambda Calculus: Its Syntax and Semantics*. North-Holland, Amsterdam.
- [7] John J. Benedetto & Wojciech Czaja (2009): *Integration and Modern Analysis*, chapter Lebesgue Measure and General Measure Theory. Birkh user Advanced Texts Basler Lehrb ucher, Birkh user Boston.

- [8] Gérard Boudol (1994): *Lambda-Calculi for (Strict) Parallel Functions*. *Information and Computation* 108(1), pp. 51–127, doi:[10.1006/inco.1994.1003](https://doi.org/10.1006/inco.1994.1003).
- [9] Olivier Bournez & Mathieu Hoyrup (2003): *Rewriting Logic and Probabilities*. In Robert Nieuwenhuis, editor: *Rewriting Techniques and Applications, Lecture Notes in Computer Science* 2706, pp. 61–75, doi:[10.1007/3-540-44881-0_6](https://doi.org/10.1007/3-540-44881-0_6). Available at <http://hal.inria.fr/inria-00099620>.
- [10] Antonio Bucciarelli, Thomas Ehrhard & Giulio Manzonetto (2012): *A Relational Semantics for Parallelism and Non-Determinism in a Functional Setting*. *Annals of Pure and Applied Logic* 163(7), pp. 918–934, doi:[10.1016/j.apal.2011.09.008](https://doi.org/10.1016/j.apal.2011.09.008). Available at hal.inria.fr/inria-00628887.
- [11] Ugo Dal Lago & Margherita Zorzi (2012): *Probabilistic operational semantics for the lambda calculus*. *RAIRO - Theoretical Informatics and Applications* 46(03), pp. 413–450, doi:[10.1051/ita/2012012](https://doi.org/10.1051/ita/2012012). Available at <http://www.arxiv.org/abs/1104.0195>.
- [12] Ugo de'Liguoro & Adolfo Piperno (1995): *Non Deterministic Extensions of Untyped λ -calculus*. *Information and Computation* 122(2), pp. 149–177, doi:[10.1006/inco.1995.1145](https://doi.org/10.1006/inco.1995.1145).
- [13] Mariangiola Dezani-Ciancaglini, Ugo de'Liguoro & Adolfo Piperno (1998): *A filter model for concurrent λ -calculus*. *SIAM Journal on Computing* 27(5), pp. 1376–1419, doi:[10.1137/S0097539794275860](https://doi.org/10.1137/S0097539794275860).
- [14] Alessandra Di Pierro, Chris Hankin & Herbert Wiklicky (2005): *Probabilistic λ -Calculus and Quantitative Program Analysis*. *Journal of Logic and Computation* 15(2), pp. 159–179, doi:[10.1093/logcom/exi008](https://doi.org/10.1093/logcom/exi008).
- [15] Alejandro Díaz-Caro & Gilles Dowek (2013): *Non determinism through type isomorphism*. In Delia Kesner & Petrucio Viana, editors: *Logical and Semantic Frameworks, with Applications, Electronic Proceedings in Theoretical Computer Science* 113, pp. 137–144, doi:[10.4204/EPTCS.113.13](https://doi.org/10.4204/EPTCS.113.13).
- [16] Alejandro Díaz-Caro & Gilles Dowek (2013): *Normalisation of a non-deterministic type isomorphic λ -calculus*. Available at <http://www.arxiv.org/abs/1306.5089>. (Under review).
- [17] Alejandro Díaz-Caro, Giulio Manzonetto & Michele Pagani (2013): *Call-by-value non-determinism in a linear logic type discipline*. In Sergei Artemov & Anil Nerode, editors: *Logical Foundations of Computer Science, Lecture Notes in Computer Science* 7734, pp. 164–178, doi:[10.1007/978-3-642-35722-0_12](https://doi.org/10.1007/978-3-642-35722-0_12).
- [18] Alejandro Díaz-Caro & Barbara Petit (2012): *Linearity in the non-deterministic call-by-value setting*. In Luke Ong & Ruy de Queiroz, editors: *Logic, Language, Information and Computation, Lecture Notes in Computer Science* 7456, pp. 216–231, doi:[10.1007/978-3-642-32621-9_16](https://doi.org/10.1007/978-3-642-32621-9_16). Available at <http://www.arxiv.org/abs/1011.3542>.
- [19] Herman Geuvers, Robbert Krebbers, James McKinna & Freek Wiedijk (2010): *Pure Type Systems without Explicit Contexts*. In Karl Crary & Marino Miculan, editors: *Logical Frameworks and Meta-languages: Theory and Practice, Electronic Proceedings in Theoretical Computer Science* 34, pp. 53–67, doi:[10.4204/EPTCS.34.6](https://doi.org/10.4204/EPTCS.34.6).
- [20] Oltea Mihaela Herescu & Catuscia Palamidessi (2000): *Probabilistic Asynchronous π -Calculus*. In Jerzy Tiuryn, editor: *Foundations of Software Science and Computation Structures, Lecture Notes in Computer Science* 1784, pp. 146–160, doi:[10.1007/3-540-46432-8_10](https://doi.org/10.1007/3-540-46432-8_10). Available at <http://www.arxiv.org/abs/cs/0109002>.
- [21] Michele Pagani & Simona Ronchi Della Rocca (2010): *Linearity, non-determinism and solvability*. *Fundamental Informaticae* 103(1–4), pp. 173–202, doi:[10.3233/FI-2010-324](https://doi.org/10.3233/FI-2010-324).
- [22] Jonghyun Park, Jeongbong Seo, Sungwoo Park & Gyesik Lee (2013): *Mechanizing Metatheory without Typing Contexts*. *Journal of Automated Reasoning*, doi:[10.1007/s10817-013-9287-4](https://doi.org/10.1007/s10817-013-9287-4).
- [23] TeReSe (2003): *Term Rewriting Systems*. *Cambridge Tracts in Theoretical Computer Science* 55, Cambridge University Press.
- [24] Lionel Vaux (2009): *The algebraic lambda calculus*. *Mathematical Structures in Computer Science* 19(5), pp. 1029–1059, doi:[10.1017/S0960129509990089](https://doi.org/10.1017/S0960129509990089). Available at <http://hal.archives-ouvertes.fr/hal-00379750>.