



# Slow oscillations of 'up' and 'down' states during general anesthesia: bi-stability in a non-linear model

Pedro Garcia, Axel Hutt

## ► To cite this version:

Pedro Garcia, Axel Hutt. Slow oscillations of 'up' and 'down' states during general anesthesia: bi-stability in a non-linear model. Bernstein Conference 2013, Sep 2013, Tuebingen, Germany. <hal-00920007>

**HAL Id: hal-00920007**

**<https://hal.inria.fr/hal-00920007>**

Submitted on 17 Dec 2013

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Slow oscillations of 'up' and 'down' states during general anesthesia: bi-stability in a non-linear model

Pedro Garcia Rodriguez, Axel Hutt

*NRIA CR Nancy - Grand Est, France*

The presence of sequences of alternating quiescent ('down') and bursting ('up') states in NREM sleep neuronal recordings is well established across mammalian species [1]. A striking related pattern of activity can also be observed in LFP and EEG recordings of anesthetized animals [2]. However, the dynamical principles underlying such phenomenon are unclear.

Some of the previous theoretical attempts to understand the neural basis of anesthesia effects have shown to be successful in replicating other remarkable characteristics of the associated EEG signals, e.g., the initial increase of the power spectrum and its subsequent decrease (in several frequency bands) during the process of the anesthesia induction [3-4]. Interestingly, such models predict the existence of notable non-linearities in the behavior of the neuronal membrane potential, with more than two stable branches through which the dynamics may evolve as the concentration of the anesthetic agent (AA) increases. Noise-induced transitions between two attractors might then occur, a theoretical framework that could explain the observed alternating sequences of 'up' and 'down' states. An example of this scenario is shown in Fig. 1, where 'p' stands for the AA level, and the oblique line is the separatrix between the two basins of attraction.

Nevertheless, the mathematical tractability of these non-linear scenarios is limited by the 2D phase space defined by the describing variables: the excitatory ( $V_e$ ) and inhibitory ( $V_i$ ) PSPs at excitatory cells. Exact solutions to these problems are only known for 1D systems and, consequently, the analysis of the existing non-linear models is typically avoided. In this work we undertake a first attempt to solve this task, by applying known techniques on stochastic phenomena theory [5] to the model proposed in [4]. We consider the time the system spends in the 'up' or 'down' states as being similar to the mean exit-time needed to escape from the corresponding attractor.

### References

- [1] Destexhe A. et al., Trends in Neurosciences 30: 334-342, 2007.
- [2] Hutt A. (ed.), Sleep and anesthesia: neural correlates in theory and experiment, Springer-Verlag, 2011. [3] Steyn-Ross M. et al., Physical Review E 60(6): 7299-7311, 1999.
- [4] Hutt A. and Longtin A., Cognitive Neurodynamics 4(1): 37-59, 2009.
- [5] Gardiner C.W., Handbook of Stochastic Methods (3rd edition), Springer-Verlag, 2004.

**Identifier:** W 110

**Conference:** Bernstein Conference 2013, Tübingen, Germany, September 24 - 27

**Citation:** Rodriguez P, Hutt A (2013) Slow oscillations of 'up' and 'down' states during general anesthesia: bi-stability in a non-linear model. *Bernstein Conference 2013*. doi: [10.12751/nncn.bc2013.0113](https://doi.org/10.12751/nncn.bc2013.0113)

**Type:** Poster