



Polarizing Double Negation Translations

Mélanie Boudard, Olivier Hermant

► **To cite this version:**

Mélanie Boudard, Olivier Hermant. Polarizing Double Negation Translations. Ken McMillan and Aart Middeldorp and Andrei Voronkov. LPAR, Dec 2013, Stellenbosch, South Africa. Springer, 8312, pp.182-197, 2013, LNCS ARCoSS. <hal-00920224v2>

HAL Id: hal-00920224

<https://hal.inria.fr/hal-00920224v2>

Submitted on 18 Dec 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Polarizing Double-Negation Translations

Mélanie Boudard¹ and Olivier Hermant²

¹ PRiSM, Univ. de Versailles-St-Quentin-en-Yvelines
CNRS, France

`melanie.boudard@prism.uvsq.fr`

² CRI, MINES ParisTech

`olivier.hermant@mines-paristech.fr`

Abstract. Double-negation translations are used to encode and decode classical proofs in intuitionistic logic. We show that, in the cut-free fragment, we can simplify the translations and introduce fewer negations. To achieve this, we consider the polarization of the formulæ and adapt those translation to the different connectives and quantifiers. We show that the embedding results still hold, using a customized version of the focused classical sequent calculus. We also prove the latter equivalent to more usual versions of the sequent calculus. This polarization process allows lighter embeddings, and sheds some light on the relationship between intuitionistic and classical connectives.

Keywords: classical logic, intuitionistic logic, double-negation translation, focusing.

1 Introduction

The relationship between different formal systems is a longstanding field of studies, and involves for instance conservativity, relative consistency or independence problems [1]. As for deductive systems, the natural question is to find a conservative encoding of formulæ. By conservative, we mean an encoding of formulæ such that a formula is provable in the first system if and only if its encoding is provable in the second system. This work was pioneered by Kolmogorov [2], Gödel [3] and Gentzen [4] for classical and intuitionistic logics. There exist several classes of sequents that are known to be classically provable if and only if they are intuitionistically provable [5].

In this paper, we refine those translations by removing a large number of unnecessary negations. Instead of focusing on invariant classes as in [5], we consider a translation on all the formulæ. A common point with this work, however, is the use of syntactic transformations. The proof systems we consider are the cut-free intuitionistic and classical sequent calculi [6]. This allows two remarks:

- the left rules of both calculi are identical; therefore it seems natural to translate them by themselves, when possible.

- In the absence of the cut rule, a formula is never active in different sides (both as an hypothesis and as a conclusion) of the turnstyle, having therefore a well-defined *polarity*. This last fact holds for all the rules except the axiom rule, which is easily dealt with, by an η -expansion-like argument, i.e. decomposing the formula by structural rules until we get axioms between atomic formula only.

In summary, we can avoid the introduction of negations on formulæ belonging to the “left” (or hypothesis) side of sequents. We also introduce further refinements, inspired by those of [3,4], to remove even more negations in the translation, based on the observation that some right-rules are also identical in the classical and intuitionistic calculi. To show conservativity by syntactic means without the cut rule, we need to impose a focusing discipline on the right-hand side of the classical sequent calculus, forced by the single-formula condition on the right-hand side of an intuitionistic sequent. We dedicate Section 4 to the study of a customized focused sequent calculus.

The price to pay of an asymmetric translation is that the result misses some modularity since we dismiss the cut rule: given a proof of a A and a proof of $A \Rightarrow B$, we cannot combine them with a cut rule. Both translations of A are not the same and so the translations of the proofs do not compose *directly*. See also the discussion in Section 6.

The paper is organized as follows. In Section 2, we give a brief overview of the background material, in particular the negative translations. In Section 3, we introduce a first polarized refinement of Kolmogorov negative translation, while Section 4 discusses the properties of the focused sequent calculus that we need in Section 5 to show that the polarized refinement of Gentzen-Gödel negative translation still has the same properties than the other translations. Section 6 concludes the paper.

2 Prerequisites

Here, we briefly recall the syntax of first-order logic, sequent calculus and the already known double-negation translations.

2.1 First-Order Logic

We assume that the reader is familiar with one-sorted first-order logic [6]: terms are composed of variables and function symbols applied to terms along their arities, and formulæ are either predicate symbols applied to terms along their arities or composed ones with the help of the conjunction (\wedge), disjunction (\vee), implication (\Rightarrow), negation (\neg) connectives and the universal (\forall) and existential (\exists) quantifiers.

To shorten the statement of our results and their proofs, we also define an operator that removes the trailing negation of formulæ, if any, and otherwise adds it.

Definition 1 (antinegation). *Let A be a formula, we let $\lrcorner A$ be:*

- B if A is equal to $\neg B$
- $\neg A$ otherwise.

Note that \lrcorner is not a connective, it is an operator, similar to Boolean complement in that $\lrcorner \lrcorner$ is the identity. In particular it has no associated rule in the sequent calculus. For instance $\lrcorner P(a)$ is the same as $\neg P(a)$ while $\lrcorner \lrcorner(A \wedge B)$ is the same as $(A \wedge B)$.

2.2 Sequent Calculi

Since they will be discussed in details in the next sections, we explicitly give the details of the classical and intuitionistic sequent calculi. A sequent is a pair of two multisets of formulæ, denoted $\Gamma \vdash \Delta$. The comma serves as a shorthand for multi-set union and Γ, A is an overloaded notation for $\Gamma, \{A\}$.

The classical sequent calculus is presented in Figure 1. The formula that is decomposed is called the *active* formula. The intuitionistic sequent calculus differs from the classical in the restriction imposed to the right-hand sides of sequents: it must be either empty, or reduced to one formula. Consequently, the following rules are modified: contr_R disappears; in the first premiss of the \Rightarrow_L rule and the axiom rule, Δ is empty; finally, the \vee_R rule splits to account for the choice of keeping A or B . For clarity, the intuitionistic sequent calculus is presented in Figure 2.

Note that, as announced, we do not consider the cut rule to be part of the calculus; so we reason in cut-free calculi.

2.3 Negative Translations

In this section, we briefly recall four existing translations [7,8]. In 1925, the first translation is published by Kolmogorov [2]. This translation involves adding a double negation in front of every subformula:

$$\begin{aligned} A^{Ko} &\equiv \neg\neg A \text{ for } A \text{ atomic} & (\neg A)^{Ko} &\equiv \neg\neg(\neg A^{Ko}) \\ (A \wedge B)^{Ko} &\equiv \neg\neg(A^{Ko} \wedge B^{Ko}) & (\forall x A)^{Ko} &\equiv \neg\neg\forall x A^{Ko} \\ (A \vee B)^{Ko} &\equiv \neg\neg(A^{Ko} \vee B^{Ko}) & (\exists x A)^{Ko} &\equiv \neg\neg\exists x A^{Ko} \\ (A \Rightarrow B)^{Ko} &\equiv \neg\neg(A^{Ko} \Rightarrow B^{Ko}) \end{aligned}$$

With the Kolmogorov's translation, A is provable using classical logic if and only if A^{Ko} is provable using intuitionistic logic.

A few years later, Gödel[3], and independently Gentzen[4], proposed a new translation, where disjunctions and existential quantifiers are replaced by a combination of negation and their De Morgan duals, respectively conjunctions and universal quantifiers:

| | |
|---|---|
| $\frac{}{\Gamma, A \vdash A, \Delta} \text{ax}$ | |
| $\frac{\Gamma, A, B \vdash \Delta}{\Gamma, A \wedge B \vdash \Delta} \wedge_L$ | $\frac{\Gamma \vdash A, \Delta \quad \Gamma \vdash B, \Delta}{\Gamma \vdash A \wedge B, \Delta} \wedge_R$ |
| $\frac{\Gamma, A \vdash \Delta \quad \Gamma, B \vdash \Delta}{\Gamma, A \vee B \vdash \Delta} \vee_L$ | $\frac{\Gamma \vdash A, B, \Delta}{\Gamma \vdash A \vee B, \Delta} \vee_R$ |
| $\frac{\Gamma \vdash A, \Delta \quad \Gamma, B \vdash \Delta}{\Gamma, A \Rightarrow B \vdash \Delta} \Rightarrow_L$ | $\frac{\Gamma, A \vdash B, \Delta}{\Gamma \vdash A \Rightarrow B, \Delta} \Rightarrow_R$ |
| $\frac{\Gamma \vdash A, \Delta}{\Gamma, \neg A \vdash \Delta} \neg_L$ | $\frac{\Gamma, A \vdash \Delta}{\Gamma \vdash \neg A, \Delta} \neg_R$ |
| $\frac{\Gamma, A[c/x] \vdash \Delta}{\Gamma, \exists x A \vdash \Delta} \exists_L$ | $\frac{\Gamma \vdash A[t/x], \Delta}{\Gamma \vdash \exists x A, \Delta} \exists_R$ |
| $\frac{\Gamma, A[t/x] \vdash \Delta}{\Gamma, \forall x A \vdash \Delta} \forall_L$ | $\frac{\Gamma \vdash A[c/x], \Delta}{\Gamma \vdash \forall x A, \Delta} \forall_R$ |
| $\frac{\Gamma, A, A \vdash \Delta}{\Gamma, A \vdash \Delta} \text{contr}_L$ | $\frac{\Gamma \vdash A, \Delta}{\Gamma \vdash A, A, \Delta} \text{contr}_R$ |
| $\frac{\Gamma \vdash \Delta}{\Gamma, A \vdash \Delta} \text{weak}_L$ | $\frac{\Gamma \vdash \Delta}{\Gamma \vdash A, \Delta} \text{weak}_R$ |

where, in \forall_L and \exists_R , c is a fresh constant and, in \forall_R and \exists_L , t is any term.

Fig. 1. Classical sequent calculus

$$\begin{aligned}
(A^{gg}) &\equiv \neg\neg A \text{ for } A \text{ atomic} & (\neg A)^{gg} &\equiv \neg A^{gg} \\
(A \wedge B)^{gg} &\equiv A^{gg} \wedge B^{gg} & (\forall x A)^{gg} &\equiv \forall x A^{gg} \\
(A \vee B)^{gg} &\equiv \neg(\neg A^{gg} \wedge \neg B^{gg}) & (\exists x A)^{gg} &\equiv \neg \forall x \neg A^{gg} \\
(A \Rightarrow B)^{gg} &\equiv A^{gg} \Rightarrow B^{gg}
\end{aligned}$$

As Kolmogorov's translation, Gödel-Gentzen's translation allows to show that A is provable using classical logic if and only if A^{gg} is provable using intuitionistic logic.

Kuroda [9] defined in 1951 a new translation:

$$\begin{aligned}
A^{Ku} &\equiv A \text{ for } A \text{ atomic} & (\neg A)^{Ku} &\equiv \neg A^{Ku} \\
(A \wedge B)^{Ku} &\equiv A^{Ku} \wedge B^{Ku} & (\forall x A)^{Ku} &\equiv \forall x \neg \neg A^{Ku} \\
(A \vee B)^{Ku} &\equiv A^{Ku} \vee B^{Ku} & (\exists x A)^{Ku} &\equiv \exists x A^{Ku} \\
(A \Rightarrow B)^{Ku} &\equiv A^{Ku} \Rightarrow B^{Ku}
\end{aligned}$$

A is provable classically if and only if $\neg\neg A^{Ku}$ is provable intuitionistically.

More recently, Krivine [10] has introduced a fourth translation:

$$\begin{aligned}
A^{Kr} &\equiv \neg A \text{ for } A \text{ atomic} & (\neg A)^{Kr} &\equiv \neg A^{Kr} \\
(A \wedge B)^{Kr} &\equiv A^{Kr} \vee B^{Kr} & (\forall x A)^{Kr} &\equiv \exists A^{Kr} \\
(A \vee B)^{Kr} &\equiv A^{Kr} \wedge B^{Kr} & (\exists x A)^{Kr} &\equiv \neg \exists x \neg A^{Kr} \\
(A \Rightarrow B)^{Kr} &\equiv \neg A^{Kr} \wedge B^{Kr}
\end{aligned}$$

| | |
|---|---|
| $\frac{}{\Gamma, A \vdash A} \text{ax}$ | |
| $\frac{\Gamma, A, B \vdash \Delta}{\Gamma, A \wedge B \vdash \Delta} \wedge_L$ | $\frac{\Gamma \vdash A \quad \Gamma \vdash B}{\Gamma \vdash A \wedge B} \wedge_R$ |
| $\frac{\Gamma, A \vdash \Delta \quad \Gamma, B \vdash \Delta}{\Gamma, A \vee B \vdash \Delta} \vee_L$ | $\frac{\Gamma \vdash A}{\Gamma \vdash A \vee B} \vee_{R1} \quad \frac{\Gamma \vdash B}{\Gamma \vdash A \vee B} \vee_{R2}$ |
| $\frac{\Gamma \vdash A \quad \Gamma, B \vdash \Delta}{\Gamma, A \Rightarrow B \vdash \Delta} \Rightarrow_L$ | $\frac{\Gamma, A \vdash B}{\Gamma \vdash A \Rightarrow B} \Rightarrow_R$ |
| $\frac{\Gamma \vdash A}{\Gamma, \neg A \vdash \Delta} \neg_L$ | $\frac{\Gamma, A \vdash}{\Gamma \vdash \neg A} \neg_R$ |
| $\frac{\Gamma, A[c/x] \vdash \Delta}{\Gamma, \exists x A \vdash \Delta} \exists_L$ | $\frac{\Gamma \vdash A[t/x]}{\Gamma \vdash \exists x A} \exists_R$ |
| $\frac{\Gamma, A[t/x] \vdash \Delta}{\Gamma, \forall x A \vdash \Delta} \forall_L$ | $\frac{\Gamma \vdash A[c/x]}{\Gamma \vdash \forall x A} \forall_R$ |
| $\frac{\Gamma, A, A \vdash \Delta}{\Gamma, A \vdash \Delta} \text{contr}_L$ | |
| $\frac{\Gamma \vdash \Delta}{\Gamma, A \vdash \Delta} \text{weak}_L$ | $\frac{\Gamma \vdash}{\Gamma \vdash A} \text{weak}_R$ |

where, in \forall_L and \exists_R , c is a fresh constant and, in \forall_R and \exists_L , t is any term.

Fig. 2. Intuitionistic sequent calculus

A is provable classically if and only if $\neg A^{Kr}$ is provable intuitionistically.

Using these existing translations, in particular Kolmogorov's and Gödel-Gentzen's translations, we propose to simplify them as described below.

3 Polarizing Kolmogorov's Translation

As in Kolmogorov's translation, let us define the polarized Kolmogorov's translation:

Definition 2. Let A, B, C and D be propositions. An occurrence of A in B is

– positive if:

- $B = A$.
- $B = C \wedge D$ and the occurrence of A is in C or in D and is positive.
- $B = C \vee D$ and the occurrence of A is in C or in D and is positive.
- $B = C \Rightarrow D$ and the occurrence of A is in C (resp. in D) and is negative (resp. positive).
- $B = \neg C$ and the occurrence of A is in C and is negative.
- $B = \forall x C$ and the occurrence of A is in C and is positive.
- $B = \exists x C$ and the occurrence of A is in C and is positive.

- negative if:
 - $B = C \wedge D$ and the occurrence of A is in C or in D and is negative.
 - $B = C \vee D$ and the occurrence of A is in C or in D and is negative.
 - $B = C \Rightarrow D$ and the occurrence of A is in C (resp. in D) and is positive (resp. negative).
 - $B = \neg C$ and the occurrence of A is in C and is positive.
 - $B = \forall x C$ and the occurrence of A is in C and is negative.
 - $B = \exists x C$ and the occurrence of A is in C and is negative.

Definition 3. Let A and B be propositions. We define by induction on the structure of propositions the positive (K^+) and negative translation (K^-):

$$\begin{array}{ll}
 A^{K^+} \equiv A \text{ if } A \text{ is atomic} & A^{K^-} \equiv A \text{ if } A \text{ is atomic} \\
 (A \wedge B)^{K^+} \equiv A^{K^+} \wedge B^{K^+} & (A \wedge B)^{K^-} \equiv \neg\neg A^{K^-} \wedge \neg\neg B^{K^-} \\
 (A \vee B)^{K^+} \equiv A^{K^+} \vee B^{K^+} & (A \vee B)^{K^-} \equiv \neg\neg A^{K^-} \vee \neg\neg B^{K^-} \\
 (A \Rightarrow B)^{K^+} \equiv \neg\neg A^{K^-} \Rightarrow B^{K^+} & (A \Rightarrow B)^{K^-} \equiv A^{K^+} \Rightarrow \neg\neg B^{K^-} \\
 (\neg A)^{K^+} \equiv \neg A^{K^-} & (\neg A)^{K^-} \equiv \neg A^{K^+} \\
 (\forall x A)^{K^+} \equiv \forall x A^{K^+} & (\forall x A)^{K^-} \equiv \forall x \neg\neg A^{K^-} \\
 (\exists x A)^{K^+} \equiv \exists x A^{K^+} & (\exists x A)^{K^-} \equiv \exists x \neg\neg A^{K^-}
 \end{array}$$

Notice how, compared to Section 2.3, we introduce double negations in front of subformulae instead of the whole formula. For instance axioms are translated by themselves, and the price to pay is, as for Kuroda’s and Krivine’s translations, a negation of the whole formula in the following theorem.

Theorem 1. If the sequent $\Gamma \vdash \Delta$ is provable in the classical sequent calculus then $\Gamma^{K^+}, \neg\Delta^{K^-} \vdash$ is provable in the intuitionistic sequent calculus.

Proof. By induction on the proof-tree. Since this theorem is not the main result of this paper, and is refined below (Theorem 3), let us process only one case. All other cases follow a similar pattern.

$$\exists_R \frac{\frac{\pi}{\Gamma \vdash \Delta, A[t/x]}}{\Gamma \vdash \Delta, \exists x A} \quad \Leftrightarrow \quad \frac{\frac{\text{IH}(\pi)}{\Gamma^{K^+}, \neg\Delta^{K^-}, \neg A^{K^-}[t/x] \vdash}}{\Gamma^{K^+}, \neg\Delta^{K^-} \vdash \neg\neg A^{K^-}[t/x]} \neg_R}{\Gamma^{K^+}, \neg\Delta^{K^-} \vdash \exists x \neg\neg A^{K^-}} \exists_R \quad \neg_L$$

where $\text{IH}(\pi)$ denotes, here and later, the proof obtained by the application of the induction hypothesis on π .

We also have the inverse translation.

Theorem 2. If the sequent $\Gamma^{K^+}, \neg\Delta^{K^-} \vdash D^{K^-}$ is provable in the intuitionistic sequent calculus, then $\Gamma \vdash \Delta, D$ is provable in the classical sequent calculus.

Proof. By a straightforward induction on the proof-tree.

We now focus on the polarization of the Gödel-Gentzen’s translation, which is lighter than the Kolmogorov’s translation, again with the idea of getting a simpler translation in both directions.

4 A Focused Sequent Calculus

Gödel-Gentzen negative translation (Definition 2.3 above) removes many negations from translations and the polarization we give in Section 5 will even more. If we want to follow the pattern of Theorem 1 to show equiprovability (in the absence of cut), we can *no longer* systematically move formulæ from the right to the left hand sides, since we lack negation on almost all connectives. Therefore, we must *constrain* our classical sequent calculus to forbid arbitrary proofs, and in particular to impose that once a rule has been applied on some formula of the right-hand side, the next rule must apply on the corresponding subformula of the premiss. Working on the same formula up to some well-chosen point is a discipline of capital importance, since we avoid to eagerly swap formulæ from right to left.

This is why we introduce a focused version of the classical sequent calculus. The resulting constraint is that we must decompose the *stoup* [11,12] formula until it gets removed from the *stoup* position. Only when the stoup becomes empty, can we apply rules on other formulæ.

Definition 4 (Focused sequent). *A focused sequent is a triple, composed of two multisets of formulæ and a distinguished set (the stoup) containing zero or one formula. It will be noted $\Gamma \vdash A; \Delta$ when the distinguished set contains a formula A , and $\Gamma \vdash .; \Delta$ when it contains no formula.*

The focused sequent calculus we define serves our particular purpose; for instance it is not optimized to maximize the so-called negative and positive phases [13]. Note also that in our paper, negative and positive has a very different meaning. The calculus is presented in Figure 3 and contains a stoup only in the right-hand side, since this is the only problematic side.

Note that all the left rules require an empty stoup, and that two new right rules, **focus** and **release**, respectively place and remove a formula of the right-hand side in the focus.

Only atomic, negated, disjunctive or existentially quantified formulæ can be removed from the stoup:

- Due to the freshness condition of the \exists -left and \forall -right rule, the \exists -right rule is the only rule that cannot be inverted (or equivalently permuted downwards). Therefore existential statements must be removable from the stoup.
- The stoup has only one place, so we cannot allow in it both subformulæ of a disjunction. This choice must be done by a subsequent call to the focus rule. More pragmatically, Gödel-Gentzen’s translation introduces negations in this case, enabling the storage of the subformulæ on the left-hand side of the sequent. As an informal translation rule, intuitionistic \neg_R rules will correspond to a lost of focus.
- The same reasoning holds for allowing atomic formulæ to be removed from the stoup. Also, if we do not allow this, the system loses completeness since the stoup becomes stuck forever.

- Allowing to remove negated formulæ from the stoup accounts for the aggressive behavior of the operator \lrcorner : to keep the statement of Theorem 3 short and close to statements of previous theorems, we must remember that \lrcorner removes the negation of negated formulæ, therefore forcing them to move on the left hand side.

As a consequence of the design constraint imposed by our translation, the rule focus cannot act on a formula which has \exists , \neg or \vee as main connective and the \exists_R , \neg_R and \vee_R rules act on formulæ that are not in the stoup (and, as mentioned, when the stoup itself is empty). The reasons become clear in the proof of Theorem 3.

Lastly, we impose the formula in the axiom rule to be atomic, which boils down to an η -expansion of the usual axiom rule.

To sum up, we consider the connectives \exists , \vee and \neg , when they appear on the right-hand side of a sequent, to have a "positive phase" in the sense of [13] and the other ones to have a negative phase.

We show that this calculus is equivalent to the usual sequent calculus of Figure 1.

Proposition 1. *Let Γ, Δ be two multisets of formulæ and A be a formula. If the sequent $\Gamma \vdash \cdot; \Delta$ (resp. $\Gamma \vdash A; \Delta$) has a proof in the focused sequent calculus, then it has a proof in the classical sequent calculus.*

Proof. Straightforward by noticing that, forgetting about the stoup (transforming the semicolon into a comma), all focused rules are instances of the classical sequent calculus rules. Both rules *focus* and *release* lose their meaning and are simply erased from the proof-tree. \square

The converse is a corollary of the slightly more general following statement. As we see below, it is crucial to have some degree of freedom to decompose arbitrarily Δ' into A and Δ in order to reason properly by induction.

Proposition 2. *Let Γ, Δ' be two multisets of formulæ. Assume that the sequent $\Gamma \vdash \Delta'$ has a proof in the classical sequent calculus. Let A be a set containing either a formula (also named A by abuse of notation) or the empty formula, and let Δ such that $\Delta' = A, \Delta$.*

Then the sequent $\Gamma \vdash A; \Delta$ has a proof in the focused sequent calculus.

Proof. The proof is a little bit more involved, but it appeals only to simple and well-known principles, in particular to Kleene's inversion lemmas [14,15], stating that inferences rules can be permuted and, therefore, gathered.

We give only a sketch of the proof, leaving out the details to the reader, for two reasons. Firstly, giving all the lengthy details would not add any insight on the structure of the proof; in the contrary they would blur the visibility of the main ideas. Secondly, similar completeness results are known for much more constrained focused proof systems; see for instance the one presented in [13].

| | |
|---|---|
| $\frac{}{\Gamma, A \vdash \cdot; A, \Delta} \text{ ax}$ | |
| $\frac{\Gamma, A, B \vdash \cdot; \Delta}{\Gamma, A \wedge B \vdash \cdot; \Delta} \wedge_L$ | $\frac{\Gamma \vdash A; \Delta \quad \Gamma \vdash B; \Delta}{\Gamma \vdash A \wedge B; \Delta} \wedge_R$ |
| $\frac{\Gamma, A \vdash \cdot; \Delta \quad \Gamma, B \vdash \cdot; \Delta}{\Gamma, A \vee B \vdash \cdot; \Delta} \vee_L$ | $\frac{\Gamma \vdash \cdot; A, B, \Delta}{\Gamma \vdash \cdot; A \vee B, \Delta} \vee_R$ |
| $\frac{\Gamma \vdash A; \Delta \quad \Gamma, B \vdash \cdot; \Delta}{\Gamma, A \Rightarrow B \vdash \cdot; \Delta} \Rightarrow_L$ | $\frac{\Gamma, A \vdash B; \Delta}{\Gamma \vdash A \Rightarrow B; \Delta} \Rightarrow_R$ |
| $\frac{\Gamma \vdash A; \Delta}{\Gamma, \neg A \vdash \cdot; \Delta} \neg_L$ | $\frac{\Gamma, A \vdash \cdot; \Delta}{\Gamma \vdash \cdot; \neg A, \Delta} \neg_R$ |
| $\frac{\Gamma, A[c/x] \vdash \cdot; \Delta}{\Gamma, \exists x A \vdash \cdot; \Delta} \exists_L$ | $\frac{\Gamma \vdash \cdot; A[t/x], \Delta}{\Gamma \vdash \cdot; \exists x A, \Delta} \exists_R$ |
| $\frac{\Gamma, A[t/x] \vdash \cdot; \Delta}{\Gamma, \forall x A \vdash \cdot; \Delta} \forall_L$ | $\frac{\Gamma \vdash A[c/x]; \Delta}{\Gamma \vdash \forall x A; \Delta} \forall_R$ |
| $\frac{\Gamma, A, A \vdash \cdot; \Delta}{\Gamma, A \vdash \cdot; \Delta} \text{ contr}_L$ | $\frac{\Gamma \vdash \cdot; A, \Delta}{\Gamma \vdash \cdot; A, A, \Delta} \text{ contr}_R$ |
| $\frac{\Gamma \vdash \cdot; \Delta}{\Gamma, A \vdash \cdot; \Delta} \text{ weak}_L$ | $\frac{\Gamma \vdash \cdot; \Delta}{\Gamma \vdash A; \Delta} \text{ weak}_R$ |
| | $\frac{\Gamma \vdash A; \Delta}{\Gamma \vdash \cdot; A, \Delta} \text{ focus}$ |
| | $\frac{\Gamma \vdash \cdot; A, \Delta}{\Gamma \vdash A; \Delta} \text{ release}$ |

where:

- the axiom rule involves only atomic formulæ,
- in \forall_L and \exists_R , c is a fresh constant,
- in \forall_R and \exists_L , t is any term,
- in release, A is either atomic or of the form $\exists x B, B \vee C$ or $\neg B$,
- in focus, A is neither atomic nor of the form $\exists x B, B \vee C$ or $\neg B$.

Fig. 3. Focused classical sequent calculus

First of all, we consider a refined version of the classical sequent calculus of Figure 1 where proofs are restricted to use the axiom and weak rules on atomic formulæ. In this way, we know [15] that Kleene's inversion lemmas [14] make the proof height *decrease* strictly. We reason by induction of the *height* of this modified proof-tree π , distinguishing the three following cases:

- A is empty, or A contains an atomic, existential, disjunctive or negated formula that is not the active formula of the last rule r of π . Then we release A , focus on the active formula if necessary, apply rule r , and we get one or two premises, on which we can apply the induction hypothesis. Let us give two instances, where, in the second case, A is empty:

$$\begin{array}{c}
\frac{\pi}{\frac{\Gamma \vdash B, A, \Delta}{\Gamma, \neg B \vdash A, \Delta}} \neg_L \quad \hookrightarrow \quad \frac{\text{IH}(\pi)}{\frac{\Gamma \vdash B; A, \Delta}{\Gamma, \neg B \vdash \cdot; A, \Delta} \neg_L} \text{release} \\
\frac{\pi_1}{\Gamma \vdash B, \Delta} \quad \frac{\pi_2}{\Gamma \vdash C, \Delta} \quad \hookrightarrow \quad \frac{\text{IH}(\pi_1)}{\Gamma \vdash B; \Delta} \quad \frac{\text{IH}(\pi_2)}{\Gamma \vdash C; \Delta} \quad \wedge_R \\
\frac{\Gamma \vdash B \wedge C, \Delta}{\Gamma \vdash B \wedge C, \Delta} \quad \hookrightarrow \quad \frac{\Gamma \vdash B \wedge C; \Delta}{\Gamma \vdash \cdot; B \wedge C, \Delta} \text{focus}
\end{array}$$

- A contains an atomic, existential, disjunctive or negated formula that is active in the last rule r of π . Then r must be one of the six rules axiom, \exists_R , \vee_R , \neg_R , weak_R or contr_R . They are direct and all the remaining cases follow a similar pattern. Here is the case for the \exists_R rule:

$$\frac{\pi}{\frac{\Gamma \vdash B[t/x], \Delta}{\Gamma \vdash \exists x B, \Delta}} \exists_R \quad \hookrightarrow \quad \frac{\text{IH}(\pi)}{\frac{\Gamma \vdash \cdot; B[t/x], \Delta}{\Gamma \vdash \cdot; \exists x B, \Delta} \exists_R} \text{release}$$

- If A is not empty and not an atomic, existential, disjunctive or negated formula then, disregarding the last rule of π , we apply Kleene's inversion lemma on A , the induction hypothesis on the premises, since the proof height has decreased, and recompose those premises to get back the corresponding component(s) of A in the stoup. Here is an example of such a rule :

$$\Rightarrow_R \frac{\pi}{\frac{\Gamma, B \vdash C, \Delta}{\Gamma \vdash B \Rightarrow C, \Delta}} \hookrightarrow \frac{\text{IH}(\pi)}{\frac{\Gamma, B \vdash C; \Delta}{\Gamma \vdash B \Rightarrow C; \Delta}} \Rightarrow_R$$

□

5 Polarizing Gödel-Gentzen's Translation

We try to reduce the number of negations. We use the polarization of propositions (Definition 2 above) and replace disjunction and existential quantifiers by conjunction and universal quantifiers, as in Gödel-Gentzen's translation.

Definition 5. *Let A and B be propositions. We define, by induction on the structure of propositions, the positive^(p) and negative⁽ⁿ⁾ translations:*

$$\begin{array}{ll}
A^p \equiv A \text{ if } A \text{ is atomic} & A^n \equiv \neg\neg A \text{ if } A \text{ is atomic} \\
(A \wedge B)^p \equiv A^p \wedge B^p & (A \wedge B)^n \equiv A^n \wedge B^n \\
(A \vee B)^p \equiv A^p \vee B^p & (A \vee B)^n \equiv \neg(\neg A^n \wedge \neg B^n) \\
(A \Rightarrow B)^p \equiv A^n \Rightarrow B^p & (A \Rightarrow B)^n \equiv A^p \Rightarrow B^n \\
(\neg A)^p \equiv \neg A^n & (\neg A)^n \equiv \neg A^p \\
(\forall x A)^p \equiv \forall x A^p & (\forall x A)^n \equiv \forall x A^n \\
(\exists x A)^p \equiv \exists x A^p & (\exists x A)^n \equiv \neg \forall x \neg A^n
\end{array}$$

Theorem 3. *Let Γ, Δ be multisets of formulae, and A be a set containing zero or one formula. If the sequent $\Gamma \vdash A; \Delta$ has a proof in the (classical) focused sequent calculus, then, in the intuitionistic sequent calculus, the sequent $\Gamma^p, \lrcorner\Delta^n \vdash A^n$ has a proof.*

Notice that \lrcorner removes the trailing negation of Δ in three cases: the negative translations of \exists , of \vee and of \neg (this last one more as a side-effect).

Proof. By induction on the proof of $\Gamma \vdash A; \Delta$, considering one by one the 19 cases from Figure 3:

- A left rule. We apply the induction hypothesis on the premises and copy the left rule. For instance, if the rule is \Rightarrow_L , then the induction hypothesis gives us proofs of the two sequents $\Gamma^p, \lrcorner\Delta^n \vdash A^n$ (since A is put in the stoup in the \Rightarrow_L rule) and $\Gamma^p, B^p, \lrcorner\Delta^n \vdash$ that can be readily combined with the (intuitionistic) \Rightarrow_L rule to yield a proof of the sequent $\Gamma^p, A^n \Rightarrow B^p, \lrcorner\Delta^n \vdash$. This is what we were looking for, since $(A \Rightarrow B)^p \equiv A^n \Rightarrow B^p$.
- A contr_R rule. It is transformed (after application of the induction hypothesis) into a contr_L rule.
- A weak_R rule. It is transformed into a weak_R rule.
- A release rule. This can occur only if A is atomic or of the form $\exists xB, B \vee C$ or $\neg B$. In all cases, we translate it as a \neg_R rule, which removes the trailing negation of A^n , turning it into the formula $\lrcorner A^n$ (see Definition 1), that integrates directly $\lrcorner\Delta^n$, so that we can readily plug the proof obtained by the application of the induction hypothesis.
- A focus rule on $A \in \Delta$. This can occur only if A is neither atomic nor of the form $\exists xB, B \vee C$ or $\neg B$. Therefore, $\lrcorner A^n = \neg A^n$, and we apply a \neg_L rule.
- An axiom rule. Since A is restricted to be atomic, we need to build an intuitionistic proof of the sequent $\Gamma^p, A, \lrcorner\neg A, \lrcorner\Delta^n \vdash$ which is a trivial two-step proof since $\lrcorner\neg A$ is $\neg A$.
- A \neg_R rule:

$$\frac{\frac{\pi}{\Gamma, B \vdash ; \Delta}}{\Gamma \vdash ; \neg B, \Delta}$$

We must find a proof of the sequent $\Gamma^p, \lrcorner\neg B^p, \lrcorner\Delta^n \vdash$. But $\lrcorner\neg B^p$ is the same as B^p , and the induction hypothesis gives us directly a proof of $\Gamma^p, B^p, \lrcorner\Delta^n \vdash$. In other words, \neg_R is not translated, thanks to the operator \lrcorner (that will soon lead us into minor considerations).

- A \vee_R rule:

$$\frac{\frac{\pi_1}{\Gamma \vdash ; B, C, \Delta}}{\Gamma \vdash ; B \vee C, \Delta}$$

We must build a proof of the sequent $\Gamma^p, \lrcorner(\neg B^n \wedge \neg C^n), \lrcorner\Delta^n \vdash$, which is equal to $\Gamma^p, \neg B^n \wedge \neg C^n, \lrcorner\Delta^n \vdash$. It is natural to try to apply the \wedge_L rule:

$$\frac{\Gamma^p, \neg B^n, \neg C^n, \lrcorner \Delta^n \vdash}{\Gamma^p, \neg B^n \wedge \neg C^n, \lrcorner \Delta^n \vdash} \wedge_L$$

We are committed to find a proof of the premiss, while the induction hypothesis gives us a proof of the following slightly different sequent:

$$\Gamma^p, \lrcorner B^n, \lrcorner C^n, \lrcorner \Delta^n \vdash$$

Therefore we must examine two subcases:

- B is an atom, an existential, disjunctive or negated formula. Then $B^n = \neg D$ for some D , and $\lrcorner B^n = D$. We build the following proof, given the proof obtained by application of the induction hypothesis:

$$\frac{\frac{\frac{\Gamma^p, D, \lrcorner C^n, \lrcorner \Delta^n \vdash}{\Gamma^p, \lrcorner C^n, \lrcorner \Delta^n \vdash \neg D} \neg_R}{\Gamma^p, \neg B^n, \lrcorner C^n, \lrcorner \Delta^n \vdash} \neg_L$$

- otherwise $\lrcorner B^n = \neg B^n$, and the induction hypothesis gives us directly a proof of the above sequent.

We do a similar case distinction on C to get from the previous proof a proof of the sequent $\Gamma^p, \neg B^n, \neg C^n, \neg \Delta^n \vdash$, which is now exactly what we were looking for.

– A \exists_R rule:

$$\frac{\frac{\pi}{\Gamma \vdash .; A[t/x], \Delta}}{\Gamma \vdash .; \exists x A, \Delta}$$

The induction hypothesis gives us a proof of the sequent $\Gamma^p, \lrcorner A[t/x]^n, \lrcorner \Delta^n \vdash$, that we turn, in the same way as in the previous case, into a proof of the sequent $\Gamma^p, \neg A[t/x]^n, \lrcorner \Delta^n \vdash$, to which we apply the \forall_L rule:

$$\forall_L \frac{\Gamma^p, \neg A[t/x]^n, \lrcorner \Delta^n \vdash}{\Gamma^p, \forall x \neg A^n, \lrcorner \Delta^n \vdash}$$

the end sequent is also equal to $\Gamma^p, \lrcorner \exists x A^n, \lrcorner \Delta^n \vdash$; so we have exhibited the proof we were looking for.

– A \wedge_R , \Rightarrow_R or \forall_R rule. Those three last cases are easy, since we are in the stoup, which corresponds to the right-hand side of the (intuitionistic) sequent. For example, let us consider the case of the \Rightarrow_R rule:

$$\Rightarrow_R \frac{\frac{\pi}{\Gamma, A \vdash B; \Delta}}{\Gamma \vdash A \Rightarrow B; \Delta} \quad \hookrightarrow \quad \frac{\text{IH}(\pi)}{\Gamma^p, A^p, \lrcorner \Delta^n \vdash B^n}}{\Gamma^p, \lrcorner \Delta^n \vdash A^p \Rightarrow B^n} \Rightarrow_R$$

□

The reverse translation is expressed with respect to the unfocused sequent calculus, which is more liberal and therefore more convenient for the reverse way. We nevertheless need to slightly generalize the statement.

Theorem 4. *Let $\Gamma, \Delta_1, \Delta_2$ be multisets of formulæ, such that Δ_1 does not contain any negated formula. Let D be at most one formula. If the sequent $\Gamma^p, \lrcorner\Delta_1^n, \neg\Delta_2^n \vdash D^n$ is provable in the intuitionistic sequent calculus, then $\Gamma \vdash \Delta_1, \Delta_2, D$ is provable in the classical sequent calculus.*

Proof. We constraint the intuitionistic proof to have a certain shape before starting the induction. First, we assume that axiom rules are restricted to atoms. It is always possible to expand the axioms that are not of this form. Second, we also assume that \neg_L rules on atomic formulæ are permuted upwards as far as they can [14], this basically induces that this \neg_L rule becomes glued either to an axiom or to a weakening rule and therefore the axiom case will be integrated to \neg_L case. This way, we avoid the presence of non-double negated axioms on the right-hand side. Unless stated otherwise we do not mention explicitly the application of the induction hypothesis, which is clear from the context.

- A contraction or a weakening on any of the formulæ of $\Gamma^p, \lrcorner\Delta_1^n, \neg\Delta_2^n$ or D^n is turned into the same rule on the corresponding formula of $\Gamma, \Delta_1, \Delta_2$ or D . Below, we now concentrate on connective and quantifier rules.
- A left-rule on Γ^p is turned into the same left-rule on Γ . The potential erasing of D^n in the two cases \Rightarrow_L and \neg_L is handled through a weakening.
- A left-rule on $\neg\Delta_2^n$ can be only a \neg_L rule. It is turned into a weakening on D^n if necessary, since we apply the induction hypothesis on the premiss $\Gamma^p, \lrcorner\Delta_1^n, \neg(\Delta_2')^n \vdash D_2^n$, with $\Delta_2 = \Delta_2', D_2$.
- A right-rule on D^n , assuming the main connective or quantifier of D is \wedge, \Rightarrow or \forall . The rule is \wedge_R, \Rightarrow_R or \forall_R , respectively. It is turned into the same right-rule on D .
- A right-rule on D^n , assuming D is an atomic, existentially quantified or disjunctive formula. The rule is \neg_R and the premiss is $\Gamma^p, \lrcorner\Delta_1^n, \lrcorner D^n, \neg\Delta_2^n \vdash$, to which we only need to apply the induction hypothesis.
- A right-rule on D^n assuming D is a negated formula $\neg D'$. In this case, the rule must be \neg_R and the premiss is of the form $\Gamma^p, \lrcorner\Delta_1^n, \lrcorner(D)^n, \neg\Delta_2^n \vdash$. But D is negated, and $\lrcorner D^n = \lrcorner\neg D'^p = D'^p$. So, to apply the induction hypothesis, we consider that the premiss is $\Gamma^p, D'^p, \lrcorner\Delta_1^n, \neg\Delta_2^n \vdash$.
- All possibilities for D have been examined. Notice in particular that a \vee_R or \exists_R rule cannot be applied on D^n , since the negative translation of Definition 5 never introduces this connective (resp. quantifier) in head position.
- A left-rule on $\lrcorner\Delta_1^n \in \lrcorner\Delta_1^n$, assuming D_1 is a disjunctive (resp. existentially quantified) formula. The rule is \wedge_L (resp. \forall_L) and is turned in an \vee_R (resp. \exists_R) rule on D_1 , making the active formula(e) of the premiss(es) move from Δ_1 to Δ_2 if necessary. Let us detail the \vee case. $D_1 = B_1 \vee C_1$, and $\lrcorner\Delta_1^n = \neg B_1^n \wedge \neg C_1^n$. The \wedge_L rule gives us the premiss :

$$\Gamma_1^p, \lrcorner(\Delta_1')^n, \neg B_1^n, \neg C_1^n, \neg\Delta_2^n \vdash D^n$$

We must distinguish according to the shapes of B_1 and C_1 . Let us discuss only B_1 , the discussion on C_1 being exactly the same:

- B_1 is an atomic, existentially quantified, disjunctive or negated formula. Then $\neg B_1^n$ is different from $\lrcorner B_1^n$, and to apply properly the induction hypothesis, B_1 must be placed into Δ_2 .
 - Otherwise $\neg B_1^n = \lrcorner B_1^n$ and it is left into Δ_1 .
- A left-rule on $\lrcorner D_1^n \in \lrcorner \Delta_1^n$, assuming that the main connective or quantifier of D_1 is \Rightarrow , \wedge or \forall . The rule is \neg_L in all cases, and we only need to weaken on D^n if necessary, before applying the induction hypothesis on the premises.
 - A left rule on $\lrcorner D_1^n \in \lrcorner \Delta_1^n$, assuming D_1 is atomic. By assumption, the next rule is a rule on D_1 . If it is a weakening, we translate both rules at once by weakening on D_1 and apply the induction hypothesis to the premiss of the weakening rule. Otherwise the next rule is an axiom. The only possibility is that D_1 belongs to Γ^p , and we translate both rules as an axiom.
 - By assumption, D_1 cannot be a negated formula, and therefore all the cases have been considered. \square

Corollary 1. *Let Γ, Δ be multisets of formulæ and D be at most one formula. If the sequent $\Gamma^p, \lrcorner \Delta^n \vdash D^n$ is provable in the intuitionistic sequent calculus, then the sequent $\Gamma \vdash \Delta, D$ is provable in the classical sequent calculus.*

Proof. Let $\neg C_1, \dots, \neg C_n$ be the negated formulæ of Δ and Δ' the other ones. We apply Theorem 4 to a Γ composed of Γ, C_1, \dots, C_n , a Δ_1 composed of Δ' , an empty Δ_2 and finally a D equal to D , which gives a proof of the sequent:

$$\Gamma, C_1, \dots, C_n \vdash \Delta', D$$

to which we apply n times the \neg_R rule to get back a proof of the wanted sequent.

6 Conclusion and Further Work

In this paper, we have shown that polarized double-negation translations still are used to navigate between intuitionistic and classical logics. They are lighter in terms of double negation, and let more statements being invariant by translation.

For instance, consider the axiom $(A \wedge B) \Rightarrow (A \vee B)$. Kolmogorov’s translation introduces 14 negations: $\neg\neg(\neg\neg(\neg\neg A \wedge \neg\neg B) \Rightarrow \neg\neg(\neg\neg A \vee \neg\neg B))$, while its (positive) polarized variant, only 10 of them: $(\neg\neg(\neg\neg A \wedge \neg\neg B) \Rightarrow (\neg\neg A \vee \neg\neg B))$. Gödel-Gentzen’s translation would be $(\neg\neg A \wedge \neg\neg B) \Rightarrow \neg(\neg\neg A \wedge \neg\neg B)$, introducing 11 negations, while its polarized version introduces only 4 of them: $(\neg\neg A \wedge \neg\neg B) \Rightarrow (A \vee B)$. Recent work from Frédéric Gilbert tends to show that it is possible to go further in the removal of double-negations by turning double negations into an operator that analyses the structure of the formula it double-negates. The polarization of Krivine’s translation remains also to be examined.

Polarized translations are particularity adapted to cut-free proofs; otherwise the same active formula may appear both in the left and the right hand sides. As the negative and positive translations of a formula usually differ, it is impossible to cut them back. The workaround can be a “manual” elimination of this cut

by reductive methods up to the point where both translations become equal, or to loosen the intuitionistic cut rule. We can also decide not to bother with cuts by eliminating them a priori. In all cases, however, we rely on a cut-elimination theorem that does not hold in the general case of the application described below.

Polarized double-negation translations has been primarily designed to fit *polarized deduction modulo* [16], an extension of first-order logic by a congruence on formulæ that is generated by *polarized* rewrite rules that apply only on a given side of the turn-style. It has already led to interesting results [17,18] in automated theorem proving within axiomatic theories. To support this approach, we must ensure the cut-elimination property of the (sequent calculus modulo the polarized) rewrite system.

One canonical way is to first show proof normalization for the natural deduction, and shift this result to the intuitionistic sequent calculus. Then, through a double-negation translation of *the rewrite system* this result can be extended to the classical sequent calculus [19]. In case of polarized rewriting, a polarized translation can be of great help for this last step, in addition to the development of normalization proofs via reducibility candidates. Another way to get cut admissibility would be to develop semantic proofs.

Lastly, it could be interesting to investigate whether, even in absence of cut admissibility as it can be the case, the modularity of our translations can be enforced, or whether cuts between two differently translated left- and right-formulæ can nevertheless be eliminated. We conjecture that this is possible, provided the rewrite relation is confluent and terminating.

Acknowledgments. The authors would like to thank P. Jouvelot for his comments and the choice of the title of this paper.

References

1. Cohen, P.J.: Set Theory and the Continuum Hypothesis. W.A. Benjamin, New York (1966)
2. Kolmogorov, A.: On the principle of the excluded middle. Mat. Sb. **32** (1925) 646–667
3. Gödel, K.: Zur intuitionistischen arithmetik und zahlentheorie. Ergebnisse eines mathematischen Kolloquiums **4** (1933) 34–38
4. Gentzen, G.: Die widerspruchsfreiheit der reinen zahlentheorie. Mathematische Annalen **112** (1936) 493:565
5. Schwichtenberg, H., Senjak, C.: Minimal from classical proofs. Annals of Pure and Applied Logic **164** (2013) 740–748
6. Schwichtenberg, H., Troelstra, A.S.: Basic Proof Theory. Cambridge University Press (1996)
7. Ferreira, G., Oliva, P.: On various negative translations. In: CL&C. (2010) 21–33
8. Troelstra, A.S., van Dalen, D.: Constructivism in Mathematics, An Introduction. North-Holland (1988)
9. Kuroda, S.: Intuitionistische untersuchungen der formalistischen logik. Nagoya Mathematical Journal **3** (1951) 35–47

10. Krivine, J.L.: Opérateurs de mise en mémoire et traduction de Gödel. *Arch. Math. Logic* **30** (1990) 241–267
11. Girard, J.Y.: On the unity of logic. *Ann. Pure Appl. Logic* **59** (1993) 201–217
12. Curién, P.L., Herbelin, H.: The duality of computation. In Odersky, M., Wadler, P., eds.: *ICFP, ACM* (2000) 233–243
13. Liang, C., Miller, D.: Focusing and polarization in linear, intuitionistic, and classical logics. *Theor. Comput. Sci.* **410** (2009) 4747–4768
14. Kleene, S.C.: Permutability of inferences in Gentzen’s calculi LK and LJ. *Memoirs of the American Mathematical Society* **10** (1952) 1–26, 27–68
15. Hermant, O.: Resolution is cut-free. *Journal of Automated Reasoning* **44** (2010) 245–276
16. Dowek, G.: Polarized deduction modulo. In: *IFIP Theoretical Computer Science*. (2010)
17. Burel, G.: Embedding deduction modulo into a prover. In Dawar, A., Veith, H., eds.: *CSL*. Volume 6247 of *Lecture Notes in Computer Science.*, Springer (2010) 155–169
18. Burel, G.: Experimenting with deduction modulo. In Bjørner, N., Sofronie-Stokkermans, V., eds.: *CADE*. Volume 6803 of *Lecture Notes in Computer Science.*, Springer (2011) 162–176
19. Dowek, G., Werner, B.: Proof normalization modulo. *The Journal of Symbolic Logic* **68** (2003) 1289–1316