

**Inverse problem in electrocardiography via factorization  
method of boundary problems : how reconstruct  
epicardial potential maps from measurements of the  
torso ?**

Julien Bouyssier, Nejib Zemzemi, Jacques Henry

► **To cite this version:**

Julien Bouyssier, Nejib Zemzemi, Jacques Henry. Inverse problem in electrocardiography via factorization method of boundary problems : how reconstruct epicardial potential maps from measurements of the torso ?. Poster Workshop Liryc, 24 et 25 Octobre 2013, Bordeaux. 2013. <hal-00926344>

**HAL Id: hal-00926344**

**<https://hal.inria.fr/hal-00926344>**

Submitted on 10 Jan 2014

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# INVERSE PROBLEM IN ELECTROCARDIOGRAPHY VIA FACTORIZATION METHOD OF BOUNDARY VALUE PROBLEMS :

How reconstruct epicardial potential maps from measurements of the torso ?

Julien Bouyssier, Nejib Zemzemi, Jacques Henry  
julien.bouyssier@inria.fr, nejib.zemzemi@inria.fr, jacques.henry@inria.fr



## Motivation and goal

**Motivation** : Solve the inverse problem in electrocardiography from measurements of the torso.

**Goal** : Use factorization method to compute epicardial potential maps.

This work is a simplified presentation of the method by considering a cylinder as geometry of our problem.

Initial problem : electrical potential  $u$  in the domain  $\Omega$  is governed by

$$(\mathcal{P}_0) \begin{cases} \Delta u = 0 \text{ in } \Omega, & \Omega : \text{cylinder} \\ u = 0 \text{ on } \Sigma, & \Sigma : \text{lateral surface} \\ u = T \text{ on } \Gamma_T, & T : \text{potential on the torso surface } \Gamma_T \\ \nabla u \cdot n = \Phi \text{ on } \Gamma_T, & \Phi : \text{normal derivative of the potential} \end{cases}$$

With  $T$  and  $\Phi$  known, find potential  $t$  and his normal derivative  $\phi$  on the heart surface  $\Gamma_H$  to complete this ill-posed Cauchy problem.

## Optimal control problem

$(\mathcal{P}_0)$  is decomposed into two sub-problems,  $\forall (\eta, \tau)$  :

$$(1) \begin{cases} \Delta u^1 = 0 \text{ in } \Omega \\ u^1 = 0 \text{ on } \Sigma \\ u^1 = T \text{ on } \Gamma_T \\ \nabla u^1 \cdot n = \eta \text{ on } \Gamma_H \end{cases} \quad \text{and} \quad (2) \begin{cases} \Delta u^2 = 0 \text{ in } \Omega \\ u^2 = 0 \text{ on } \Sigma \\ u^2 = \tau \text{ on } \Gamma_H \\ \nabla u^2 \cdot n = \Phi \text{ on } \Gamma_T \end{cases}$$

Solve  $(\mathcal{P}_0)$  :

$\Rightarrow$  Define the cost function  $E(\eta, \tau) = \int_{\Omega} (\nabla u^1(\eta) - \nabla u^2(\tau))^2$

$\Rightarrow$  Find  $(\phi, t)$  : minimize  $E(\eta, \tau)$

**New approach** : the factorization method by invariant embedding

## Principle of invariant embedding

**Principle** : transport potential data from torso surface to heart surface

$\Rightarrow$  Boundary value problems (1) and (2) are embedded into a family of similar problems on subdomains  $\Omega_S$

$\Rightarrow \Omega_S$  are bounded by a moving boundary  $\Gamma_S$  defined at  $x = s$  for  $x = 0 \rightarrow x = a$

$\Rightarrow$  At each position  $x = s$ , we impose a Neumann boundary condition  $\frac{\partial u_S^1}{\partial x}|_{\Gamma_S} = \alpha$  for (1) and a Dirichlet boundary condition  $(u_S^2)|_{\Gamma_S} = \beta$  for (2) :

$$(\mathcal{P}_S^1) \begin{cases} \Delta u_S^1 = 0 \text{ in } \Omega_S \\ u_S^1 = 0 \text{ on } \Sigma_S \\ u_S^1 = T \text{ on } \Gamma_T \\ \nabla u_S^1 \cdot n = \alpha \text{ on } \Gamma_S \end{cases} \quad \text{and} \quad (\mathcal{P}_S^2) \begin{cases} \Delta u_S^2 = 0 \text{ in } \Omega_S \\ u_S^2 = 0 \text{ on } \Sigma_S \\ u_S^2 = \beta \text{ on } \Gamma_S \\ \nabla u_S^2 \cdot n = \Phi \text{ on } \Gamma_T \end{cases}$$




FIGURE 1: Illustration of the moving boundary. The domain  $\Omega$  is a cylinder of a length  $a$  with a section  $\mathcal{O}$  and a lateral surface  $\Sigma$ . The section  $\mathcal{O} = \Gamma_T$  at  $x = 0$  represents the torso surface and the section  $\mathcal{O} = \Gamma_H$  at  $x = a$  represents the heart surface. The moving boundary  $\mathcal{O} = \Gamma_S$  at  $x = s$  is defined between these two surfaces and moving along the axis of revolution for  $x \in [0; a]$ . For each position  $s$ , a cylinder with a section  $\mathcal{O}$ , a lateral surface  $\Sigma_S$  and a length  $s$  is defined and forms a new subdomain  $\Omega_S$ .

## Resolution method

At each position  $x = s$  of  $\Gamma_S$ , we define two linear operators :

Neumann-to-Dirichlet application  $Q(s) : \alpha \mapsto u_S^1|_{\Gamma_S}$ , associated to  $(\mathcal{P}_S^1)$

Dirichlet-to-Neumann application  $P(s) : \beta \mapsto \frac{\partial u_S^2}{\partial x}|_{\Gamma_S}$ , associated to  $(\mathcal{P}_S^2)$

$\Rightarrow P$  and  $Q$  depend on  $s$  : variable that describes the axis of evolution

$\Rightarrow P$  and  $Q$  act on functions defined on section  $\mathcal{O}$

Let  $w_1$  and  $w_2$  : residual functions, defined on  $\mathcal{O}$ .  $\forall x \in [0; a]$ , we have :

$$(3) \begin{cases} u_S^1(x) = Q(x)\alpha + w_1(x) \\ \text{with } Q(0) = 0 \text{ and } w_1(0) = T \end{cases} \quad \text{and} \quad (4) \begin{cases} \frac{\partial u_S^2}{\partial x}(x) = P(x)\beta + w_2(x) \\ \text{with } P(0) = 0 \text{ and } w_2(0) = -\Phi \end{cases}$$

Let  $\Delta_y$  : laplacian operator, defined on the section  $\mathcal{O}$ .  $\forall x \in [0; a]$ , we have :

$$(5) \begin{cases} \frac{dP}{dx} + P^2 = -\Delta_y, & P(0) = 0 \\ \frac{dw_2}{dx} + Pw_2 = 0, & w_2(0) = -\Phi \end{cases} \quad \text{and} \quad (6) \begin{cases} \frac{dQ}{dx} - Q\Delta_y Q = I, & Q(0) = 0 \\ \frac{dw_1}{dx} - Q\Delta_y w_1 = 0, & w_1(0) = T \end{cases}$$

$\Rightarrow$  First solve Riccati equations for  $P$  and  $Q$  for  $x = 0 \rightarrow x = a$

$\Rightarrow$  Then solve equations for  $w_1$  and  $w_2$  for  $x = 0 \rightarrow x = a$

$\Rightarrow$  Compute operators and residuals at  $x = a$ .

Rename  $P(a) = P$ ,  $Q(a) = Q$ ,  $w_1(a) = w_1$ ,  $w_2(a) = w_2$

Define the matrix  $A$  as :  $A = \begin{pmatrix} Q & -QP \\ -PQ & P \end{pmatrix}$

We can rewrite  $E(\eta, \tau)$  as :  $E(\eta, \tau) = C + [\eta, \tau]A[\eta, \tau]' - 2 \int_{\Gamma_H} (w_1 P \tau + Q w_2 \eta)$

Finally :

$$(\phi, t) = \arg \min E(\eta, \tau) \iff A[\phi, t]' = [Qw_2, Pw_1]'$$

$\Rightarrow$  Find  $(\phi, t)$  : regularize previous system and inverse  $A$

## Conclusions and perspectives

**Conclusions** :

Direct optimal estimation of  $t$  and  $\phi$  before using any discretisation :

$\Rightarrow$  Analyse ill-posedness and propose a better regularization and discretization

Equations for  $P$  and  $Q$  depend only of the geometry :

$\Rightarrow$  Not necessary to repeat resolution at every time step of cardiac cycle

**Perspectives** :

Apply the method to 3D case where the moving boundary  $\Gamma_S$  will be a deformed surface :

$\Rightarrow$  First : model of spheres

$\Rightarrow$  Then : realistic geometries : how compute 3D surfaces ? + numerical cost ?




FIGURE 2: Illustration of  $\Gamma_S$  for the case of spheres, during the depolarization phase (left) and the repolarization phase (right). The smaller sphere represents the heart surface and the bigger one the torso surface. The heart was stimulated and direct problem was solved. We use computed potential data on the torso surface to apply factorization method and recover potential at heart surface. Spheres between torso and heart represent successive positions of  $\Gamma_S$  during invariant embedding. We also represent the potential recovered by the method.

**References**

- [1] Jacques Henry and Angel Manuel Ramos, *La méthode de factorisation des problèmes aux limites*, book (in preparation), (2013)
- [2] Fadhel Jday, *Complétion de données frontières : la méthode de plongement invariant*, PHD thesis, (2012)