

Asymptotic description of neural networks with correlated synaptic weights

Olivier Faugeras, James Maclaurin

► **To cite this version:**

Olivier Faugeras, James Maclaurin. Asymptotic description of neural networks with correlated synaptic weights. Entropy, MDPI, 2015, Special Issue Entropic Aspects in Statistical Physics of Complex Systems, 17(7), 4701-4743 (7), pp.4701-4743. hal-00955770

HAL Id: hal-00955770

<https://hal.inria.fr/hal-00955770>

Submitted on 6 Mar 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Asymptotic description of neural networks with correlated synaptic weights

Olivier Faugeras, James Maclaurin

**RESEARCH
REPORT**

N° 8495

March 2014

Project-Team NeuroMathComp



Asymptotic description of neural networks with correlated synaptic weights

Olivier Faugeras*, James Maclaurin*

Project-Team NeuroMathComp

Research Report n° 8495 — March 2014 — 44 pages

Abstract: We study the asymptotic law of a network of interacting neurons when the number of neurons becomes infinite. Given a completely connected network of neurons in which the synaptic weights are Gaussian correlated random variables, we describe the asymptotic law of the network when the number of neurons goes to infinity. All previous works assumed that the weights were i.i.d. random variables, thereby making the analysis much simpler. This hypothesis is not realistic from the biological viewpoint. In order to cope with this extra complexity we introduce the process-level empirical measure of the trajectories of the solutions to the equations of the finite network of neurons and the averaged law (with respect to the synaptic weights) of the trajectories of the solutions to the equations of the network of neurons. The main result of this article is that the image law through the empirical measure satisfies a large deviation principle with a good rate function which is shown to have a unique global minimum. Finally, our analysis of the rate function allows us also to describe this minimum as a stationary Gaussian measure which completely characterizes the activity of the infinite size network.

Key-words: Large Deviations, Good Rate Function, Stationary Gaussian Processes, Stationary Measures, Spectral Representations, Neural Networks, Firing Rate Neurons, Correlated Synaptic Weights.

AMS Mathematics Subject Classification:

28C20, 34F05, 34K50, 37L55, 60B11, 60F10, 60G10, 60G15, 60G57, 60G60, 60H10, 62M45, 92C20.

* NeuroMathComp Laboratory, INRIA Sophia-Antipolis, 2004 route des Lucioles, B.P. 93, 06902 Sophia-Antipolis, France. Email: `firstname.name@inria.fr`

**RESEARCH CENTRE
SOPHIA ANTIPOLIS – MÉDITERRANÉE**

2004 route des Lucioles - BP 93
06902 Sophia Antipolis Cedex

Description asymptotique de réseaux de neurones avec poids synaptiques corrélés

Résumé : Nous étudions la loi asymptotique décrivant un réseau de neurones interconnectés lorsque le nombre de neurones tend vers l'infini. Étant donné un réseau complètement connecté de neurones dans lequel les poids synaptiques sont des variables aléatoires gaussiennes corrélées, nous caractérisons la loi asymptotique de ce réseau lorsque le nombre de neurones tend vers l'infini. Tous les travaux précédents faisaient l'hypothèse de poids indépendants. Cette hypothèse n'est pas réaliste d'un point de vue biologique mais elle simplifie considérablement l'analyse mathématique du problème. Pour faire face à cette difficulté supplémentaire nous introduisons la mesure empirique sur l'espace des trajectoires solutions des équations du réseau de neurones de taille finie et la loi moyennée (par rapport aux poids synaptiques) des trajectoires de ces solutions. Notre résultat principal est que la loi image de cette loi par la mesure empirique satisfait un principe de grandes déviations dont nous montrons que la bonne fonction de taux admet un minimum global unique. Notre analyse de la fonction de taux nous permet enfin de décrire ce minimum comme une mesure gaussienne stationnaire nous permettant ainsi de caractériser complètement l'activité du réseau de taille infinie.

Mots-clés : Grandes Déviations, Bonne Fonction de Taux, Processus Gaussiens Stationnaires, Mesures Stationnaires, Représentations Spectrales, Réseaux de Neurones, Neurones à Taux de Décharge, Poids Synaptiques Corrélés.

Contents

1	Introduction	4
2	Frequently used notations	6
3	The neural network model	8
3.1	The model equations	9
3.2	The laws of the uncoupled and coupled processes	10
3.2.1	Preliminaries	10
3.2.2	Coupled and uncoupled processes	12
4	The good rate function	14
4.1	Gaussian processes	14
4.2	Convergence of Gaussian Processes	16
4.3	Definition of the functional Γ	18
4.3.1	Γ_1	18
4.3.2	Γ_2	19
4.4	The Radon-Nikodym derivative	21
5	The large deviation principle	22
5.1	Lower bound on the open sets	24
5.2	Exponential Tightness of Π^N	24
5.3	Upper Bound on the Compact Sets	25
5.3.1	An LDP for a Gaussian measure	25
5.3.2	An upper bound for Π^N over compact sets	27
5.4	End of the proof of theorem 3.8	30
6	Characterization of the unique minimum of the rate function	31
7	Some important consequences of theorem 3.8	33
8	Possible extensions	34
9	Conclusion	35
A	Two useful lemmas	35
B	Proof of proposition 4.6	36
C	Existence of a lower bound for $\Phi^N(\mu, v)$	37
D	Proof of lemmas 5.4 and 5.14	38
E	Proof of lemma 5.11	41

1 Introduction

The goal of this paper is to study the asymptotic behaviour and large deviations of a network of interacting neurons when the number of neurons becomes infinite. Our network may be thought of as a network of weakly-interacting diffusions: thus before we begin we briefly overview other asymptotic analyses of such systems. In particular, a lot of work has been done on spin glass dynamics, including Ben Arous and Guionnet on the mathematical side [33, 3, 4, 34] and Sompolinsky and his co-workers on the theoretical physics side [44, 45, 13, 14]. Furthermore the large deviations of weakly interacting diffusions has been extensively studied by Dawson and Gartner [17, 18], and more recently Budhiraja, Dupuis and Fischer [6, 28]. More references to previous work on this particular subject can be found in these references.

Because the dynamics of spin glasses is not too far from that of networks of interacting neurons, Sompolinsky also successfully explored this particular topic [43] for fully connected networks of rate neurons, i.e. neurons represented by the time variation of their firing rates (the number of spikes they emit per unit of time), as opposed to spiking neurons, i.e. neurons represented by the time variation of their membrane potential (including the individual spikes). For an introduction to these notions, the interested reader is referred to such textbooks as [30, 35, 26]. In his study of the continuous time dynamics of networks of rate neurons, Sompolinsky and his colleagues assumed, as in the work on spin glasses, that the coupling coefficients, called the synaptic weights in neuroscience, were random variables i.i.d. with zero mean Gaussian laws. The main result obtained by Ben Arous and Guionnet for spin glass networks using a large deviations approach (resp. by Sompolinsky and his colleagues for networks of rate neurons using the local chaos hypothesis) under the previous hypotheses is that the averaged law of Langevin spin glass (resp. rate neurons) dynamics is chaotic in the sense that the averaged law of a finite number of spins (resp. of neurons) converges to a product measure as the system gets very large.

The next theoretical efforts in the direction of understanding the averaged law of rate neurons are those of Cessac, Moynot and Samuelides [10, 39, 40, 11, 42]. From the technical viewpoint, the study of the collective dynamics is done in discrete time, assuming no leak (this term is explained below) in the individual dynamics of each of the rate neurons. Moynot and Samuelides obtained a large deviation principle and were able to describe in detail the limit averaged law that had been obtained by Cessac using the local chaos hypothesis and to prove rigorously the propagation of chaos property. Moynot extended these results to the more general case where the neurons can belong to two populations, the synaptic weights are non-Gaussian (with some restrictions) but still i.i.d., and the network is not fully connected (with some restrictions) [39].

One of the next challenges is to incorporate in the network model the fact that the synaptic weights are not independent and in effect often highly correlated. One of the reasons for this is the plasticity processes at work at the levels of the synaptic connections between neurons; see for example [36] for a biological viewpoint, and [19, 30, 26] for a more computational and mathematical account of these phenomena.

The problem we solve in this paper is the following. Given a completely connected network of firing rate neurons in which the synaptic weights are Gaussian correlated random variables, we describe the asymptotic law of the network when the number of neurons goes to infinity. Like in [39, 40] we study a discrete time dynamics but unlike these authors we cope with more complex intrinsic dynamics of the neurons, in particular we allow for a leak (to be explained in more detail below).

To be complete, let us mention the fact that this problem has already partially been explored in Physics by Sompolinsky and Zippelius [44, 45] and in Mathematics by Alice Guionnet [34] who analysed symmetric spin glass dynamics, i.e. the case where the matrix of the coupling coefficients (the synaptic weights in our case) is symmetric. This is a very special case of correlation. The

work in [15] is also an important step forward in the direction of understanding the spin glass dynamics when more general correlations are present.

Let us also mention very briefly another class of approaches toward the description of very large populations of neurons where the individual spikes generated by the neurons are considered. The model for individual neurons is usually of the class of Integrate and Fire (IF) neurons [37] and the underlying mathematical tools are those of the theory of point-processes [16]. Important results have been obtained in this framework by Gerstner and his collaborators, e.g. [31, 29] in the case of deterministic synaptic weights. Related to this approach but from a more mathematical viewpoint, important results on the solutions of the mean-field equations have been obtained in [9]. In the case of spiking neurons but with a continuous dynamics (unlike that of IF neurons), the first author and collaborators have recently obtained some limit equations that describe the asymptotic dynamics of fully connected networks of neurons [1] with independent synaptic weights.

Because of the correlation of the synaptic weights, the natural space to work in is the infinite dimensional space of the trajectories, noted \mathcal{T}^Z , of a countably-infinite set of neurons and the set of stationary probability measures defined on this set, noted $\mathcal{M}_{1,S}^+(\mathcal{T}^Z)$.

We introduce the process-level empirical measure, noted $\hat{\mu}_N$, of the N trajectories of the solutions to the equations of the network of N neurons and the averaged (with respect to the synaptic weights) law Q^N of the N trajectories of the solutions to the equations of the network of N neurons. The first result of this article (theorem 3.8) is that the image law Π^N of Q^N through $\hat{\mu}^N$ satisfies a large deviation principle (LDP) with a good rate function H which is shown to have a unique global minimum, μ_e . Thus, with respect to the measure Π^N on $\mathcal{M}_{1,S}^+(\mathcal{T}^Z)$, if the set X contains the measure δ_{μ_e} , then $\Pi^N(X) \rightarrow 1$ as $N \rightarrow \infty$, whereas if δ_{μ_e} is not in the closure of X , $\Pi^N(X) \rightarrow 0$ as $N \rightarrow \infty$ exponentially fast and the constant in the exponential rate is determined by the rate function. Our analysis of the rate function allows us also, and this is our second result (theorem 6.4), to characterize the limit measure μ_e as the image of a stationary Gaussian measure $\underline{\mu}_e$ defined on a transformed set of trajectories \mathcal{T}^Z . This is potentially very useful for applications since $\underline{\mu}_e$ can be completely characterized by its mean and spectral density. Furthermore the rate function allows us to quantify the probability of finite-size effects. Theorems 3.8 and 6.4 allows us to characterize the average (over the synaptic weights) behaviour of the network. We also derive, and this is our third result, some properties of the infinite-size network that are true for almost all realizations of the synaptic weights (theorems 7.1 and 7.4).

The paper is organized as follows. In section 2 we provide a glossary of some frequently used notations, in section 3 we describe the equations of our network of neurons, the type of correlation between the synaptic weights, define the proper state spaces and introduce the different probability measures that are necessary for establishing our results, in particular the process-level empirical measure, $\hat{\mu}_N$, Π^N and the image R^N through $\hat{\mu}_N$ of the law of the uncoupled neurons. We state the principle result of this paper in Theorem 3.8.

In section 4 we introduce a certain Gaussian process attached to a given measure in $\mathcal{M}_{1,S}^+(\mathcal{T}^Z)$ and $\mathcal{M}_{1,S}^+(\mathcal{T}^N)$ and motivate this introduction by showing that the Radon-Nikodym derivative of Q^N with respect to the law of the uncoupled neurons can be expressed by the Gaussian process corresponding to the empirical measure $\hat{\mu}_N$. This allows us to compute the Radon-Nikodym derivative of Π^N with respect to R^N for any measure in $\mathcal{M}_{1,S}^+(\mathcal{T}^Z)$. Using these results, section 5 is dedicated to the proof of the existence of a strong LDP for the measure Π^N . In section 6 we show that the good rate function obtained in the previous section has a unique global minimum and we characterize it as the image of a stationary Gaussian measure. Section 7 is dedicated to drawing some important consequences of our first main theorem, in particular some quenched results. Section 8 explores some possible extensions of our work and we conclude with section 9.

2 Frequently used notations

$\langle \cdot, \cdot \rangle, \ \cdot \ $	The usual Euclidean inner product and corresponding norm in \mathbb{R}^T , in effect $\mathcal{T}_{1,T}$.
*	Complex conjugation.
$\mathbf{1}_M$	The M -dimensional vector (M positive integer) with coordinates equal to 1.
$\mathbf{1}_X$	The indicator function of the set X .
$A_{[N]}^\mu$	The $NT \times NT$ symmetric block-circulant matrix equal to $K_{[N]}^\mu(\sigma^2 \text{Id}_{NT} + K_{[N]}^\mu)^{-1}$ (page 16).
$A_{[N]}^{\mu,k}$	The k th $T \times T$ block of $A_{[N]}^\mu$, $k = -n \cdots n$ (page 16).
$\tilde{A}_{[N]}^{\mu,l}$	The Discrete Fourier transform of the sequence $A_{[N]}^{\mu,k}$, $k, l = -n \cdots n$ (equation (4.12)).
$\tilde{A}^\mu(\theta)$	Defined by equation (4.13) from $\tilde{K}^\mu(\theta)$.
$\tilde{A}_{[N]}^\mu(\theta)$	Defined right after equation (4.14).
$A^{\mu,k}$	The k th Fourier coefficient of the periodic matrix-valued function $\tilde{A}^\mu(\theta)$, see equation (4.14).
$\tilde{A}^{\mu,N,l}$	The l th Fourier coefficient of the sequence $(A^{\mu,k})$, $k = -n \cdots n$, beginning of section 5.3.2.
${}^\dagger B$	The transpose of the matrix or vector B .
$\mathcal{B}(\mathcal{X})$	The set of Borelian sets of the topological set \mathcal{X} .
c^μ	The T -dimensional mean vector of $G^{\mu,k}$, $k \in \mathbb{Z}$, for $\mu \in \mathcal{M}_{1,S}^+(\mathcal{T}^{\mathbb{Z}})$ (equation (4.1)).
DFT	Discrete Fourier Transform (page 10).
\mathcal{E}_2	subset of $\mathcal{M}_{1,S}^+(\mathcal{T}^{\mathbb{Z}})$, such that $\pi^1(\pi_{1,T}(v))$ is square-integrable, see definition 4.
f	Monotonically increasing bijection $\mathbb{R} \rightarrow]0, 1[$, page 9.
\mathcal{F}	Short notation for $\mathcal{B}(\mathcal{T}^{\mathbb{Z}})$, see page 11.
$\mathcal{F}_{s,t}$	$\pi_{s,t}(\mathcal{F})$
G^μ	The T -dimensional spatially stationary Gaussian process taking its values in $\mathcal{T}_{1,T}^{\mathbb{Z}}$ and defined by the measure $\mu \in \mathcal{M}_{1,S}^+(\mathcal{T}^{\mathbb{Z}})$ (section 4.1).
$G_{[N]}^\mu$	The T -dimensional spatially shift-invariant (modulo N) Gaussian process taking its values in $\mathcal{T}_{1,T}^N$ and defined by the measure $\mu^N \in \mathcal{M}_{1,S}^+(\mathcal{T}^N)$ (section 4.1).

$\Gamma_{[N],1}, \Gamma_1$	Section 4.3.1.
$\Gamma_{[N],2}, \Gamma_2$	Section 4.3.2.
Γ^ν, Γ_2^ν	Linear approximations of Γ and Γ_2 , section 5.3.1.
$\Gamma_{[N]}^\nu, \Gamma_1^N, \Gamma_{[N],2}^\nu$	Section 5.3.1.
$I^{(2)}(\mu^N, \nu^N)$	Relative entropy of μ^N with respect to ν^N , $\mu^N, \nu^N \in \mathcal{M}_1^+(\mathcal{T}^N)$ (equation (5.1)).
$i \bmod N$	lies between $-n$ and n , $N = 2n + 1$, see page 9
$I^{(3)}(\mu, \nu^Z)$	Process-level entropy of μ with respect to $\nu \in \mathcal{M}_1^+(\mathcal{T})$, $\mu \in \mathcal{M}_{1,S}^+(\mathcal{T}^Z)$, (equation (5.3)).
J_{ij}	The (random) synaptic weight from neuron j to neuron i .
J^N	The $N \times N$ matrix of the synaptic weights (page 9).
\bar{J}	Determines the mean of the random synaptic weights (page 9).
Γ	Mapping from \mathcal{T}^Z or \mathcal{T}^N to $\mathbb{R} \cup \{+\infty\}$ (section 4.3). One has $\Gamma = \Gamma_1 + \Gamma_2$.
k_f	Lipschitz constant of f , page 9 and equation (3.11).
$K^{\mu,k}$	The $T \times T$ covariance matrix of $G^{\mu,i}$ and $G^{\mu,i+k}$, $i, k \in \mathbb{Z}$ and $\mu \in \mathcal{M}_{1,S}^+(\mathcal{T}^Z)$ (equation (4.6)).
$K_{[N]}^{\mu,k}$	The $T \times T$ covariance matrix of $G_{[N]}^{\mu,i}$ and $G_{[N]}^{\mu,i+k}$, $i, k = -n \cdots n$, $i+k$ modulo N and $\mu \in \mathcal{M}_{1,S}^+(\mathcal{T}^Z)$ (equation (4.11)).
$K_{[N]}^\mu$	The $(NT \times NT)$ covariance matrix of the sequence of Gaussian random variables $(G_{[N]}^{\mu,-n}, \dots, G_{[N]}^{\mu,n})$ (page 16).
$K^{\mu,N}$	The $(NT \times NT)$ covariance matrix of the sequence of Gaussian random variables, see Section 5.3.1. $(G^{\mu,0}, G^{\mu,1}, \dots, G^{\mu,n}, G^{\mu,-n}, G^{\mu,-n+1}, \dots, G^{\mu,-1})$ (equation (5.13)).
$\tilde{K}^\mu(\theta), (K_{[N]}^\mu(\theta))$	The Fourier transform of the sequence $K^{\mu,k}$, $k \in \mathbb{Z}$, $K_{[N]}^{\mu,k}$, $k = -n \cdots n$, (proposition 4.1 (page 16 and Section 4.2)
$\tilde{K}_{[N]}^{\mu,l}$	The Discrete Fourier transform of the sequence $K_{[N]}^{\mu,k}$, $k, l = -n \cdots n$ (page 15).
$\Lambda(k, l)$	Describes the covariance matrix of the synaptic weights, see (3.2).
$\tilde{\Lambda}(\theta_1, \theta_2)$	The Fourier transform of $(\Lambda(k, l))_{k,l \in \mathbb{Z}}$, (proposition 3.2).
Λ^{min}	The sum of the absolutely convergent series $(\Lambda(k, l))_{k,l \in \mathbb{Z}}$ (page 10).
Λ^{sum}	The sum of the convergent series $(\Lambda(k, l))_{k,l \in \mathbb{Z}}$ (equation (3.4)).
LDP	Large Deviation Principle (theorem 3.8).
$M^{\mu,k}$	The $T \times T$ matrix measuring the correlation between the vectors $f(\pi_{0,T-1}(u^0))$ and $f(\pi_{0,T-1}(u^k))$, $k \in \mathbb{Z}$ for $\mu \in \mathcal{M}_{1,S}^+(\mathcal{T}^Z)$ (equation 4.2).
$\mathcal{M}_1^+(\mathcal{T}^N), \mathcal{M}_1^+(\mathcal{T}^Z)$	Sets of probability measures on $\mathcal{T}^N, \mathcal{T}^Z$ (equation (4.2)).
$\mathcal{M}_{1,S}^+(\mathcal{T}^Z)$	Set of stationary probability measures on \mathcal{T}^Z , see section 3.2.1
$\mathcal{M}_{1,S}^+(\mathcal{T}^N)$	Set of stationary probability measures on \mathcal{T}^N , see section 3.2.2
μ^N	The N -dimensional spatial marginal $\mu \circ (\pi^N)^{-1}$ of $\mu \in \mathcal{M}_{1,S}^+(\mathcal{T}^Z)$ (page 11).

$\mu_{s,t}$	The $t - s + 1$ -dimensional ($0 \leq s \leq t \leq T$) temporal marginal $\mu \circ \pi_{s,t}^{-1}$ of $\mu \in \mathcal{M}_{1,S}^+(\mathcal{T}^Z)$ (page 11).
$\underline{\mu}$ $\hat{\mu}_N(u^{-n}, \dots, u^n)$	For each measure $\mu \in \mathcal{M}_1^+(\mathcal{T}^N)$ or $\mathcal{M}_{1,S}^+(\mathcal{T}^Z)$, $\mu \circ \Psi^{-1}$ (definition 1). The process level empirical measure of the element (u^{-n}, \dots, u^n) of \mathcal{T}^N : an element of $\mathcal{M}_{1,S}^+(\mathcal{T}^Z)$, see (3.10).
μ_I	The law of the initial condition of each neuron, an element of $\mathcal{M}_1^+(\mathbb{R})$, see section 3.2.2.
μ_{u_0}	The conditional probability distribution of μ , given the initial conditions u_0^j , $j \in Z$, see page 11.
N	The number of neurons in the finite network, assumed to be odd, $N = 2n + 1$, $n \geq 0$.
$\mathcal{N}_p(m, \Sigma)$	The neurons are indexed from $-n$ to n . The law of the p -dimensional Gaussian variable with mean m and covariance matrix Σ , page 9.
P	The law of the solution to (3.13) (in $\mathcal{M}_1^+(\mathcal{T})$).
$\pi_{s,t}$	The temporal projection from \mathcal{T} to $\mathcal{T}_{s,t}$ (section 3.2.1).
π^N	The spatial projection from \mathcal{T}^Z to \mathcal{T}^N , page 9.
ϕ^N	Mapping from $\mathcal{M}_{1,S}^+(\mathcal{T}^Z) \times \mathcal{T}_{1,T}^N$ to \mathbb{R} used in the definition of $\Gamma_{[N],2}$, see equation (4.26).
ϕ_∞^N	Mapping from $\mathcal{M}_{1,S}^+(\mathcal{T}^Z) \times \mathcal{T}_{1,T}^N$ to \mathbb{R} used in the definition of $\Gamma_{[N],2}^\nu$, see equation (5.12).
π^N	The spatial projection from \mathcal{T}^Z onto \mathcal{T}^N such that $\pi^N(u) = (u^{-n}, \dots, u^n)$ (page 10)
Π_N	The image law of Q^N through $\hat{\mu}_N$ (definition 2).
Ψ	A bijection from \mathcal{T} to \mathcal{T} (and by extension from \mathcal{T}^N to \mathcal{T}^N and from \mathcal{T}^Z to \mathcal{T}^Z), see equation (3.14).
$\Psi_{1,T}$	A mapping from \mathcal{T} to $\mathcal{T}_{1,T}$, page 13.
$Q^N(J^N)$	The law of the solution to (3.1) (in $\mathcal{M}_1^+(\mathcal{T}^N)$)
Q^N	The averaged law of $Q^N(J^N)$ w.r.t. to the synaptic weights J^N : an element of $\mathcal{M}_{1,S}^+(\mathcal{T}^N)$.
R_N	The image law of $P^{\otimes N}$ through $\hat{\mu}_N$ (definition 2).
S	The shift operator on \mathcal{T}^Z , see (3.6).
T	The maximum time in the study of the dynamics described by (3.1).
\mathcal{T}	Set of trajectories from time 0 to time T , elements of $\mathbb{R}^{[0 \dots T]}$ (section 3.2.1).
$\mathcal{T}_{s,t}$	Set of trajectories from time s to time $0 \leq s \leq t \leq T$, elements of $\mathbb{R}^{[s \dots t]}$ (section 3.2.1).
$\mathcal{T}^Z, \mathcal{T}_{s,t}^Z$	Set of doubly infinite sequences of elements of \mathcal{T} , (section 3.2.1), of elements of $\mathcal{T}_{s,t}$.

3 The neural network model

We consider a fully connected network of N neurons. Not all sets of neurons are fully connected but many are, e.g. within the same cortical column. One of the major aims of this article is to quantify how quickly the system converges to its limit, so the rate function gives us a means of assessing whether the number of neurons in a cortical column is sufficiently high for the mean

field equations to be accurate. For simplicity but without loss of generality, we assume N odd¹ and write $N = 2n + 1$, $n \geq 0$. The state of the neurons is described by the variables (U_t^j) , $j = -n \cdots n$, $t = 0 \cdots T$ which represent the values of the neurons membrane potentials.

3.1 The model equations

The equation describing the time variation of the membrane potential U_t^j of the j th neuron writes

$$U_t^j = \gamma U_{t-1}^j + \sum_{i=-n}^n J_{ji}^N f(U_{t-1}^i) + B_{t-1}^j, \quad j = -n, \dots, n \quad t = 1, \dots, T. \quad (3.1)$$

$f : \mathbb{R} \rightarrow]0, 1[$ is a monotonically increasing bijection which we assume to be Lipschitz continuous. Its Lipschitz constant is noted k_f . We could for example employ $f(x) = (1 + \tanh(gx))/2$, where the parameter g can be used to control the slope of the ‘‘sigmoid’’ f at the origin $x = 0$.

This equation involves the parameters γ , J_{ij}^N , and the time processes B_t^j , $i, j = -n, \dots, n$, $t = 0, \dots, T - 1$. The initial conditions are discussed at the beginning of section 3.2.2.

γ is in $[0, 1)$ and determines the time scale of the intrinsic dynamics, i.e. without interactions, of the neurons. If $\gamma = 0$ the dynamics is said to have no leak.

The B_t^j s represent random fluctuations of the membrane potential of neuron j . They are independent *random processes* with the same law. We assume that at each time instant t , the B_t^j s are i.i.d. random variables distributed as $\mathcal{N}_1(0, \sigma^2)^2$.

The J_{ij}^N s are the synaptic weights. J_{ij}^N represents the strength with which the ‘presynaptic’ neuron j influences the ‘postsynaptic’ neuron i . They are *Gaussian random variables*, independent of the membrane fluctuations, whose mean is given by

$$\mathbb{E}[J_{ij}^N] = \frac{\bar{J}}{N},$$

where \bar{J} is some number independent of N .

We note J^N the $N \times N$ matrix of the synaptic weights, $J^N = (J_{ij}^N)_{i,j=-n,\dots,n}$. Their covariance is assumed to satisfy the following shift invariance property,

$$\text{cov}(J_{ij}^N J_{kl}^N) = \text{cov}(J_{i+m,j+n}^N J_{k+m,l+n}^N)$$

for all indexes $i, j, k, l = -n, \dots, n$ and all integers m and n , the indexes being taken modulo N . Here, and throughout this paper, $i \bmod N$ is taken to lie between $-n$ and n .

Remark 3.1. *This shift invariance property is technically useful since it allows us to use the tools of Fourier analysis. In terms of the neural population it means that the neurons “live” on a circle. Therefore, unlike in the uncorrelated case studied in the papers cited in the introduction, we have to indirectly introduce a notion of space.*

We stipulate the covariances through a covariance function $\Lambda : \mathbb{Z}^2 \rightarrow \mathbb{R}$ and assume that they scale as $1/N$. We write

$$\text{cov}(J_{ij}^N J_{kl}^N) = \frac{1}{N} \Lambda((k - i) \bmod N, (l - j) \bmod N). \quad (3.2)$$

The function Λ is even:

$$\Lambda(-k, -l) = \Lambda(k, l), \quad (3.3)$$

¹When N is even the formulae are slightly more complicated but all the results we prove below in the case N odd are still valid.

²We note $\mathcal{N}_p(m, \Sigma)$ the law of the p -dimensional Gaussian variable with mean m and covariance matrix Σ .

corresponding to the simultaneous exchange of the two presynaptic and postsynaptic neurons ($cov(J_{ij}^N J_{kl}^N) = cov(J_{kl}^N J_{ij}^N)$).

We must make further assumptions on Λ to ensure that the system is well-behaved as the number of neurons N asymptotes to infinity. We assume that the series $(\Lambda(k, l))_{k, l \in \mathbb{Z}}$ is absolutely convergent, i.e.

$$\Lambda^{sum} = \sum_{k, l = -\infty}^{\infty} |\Lambda(k, l)| < \infty, \quad (3.4)$$

and furthermore that

$$\Lambda^{min} = \sum_{k, l = -\infty}^{\infty} \Lambda(k, l) > 0$$

We let Λ^N be the restriction of Λ to $[-n, n]^2$, i.e. $\Lambda^N(i, j) = \Lambda(i, j)$ for $-n \leq i, j \leq n$.

We next introduce the spectral properties of Λ that are crucial for the results in this paper. We use throughout the paper the notation that if x is some quantity, \tilde{x} represents its Fourier transform in a sense that depends on the particular space where x is defined. For example $\tilde{\Lambda}$ is the 2π doubly periodic Fourier transform of the function Λ whose properties are described in the next proposition. Similarly, $\tilde{\Lambda}^N$ is the two-dimensional Discrete Fourier Transform (DFT) of the doubly periodic sequence Λ^N . The proof of the following proposition is obvious.

Proposition 3.2. *The sum $\tilde{\Lambda}(\theta_1, \theta_2)$ of the absolutely convergent series $(\Lambda(k, l)e^{-i(k\theta_1 + l\theta_2)})_{k, l \in \mathbb{Z}}$ is continuous on $[-\pi, \pi]^2$ and positive. The covariance function Λ is recovered from the inverse Fourier transform of $\tilde{\Lambda}$:*

$$\Lambda(k, l) = \frac{1}{(2\pi)^2} \int_{-\pi}^{\pi} \int_{-\pi}^{\pi} \tilde{\Lambda}(\theta_1, \theta_2) e^{i(k\theta_1 + l\theta_2)} d\theta_1 d\theta_2$$

Moreover there exists $\tilde{\Lambda}^{min} > 0$ such that

$$\tilde{\Lambda}^N(0, 0) \geq \tilde{\Lambda}^{min} > 0, \quad (3.5)$$

for N large enough.

3.2 The laws of the uncoupled and coupled processes

3.2.1 Preliminaries

Sets of trajectories, temporal and spatial projections

The time evolution of one membrane potential is represented by the set $\mathbb{R}^{[0 \dots T]} := \mathcal{T}$ of finite sequences $(u_t)_{t=0, \dots, T}$ of length $T + 1$ of real numbers. \mathcal{T}^N is the set of sequences (u^{-n}, \dots, u^n) ($N = 2n + 1$) of elements of \mathcal{T} that we use to describe the solutions to (3.1). Similarly we note $\mathcal{T}^{\mathbb{Z}}$ the set of doubly infinite sequences of elements of \mathcal{T} . If u is in $\mathcal{T}^{\mathbb{Z}}$ we note u^i , $i \in \mathbb{Z}$, its i th coordinate, an element of \mathcal{T} . Hence $u = (u^i)_{i=-\infty \dots \infty}$.

Given the integers s and t , $0 \leq s \leq t \leq T$, we define the temporal projection $\pi_{s,t} : \mathcal{T} \rightarrow \mathbb{R}^{[s \dots t]} := \mathcal{T}_{s,t}$ as the set of finite sequences of length $t - s + 1$ of real numbers such that $\pi_{s,t}(u) = (u_r)_{r=s \dots t} := u_{s,t}$. When $s = t$ we note π_t and \mathcal{T}_t rather than $\pi_{t,t}$ and $\mathcal{T}_{t,t}$. The temporal projection $\pi_{s,t}$ extends in a natural way to \mathcal{T}^N and $\mathcal{T}^{\mathbb{Z}}$: for example $\pi_{s,t}$ maps \mathcal{T}^N to $\mathcal{T}_{s,t}^N$. We define the spatial projection $\pi^N : \mathcal{T}^{\mathbb{Z}} \rightarrow \mathcal{T}^N$ ($N = 2n + 1$) to be $\pi^N(u) = (u^{-n}, \dots, u^n)$. Temporal and spatial projections commute, i.e. $\pi^N \circ \pi_{s,t} = \pi_{s,t} \circ \pi^N$.

The shift operator $\mathcal{S} : \mathcal{T}^{\mathbb{Z}} \rightarrow \mathcal{T}^{\mathbb{Z}}$ is defined by

$$(\mathcal{S}u)^i = u^{i+1}, \quad i \in \mathbb{Z}. \quad (3.6)$$

Given the element $u = (u^{-n}, \dots, u^n)$ of \mathcal{T}^N we form the doubly infinite periodic sequence

$$p_N(u) = (\dots, u^{n-1}, u^n, u^{-n}, \dots, u^n, u^{-n}, u^{-n+1}, \dots) \quad (3.7)$$

which is an element of $\mathcal{T}^{\mathbb{Z}}$. We have $(p_N(u))^i = u^{(i \bmod N)}$. p_N is a mapping $\mathcal{T}^N \rightarrow \mathcal{T}^{\mathbb{Z}}$. With a slight abuse of notation we also note \mathcal{S} the shift operator induced by \mathcal{S} on \mathcal{T}^N through the function p_N :

$$\mathcal{S}u = \pi^N(\mathcal{S}p_N(u)) \quad u \in \mathcal{T}^N \quad (3.8)$$

Topologies on the sets of trajectories

We equip $\mathcal{T}^{\mathbb{Z}}$ with the projective topology, i.e. the topology generated by the following metric. For $u, v \in \mathcal{T}^N$, we define their distance $d_N(u, v)$ to be

$$d_N(u, v) = \sup_{|j| \leq n, 0 \leq s \leq T} |u_s^j - v_s^j|.$$

This allows us to define the following metric over $\mathcal{T}^{\mathbb{Z}}$, whereby if $u, v \in \mathcal{T}^{\mathbb{Z}}$, then

$$d(u, v) = \sum_{N=1}^{\infty} 2^{-N} (d_N(\pi^N u, \pi^N v) \wedge 1). \quad (3.9)$$

Equipped with this topology, $\mathcal{T}^{\mathbb{Z}}$ is Polish (a complete, separable metric space).

The metric d generates the Borelian σ -algebra $\mathcal{B}(\mathcal{T}^{\mathbb{Z}}) := \mathcal{F}$. It is generated by the coordinate functions $(u_t^i)_{i \in \mathbb{Z}, t=0 \dots T}$. The spatial and temporal projections defined above can be used to define the corresponding σ -algebras on the sets $\mathcal{T}_{s,t}^N$, e.g. $\mathcal{F}_{s,t}^N = \pi^N(\pi_{s,t}(\mathcal{F}))$, $0 \leq s \leq t \leq T$.

Probability measures on the sets of trajectories

We note $\mathcal{M}_1^+(\mathcal{T}^{\mathbb{Z}})$ (resp. $\mathcal{M}_1^+(\mathcal{T}^N)$) the set of probability measures on $(\mathcal{T}^{\mathbb{Z}}, \mathcal{F})$ (resp. $(\mathcal{T}^N, \mathcal{F}^N)$).

For $\mu \in \mathcal{M}_1^+(\mathcal{T}^{\mathbb{Z}})$, we denote its marginal distribution at time t by $\mu_t = \mu \circ \pi_t^{-1}$. Similarly, $\mu_{s,t}^N$ is its N -dimensional spatial, $t-s+1$ -dimensional time marginal $\mu \circ (\pi^N)^{-1} \circ \pi_{s,t}^{-1}$.

We denote the conditional probability distribution of μ , given $U_0^j = u_0^j$ (for all j), by μ_{u_0} . This is understood to be a probability measure over $\mathcal{B}(\mathcal{T}_{1,T}^{\mathbb{Z}}) = \mathcal{F}_{1,T}$.

We note $\mathcal{M}_{1,\mathcal{S}}^+(\mathcal{T}^{\mathbb{Z}})$ the set of stationary probability measures on $\mathcal{T}^{\mathbb{Z}}$. Given a random variable u with values in $\mathcal{T}^{\mathbb{Z}}$ governed by μ in $\mathcal{M}_{1,\mathcal{S}}^+(\mathcal{T}^{\mathbb{Z}})$, so is the random variable $\mathcal{S}u$, with \mathcal{S} the shift operator defined by (3.6) (equivalently $\mu \circ \mathcal{S}^{-1} = \mu$). With a slight abuse of notation, we define $\mathcal{M}_{1,\mathcal{S}}^+(\mathcal{T}^N)$ to be the set of all $\mu \in \mathcal{M}_1(\mathcal{T}^N)$ satisfying the following property. If (u^{-n}, \dots, u^n) are random variables governed by μ^N , then for all $|m| \leq n$, $(u^{m-n}, \dots, u^{m+n})$ has the same law as (u^{-n}, \dots, u^n) (recall that the indexing is taken modulo N), or equivalently that $\mu^N \circ \mathcal{S}^{-1} = \mu^N$ (remember (3.8)).

Remark 3.3. *Note that the stationarity discussed here is a spatial stationarity.*

Process-level empirical measure

We next introduce the following process-level empirical measure. Given an element $u = (u^{-n}, \dots, u^n)$ in \mathcal{T}^N we associate with it the measure, noted $\hat{\mu}_N(u^{-n}, \dots, u^n)$, in $\mathcal{M}_{1,\mathcal{S}}^+(\mathcal{T}^{\mathbb{Z}})$ defined by

$$\hat{\mu}_N : \mathcal{T}^N \rightarrow \mathcal{M}_{1,\mathcal{S}}^+(\mathcal{T}^{\mathbb{Z}}) \quad \text{such that} \quad d\hat{\mu}_N(u^{-n}, \dots, u^n)(y) = \frac{1}{N} \sum_{i=-n}^n \delta_{\mathcal{S}^i p_N(u)}(y). \quad (3.10)$$

\mathcal{S} is the shift operator defined in (3.6).

Remark 3.4. *This is a significant difference with previous work dealing with uncorrelated weights (e.g. [40]) where the N processes are coupled through the "usual" empirical measure $d\hat{\mu}_N(u^{-n}, \dots, u^n)(y) = \frac{1}{N} \sum_{i=-n}^n \delta_{u^i}(y)$ which is a measure on \mathcal{T} . In our case, because of the correlations and as shown in section 4.4 the processes are coupled through the process-level empirical measure (3.10) which is a probability measure on $\mathcal{T}^{\mathbb{Z}}$. This makes our analysis more biologically realistic, since we know that correlations between the synaptic weights do exist, but technically more involved.*

Topology on sets of measures

We next equip $\mathcal{M}_1^+(\mathcal{T}^{\mathbb{Z}})$ with the topology of weak convergence, as follows. For $\mu^N, \nu^N \in \mathcal{M}_1^+(\mathcal{T}^{\mathbb{Z}})$, we note the Wasserstein distance induced by the metric $k_f d_N(u, v) \wedge 1$,

$$D_N(\mu^N, \nu^N) = \inf_{\mathcal{L} \in \mathcal{J}} \{ \mathbb{E}^{\mathcal{L}}(k_f d_N(u, v) \wedge 1) \}, \quad (3.11)$$

where k_f is a positive constant defined at the start of section 3.1 and \mathcal{J} is the set of all measures in $\mathcal{M}_1^+(\mathcal{T}^{\mathbb{Z}} \times \mathcal{T}^{\mathbb{Z}})$ with N -dimensional marginals μ^N and ν^N .

Remark 3.5. *The use of k_f in (3.11) is technical and used to simplify the proof of proposition 4.5.*

For $\mu, \nu \in \mathcal{M}_1^+(\mathcal{T}^{\mathbb{Z}})$, we define

$$D(\mu, \nu) = 2 \sum_{n=0}^{\infty} \kappa_n D_N(\mu^N, \nu^N), \quad (3.12)$$

where $N = 2n + 1$. Here $\kappa_n = \max(\lambda_n, 2^{-N})$ and $\lambda_n = \sum_{k=-\infty}^{\infty} |\Lambda(k, n)|$. We note that this metric is well-defined because $D_N(\mu^N, \nu^N) \leq 1$ and $\sum_{n=0}^{\infty} \kappa_n < \infty$. It can be shown that $\mathcal{M}_1^+(\mathcal{T}^{\mathbb{Z}})$ equipped with this metric is Polish.

3.2.2 Coupled and uncoupled processes

We specify the initial conditions for (3.1) as N i.i.d. random variables $(U_0^j)_{j=-n, \dots, n}$. Let μ_I be the individual law on \mathbb{R} of U_0^j ; it follows that the joint law of the variables is $\mu_I^{\otimes N}$ on \mathbb{R}^N . We note P the law of the solution to one of the uncoupled equations (3.1) where we take $J_{ij}^N = 0$, $i, j = -n, \dots, n$. P is the law of the solution to the following stochastic difference equation:

$$U_t = \gamma U_{t-1} + B_{t-1}, \quad t = 1, \dots, T \quad (3.13)$$

the law of the initial condition being μ_I . This process can be characterized exactly, as follows.

Let $\Psi : \mathcal{T} \rightarrow \mathcal{T}$ be the following bicontinuous bijection. Writing $v = \Psi(u)$, we define

$$\begin{cases} v_0 &= \Psi_0(u) = u_0 \\ v_s &= \Psi_s(u) = u_s - \gamma u_{s-1} \quad s = 1, \dots, T. \end{cases} \quad (3.14)$$

The following proposition is evident from equations (3.13) and (3.14).

Proposition 3.6. *The law P of the solution to (3.13) writes*

$$P = (\mathcal{N}_T(0_T, \sigma^2 \text{Id}_T) \otimes \mu_I) \circ \Psi,$$

where 0_T is the T -dimensional vector of coordinates equal to 0 and Id_T is the T -dimensional identity matrix.

We later employ the convention that if $u = (u^{-n}, \dots, u^n) \in \mathcal{T}^N$ then $\Psi(u) = (\Psi(u^{-n}), \dots, \Psi(u^n))$. A similar convention applies if $u \in \mathcal{T}^Z$. We also use the notation $\Psi_{1,T}$ for the mapping $\mathcal{T} \rightarrow \mathcal{T}_{1,T}$ such that $\Psi_{1,T} = \pi_{1,T} \circ \Psi$.

Reintroducing the coupling between the neurons, we note $Q^N(J^N)$ the element of $\mathcal{M}_1^+(\mathcal{T}^N)$ which is the law of the solution to (3.1) conditioned on J^N . We let $Q^N = \mathbb{E}^J[Q^N(J^N)]$ be the law averaged with respect to the weights. The reason for this is as follows. We want to study the empirical measure $\hat{\mu}_N$ on path space. There is no reason for this to be a simple problem since for a fixed interaction J^N , the variables (U^{-n}, \dots, U^n) are not exchangeable. So we first study the law of $\hat{\mu}_N$ averaged over the interaction before we prove in section 7 some almost sure properties of this law. Q^N is a common construction in the physics of interacting particle systems and is known as the *annealed law* [38].

We may thus infer that

Lemma 3.7. $P^{\otimes N}$, Q^N and $\hat{\mu}_N^N$ (the N -dimensional marginal of $\hat{\mu}_N$) are in $\mathcal{M}_{1,S}^+(\mathcal{T}^N)$.

Since the application Ψ defined in (3.14) plays a central role in the sequel we introduce the following definition.

Definition 1. For each measure $\mu \in \mathcal{M}_1^+(\mathcal{T}^N)$ or $\mathcal{M}_{1,S}^+(\mathcal{T}^Z)$ we define $\underline{\mu}$ to be $\mu \circ \Psi^{-1}$.

In particular, note that

$$\underline{P} = \mathcal{N}_T(0_T, \sigma^2 \text{Id}_T) \otimes \mu_I. \quad (3.15)$$

Finally we introduce the image laws in terms of which the principal results of this paper are formulated.

Definition 2. Let Π^N (respectively R^N) be the image law of Q^N (respectively $P^{\otimes N}$) through the function $\hat{\mu}_N : \mathcal{T}^N \rightarrow \mathcal{M}_{1,S}^+(\mathcal{T}^Z)$ defined by (3.10).

The central result of this paper is in the next theorem.

Theorem 3.8. Π^N is governed by a large deviation principle (LDP) with a good rate function H (to be found in definition 5). That is, if F is a closed set in $\mathcal{M}_{1,S}^+(\mathcal{T}^Z)$, then

$$\overline{\lim}_{N \rightarrow \infty} N^{-1} \log \Pi^N(F) \leq - \inf_{\mu \in F} H(\mu). \quad (3.16)$$

Conversely, for all open sets O in $\mathcal{M}_{1,S}^+(\mathcal{T}^Z)$,

$$\underline{\lim}_{N \rightarrow \infty} N^{-1} \log \Pi^N(O) \geq - \inf_{\mu \in O} H(\mu). \quad (3.17)$$

Remark 3.9. We recall that the above LDP is also called a strong LDP.

Our proof of theorem 3.8 will occur in several steps. We prove in sections 5.1 and 5.3 that Π^N satisfies a weak LDP i.e. that it satisfies (3.16) when F is compact and (3.17) for all open O . We also prove in section 5.2 that $\{\Pi^N\}$ is exponentially tight, and we prove in section 5.4 that H is a good rate function. It directly follows from these results that Π^N satisfies a strong LDP with good rate function H [20]. Finally, in section 6 we prove that H has a unique minimum μ_e which $\hat{\mu}_N$ converges to weakly as $N \rightarrow \infty$. This minimum is a (stationary) Gaussian measure which we describe in detail in theorem 6.4.

4 The good rate function

In the sections to follow we will obtain an LDP for the process with correlations (Π^N) via the (simpler) process without correlations (R^N) . However to do this we require an expression for the Radon-Nikodym derivative of Π^N with respect to R^N , which is the main result of this section. The derivative will be expressed in terms of a function $\Gamma_{[N]} : \mathcal{M}_{1,S}^+(\mathcal{T}^N) \rightarrow \mathbb{R}$. We will firstly define $\Gamma_{[N]}(\mu)$, demonstrating that it may be expressed in terms of a Gaussian process $G_{[N]}^\mu$ (to be defined below), and then use this to determine the Radon-Nikodym derivative of Π^N with respect to R^N .

4.1 Gaussian processes

Given μ in $\mathcal{M}_{1,S}^+(\mathcal{T}^Z)$ we define a stationary Gaussian process G^μ . with values in $\mathcal{T}_{1,T}^Z$.

For all i the mean of $G_t^{\mu,i}$ is given by c_t^μ , where

$$c_t^\mu = \bar{J} \int_{\mathcal{T}^Z} f(u_{t-1}^i) d\mu(u), \quad t = 1, \dots, T, i \in Z, \quad (4.1)$$

the above integral being well-defined because of the definition of f and independent of i due to the stationarity of μ .

We now define the covariance of G^μ . We first define the following matrix-valued process.

Definition 3. Let $M^{\mu,k}$, $k \in Z$ be the $T \times T$ matrix defined by (for $s, t \in [1, T]$),

$$M_{st}^{\mu,k} = \int_{\mathcal{T}^Z} f(u_{s-1}^0) f(u_{t-1}^k) d\mu(u), \quad (4.2)$$

the above integral being well-defined because of the definition of f .

These matrixes satisfy

$$\dagger M^{\mu,k} = M^{\mu,-k}, \quad (4.3)$$

because of the stationarity of μ . Furthermore, they feature a spectral representation, i.e. there exists a $T \times T$ matrix-valued measure $\tilde{M}^\mu = (\tilde{M}^\mu)_{s,t=1,\dots,T}$ with the following properties. Each \tilde{M}_{st}^μ is a complex measure on $[-\pi, \pi[$ of finite total variation and such that

$$M^{\mu,k} = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{ik\theta} \tilde{M}^\mu(d\theta). \quad (4.4)$$

Relations (4.3) and (4.4) imply the following relations, for all Borelian sets $\mathcal{A} \subset [-\pi, \pi[$,

$$\tilde{M}^\mu(-\mathcal{A}) = \dagger \tilde{M}^\mu(\mathcal{A}) = \tilde{M}^\mu(\mathcal{A})^*, \quad (4.5)$$

where $*$ indicates complex conjugation. We may infer from this that \tilde{M}^μ is Hermitian-valued. The spectral representation means that for all vectors $W \in \mathbb{R}^T$, $\dagger W \tilde{M}^\mu(d\theta) W$ is a positive measure on $[-\pi, \pi[$.

The covariance between the Gaussian vectors $G^{\mu,i}$ and $G^{\mu,i+k}$ is defined to be

$$K^{\mu,k} = \sum_{l=-\infty}^{\infty} \Lambda(k,l) M^{\mu,l}. \quad (4.6)$$

We note that the above summation converges for all $k \in Z$ since the series $(\Lambda(k,l))_{k,l \in Z}$ is absolutely convergent and the elements of $M^{\mu,l}$ are bounded by 1 for all $l \in Z$.

It follows immediately from the definition that for $\mu \in \mathcal{M}_{1,S}^+(\mathcal{T}^{\mathbb{Z}})$ and $k \in \mathbb{Z}$ we have

$$\dagger K^{\mu,k} = K^{\mu,-k}. \quad (4.7)$$

This is necessary for the covariance function to be well-defined. The following proposition may be easily proved from the above definitions.

Proposition 4.1. *The sequence $(K^{\mu,k})_{k \in \mathbb{Z}}$ has spectral density \tilde{K}^μ given by*

$$\tilde{K}^\mu(\theta) = \frac{1}{2\pi} \int_{-\pi}^{\pi} \tilde{\Lambda}(\theta, -\varphi) \tilde{M}^\mu(d\varphi).$$

That is, \tilde{K}^μ is Hermitian positive and satisfies $\tilde{K}^\mu(-\theta) = \dagger \tilde{K}^\mu(\theta)$ and $K^{\mu,k} = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{ik\theta} \tilde{K}^\mu(\theta) d\theta$.

Proof. The proof essentially consists of demonstrating that the matrix function

$$\tilde{K}^\mu(\theta) = \sum_{k=-\infty}^{\infty} K^{\mu,k} e^{-ik\theta} \quad (4.8)$$

is well-defined on $[-\pi, \pi[$ and is equal to the expression in the statement of the proposition. Afterwards, we will prove that \tilde{K}^μ is positive.

From (4.6) we obtain that, for all $s, t \in [1 \cdots T]$,

$$|K_{st}^{\mu,k}| \leq \sum_{l=-\infty}^{\infty} |\Lambda(k, l)|. \quad (4.9)$$

This shows that, because by (3.4) the series $(\Lambda(k, l))_{k, l \in \mathbb{Z}}$ is absolutely convergent, $\tilde{K}^\mu(\theta)$ is well-defined on $[-\pi, \pi[$. The fact that $\tilde{K}^\mu(\theta)$ is Hermitian follows from (4.8) and (4.7).

Combining (4.4), (4.6) and (4.8) we write

$$\tilde{K}^\mu(\theta) = \frac{1}{2\pi} \int_{-\pi}^{\pi} \left(\sum_{m=-\infty}^{\infty} \sum_{k=-\infty}^{\infty} \Lambda(k, m) e^{-i(k\theta - m\varphi)} \right) \tilde{M}^\mu(d\varphi).$$

This can be rewritten in terms of the spectral density $\tilde{\Lambda}$ of Λ

$$\tilde{K}^\mu(\theta) = \frac{1}{2\pi} \int_{-\pi}^{\pi} \tilde{\Lambda}(\theta, -\varphi) \tilde{M}^\mu(d\varphi).$$

We note that $\tilde{K}^\mu(\theta)$ is positive, because for all vectors W of \mathbb{R}^T ,

$$\dagger W \tilde{K}^\mu(\theta) W = \frac{1}{2\pi} \int_{-\pi}^{\pi} \tilde{\Lambda}(\theta, -\varphi) \left(\dagger W \tilde{M}^\mu(d\varphi) W \right), \quad (4.10)$$

the spectral density $\tilde{\Lambda}$ is positive and the measure $\dagger W \tilde{M}^\mu(d\varphi) W$ is positive. The identity $\tilde{K}^\mu(-\theta) = \dagger \tilde{K}^\mu(\theta)$ follows from (4.7). \square

We may also define the N -dimensional Gaussian process $G_{[N]}^\mu$ with values in $\mathcal{T}_{1,T}^N$ as follows. The mean of $G_{[N]}^{\mu,i}$, $i = -n, \dots, n$ is given by (4.1) (or rather its finite dimensional analog) and the covariance between $G_{[N]}^{\mu,i}$ and $G_{[N]}^{\mu,i+k}$ is given by

$$K_{[N]}^{\mu,k} = \sum_{m=-n}^n \Lambda(k, m) M^{\mu,m}, \quad (4.11)$$

for $k = -n, \dots, n$. Equation (4.7) holds for $K_{[N]}^{\mu,k}$, $k = -n \cdots n$. This finite sequence has a Hermitian positive Discrete Fourier Transform denoted by $\tilde{K}_{[N]}^{\mu,l}$, for $l = -n \cdots n$.

4.2 Convergence of Gaussian Processes

The finite-dimensional system ‘converges’ to the infinite-dimensional system in the following sense. In what follows, and throughout the paper, we use the Frobenius norm on the $T \times T$ matrices. We write $\tilde{K}_{[N]}^\mu(\theta) = \sum_{k=-n}^n K_{[N]}^{\mu,k} \exp(-ik\theta)$. Note that for $|j| \leq n$, $\tilde{K}_{[N]}^\mu(2\pi j/N) = \tilde{K}_{[N]}^{\mu,j}$. The lemma below follows directly from the absolute convergence of $\sum_{j,k} |\Lambda(j,k)|$.

Lemma 4.2. *Fix $\mu \in \mathcal{M}_{1,S}^+(\mathcal{T}^{\mathbb{Z}})$. For all ε , there exists an N such that for all $M > N$ and all j such that $2|j| + 1 \leq M$, $\|K_{[M]}^{\mu,j} - K^{\mu,j}\| < \varepsilon$ and for all $\theta \in [-\pi, \pi[$, $\|\tilde{K}_{[M]}^\mu(\theta) - \tilde{K}^\mu(\theta)\| \leq \varepsilon$.*

Lemma 4.3. *The eigenvalues of $\tilde{K}_{[N]}^{\mu,l}$ and $\tilde{K}^\mu(\theta)$ are upperbounded by $\rho_K \stackrel{\text{def}}{=} T\Lambda^{\text{sum}}$, where Λ^{sum} is defined in (3.4).*

Proof. Let $W \in \mathbb{R}^T$. We find from proposition 4.1, and (3.4) that

$$\dagger W \tilde{K}^\mu(\theta) W \leq \frac{\Lambda^{\text{sum}}}{2\pi} \int_{-\pi}^{\pi} \dagger W \tilde{M}^\mu(d\varphi) W = \Lambda^{\text{sum}} \dagger W M^{\mu,0} W.$$

The eigenvalues of $M^{\mu,0}$ are all positive (since it is a correlation matrix), which means that each eigenvalue is upperbounded by the trace, which in turn is upperbounded by T . The proof in the finite dimensional case follows similarly. \square

We note $K_{[N]}^\mu$ the $(NT \times NT)$ covariance matrix of the sequence of Gaussian random variables $(G_{[N]}^{\mu,-n}, \dots, G_{[N]}^{\mu,n})$. Because of the properties of the matrixes $K_{[N]}^{\mu,k}$, $k = -n \dots n$, this is a symmetric block circulant matrix. It is also positive, being a covariance matrix.

We let $A_{[N]}^\mu = K_{[N]}^\mu(\sigma^2 \text{Id}_{NT} + K_{[N]}^\mu)^{-1}$. This is well-defined because $K_{[N]}^\mu$ is diagonalizable (being symmetric and real) and has positive eigenvalues (being a covariance matrix). It follows from lemma A.2 in appendix A that this is a symmetric block circulant matrix, with blocks $A_{[N]}^{\mu,k}$ ($k = -n, \dots, n$) such that

$$A_{[N]}^{\mu,-k} = \dagger A_{[N]}^{\mu,k}$$

and that the matrixes

$$\tilde{A}_{[N]}^{\mu,l} = \sum_{k=-n}^n A_{[N]}^{\mu,k} e^{-\frac{2\pi i k l}{N}} = \tilde{K}_{[N]}^{\mu,l} (\sigma^2 \text{Id}_T + \tilde{K}_{[N]}^{\mu,l})^{-1}. \quad (4.12)$$

are Hermitian positive.

In the limit $N \rightarrow \infty$ we may define

$$\tilde{A}^\mu(\theta) = \tilde{K}^\mu(\theta) (\sigma^2 \text{Id}_T + \tilde{K}^\mu(\theta))^{-1}. \quad (4.13)$$

The Fourier series of \tilde{A}^μ is absolutely convergent as a consequence of Wiener’s Theorem. We thus find that, for $l \in \mathbb{Z}$,

$$A^{\mu,l} = \frac{1}{2\pi} \int_{-\pi}^{\pi} \tilde{A}^\mu(\theta) e^{il\theta} d\theta = \lim_{N \rightarrow \infty} A_{[N]}^{\mu,l}, \quad (4.14)$$

and $\tilde{A}^\mu(\theta) = \sum_{l=-\infty}^{\infty} A^{\mu,l} e^{-il\theta}$. Let $\tilde{A}_{[N]}^\mu(\theta) = \sum_{k=-n}^n A_{[N]}^{\mu,k} \exp(-ik\theta)$ and note that for $|j| \leq n$, $\tilde{A}_{[N]}^\mu(2\pi j/N) = \tilde{A}_{[N]}^{\mu,j}$.

Lemma 4.4. *The map $B \rightarrow B(\sigma^2 \text{Id}_T + B)^{-1}$ is Lipschitz continuous over the set $\Delta = \{\tilde{K}_{[N]}^\mu(\theta), \tilde{K}^\mu(\theta) : \mu \in \mathcal{M}_{1,S}^+(\mathcal{T}^Z), N > 0, \theta \in [-\pi, \pi[]\}$.*

Proof. The proof is straightforward using the boundedness of the eigenvalues of the matrixes in Δ . \square

The following lemma is a consequence of lemmas 4.2 and 4.4.

Lemma 4.5. *Fix $\mu \in \mathcal{M}_{1,S}^+(\mathcal{T}^Z)$. For all ε , there exists an N such that for all $M > N$ and all $\theta \in [-\pi, \pi[$, $\|\tilde{A}_{[M]}^\mu(\theta) - \tilde{A}^\mu(\theta)\| \leq \varepsilon$.*

The above-defined matrices have the following ‘uniform convergence’ properties.

Proposition 4.6. *Fix $\nu \in \mathcal{M}_{1,S}^+(\mathcal{T}^Z)$. For all $\varepsilon > 0$, there exists an open neighbourhood $V_\varepsilon(\nu)$ such that for all $\mu \in V_\varepsilon(\nu)$, all $s, t \in [1, T]$ and all $\theta \in [-\pi, \pi[$,*

$$\left| \tilde{K}_{st}^\nu(\theta) - \tilde{K}_{st}^\mu(\theta) \right| \leq \varepsilon, \quad (4.15)$$

$$\left| \tilde{A}_{st}^\nu(\theta) - \tilde{A}_{st}^\mu(\theta) \right| \leq \varepsilon, \quad (4.16)$$

$$|c_s^\nu - c_s^\mu| \leq \varepsilon, \quad (4.17)$$

and for all $N > 0$, and for all k such that $|k| \leq n$,

$$\left| \tilde{K}_{[N],st}^{\nu,k} - \tilde{K}_{[N],st}^{\mu,k} \right| \leq \varepsilon, \quad (4.18)$$

and

$$\left| \tilde{A}_{[N],st}^{\nu,k} - \tilde{A}_{[N],st}^{\mu,k} \right| \leq \varepsilon. \quad (4.19)$$

Proof. The proof is found in appendix B. \square

Before we close this section we define a subset of $\mathcal{M}_{1,S}^+(\mathcal{T}^Z)$ which appears naturally, i.e. it is the subset of $\mathcal{M}_{1,S}^+(\mathcal{T}^Z)$ where the rate function (to be defined) is not infinite, see section 4.3.2 and lemma 5.2.

Definition 4. Let \mathcal{E}_2 be the subset of $\mathcal{M}_{1,S}^+(\mathcal{T}^Z)$ defined by

$$\mathcal{E}_2 = \{\mu \in \mathcal{M}_{1,S}^+(\mathcal{T}^Z) \mid \mathbb{E}^{\mu_{1,T}}[\|v^0\|^2] < \infty\},$$

where $v \in \mathcal{T}_{1,T}^Z$ and

$$\mathbb{E}^{\mu_{1,T}}[\|v^0\|^2] = \int_{\mathcal{T}_{1,T}^Z} \|\pi^1(v)\|^2 d\mu_{1,T}(v) = \int_{\mathcal{T}_{1,T}} \|v^0\|^2 \mu_{1,T}^1(dv^0).$$

For this set of measures, we may define the stationary process $(v^k)_{k \in \mathbb{Z}}$ in $\mathcal{T}_{1,T}^Z$, where $v_s^k = \Psi_s(u^k)$, $s = 1, \dots, T$. This has a finite mean $\mathbb{E}^{\mu_{1,T}}[v^0]$, noted \bar{v}^μ . It admits the following spectral density measure, noted \tilde{v}^μ , such that

$$\mathbb{E}^{\mu_{1,T}}[v^{0\dagger} v^k] = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{ik\theta} \tilde{v}^\mu(d\theta). \quad (4.20)$$

4.3 Definition of the functional Γ

In this section we define and study a functional $\Gamma_{[N]} = \Gamma_{[N],1} + \Gamma_{[N],2}$, which will be used to characterise the Radon-Nikodym derivative of Π^N with respect to R^N . Let $\mu \in \mathcal{M}_{1,S}^+(\mathcal{T}^Z)$, and let $(\mu^N)_{N \geq 1}$ be the N -dimensional marginals of μ (for $N = 2n + 1$ odd).

4.3.1 Γ_1

We define

$$\Gamma_{[N],1}(\mu) = -\frac{1}{2N} \log \left(\det \left(\text{Id}_{NT} + \frac{1}{\sigma^2} K_{[N]}^\mu \right) \right). \quad (4.21)$$

Because of the remarks after lemma 4.3 the spectrum of $K_{[N]}^\mu$ is positive, that of $\text{Id}_{NT} + \frac{1}{\sigma^2} K_{[N]}^\mu$ is strictly positive (in effect larger than 1) and the above expression has a sense. Moreover, $\Gamma_{[N],1}(\mu) \leq 0$.

We now define $\Gamma_1(\mu) = \lim_{N \rightarrow \infty} \Gamma_{[N],1}(\mu)$. The following lemma indicates that this is well-defined.

Lemma 4.7. *When N goes to infinity the limit of (4.21) is given by*

$$\Gamma_1(\mu) = -\frac{1}{4\pi} \int_{-\pi}^{\pi} \log \left(\det \left(\text{Id}_T + \frac{1}{\sigma^2} \tilde{K}^\mu(\theta) \right) \right) d\theta \quad (4.22)$$

for all $\mu \in \mathcal{M}_{1,S}^+(\mathcal{T}^Z)$.

Proof. Through lemma A.2 in appendix A, we have that

$$\Gamma_{[N],1}(\mu) = -\frac{1}{2N} \sum_{l=-n}^n \log \left(\det \left(\text{Id}_T + \frac{1}{\sigma^2} \tilde{K}_{[N]}^\mu \left(\frac{2\pi l}{N} \right) \right) \right), \quad (4.23)$$

where we recall that $\tilde{K}_{[N]}^\mu \left(\frac{2\pi l}{N} \right) = \tilde{K}_{[N]}^{\mu,l}$. Since, by lemma 4.2, $\tilde{K}_{[N]}^\mu(\theta)$ converges uniformly to $\tilde{K}^\mu(\theta)$, it is evident that the above expression converges to the desired result. \square

Proposition 4.8. $\Gamma_{[N],1}$ and Γ_1 are bounded below and continuous on $\mathcal{M}_{1,S}^+(\mathcal{T}^Z)$.

Proof. Applying lemma A.1 in the case of $Z = (G_{[N]}^{\mu,-n} - c^\mu, \dots, G_{[N]}^{\mu,n} - c^\mu)$, $a = 0$, $b = \sigma^{-2}$, we write

$$\Gamma_{[N],1}(\mu) = \frac{1}{N} \log \mathbb{E} \left[\exp \left(-\frac{1}{2\sigma^2} \sum_{k=-n}^n \|G_{[N]}^{\mu,k} - c^\mu\|^2 \right) \right].$$

Using Jensen's inequality we have

$$\Gamma_{[N],1}(\mu) \geq -\frac{1}{2N\sigma^2} \mathbb{E} \left[\sum_{k=-n}^n \|G_{[N]}^{\mu,k} - c^\mu\|^2 \right] = -\frac{1}{2\sigma^2} \mathbb{E} \left[\|G_{[N]}^{\mu,0} - c^\mu\|^2 \right].$$

By definition of $K_{[N]}^{\mu,0}$, the right-hand side is equal to $-\frac{1}{2\sigma^2} \text{Trace}(K_{[N]}^{\mu,0})$. From (4.11), we find that

$$\text{Trace}(K_{[N]}^{\mu,0}) = \sum_{m=-n}^n \Lambda(0, m) \text{Trace}(M^{\mu,m}).$$

It follows from equation (4.2) that $0 \leq \text{Trace}(M^{\mu,m}) \leq T$. Hence $\Gamma_{[N],1}(\mu) \geq -\beta_1$, where

$$\beta_1 = \frac{T\Lambda^{sum}}{2\sigma^2}. \quad (4.24)$$

It follows from lemma 4.7 that $-\beta_1$ is a lower bound for $\Gamma_1(\mu)$ as well.

The continuity of both $\Gamma_{[N],1}$ and Γ_1 follows from the expressions (4.21) and (4.22), continuity of the applications $\mu \rightarrow \tilde{K}_{[N]}^\mu$ and $\mu \rightarrow \tilde{K}^\mu$ (proposition 4.6) and the continuity of the determinant. \square

4.3.2 Γ_2

For $\mu \in \mathcal{M}_{1,S}^+(\mathcal{T}^Z)$ we define

$$\Gamma_{[N],2}(\mu) = \int_{\mathcal{T}_{1,T}^N} \phi^N(\mu, v) \underline{\mu}_{1,T}^N(dv) \quad (4.25)$$

where $\phi^N : \mathcal{M}_{1,S}^+(\mathcal{T}^Z) \times \mathcal{T}_{1,T}^N \rightarrow \mathbf{R}$ is defined by

$$\phi^N(\mu, v) = \frac{1}{2\sigma^2} \left(\frac{1}{N} \sum_{j,k=-n}^n \dagger(v^j - c^\mu) A_{[N]}^{\mu,k}(v^{k+j} - c^\mu) + \frac{2}{N} \sum_{j=-n}^n \langle c^\mu, v^j \rangle - \|c^\mu\|^2 \right). \quad (4.26)$$

$\Gamma_{[N],2}(\mu)$ is finite in the subset \mathcal{E}_2 of $\mathcal{M}_{1,S}^+(\mathcal{T}^Z)$ defined in definition 4. If $\mu \notin \mathcal{E}_2$, then we set $\Gamma_{[N],2}(\mu) = \infty$.

We define $\Gamma_2(\mu) = \lim_{N \rightarrow \infty} \Gamma_{[N],2}(\mu)$. The following proposition indicates that $\Gamma_2(\mu)$ is well-defined.

Proposition 4.9. *If the measure μ is in \mathcal{E}_2 , i.e. if $\mathbb{E}^{\mu_{1,T}}[\|v^0\|^2] < \infty$, then $\Gamma_2(\mu)$ is finite and writes*

$$\Gamma_2(\mu) = \frac{1}{2\sigma^2} \left(\frac{1}{2\pi} \int_{-\pi}^{\pi} \tilde{A}^\mu(-\theta) : \tilde{v}^\mu(d\theta) + \dagger c^\mu (\tilde{A}^\mu(0) - \text{Id}_T) c^\mu + 2\mathbb{E}^{\mu_{1,T}} \left[\dagger v^0 (\text{Id}_T - \tilde{A}^\mu(0)) c^\mu \right] \right).$$

The “:” symbol indicates the double contraction on the indexes. One also has

$$\Gamma_2(\mu) = \frac{1}{2\sigma^2} \left(\lim_{n \rightarrow \infty} \sum_{k=-n}^n \int_{\mathcal{T}_{1,T}^Z} \dagger(v^0 - c^\mu) A^{\mu,k}(v^k - c^\mu) d\underline{\mu}_{1,T}(v) + 2\mathbb{E}^{\mu_{1,T}}[\langle c^\mu, v^0 \rangle] - \|c^\mu\|^2 \right).$$

Proof. Using (4.20), (4.25) the stationarity of μ and the fact that $\sum_{k=-n}^n A_{[N]}^{\mu,k} = \tilde{A}_{[N]}^\mu(0)$, we have

$$\begin{aligned} \Gamma_{[N],2}(\mu) &= \frac{1}{4\pi\sigma^2} \int_{-\pi}^{\pi} \sum_{k=-n}^n \exp(ik\theta) A_{[N]}^{\mu,k} : \tilde{v}^\mu(d\theta) \\ &\quad + \frac{1}{\sigma^2} \int_{\mathcal{T}_{1,T}^N} \langle c^\mu, v^0 \rangle - \dagger c^\mu \tilde{A}_{[N]}^\mu(0) v^0 d\underline{\mu}_{1,T}^N(v) + \frac{1}{2\sigma^2} \dagger c^\mu \left(\text{Id}_T - \tilde{A}_{[N]}^\mu(0) \right) c^\mu. \end{aligned} \quad (4.27)$$

From the spectral representation of $A_{[N]}^\mu$ we find that

$$\begin{aligned} \Gamma_{[N],2}(\mu) &= \frac{1}{4\pi\sigma^2} \int_{-\pi}^{\pi} \tilde{A}_{[N]}^\mu(-\theta) : \tilde{v}^\mu(d\theta) \\ &\quad + \frac{1}{\sigma^2} \mathbb{E}^{\mu_{1,T}} \left[\dagger v^0 (\text{Id}_T - \tilde{A}_{[N]}^\mu(0)) c^\mu \right] + \frac{1}{2\sigma^2} \dagger c^\mu \left(\text{Id}_T - \tilde{A}_{[N]}^\mu(0) \right) c^\mu. \end{aligned} \quad (4.28)$$

Since (according to proposition 4.5) $\tilde{A}_{[N]}^\mu(\theta)$ converges uniformly to $\tilde{A}^\mu(\theta)$ as $N \rightarrow \infty$, it follows by dominated convergence that $\Gamma_{[N],2}(\mu)$ converges to the expression in the proposition.

The second expression for $\Gamma_2(\mu)$ follows analogously, although this time we make use of the fact that the partial sums of the Fourier Series of \tilde{A}^μ converge uniformly to \tilde{A}^μ (because the Fourier Series is absolutely convergent). \square

We next obtain more information about the eigenvalues of the matrices $\tilde{A}_{[N]}^{\mu,k} = \tilde{A}_{[N]}^\mu(\frac{2k\pi}{N})$ (where $k = -n, \dots, n$) and $\tilde{A}^\mu(\theta)$.

Lemma 4.10. *There exists $0 < \alpha < 1$, such that for all N and μ , the eigenvalues of $\tilde{A}_{[N]}^{\mu,k}$, $\tilde{A}^\mu(\theta)$ and $A_{[N]}^\mu$ are less than or equal to α .*

Proof. By lemma 4.3, the eigenvalues of $\tilde{K}^\mu(\theta)$ are positive and upperbounded by ρ_K . Since $\tilde{K}^\mu(\theta)$ and $(\sigma^2 \text{Id}_T + \tilde{K}^\mu(\theta))^{-1}$ are coaxial (because \tilde{K}^μ is Hermitian and therefore diagonalisable), we may take

$$\alpha = \frac{\rho_K}{\sigma^2 + \rho_K}.$$

This upperbound also holds for $\tilde{A}_{[N]}^{\mu,k}$, and for the eigenvalues of $A_{[N]}^\mu$, because of lemma A.2. \square

We wish to prove that $\Gamma_{[N],2}(\mu)$ is lower semicontinuous. A consequence of this will be that $\Gamma_{[N],2}(\mu)$ is measurable with respect to $\mathcal{B}(\mathcal{M}_{1,S}^+(\mathcal{T}^Z))$. In effect, we prove in appendix C that $\phi^N(\mu, v)$ defined by (4.26) satisfies

$$\phi^N(\mu, v) \geq -\beta_2$$

for some positive constant β_2 defined in equation (C.5) in appendix C.

We then have the following proposition.

Proposition 4.11. *$\Gamma_{[N],2}(\mu)$ is lower-semicontinuous.*

Proof. We define $\phi^{N,M}(\mu, v) = 1_{B_M}(v) (\phi^N(\mu, v) + \beta_2)$, where 1_{B_M} is the indicator of B_M and $v \in B_M$ if $N^{-1} \sum_{j=-n}^n \|v^j\|^2 \leq M$. We have just seen that $\phi^{N,M} \geq 0$. We also define

$$\Gamma_{[N],2}^M(\mu) = \int_{\mathcal{T}_{1,T}^N} \phi^{N,M}(\mu, v) \underline{\mu}_{1,T}^N(dv) - \beta_2.$$

Suppose that $\nu_n \rightarrow \mu$ with respect to the weak topology in $\mathcal{M}_{1,S}^+(\mathcal{T}^Z)$. Observe that

$$\begin{aligned} \left| \Gamma_{[N],2}^M(\mu) - \Gamma_{[N],2}^M(\nu_n) \right| &\leq \left| \int_{\mathcal{T}_{1,T}^N} \phi^{N,M}(\mu, v) \underline{\mu}_{1,T}^N(dv) - \int_{\mathcal{T}_{1,T}^N} \phi^{N,M}(\mu, v) \underline{\nu}_{n,1,T}^N(dv) \right| \\ &\quad + \left| \int_{\mathcal{T}_{1,T}^N} \phi^{N,M}(\mu, v) \underline{\nu}_{n,1,T}^N(dv) - \int_{\mathcal{T}_{1,T}^N} \phi^{N,M}(\nu_n, v) \underline{\nu}_{n,1,T}^N(dv) \right|. \end{aligned}$$

We may infer from the above expression that $\Gamma_{2,[N]}^M(\mu)$ is continuous (with respect to μ) for the following reasons. The first term on the right hand side converges to zero because $\phi^{N,M}$ is continuous and bounded (with respect to v). The second term converges to zero because $\phi^{N,M}(\mu, v)$ is a continuous function of μ , see proposition 4.6.

Since $\Gamma_{[N],2}^M(\mu)$ grows to $\Gamma_{[N],2}(\mu)$ as $M \rightarrow \infty$, we may conclude that $\Gamma_{[N],2}(\mu)$ is lower semicontinuous with respect to μ . \square

We define $\Gamma_{[N]}(\mu) = \Gamma_{[N],1}(\mu) + \Gamma_{[N],2}(\mu)$. We may conclude from propositions 4.8 and 4.11 that $\Gamma_{[N]}$ is measurable.

4.4 The Radon-Nikodym derivative

In this section we determine the Radon-Nikodym derivative of Π^N with respect to R^N . However in order for us to do this, we must first compute the Radon-Nikodym derivative of Q^N with respect to $P^{\otimes N}$. We do this in the next proposition.

Proposition 4.12. *The Radon-Nikodym derivative of Q^N with respect to $P^{\otimes N}$ is given by the following expression.*

$$\frac{dQ^N}{dP^{\otimes N}}(u^{-n}, \dots, u^n) = \mathbb{E} \left[\exp \left(\frac{1}{\sigma^2} \left(\sum_{j=-n}^n \langle \Psi_{1,T}(u^j), G^j \rangle - \frac{1}{2} \|G^j\|^2 \right) \right) \right], \quad (4.29)$$

the expectation being taken against the N T -dimensional Gaussian processes (G^i) , $i = -n, \dots, n$ given by

$$G_t^i = \sum_{j=-n}^n J_{ij}^N f(u_{t-1}^j), \quad t = 1, \dots, T, \quad (4.30)$$

and the function Ψ being defined by (3.14).

Proof. For fixed J^N , we let $R_{J^N} : \mathbb{R}^{N(T+1)} \rightarrow \mathbb{R}^{N(T+1)}$ be the mapping $u \rightarrow y$, i.e. $R_{J^N}(u^{-n}, \dots, u^n) = (y^{-n}, \dots, y^n)$, where for $j = -n, \dots, n$,

$$\begin{cases} y_0^j &= u_0^j \\ y_t^j &= u_t^j - \gamma u_{t-1}^j - G_t^j \quad t = 1, \dots, T. \end{cases}$$

The determinant of the Jacobian of R_{J^N} is 1 for the following reasons. Since $\frac{dy_s^j}{du_t^k} = 0$ if $t > s$, the determinant is $\prod_{s=0}^T D_s$, where D_s is the Jacobian of the map $(u_s^{-n}, \dots, u_s^n) \rightarrow (y_s^{-n}, \dots, y_s^n)$ induced by R_{J^N} . However D_s is evidently 1. Similar reasoning implies that R_{J^N} is a bijection.

It may be seen that the random vector $Y = R_{J^N}(U)$, U solution of (3.1), is such that $Y_0^j = U_0^j$ and $Y_t^j = B_{t-1}^j$ where $|j| \leq n$ and $t = 1, \dots, T$. Therefore

$$Y^j \simeq \mathcal{N}_T(0, \sigma^2 \text{Id}_T) \otimes \mu_I, \quad j = -n, \dots, n.$$

Since the determinant of the Jacobian of R_{J^N} is one, we obtain the law of $Q^N(J^N)$ by applying the inverse of R_{J^N} to the above distribution, i.e.

$$Q^N(J^N)(du) = (2\pi\sigma^2)^{-\frac{NT}{2}} \exp \left(-\frac{1}{2\sigma^2} \|R_{J^N}(u)\|^2 \right) \prod_{j=-n}^n \mu_I(du_0^j) \prod_{t=1}^T du_t^j.$$

Note that, exceptionally, $\|\cdot\|$ is the Euclidean norm in $\mathbb{R}^{N(T+1)}$ or \mathcal{T}^N .

Recalling that $P^{\otimes N} = Q^N(0)$, we therefore find that

$$\frac{dQ^N(J^N)}{dP^{\otimes N}}(u) = \exp \left(-\frac{1}{2\sigma^2} (\|R_{J^N}(u)\|^2 - \|R_0(u)\|^2) \right).$$

Taking the expectation of this with respect to J^N yields the result. \square

In fact, as stated in the proposition below, the Gaussian system $(G_s^i)_{i=-n, \dots, n, s=1, \dots, T}$ has the same law as the system $G_{[N]}^{\mu_N}$, as defined in (4.11) and afterwards.

Proposition 4.13. Fix $u \in \mathcal{T}^N$. The covariance of the Gaussian system (G_s^i) , where $i = -n, \dots, n$ and $s = 1, \dots, T$ writes $K_{[N]}^{\hat{\mu}_N(u)}$. For each i , the mean of G^i is $e^{\hat{\mu}_N(u)}$.

The proof of this proposition is an easy verification left to the reader. We obtain an alternative expression for the Radon-Nikodym derivative in (4.29) by applying lemma A.1 in appendix A. That is, we substitute $Z = (G^{-n}, \dots, G^n)$, $a = \frac{1}{\sigma^2}(v^{-n}, \dots, v^n)$, and $b = \frac{1}{\sigma^2}$ into the formula in lemma A.1. After noting proposition 4.13 we thus find that

Proposition 4.14. The Radon-Nikodym derivatives write as

$$\frac{dQ^N}{dP^{\otimes N}}(u^{-n}, \dots, u^n) = \exp(N\Gamma_{[N]}(\hat{\mu}_N(u^{-n}, \dots, u^n))),$$

$$\frac{d\Pi^N}{dR^N}(\mu) = \exp(N\Gamma_{[N]}(\mu)).$$

Here $\mu \in \mathcal{M}_{1,S}^+(\mathcal{T}^Z)$, $\Gamma_{[N]}(\mu) = \Gamma_{[N],1}(\mu) + \Gamma_{[N],2}(\mu)$ and the expressions for $\Gamma_{[N],1}$ and $\Gamma_{[N],2}$ have been defined in equations (4.21) and (4.25).

The second expression in the above proposition follows from the first one because $\Gamma_{[N]}$ is measurable.

Remark 4.15. Proposition 4.14 shows that the process solutions of (3.1) are coupled through the process-level empirical measure unlike in the case of independent weights where they are coupled through the usual empirical measure. As mentioned in the remark 3.4 this complicates significantly the mathematical analysis.

5 The large deviation principle

In this section we prove the principal result of this paper (Theorem 3.8), that the image laws Π^N satisfy an LDP with good rate function H (to be defined below). We do this by firstly establishing an LDP for the image law with uncoupled weights (R^N), see definition 2, and then use the Radon-Nikodym derivative of corollary 4.14 to establish the full LDP for Π^N . Therefore our first task is to write the LDP governing R^N .

Let μ, ν be probability measures over a Polish Space Ω equipped with its Borelian σ -algebra. The Kullback-Leibler divergence of μ relative to ν (also called the relative entropy) is

$$I^{(2)}(\mu, \nu) = \int_{\Omega} \log \left(\frac{d\mu}{d\nu} \right) d\mu \quad (5.1)$$

if μ is absolutely continuous with respect to ν , and $I^{(2)}(\mu, \nu) = \infty$ otherwise. It is a standard result that

$$I^{(2)}(\mu, \nu) = \sup_{f \in C_b(\Omega)} \{ \mathbb{E}^{\mu}[f] - \log \mathbb{E}^{\nu}[\exp(f)] \}, \quad (5.2)$$

the supremum being taken over all bounded continuous functions.

For $\mu \in M_{1,S}^+(\Omega^Z)$ and $\nu \in M_1^+(\Omega)$, the process-level entropy of μ with respect to ν^Z is defined to be

$$I^{(3)}(\mu, \nu^Z) = \lim_{N \rightarrow \infty} \frac{1}{N} I^{(2)}(\mu^N, \nu^{\otimes N}). \quad (5.3)$$

See [25, Lemma IX.2.4] for a proof that this (possibly infinite) limit always exists (the superadditivity of the sequence $N^{-1}I^{(2)}(\mu^N)$ follows from (5.2)).

Theorem 5.1. R^N is governed by a large deviation principle with good rate function [23, 2]

$$I^{(3)}(\mu, P^Z) = I^{(3)}(\mu_0, \mu_I^Z) + \int_{\mathbb{R}^\infty} I^{(3)}(\mu_{u_0}, P_{u_0}^Z) d\mu_0(u_0), \quad (5.4)$$

where $\mu_0 = \mu \circ \pi_0^{-1}$ is the time marginal of μ at time 0 and $\mu_{u_0} \in \mathcal{M}_{1,S}^+(\mathcal{T}_{1,T}^Z)$ is the conditional probability distribution of μ given u_0 in \mathbb{R}^Z . In addition, the set of measures $\{R^N\}$ is exponentially tight.

Proof. R^N satisfies an LDP with good rate function $I^{(3)}(\mu, P^Z)$ [25]. In turn, a sequence of probability measures (such as $\{R^N\}$) over a Polish Space satisfying a large deviations upper bound with a good rate function is exponentially tight [20].

It is an identity in [22] that

$$I^{(2)}(\mu^N, P^{\otimes N}) = I^{(2)}(\mu_0^N, \mu_I^{\otimes N}) + \int_{\mathbb{R}^N} I^{(2)}(\mu_{u_0}^N, P_{u_0}^{\otimes N}) d\mu_0^N(u_0). \quad (5.5)$$

It follows directly from the variational expression (5.2) that

$$I^{(2)}(\mu_{u_0}^{2N}, P_{u_0}^{\otimes 2N}) \geq 2I^{(2)}(\mu_{u_0}^N, P_{u_0}^{\otimes N}). \quad (5.6)$$

Note that although our convention throughout this paper is for N to be odd, the limit in (5.3) exists for any sequence of integers going to ∞ . We divide (5.5) by N and consider the subsequence of all N of the form $N = 2^k$ for $k \in \mathbb{Z}^+$. It follows from (5.6) that $N^{-1}I^{(2)}(\mu_{u_0}^N, P_{u_0}^{\otimes N})$ is strictly nondecreasing as $N = 2^k \rightarrow \infty$ (for all u_0), so that (5.4) follows by the monotone convergence theorem. \square

Because Ψ is bijective and bicontinuous, it may be easily shown that

$$I^{(2)}(\mu^N, P^{\otimes N}) = I^{(2)}(\underline{\mu}^N, \underline{P}^{\otimes N}) \quad (5.7)$$

$$I^{(3)}(\mu, P^Z) = I^{(3)}(\underline{\mu}, \underline{P}^Z). \quad (5.8)$$

Before we move to a statement of the LDP governing Π^N , we prove the following relationship between the set \mathcal{E}_2 (see definition 4) and the set of stationary measures which have a finite Kullback-Leibler divergence or process-level entropy with respect to P^Z .

Lemma 5.2.

$$\{\mu \in \mathcal{M}_{1,S}^+(\mathcal{T}^Z), I^{(3)}(\mu, P^Z) < \infty\} \subset \mathcal{E}_2.$$

See [27, Lemma 10] for a proof. We are now in a position to define what will be the rate function of the LDP governing Π^N .

Definition 5. Let H be the function $\mathcal{M}_{1,S}^+(\mathcal{T}^Z) \rightarrow \mathbb{R} \cup \{+\infty\}$ defined by

$$H(\mu) = \begin{cases} +\infty & \text{if } I^{(3)}(\mu, P^Z) = \infty \\ I^{(3)}(\mu, P^Z) - \Gamma(\mu) & \text{otherwise.} \end{cases}$$

Here $\Gamma(\mu) = \Gamma_1(\mu) + \Gamma_2(\mu)$ and the expressions for Γ_1 and Γ_2 have been defined in lemma 4.7 and proposition 4.9. Note that because of proposition 4.9 and lemma 5.2, whenever $I^{(3)}(\mu, P^Z)$ is finite, so is $\Gamma(\mu)$.

5.1 Lower bound on the open sets

We prove the second half of proposition 3.8.

Lemma 5.3. *For all open sets O , equation (3.17)*

$$\underline{\lim}_{N \rightarrow \infty} N^{-1} \log \Pi^N(O) \geq - \inf_{\mu \in O} H(\mu),$$

holds.

Proof. From the expression for the Radon-Nikodym derivative in proposition 4.14 we have

$$\Pi^N(O) = \int_O \exp(N\Gamma_{[N]}(\mu)) dR^N(\mu).$$

If $\mu \in O$ is such that $I^{(3)}(\mu, P^Z) = \infty$, then $H(\mu) = \infty$ and evidently (3.17) holds. We now prove (3.17) for all $\mu \in O$ such that $I^{(3)}(\mu, P^Z) < \infty$. Let $\varepsilon > 0$ and $Z_\varepsilon^N(\mu) \subset O$ be an open neighbourhood containing μ such that

$\inf_{\nu \in Z_\varepsilon^N(\mu)} \Gamma_{[N]}(\nu) \geq \Gamma_{[N]}(\mu) - \varepsilon$. Such $\{Z_\varepsilon^N(\mu)\}$ exist for all N because of the lower semi-continuity of $\Gamma_{[N]}(\mu)$ (see propositions 4.8 and 4.11). Then

$$\begin{aligned} \underline{\lim}_{N \rightarrow \infty} N^{-1} \log \Pi^N(O) &= \underline{\lim}_{N \rightarrow \infty} N^{-1} \log \int_O \exp(N\Gamma_{[N]}(\nu)) dR^N(\nu) \\ &\geq \underline{\lim}_{N \rightarrow \infty} N^{-1} \log \left(R^N(Z_\varepsilon^N(\mu)) \times \inf_{\nu \in Z_\varepsilon^N(\mu)} \exp(N\Gamma_{[N]}(\nu)) \right) \\ &\geq -I^{(3)}(\mu, P^Z) + \underline{\lim}_{N \rightarrow \infty} \inf_{\nu \in Z_\varepsilon^N(\mu)} \Gamma_{[N]}(\nu) \quad \text{because of theorem 5.1} \\ &\geq -I^{(3)}(\mu, P^Z) + \underline{\lim}_{N \rightarrow \infty} \Gamma_{[N]}(\mu) - \varepsilon \\ &= -I^{(3)}(\mu, P^Z) + \Gamma(\mu) - \varepsilon. \end{aligned}$$

The last equality follows from lemma 4.7 and proposition 4.9. Since ε is arbitrary, we may take the limit as $\varepsilon \rightarrow 0$ to obtain (3.17). Since (3.17) is true for all $\mu \in O$ the lemma is proved. \square

5.2 Exponential Tightness of Π^N

We begin with the following technical lemma, the proof of which can be found in appendix D.

Lemma 5.4. *There exist positive constants $c > 0$ and $a > 1$ such that, for all N ,*

$$\int_{\mathcal{T}^N} \exp(aN\phi^N(\hat{\mu}_N(u), \Psi_{1,T}(u))) P^{\otimes N}(du) \leq \exp(Nc),$$

where ϕ^N is defined in (4.26).

This lemma allows us to prove the exponential tightness:

Proposition 5.5. *The family $\{\Pi^N\}$ is exponentially tight.*

Proof. Let $B \in \mathcal{B}(\mathcal{M}_{1,S}^+(\mathcal{T}^Z))$. We have from proposition 4.14

$$\Pi^N(B) = \int_{(\hat{\mu}_N)^{-1}(B)} \exp N\Gamma_{[N]}(\hat{\mu}_N(u)) P^{\otimes N}(du).$$

Through Hölder's Inequality, we find that for any $a > 1$:

$$\Pi^N(B) \leq R^N(B)^{(1-\frac{1}{a})} \left(\int_{(\hat{\mu}_N)^{-1}(B)} \exp(aN\Gamma_{[N]}(\hat{\mu}_N(u))) P^{\otimes N}(du) \right)^{\frac{1}{a}}.$$

Now it may be observed that

$$\begin{aligned} & \int_{\mathcal{T}^N} \exp(aN\Gamma_{[N]}(\hat{\mu}_N(u))) P^{\otimes N}(du) \\ &= \int_{\mathcal{T}^N} \exp(aN\phi^N(\hat{\mu}_N(u), \Psi_{1,T}(u)) + aN\Gamma_{[N],1}(\hat{\mu}_N(u))) P^{\otimes N}(du). \end{aligned} \quad (5.9)$$

Since $\Gamma_1 \leq 0$, it follows from lemma 5.4 that

$$\Pi^N(B) \leq R^N(B)^{(1-\frac{1}{a})} \exp\left(\frac{Nc}{a}\right). \quad (5.10)$$

By the exponential tightness of $\{R^N\}$ (as stated in theorem 5.1), for each $L > 0$, there exists a compact set K_L such that $\overline{\lim}_{N \rightarrow \infty} N^{-1} \log(R^N(K_L^c)) \leq -L$. Thus if we choose compact K_{Π} such that $K_{\Pi,L} = K_{\frac{a}{a-1}(L+\frac{\epsilon}{a})}$, then for all $L > 0$, $\overline{\lim}_{N \rightarrow \infty} N^{-1} \log(\Pi^N(K_{\Pi,L}^c)) \leq -L$. \square

5.3 Upper Bound on the Compact Sets

In this section we obtain an upper bound on the compact sets, i.e. the first half of theorem 3.8 for F compact. Our method is to obtain an LDP for a simplified Gaussian system (with fixed A^ν and c^ν), and then prove that this converges to the required bound as $\nu \rightarrow \mu$.

5.3.1 An LDP for a Gaussian measure

We linearise Γ in the following manner. Fix $\nu \in \mathcal{M}_{1,S}^+(\mathcal{T}^Z)$ and assume for the moment that $\mu \in \mathcal{E}_2$. Let

$$\Gamma_{[N],2}^\nu(\mu) = \int_{\mathcal{T}_{1,T}^N} \phi_\infty^N(\nu, v) d\mu_{1,T}^N(v), \quad \text{where} \quad (5.11)$$

$\phi_\infty^N : \mathcal{M}_{1,S}^+(\mathcal{T}^Z) \times \mathcal{T}_{1,T}^N \rightarrow \mathbb{R}$ is defined by

$$\phi_\infty^N(\nu, v) = \frac{1}{2\sigma^2} \left(\frac{1}{N} \sum_{j,k=-n}^n \dagger(v^j - c^\nu) A^{\nu,k} (v^{k+j} - c^\nu) + \frac{2}{N} \sum_{j=-n}^n \langle c^\nu, v^j \rangle - \|c^\nu\|^2 \right). \quad (5.12)$$

Remark 5.6. Note the subtle difference with the definition of Φ^N in (4.26): we use $A^{\nu,k}$ instead of $A_{[N]}^{\nu,k}$. When turning to spectral representations of Φ_∞^N this will bring in the matrixes $\tilde{A}^{\nu,N,l}$, $l = -n \dots n$ defined at the beginning of section 5.3.2.

Let us also define

$$\Gamma_1^N(\nu) = -\frac{1}{2N} \log \det \left(\text{Id}_{NT} + \frac{1}{\sigma^2} K^{\nu,N} \right), \quad (5.13)$$

where $K^{\nu,N}$ is the $NT \times NT$ matrix with $T \times T$ blocks noted $K^{\nu,N,l}$. We define

$$\Gamma_{[N]}^\nu(\mu) = \Gamma_1^N(\nu) + \Gamma_{[N],2}^\nu(\mu), \quad \text{and} \quad (5.14)$$

$\Gamma_2^\nu(\mu) = \lim_{N \rightarrow \infty} \Gamma_{[N],2}^\nu(\mu)$. We find, using the first identity in proposition 4.9, that

$$\Gamma_2^\nu(\mu) = \frac{1}{2\sigma^2} \left(\frac{1}{2\pi} \int_{-\pi}^{\pi} \tilde{A}^\nu(-\theta) : \tilde{v}^\mu(d\theta) - 2 \dagger c^\nu \tilde{A}^\nu(0) \bar{v}^\mu + \dagger c^\nu \tilde{A}^\nu(0) c^\nu + 2 \langle c^\nu, \bar{v}^\mu \rangle - \|c^\nu\|^2 \right), \quad (5.15)$$

where $\bar{v}^\mu = \mathbb{E}^{\mu_{1,T}}[v^0]$, and \tilde{v}^μ is the spectral measure defined in (4.20). We recall that $:$ denotes double contraction on the indices.

Similarly to lemma 4.7, we find that

$$\lim_{N \rightarrow \infty} \Gamma_1^N(\nu) = -\frac{1}{4\pi} \int_{-\pi}^{\pi} \left(\log \det \left(\text{Id}_T + \frac{1}{\sigma^2} \tilde{K}^\nu(\theta) \right) \right) d\theta = \Gamma_1(\nu). \quad (5.16)$$

For $\mu \in \mathcal{E}_2$, we define $H^\nu(\mu) = I^{(3)}(\mu, P^{\mathbb{Z}}) - \Gamma^\nu(\mu)$; for $\mu \notin \mathcal{E}_2$, we define $\Gamma_2^\nu(\mu) = \Gamma^\nu(\mu) = \infty$ and $H^\nu(\mu) = \infty$.

Definition 6. Let $\underline{Q}^\nu \in \mathcal{M}_{1,S}^+(\mathcal{T}^{\mathbb{Z}})$ with N -dimensional marginals $\underline{Q}^{\nu,N}$ given by

$$\underline{Q}^{\nu,N}(B) = \int_B \exp \left(N \Gamma_{[N]}^\nu(\hat{\mu}_N(v)) \right) \underline{P}^{\otimes N}(dv), \quad (5.17)$$

where $B \in \mathcal{B}(\mathcal{T}^N)$. This defines a law $Q^\nu \in \mathcal{M}_{1,S}^+(\mathcal{T}^{\mathbb{Z}})$ according to the correspondence in definition 1.

We have the following lemma.

Lemma 5.7. $\underline{Q}_{1,T}^\nu$ is a stationary Gaussian process of mean c^ν . Its N -dimensional spatial, T -dimensional temporal, marginals $\underline{Q}_{1,T}^{\nu,N}$ are in $\mathcal{M}_{1,S}^+(\mathcal{T}_{1,T}^N)$ and have covariance $\sigma^2 \text{Id}_{NT} + K^{\nu,N}$. The spectral density of $\underline{Q}_{1,T}^\nu$ is $\sigma^2 \text{Id}_T + \tilde{K}^\nu(\theta)$ and in addition,

$$\underline{Q}_0^\nu = \mu_I^{\mathbb{Z}}. \quad (5.18)$$

Proof. In effect we find that

$$\begin{aligned} \underline{Q}_{1,T}^{\nu,N}(B_N) &= \left(\det \left(\text{Id}_{NT} + \frac{1}{\sigma^2} K^{\nu,N} \right) \right)^{-\frac{1}{2}} \times \\ &\int_{B_N} \exp \frac{1}{2\sigma^2} \left(\sum_{j,k=-n}^n {}^t(v^j - c^\nu) A^{\nu,k} (v^{k+j} - c^\nu) + \right. \\ &\quad \left. 2 \sum_{j=-n}^n \langle c^\nu, v^j \rangle - N \|c^\nu\|^2 \right) \underline{P}_{1,T}^{\otimes N}(dv), \end{aligned} \quad (5.19)$$

for each Borelian $B_N \in \mathcal{B}(\mathcal{T}_{1,T}^N)$. We note $c^{\nu,N}$ the NT -dimensional vector obtained by concatenating N times the vector c^ν . We also have that

$$\frac{1}{\sigma^2} (\text{Id}_{NT} - A^{\nu,N}) = (\sigma^2 \text{Id}_{NT} + K^{\nu,N})^{-1}.$$

Thus, through proposition 3.6, we find that

$$\begin{aligned} \underline{Q}_{1,T}^{\nu,N}(B_N) &= (2\pi)^{-\frac{NT}{2}} \left(\det \left(\frac{1}{\sigma^2} (\text{Id}_{NT} - A^{\nu,N}) \right)^{-1} \right)^{-\frac{1}{2}} \\ &\int_{B_N} \exp -\frac{1}{2\sigma^2} \dagger (v - c^{\nu,N}) (\text{Id}_{NT} - A^{\nu,N}) (v - c^{\nu,N}) \prod_{j=-n}^n \prod_{t=1}^T dv_t^j. \end{aligned} \quad (5.20)$$

It is seen that $\underline{Q}_{1,T}^{\nu,N}$ is an NT -dimensional Gaussian measure with mean $c^{\nu,N}$, inverse covariance matrix $\frac{1}{\sigma^2}(\text{Id}_{NT} - A^{\nu,N})$, and covariance matrix $\sigma^2 \text{Id}_{NT} + K^{\nu,N}$. Hence $\underline{Q}_{1,T}^{\nu,N}$ is in $\mathcal{M}_{1,S}^+(\mathcal{T}_{1,T}^N)$, and

$$\underline{Q}^{\nu,N} = \underline{Q}_{1,T}^{\nu,N} \otimes \mu_I^N \quad (5.21)$$

is in $\mathcal{M}_{1,S}^+(\mathcal{T}^N)$. It follows also that the spectral density of $\underline{Q}_{1,T}^{\nu}$ is $\sigma^2 \text{Id}_T + \tilde{K}^{\nu}(\theta)$. \square

We may thus define the measure \underline{Q}^{ν} of a stationary process over the variables $\{v_s^j\}_{j \in \mathbb{Z}, s=0, \dots, T}$, with N -dimensional marginals given by (5.20) and (5.21).

Definition 7. Let $\underline{\Pi}^{\nu,N}$ be the image law of $\underline{Q}^{\nu,N}$ under $\hat{\mu}_N$, i.e. for $B \in \mathcal{B}(\mathcal{M}_{1,S}^+(\mathcal{T}^{\mathbb{Z}}))$,

$$\underline{\Pi}^{\nu,N}(B) = \underline{Q}^{\nu,N}(\hat{\mu}_N \in B).$$

The point is that it can be shown that the image $\underline{\Pi}^{\nu,N}$ of the measure $\underline{Q}^{\nu,N}$ satisfies a strong LDP (see next lemma) and that this LDP can be transferred to Π^N , see proposition 5.12. We begin with the following lemma which is a generalization of the result in [23].

Lemma 5.8. *The image law $\underline{\Pi}^{\nu,N}$ satisfies a strong LDP (in the manner of Theorem 3.8) with good rate function*

$$\underline{H}^{\nu}(\underline{\mu}) = I^{(3)}(\underline{\mu}, \underline{P}^{\mathbb{Z}}) - \Gamma^{\nu}(\underline{\mu}). \quad (5.22)$$

Proof. We have found an LDP for a Gaussian process in [27]. Since Q^{ν} may be separated in the manner of (5.21), we may use the expression in Theorem 5.1 to obtain the result. \square

For $B \in \mathcal{B}(\mathcal{M}_{1,S}^+(\mathcal{T}^{\mathbb{Z}}))$, we define the image law

$$\Pi^{\nu,N}(B) = Q^{\nu,N}(\hat{\mu}_N \in B) = \underline{Q}^{\nu,N}(\hat{\mu}_N \in B).$$

It follows from the contraction principle that if we write $H^{\nu}(\mu) := \underline{H}^{\nu}(\underline{\mu})$, then

Corollary 5.9. *The image law $\Pi^{\nu,N}$ satisfies a strong LDP with good rate function*

$$H^{\nu}(\mu) = I^{(3)}(\mu, P^{\mathbb{Z}}) - \Gamma^{\nu}(\mu). \quad (5.23)$$

5.3.2 An upper bound for Π^N over compact sets

In this section we derive an upper bound for Π^N over compact sets using the LDP of the previous section. Before we do this, we require two lemmas governing the ‘distance’ between Γ^{ν} and Γ . Let $\tilde{K}^{\mu,N}$ be the DFT of $(K^{\mu,j})_{j=-n}^n$, and similarly $\tilde{A}^{\mu,N}$ is the DFT of $(A^{\mu,j})_{j=-n}^n$. We define

$$C_N^{\nu} = \sup_{M \geq N, (2|l|+1) \leq M} \{\|\tilde{A}_{[M]}^{\nu,l} - \tilde{A}^{\nu,M,l}\|, \|\tilde{K}_{[M]}^{\nu,l} - \tilde{K}^{\nu,M,l}\|\}. \quad (5.24)$$

Lemma 5.10. *For all $\nu \in \mathcal{M}_{1,S}^+(\mathcal{T}^{\mathbb{Z}})$, C_N^{ν} is finite and*

$$C_N^{\nu} \rightarrow 0 \text{ as } N \rightarrow \infty.$$

Proof. We recall from proposition 4.6 that $\tilde{K}_{[M],st}^{\nu}(\theta)$ converges uniformly (in θ) to $\tilde{K}_{st}^{\nu}(\theta)$. The same holds for $\tilde{K}_{st}^{\nu,M,l}$, because this represents the partial summation of an absolutely converging Fourier Series. That is, for fixed $\theta = 2\pi l_M/M$, $\tilde{K}_{st}^{\nu,M,l_M} \rightarrow \tilde{K}_{st}^{\nu}(\theta)$ as $M \rightarrow \infty$. The result then follows from the equivalence of matrix norms. The proof for \tilde{A}^{ν} is analogous. \square

The second lemma, the proof of which can be found in appendix E, goes as follows.

Lemma 5.11. *There exists a constant C_0 such that for all ν in $\mathcal{M}_{1,S}^+(\mathcal{T}^Z)$, all $\varepsilon > 0$ and all $\mu \in V_\varepsilon(\nu) \cap \mathcal{E}_2$,*

$$\left| \Gamma_{[N]}(\mu) - \Gamma_{[N]}^\nu(\mu) \right| \leq C_0(C_N^\nu + \varepsilon)(1 + \mathbb{E}^{\underline{\mu}, T}[\|v^0\|^2]).$$

Here $V_\varepsilon(\nu)$ is the open neighbourhood defined in proposition 4.6, and $\underline{\mu}$ is given in definition 1.

We are now ready to begin the proof of the upper bound on compact sets for which we follow the ideas in [34]

Proposition 5.12. *Let \mathcal{K} be a compact subset of $\overline{\mathcal{M}}_{1,S}(\mathcal{T}^Z)$. Then*

$$\overline{\lim}_{N \rightarrow \infty} N^{-1} \log(\Pi^N(\mathcal{K})) \leq -\inf_{\mathcal{K}} H.$$

Proof. Fix $\varepsilon > 0$. Let $V_\varepsilon(\nu)$ be the open neighbourhood of ν defined in proposition 4.6, and let $\bar{V}_\varepsilon(\nu)$ be its closure. Since \mathcal{K} is compact and $\{V_\varepsilon(\nu)\}_{\nu \in \mathcal{K}}$ is an open cover, there exists an r and $\{\nu_i\}_{i=1}^r$ such that $\mathcal{K} \subset \bigcup_{i=1}^r V_\varepsilon(\nu_i)$. We find that

$$\overline{\lim}_{N \rightarrow \infty} N^{-1} \log \left(\Pi^N \left(\bigcup_{i=1}^r V_\varepsilon(\nu_i) \cap \mathcal{K} \right) \right) \leq \sup_{1 \leq i \leq r} \overline{\lim}_{N \rightarrow \infty} N^{-1} \log(\Pi^N(\bar{V}_\varepsilon(\nu_i) \cap \mathcal{K})).$$

It follows from the fact that $\hat{\mu}_N \in \mathcal{E}_2$, proposition 4.14 and lemma 5.11 that

$$\begin{aligned} \Pi^N(\bar{V}_\varepsilon(\nu_i) \cap \mathcal{K}) &\leq \int_{\hat{\mu}_N(u) \in \bar{V}_\varepsilon(\nu_i) \cap \mathcal{K}} \exp \left(N \Gamma_{(N)}^{\nu_i}(\hat{\mu}_N(u)) + \right. \\ &\quad \left. N C_0(\varepsilon + C_N^{\nu_i}) \left(1 + \frac{1}{N} \sum_{j=-n}^n \|\Psi_{1,T}(u^j)\|^2 \right) \right) P_{1,T}^{\otimes N}(du). \end{aligned} \quad (5.25)$$

From the definition of $Q^{\nu,N}$ in (5.17) and Hölder's Inequality, for p, q such that $\frac{1}{p} + \frac{1}{q} = 1$, we have

$$\Pi^N(\bar{V}_\varepsilon(\nu_i) \cap \mathcal{K}) \leq (Q^{\nu_i, N}(\hat{\mu}_N(u) \in \bar{V}_\varepsilon(\nu_i) \cap \mathcal{K}))^{\frac{1}{p}} D^{\frac{1}{q}}, \quad (5.26)$$

where

$$\begin{aligned} D &= \int_{\hat{\mu}_N(u) \in \bar{V}_\varepsilon(\nu_i) \cap \mathcal{K}} \exp \left(q N C_0(\varepsilon + C_N^{\nu_i}) \left(1 + \frac{1}{N} \sum_{j=-n}^n \|\Psi_{1,T}(u^j)\|^2 \right) \right) Q_{1,T}^{\nu_i, N}(du) \\ &= \exp q N C_0(\varepsilon + C_N^{\nu_i}) \times \int_{\underline{\mu}^N(v) \in \Psi(\bar{V}_\varepsilon(\nu_i) \cap \mathcal{K})} \exp \left(q C_0(\varepsilon + C_N^{\nu_i}) \left(\sum_{j=-n}^n \|v^j\|^2 \right) \right) \underline{Q}_{1,T}^{\nu_i, N}(dv). \end{aligned}$$

We note from lemma 4.3 that the eigenvalues of the covariance of $\underline{Q}_{1,T}^{\nu_i, N}$ are upperbounded by $\sigma^2 + \rho_K$. Thus for this integral to converge it is sufficient that

$$q C_0(\varepsilon + C_N^{\nu_i}) \leq \frac{1}{2(\sigma^2 + \rho_K)}. \quad (5.27)$$

This condition will always be satisfied for sufficiently small ε and sufficiently large N (since $C_N^{\nu_i} \rightarrow 0$ as $N \rightarrow \infty$ by lemma 5.10). Considering (5.26), by corollary 5.9,

$$\overline{\lim}_{N \rightarrow \infty} N^{-1} \log(Q^{\nu_i, N}(\hat{\mu}_N(u) \in \bar{V}_\varepsilon(\nu_i) \cap \mathcal{K})) \leq - \inf_{\mu \in \bar{V}_\varepsilon(\nu_i) \cap \mathcal{K}} H^{\nu_i}(\mu), \quad (5.28)$$

We next find an upper bound for the integral appearing in the definition of the quantity D . We apply lemma A.1 in appendix A to find

$$\int_{\underline{\mu}^N(v) \in \Psi(\bar{V}_\varepsilon(\nu_i) \cap \mathcal{K})} \exp qC_0(\varepsilon + C_N^{\nu_i}) \left(\sum_{j=-n}^n \|v^j\|^2 \right) Q_{1,T}^{\nu_i, N}(dv) \leq \\ (\det((1 - 2qC_0(\varepsilon + C_N^{\nu_i})\sigma^2)\text{Id}_{NT} - 2qC_0(\varepsilon + C_N^{\nu_i})K^{\nu_i, N}))^{-\frac{1}{2}} \times \\ \exp(2C_0^2q^2((\varepsilon + C_N^{\nu_i})^2)^\dagger(1_{NT}c^{\nu_i})B((\text{Id}_T \otimes 1_N)c^{\nu_i}) + NqC_0(\varepsilon + C_N^{\nu_i})\|c^{\nu_i}\|^2)$$

where $\text{Id}_T \otimes 1_N$ is the $NT \times T$ block matrix with each block Id_T and

$$B = (\sigma^2\text{Id}_{NT} + K^{\nu_i, N})((1 - 2C_0q(\varepsilon + C_N^{\nu_i})\sigma^2)\text{Id}_{NT} - 2C_0q(\varepsilon + C_N^{\nu_i})K^{\nu_i, N})^{-1}$$

is a symmetric block circulant matrix. We note B^k , $k = -n, \dots, n$ its $T \times T$ blocks. We have

$$^\dagger((\text{Id}_T \otimes 1_N)c^{\nu_i})B((\text{Id}_T \otimes 1_N)c^{\nu_i}) = N^\dagger c^{\nu_i} \left(\sum_{k=-n}^n B^k \right) c^{\nu_i} = N^\dagger c^{\nu_i} \tilde{B}^0 c^{\nu_i},$$

where \tilde{B}^0 is the 0th component of the spectral representation of the sequence $(B^k)_{k=-n, \dots, n}$. Let v_m be the largest eigenvalue of B . Since (by lemma A.2) the eigenvalues of \tilde{B}^0 are a subset of the eigenvalues of B , we have

$$^\dagger((\text{Id}_T \otimes 1_N)c^{\nu_i})B((\text{Id}_T \otimes 1_N)c^{\nu_i}) \leq Nv_m\|c^{\nu_i}\|^2.$$

From the definition of B and through lemma 4.3 we have $v_m \leq \frac{\sigma^2 + \rho_K}{1 - 2C_0q(\varepsilon + C_N^{\nu_i})(\sigma^2 + \rho_K)}$. Hence we have, since $\|c^{\nu_i}\|^2 \leq T\bar{J}^2$

$$\exp(2C_0^2(q^2(\varepsilon + C_N^{\nu_i})^2)^\dagger((\text{Id}_T \otimes 1_N)c^{\nu_i})B((\text{Id}_T \otimes 1_N)c^{\nu_i})) \leq \\ \exp\left(NT \times \frac{2C_0^2q^2(\varepsilon + C_N^{\nu_i})^2(\sigma^2 + \rho_K)\bar{J}^2}{1 - 2C_0q(\varepsilon + C_N^{\nu_i})(\sigma^2 + \rho_K)}\right).$$

Since the determinant is the product of the eigenvalues, we similarly find that

$$(\det((1 - 2C_0q(\varepsilon + C_N^{\nu_i})\sigma^2)\text{Id}_{NT} - 2C_0q(\varepsilon + C_N^{\nu_i})K^{\nu_i, N}))^{-\frac{1}{2}} \leq \\ (1 - 2C_0q(\varepsilon + C_N^{\nu_i})(\sigma^2 + \rho_K))^{-\frac{NT}{2}}.$$

Upon collecting the above inequalities, and noting that $\|c^\nu\|^2 \leq T\bar{J}^2$, we find that

$$D \leq \exp(Ns_N^{\nu_i}(q, \varepsilon)) \tag{5.29}$$

where

$$s_N^{\nu_i}(q, \varepsilon) = T \left(-\frac{1}{2} \log(1 - 2C_0q(\varepsilon + C_N^{\nu_i})(\sigma^2 + \rho_K)) \right. \\ \left. + \frac{2C_0^2q^2(\varepsilon + C_N^{\nu_i})^2(\sigma^2 + \rho_K)\bar{J}^2}{1 - 2C_0q(\varepsilon + C_N^{\nu_i})(\sigma^2 + \rho_K)} + qC_0(\varepsilon + C_N^{\nu_i}) \left(\frac{1}{T} + \bar{J}^2 \right) \right).$$

We let $s(q, \varepsilon) = \overline{\lim}_{N \rightarrow \infty} s_N^{\nu_i}(q, \varepsilon)$, and find through lemma 5.10 that

$$s(q, \varepsilon) = T \left(-\frac{1}{2} \log(1 - 2C_0 q \varepsilon (\sigma^2 + \rho_K)) + \frac{2C_0^2 q^2 \varepsilon^2 (\sigma^2 + \rho_K) \bar{J}^2}{1 - 2C_0 q \varepsilon (\sigma^2 + \rho_K)} + q C_0 \varepsilon \left(\frac{1}{T} + \bar{J}^2 \right) \right).$$

Notice that $s(q, \varepsilon)$ is independent of ν_i and that $s(q, \varepsilon) \rightarrow 0$ as $\varepsilon \rightarrow 0$. Using (5.26), (5.28) and (5.29) we thus find that

$$\overline{\lim}_{N \rightarrow \infty} N^{-1} \log(\Pi^N(\mathcal{K})) \leq \sup_{1 \leq i \leq r} -\frac{1}{p} \inf_{\mu \in \mathcal{K} \cap \bar{V}_\varepsilon(\nu_i)} H^{\nu_i}(\mu) - \frac{1}{q} s(q, \varepsilon).$$

Recall that $H^\nu(\mu) = \infty$ for all $\mu \notin \mathcal{E}_2$. Thus if $\mathcal{K} \cap \mathcal{E}_2 = \emptyset$, we may infer that $\overline{\lim}_{N \rightarrow \infty} N^{-1} \log(\Pi^N(\mathcal{K})) = -\infty$ and the proposition is evident. Thus we may assume without loss of generality that $\inf_{\mu \in \mathcal{K}} H^{\nu_i}(\mu) = \inf_{\mu \in \mathcal{K} \cap \mathcal{E}_2} H^{\nu_i}(\mu)$. Furthermore it follows from proposition 5.13 (below) that there exists a constant C_I such that for all $\mu \in \bar{V}_\varepsilon(\nu_i) \cap \mathcal{E}_2$,

$$H^{\nu_i}(\mu) \geq I^{(3)}(\mu, P^Z) - \Gamma(\mu) - C_I \varepsilon (1 + I^{(3)}(\mu, P^Z)).$$

We thus find that

$$\overline{\lim}_{N \rightarrow \infty} N^{-1} \log(\Pi^N(\mathcal{K})) \leq -\frac{1}{p} \inf_{\mathcal{K} \cap \mathcal{E}_2} \left(I^{(3)}(\mu, P^Z) (1 - C_I \varepsilon) - \Gamma(\mu) \right) - \frac{s(q, \varepsilon)}{q} + \frac{\varepsilon}{p} C_I.$$

We take $\varepsilon \rightarrow 0$ and find, through the use of lemma 5.14, that

$$\overline{\lim}_{N \rightarrow \infty} N^{-1} \log(\Pi^N(\mathcal{K})) \leq -\frac{1}{p} \inf_{\mathcal{K}} \left(I^{(3)}(\mu, P^Z) - \Gamma(\mu) \right).$$

The proof may thus be completed by taking $p \rightarrow 1$. \square

Proposition 5.13. *There exists a positive constant C_I such that, for all ν in $\mathcal{M}_{1,S}^+(\mathcal{T}^Z) \cap \mathcal{E}_2$, all $\varepsilon > 0$ and all $\mu \in \bar{V}_\varepsilon(\nu) \cap \mathcal{E}_2$ (where $\bar{V}_\varepsilon(\nu)$ is the neighbourhood defined in proposition 4.6),*

$$|\Gamma^\nu(\mu) - \Gamma^\mu(\mu)| \leq C_I \varepsilon \left(1 + I^{(3)}(\mu, P^Z) \right). \quad (5.30)$$

The proof is very similar to that of lemma 5.11 and we leave it to the reader. We end this section with lemma 5.14 whose proof can be found in appendix D

Lemma 5.14. *There exist constants $a > 1$ and $c > 0$ such that for all $\mu \in \mathcal{M}_{1,S}^+(\mathcal{T}^Z) \cap \mathcal{E}_2$,*

$$\Gamma(\mu) \leq \frac{(I^{(3)}(\mu, P^Z) + c)}{a}.$$

5.4 End of the proof of theorem 3.8

Lemma 5.15. *$H(\mu)$ is lower-semi-continuous.*

The proof is very similar to that in [27]. Because $\{\Pi^N\}$ is exponentially tight and satisfies the weak LDP with rate function $H(\mu)$, the following corollary is immediate [21, Lemma 2.1.5].

Corollary 5.16. *$H(\mu)$ is a good rate function, i.e. the sets $\{\mu : H(\mu) \leq \delta\}$ are compact for all $\delta \in \mathbb{R}^+$, and it satisfies the first condition of theorem 3.8.*

This allows us to complete the proof of theorem 3.8:

Proof. By combining lemmas 5.15 and 5.3, proposition 5.5, and corollary 5.16, we complete the proof of theorem 3.8. \square

6 Characterization of the unique minimum of the rate function

We prove that there exists a unique minimum μ_e of the rate function. and provide explicit equations for μ_e which would facilitate its numerical simulation. We start with the following lemma.

Lemma 6.1. *For $\mu, \nu \in \mathcal{M}_{1,S}^+(\mathcal{T}^{\mathbb{Z}})$, $H^\nu(\mu) = 0$ if and only if $\mu = Q^\nu$.*

Proof. This is a straightforward consequence of [27, Theorem 1] and Theorem 5.1. \square

Proposition 6.2. *There is a unique distribution $\mu_e \in \mathcal{M}_{1,S}^+(\mathcal{T}^{\mathbb{Z}})$ which minimises H . This distribution satisfies $H(\mu_e) = 0$.*

Proof. By the previous lemma, it suffices to prove that there is a unique μ_e such that

$$Q^{\mu_e} = \mu_e. \quad (6.1)$$

We define the mapping $L : \mathcal{M}_{1,S}^+(\mathcal{T}^{\mathbb{Z}}) \rightarrow \mathcal{M}_{1,S}^+(\mathcal{T}^{\mathbb{Z}})$ by

$$\mu \rightarrow L(\mu) = Q^\mu.$$

It follows from (5.18) that

$$Q_0^\mu = \mu_I^{\mathbb{Z}}, \quad (6.2)$$

which is independent of μ .

It may be inferred from the definitions in Section 4.1 that the marginal of $L(\mu) = Q^\mu$ over $\mathcal{F}_{0,t}$ only depends upon the marginal of μ over $\mathcal{F}_{0,t-1}$, $t \geq 1$. This follows from the fact that $Q_{1,t}^\mu$ (which determines $Q_{0,t}^\mu$) is completely determined by the means $\{c_s^\mu; s = 1, \dots, t\}$ and covariances $\{K_{uv}^{\mu,j}; j \in \mathbb{Z}, u, v \in [1, t]\}$. In turn, it may be observed from (4.1) and (4.6) that these variables are determined by $\mu_{0,t-1}$. Thus for any $\mu, \nu \in \mathcal{M}_{1,S}^+(\mathcal{T}^{\mathbb{Z}})$ and $t \in [1, T]$, if

$$\mu_{0,t-1} = \nu_{0,t-1},$$

then

$$L(\mu)_{0,t} = L(\nu)_{0,t}.$$

It follows from repeated application of the above identity that for any ν satisfying $\nu_0 = \mu_I^{\mathbb{Z}}$,

$$L^T(\nu)_{0,T} = L(L^T(\nu))_{0,T}. \quad (6.3)$$

Defining

$$\mu_e = L^T(\nu), \quad (6.4)$$

it follows from (6.3) that μ_e satisfies (6.1).

Conversely if $\mu = L(\mu)$ for some μ , then we have that $\mu = L^2(\nu)$ for any ν such that $\nu_{0,T-2} = \mu_{0,T-2}$. Continuing this reasoning, we find that $\mu = L^T(\nu)$ for any ν such that $\nu_0 = \mu_0$. But by (6.2), since $Q^\mu = \mu$, we have $\mu_0 = \mu_I^{\mathbb{Z}}$. But we have just seen that any μ satisfying $\mu = L^T(\nu)$, where $\nu_0 = \mu_I^{\mathbb{Z}}$, is uniquely defined by (6.4), which means that $\mu = \mu_e$. \square

We may use the proof of proposition 6.2 to characterize the unique measure μ_e such that $\mu_e = Q^{\mu_e}$ in terms of its image $\underline{\mu}_e$. This characterization allows one to directly numerically calculate μ_e . We characterize $\underline{\mu}_e$ recursively (in time), by providing a method of determining $\underline{\mu}_{e,0,t}$ in terms of $\underline{\mu}_{e,0,t-1}$. However we must firstly outline explicitly the bijective correspondence

between $\mu_{e0,t}$ and $\underline{\mu}_{e0,t}$, as follows. For $v \in \mathcal{T}$, we write $\Psi^{-1}(v) = (\Psi^{-1}(v)_0, \dots, \Psi^{-1}(v)_T)$. We recall from (3.14) that $\Psi^{-1}(v)_0 = v_0$. The coordinate $\Psi^{-1}(v)_t$ is the affine function of v_s , $s = 0 \dots t$ obtained from equations (3.14)

$$\Psi^{-1}(v)_t = \sum_{i=0}^t \gamma^i v_{t-i}, \quad t = 0, \dots, T.$$

Let $K_{(t-1,s-1)}^{\mu_e,l}$ be the $(t-1) \times (s-1)$ submatrix of $K^{\mu_e,l}$ composed of the rows from times 1 to $(t-1)$ and the columns from times 1 to $(s-1)$, and

$$c_{(t-1)}^{\mu_e} = \dagger(c_1^{\mu_e}, \dots, c_{t-1}^{\mu_e}).$$

Let the measures $\underline{\mu}_{e0,t}^1$ and $\underline{\mu}_{e,t,s}^{(0,l)}$ be given by

$$\begin{aligned} \underline{\mu}_{e0,t}^1(dv) &= \mu_I(dv_0) \otimes \mathcal{N}_t(c_{(t)}^{\mu_e}, \sigma^2 \text{Id}_t + K_{(t,t)}^{\mu_e,0}) dv_1 \dots dv_{t-1}. \\ \underline{\mu}_{e(t,s)}^{(0,l)}(dv^0 dv^l) &= \mu_I(dv_0^0) \otimes \mu_I(dv_0^l) \otimes \mathcal{N}_{t+s}((c_{(t)}^{\mu_e}, c_{(s)}^{\mu_e}), \sigma^2 \text{Id}_{t+s} + K_{(t,s)}^{\mu_e,(0,l)}) dv_1^0 \dots dv_t^0 dv_1^l \dots dv_s^l, \end{aligned}$$

where

$$K_{(t,s)}^{\mu_e,(0,l)} = \begin{bmatrix} K_{(t,t)}^{\mu_e,0} & K_{(t,s)}^{\mu_e,l} \\ \dagger K_{(t,s)}^{\mu_e,l} & K_{(s,s)}^{\mu_e,0} \end{bmatrix}.$$

The lemma below is evident from the definitions above.

Lemma 6.3. *For any $t \in [1, T]$, the variables $\{c_s^{\mu_e}, K_{rs}^{\mu_e,j} : 1 \leq r, s \leq t, j \in \mathbb{Z} \leq\}$ are necessary and sufficient to completely characterise the measures $\{\underline{\mu}_{e0,t}^1, \underline{\mu}_{e(0,t)}^{(0,l)} : l \in \mathbb{Z}\}$. In turn, the measures $\{\underline{\mu}_{e0,t}^1, \underline{\mu}_{e(0,t)}^{(0,l)} : l \in \mathbb{Z}\}$ are necessary and sufficient to characterise $\underline{\mu}_{e0,t}$.*

The inductive method for calculating $\underline{\mu}_e$ is outlined in the theorem below.

Theorem 6.4. *We may characterise $\underline{\mu}_e$ inductively as follows. Initially $\underline{\mu}_{e0} = \mu_I^{\mathbb{Z}}$. Given that we have a complete characterisation of*

$$\left\{ \underline{\mu}_{e(0,t-1)}^{(0,l)}, \underline{\mu}_{e0,t-1}^1 : l \in \mathbb{Z} \right\},$$

we may characterise

$$\left\{ \underline{\mu}_{e(0,t)}^{(0,l)}, \underline{\mu}_{e0,t}^1 : l \in \mathbb{Z} \right\}$$

according to the following identities. For $s \in [1, t]$,

$$c_s^{\mu_e} = \bar{J} \int_{\mathbb{R}^t} (f(\Psi^{-1}(v)_{s-1})) \underline{\mu}_{e0,s-1}^1(dv).$$

For $1 \leq r, s \leq t$, $K_{rs}^{\mu_e,k} = \sum_{l=-\infty}^{\infty} \Lambda(k, l) M_{rs}^{\mu_e,l}$. Here, for $p = \max(r-1, s-1)$,

$$M_{rs}^{\mu_e,0} = \int_{\mathbb{R}^{p+1}} (f(\Psi^{-1}(v)_{r-1})) \times (f(\Psi^{-1}(v)_{s-1})) \underline{\mu}_{e0,p}^1(dv),$$

and for $l \neq 0$

$$M_{rs}^{\mu_e,l} = \int_{\mathbb{R}^r \times \mathbb{R}^s} (f(\Psi^{-1}(v^0)_{r-1})) \times (f(\Psi^{-1}(v^l)_{s-1})) \underline{\mu}_{e(r-1,s-1)}^{(0,l)}(dv^0 dv^l).$$

Of course the measure μ_e may be determined from $\underline{\mu}_e$ since $\mu_e = \underline{\mu}_e \circ \Psi$.

7 Some important consequences of theorem 3.8

We state some important consequences of our results including some which are valid J almost surely (quenched results). We recall that $Q^N(J^N)$ is the conditional law of N neurons for given J^N .

Theorem 7.1. Π^N converges weakly to δ_{μ_e} , i.e., for all $\Phi \in \mathcal{C}_b(\mathcal{M}_{1,S}^+(\mathcal{T}^{\mathbb{Z}}))$,

$$\lim_{N \rightarrow \infty} \int_{\mathcal{T}^N} \Phi(\hat{\mu}_N(u)) Q^N(du) = \Phi(\mu_e).$$

Similarly,

$$\lim_{N \rightarrow \infty} \int_{\mathcal{T}^N} \Phi(\hat{\mu}_N(u)) Q^N(J^N)(du) = \Phi(\mu_e) \quad J \text{ almost surely}$$

Proof. The proof of the first result follows directly from the existence of an LDP for the measure Π^N , see theorem 3.8, and is a straightforward adaptation of the one in [39, Theorem 2.5.1]. The proof of the second result uses the same method, making use of theorem 7.2 below. \square

We can in fact obtain the following quenched convergence analogue of (3.16).

Theorem 7.2. For each closed set F of $\mathcal{M}_{1,S}^+(\mathcal{T}^{\mathbb{Z}})$ and for almost all J

$$\overline{\lim}_{N \rightarrow \infty} \frac{1}{N} \log [Q^N(J^N)(\hat{\mu}_N \in F)] \leq - \inf_{\mu \in F} H(\mu).$$

Proof. The proof is a combination of Tchebyshev's inequality and the Borel-Cantelli lemma and is a straightforward adaptation of the one in [39, Theorem 2.5.4, Corollary 2.5.6]. \square

We define $\check{Q}^N(J^N) = \frac{1}{N} \sum_{j=-n}^n Q^N(J^N) \circ S^{-j}$, where we recall the shift operator S defined by (3.8). Clearly $\check{Q}^N(J^N)$ is in $\mathcal{M}_{1,S}^+(\mathcal{T}^{\mathbb{Z}})$.

Corollary 7.3. Fix M and let $N > M$. For almost every J and all $h \in \mathcal{C}_b(\mathcal{T}^M)$,

$$\begin{aligned} \lim_{N \rightarrow \infty} \int_{\mathcal{T}^M} h(u) \check{Q}^{N,M}(J^N)(du) &= \int_{\mathcal{T}^M} h(u) \mu_e^M(du). \\ \lim_{N \rightarrow \infty} \int_{\mathcal{T}^M} h(u) Q^{N,M}(du) &= \int_{\mathcal{T}^M} h(u) \mu_e^M(du). \end{aligned}$$

That is, the M^{th} marginals $\check{Q}^{N,M}(J^N)$ and $Q^{N,M}$ converge weakly to μ_e^M as $N \rightarrow \infty$.

Proof. It is sufficient to apply theorem 7.1 in the case where Φ in $\mathcal{C}_b(\mathcal{M}_{1,S}^+(\mathcal{T}^{\mathbb{Z}}))$ is defined by

$$\Phi(\mu) = \int_{\mathcal{T}^M} h d\mu^M$$

and to use the fact that $Q^N, \check{Q}^N(J) \in \mathcal{M}_{1,S}^+(\mathcal{T}^{\mathbb{Z}})$ (lemma 3.7 and above remark). \square

We now prove the following ergodic-type theorem. We may represent the ambient probability space by \mathfrak{W} , where $\omega \in \mathfrak{W}$ is such that $\omega = (J_{ij}, B_t^j, u_0^j)$, where $i, j \in \mathbb{Z}$ and $0 \leq t \leq T-1$, recall (3.1). We denote the probability measure governing ω by \mathfrak{P} . Let $u^{(N)}(\omega) \in \mathcal{T}^N$ be defined by (3.1). As an aside, we may then understand $Q^N(J^N)$ to be the conditional law of \mathfrak{P} on $u^{(N)}(\omega)$, for given J^N .

Theorem 7.4. Fix $M > 0$ and let $h \in C_b(\mathcal{T}^M)$. For $u^{(N)}(\omega) \in \mathcal{T}^N$ (where $N > M$) and $|j| \leq n$. Then \mathfrak{P} almost surely,

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{j=-n}^n h\left(\pi_M(\mathcal{S}^j u^{(N)}(\omega))\right) = \int_{\mathcal{T}^M} h(u) d\mu_e^M(u), \quad (7.1)$$

where $\mathcal{S}^j u^{(N)}(\omega)$ is defined in (3.8). Hence $\hat{\mu}_N(u^{(N)}(\omega))$ converges \mathfrak{P} -almost-surely to μ_e .

Proof. Our proof is an adaptation of [39]. We may suppose without loss of generality that $\int_{\mathcal{T}^M} h(u) d\mu_e^M(u) = 0$. For $p > 1$ let

$$F_p = \left\{ \mu \in \mathcal{M}_{1,S}^+(\mathcal{T}^Z) \mid \left| \int_{\mathcal{T}^M} h(u) \mu^M(du) \right| \geq \frac{1}{p} \right\}.$$

Since $\mu_e \notin F_p$, but it is the unique zero of H , it follows that $\inf_{F_p} H = m > 0$. Thus by theorem 3.8 there exists an N_0 , such that for all $N > N_0$,

$$Q^N(\hat{\mu}_N \in F_p) \leq \exp(-mN).$$

However

$$\mathfrak{P}\left(\omega \mid \hat{\mu}_N(u^{(N)}(\omega)) \in F_p\right) = Q^N(u \mid \hat{\mu}_N(u) \in F_p).$$

Thus

$$\sum_{N=1}^{\infty} \mathfrak{P}\left(\omega \mid \hat{\mu}_N(u^{(N)}(\omega)) \in F_p\right) < \infty.$$

We may thus conclude from the Borel-Cantelli Lemma that \mathfrak{P} almost surely, for every $\omega \in \mathfrak{W}$, there exists N_p such that for all $N \geq N_p$,

$$\left| \frac{1}{N} \sum_{j=-n}^n h\left(\pi_M \mathcal{S}^j u^{(N)}(\omega)\right) \right| \leq \frac{1}{p}.$$

This yields (7.1) because p is arbitrary. The convergence of $\hat{\mu}_N(u^{(N)}(\omega))$ is a direct consequence of (7.1), since this means that each of the M^{th} marginals converge. \square

8 Possible extensions

Our results hold true if we assume that equation (3.1) is replaced by the more general equation

$$U_t^j = \sum_{k=1}^l \gamma_k U_{t-k}^j + \sum_{i=-n}^n J_{ji}^N f(U_{t-1}^i) + \theta_j + B_{t-1}^j, \quad j = -n, \dots, n \quad t = l, \dots, T,$$

where l is a positive integer strictly less than T (in practice much smaller) and the θ_j s are i.i.d. random variables independent of the synaptic weights J and the random processes B^j . They can be thought of as external stimuli imposed on the neurons. This equation accounts for a more complicated "intrinsic" dynamics of the neurons, i.e. when they are uncoupled. The parameters γ_k , $k = 1 \dots l$ must satisfy some conditions to ensure stability of the uncoupled dynamics.

This result can be straightforwardly extended to the case when the noise is correlated but stationary Gaussian, that is $\text{cov}(B_s^j, B_t^k)$ is some function of s, t and $(k - j)$. It can also be

easily extended to the case that the initial distribution is correlated but mixing, using the Large Deviation Principle in [12].

The hypothesis that the synaptic weights are Gaussian is somewhat unrealistic from the biological viewpoint. In his PhD thesis [39], Moynot has obtained some preliminary results in the case of uncorrelated weights. We think that this is also a promising avenue.

Moynot again, in his thesis, has extended the uncorrelated weights case, to include two populations with different (Gaussian) statistics for each population. This is also an important practical problem in neuroscience. Extending Moynot's result to the correlated case is probably a low hanging fruit.

Last but not least, the solutions of the equations for the mean and covariance operator of the measure minimizing the rate function derived in section 6 and their numerical simulation are very much worth investigating and their predictions confronted to biological measurements.

9 Conclusion

In recent years there has been a lot of effort to mathematically justify neural-field models, through some sort of asymptotic analysis of finite-size neural networks. Many, if not most, of these models assume / prove some sort of thermodynamic limit, whereby if one isolates a particular population of neurons in a localised area of space, they are found to fire increasingly asynchronously as the number in the population asymptotes to infinity.³ Indeed this was the result of Moynot and Samuelides. However our results imply that there are system-wide correlations between the neurons, even in the asymptotic limit. The key reason why we do not have propagation of chaos is that the Radon-Nikodym derivative $\frac{dQ^N}{dP^N}$ of the average laws in proposition 4.12 cannot be tensored into N i.i.d. processes; whereas the simpler assumptions on the weight function Λ in Moynot and Samuelides allow the Radon-Nikodym derivative to be tensored. A very important implication of our result is that the mean-field behaviour is insufficient to characterise the behaviour of a population. Our limit process μ_e is system-wide and ergodic. Our work challenges the assumption held by some that one cannot have a 'concise' macroscopic description of a neural network without an assumption of asynchronicity at the local population level.

It would be of interest to compare our LDP with other analyses of the rate of convergence of neural networks to their limits as the size asymptotes to infinity. This includes the system-size expansion of Bressloff [5], the path-integral formulation of Buice and Cowan [7] and the systematic expansion of the moments by (amongst others) [32, 24, 8].

Acknowledgements

Many thanks to Bruno Cessac whose suggestion to look at process-level empirical measures and entropies has been very useful and whose insights into the physical interpretations of our results have been very stimulating.

This work was partially supported by the European Union Seventh Framework Programme (FP7/2007- 2013) under grant agreement no. 269921 (BrainScaleS), no. 318723 (MathemacS), and by the ERC advanced grant NerVi no. 227747.

A Two useful lemmas

The following lemma from Gaussian calculus [39, 41] which we recall for completeness is used several times throughout the paper.

³We noted in the introduction that this is termed propagation of chaos by some.

Lemma A.1. *Let Z be a Gaussian vector of \mathbb{R}^p with mean c and covariance matrix K . If $a \in \mathbb{R}^p$ and $b \in \mathbb{R}$ is such that for all eigenvalues α of K the relation $\alpha b > -1$ holds, we have*

$$\mathbb{E} \left[\exp \left({}^t a Z - \frac{b}{2} \|Z\|^2 \right) \right] = \frac{1}{\sqrt{\det(\text{Id}_p + bK)}} \times \exp \left({}^t a c - \frac{b}{2} \|c\|^2 + \frac{1}{2} {}^t (a - bc) K (\text{Id}_p + bK)^{-1} (a - bc) \right)$$

Block-circulant matrices may be diagonalised using DFT's as follows.

Lemma A.2. *Let B be a symmetric block-circulant matrix with the (j, k) $T \times T$ block given by $(B^{(j-k) \bmod N})$, $j, k = -n, \dots, n$. Let $W^{(N)}$ be the $N \times N$ Unitary matrix with elements $W_{jk}^{(N)} = \frac{1}{\sqrt{N}} \exp(\frac{2\pi i j k}{N})$, $j, k = -n, \dots, n$. Then B may be 'block'-diagonalised in the follow manner (where \otimes is the Kronecker Product and $*$ the complex conjugate),*

$$B = (W^{(N)} \otimes \text{Id}_T) \text{diag} \left(\tilde{B}^{-n}, \dots, \tilde{B}^n \right) (W^{(N)} \otimes \text{Id}_T)^*.$$

Here \tilde{B}^j is a $T \times T$ Hermitian matrix and is the DFT defined in (D.1). We observe also that λ is an eigenvalue of B if and only if λ is an eigenvalue of \tilde{B}^k for some k .

B Proof of proposition 4.6

We first recall proposition 4.6:

Proposition

Fix $\nu \in \mathcal{M}_{1,S}^+(\mathcal{T}^Z)$. For all $\varepsilon > 0$, there exists an open neighbourhood $V_\varepsilon(\nu)$ such that for all $\mu \in V_\varepsilon(\nu)$, all $s, t \in [1, T]$ and all $\theta \in [-\pi, \pi[$,

$$\left| \tilde{K}_{st}^\nu(\theta) - \tilde{K}_{st}^\mu(\theta) \right| \leq \varepsilon,$$

$$\left| \tilde{A}_{st}^\nu(\theta) - \tilde{A}_{st}^\mu(\theta) \right| \leq \varepsilon,$$

$$|c_s^\nu - c_s^\mu| \leq \varepsilon,$$

and for all $N > 0$, and for all k such that $|k| \leq n$,

$$\left| \tilde{K}_{[N],st}^{\nu,k} - \tilde{K}_{[N],st}^{\mu,k} \right| \leq \varepsilon,$$

and

$$\left| \tilde{A}_{[N],st}^{\nu,k} - \tilde{A}_{[N],st}^{\mu,k} \right| \leq \varepsilon.$$

Proof. Let μ be in $\mathcal{M}_{1,S}^+(\mathcal{T}^Z)$ and $\theta \in [-\pi, \pi[$. We have

$$\tilde{K}_{st}^\mu(\theta) - \tilde{K}_{st}^\nu(\theta) = \sum_{k=-\infty}^{\infty} (K_{st}^{\mu,k} - K_{st}^{\nu,k}) e^{-ik\theta}.$$

Using (4.6) we have

$$\tilde{K}_{st}^\mu(\theta) - \tilde{K}_{st}^\nu(\theta) = \sum_{k,l=-\infty}^{\infty} \Lambda(k,l) (M_{st}^{\mu,l} - M_{st}^{\nu,l}) e^{-ik\theta},$$

hence

$$\left| \tilde{K}_{st}^\mu(\theta) - \tilde{K}_{st}^\nu(\theta) \right| \leq \sum_{k,l=-\infty}^{\infty} |\Lambda(k,l)| \inf_{\mathcal{L}^{2L}} \int_{\mathcal{T}^L \times \mathcal{T}^L} |f(u_{s-1}^0) f(u_{t-1}^l) - f(v_{s-1}^0) f(v_{t-1}^l)| \mathcal{L}^{2L}(du, dv),$$

where $L = 2|l| + 1$ and \mathcal{L}^{2L} has marginals μ^L and ν^L . Since $|f(u_{s-1}^0) f(u_{t-1}^l) - f(v_{s-1}^0) f(v_{t-1}^l)| \leq 2(k_f d_L(\pi_L u, \pi_L v) \wedge 1)$, where k_f is the Lipschitz constant of the function f , we find (through (3.12)) that

$$\left| \tilde{K}_{st}^\mu(\theta) - \tilde{K}_{st}^\nu(\theta) \right| \leq 2D(\mu, \nu).$$

Thus for (4.15) to be satisfied, it suffices for us to stipulate that $V_\varepsilon(\nu)$ is a ball of radius less than $\frac{\varepsilon}{2}$ (with respect to the distance metric in (3.12)). Similar reasoning dictates that (4.18) is satisfied too.

However in light of lemma 4.4, it is evident that we may take the radius of $V_\varepsilon(\nu)$ to be sufficiently small that (4.15), (4.18) and (4.19) are satisfied. In fact (4.16) is also satisfied, as it may be obtained by taking the limit as $N \rightarrow \infty$ of (4.19). Since c^μ is determined by the one-dimensional spatial marginal of μ , it follows from the definition of the metric in (3.12) that we may take the radius of $V_\varepsilon(\nu)$ to be sufficiently small that (4.17) is satisfied too. \square

C Existence of a lower bound for $\Phi^N(\mu, \nu)$

In order to prove that $\phi^N(\mu, \nu)$ defined in (4.25) possesses a lower bound, we use a spectral representation and let $\tilde{w}^j = \tilde{v}^j$ for all j , except that $\tilde{w}^0 = \tilde{v}^0 - Nc^\mu$. We may then write

$$\phi^N(\mu, \nu) = \frac{1}{2N^2\sigma^2} \sum_{l=-n}^n \dagger \tilde{w}^{l,*} \tilde{A}_{[N]}^{\mu,-l} \tilde{w}^l + \frac{1}{N\sigma^2} \langle c^\mu, \tilde{w}^0 \rangle + \frac{1}{2\sigma^2} \|c^\mu\|^2. \quad (\text{C.1})$$

Thus in order that the integrand possesses a lower bound, it suffices to prove, since the matrixes $\tilde{A}_{[N]}^{\mu,l}$ are Hermitian positive, that there exists a lower bound for

$$\frac{1}{N^2} \dagger \tilde{w}^0 \tilde{A}_{[N]}^{\mu,0} \tilde{w}^0 + \frac{2}{N} \langle \tilde{w}^0, c^\mu \rangle, \quad (\text{C.2})$$

We have made use of the fact that \tilde{w}^0 and $\tilde{A}_{[N]}^{\mu,0}$ are real (since they are each a sum of real variables). Let $\tilde{K}_{[N]}^{\mu,0} = O_{[N]}^\mu D_{[N]}^\mu \dagger O_{[N]}^\mu$, where $D_{[N]}^\mu$ is diagonal and $O_{[N]}^\mu$ is orthonormal. We define $X = \dagger O_{[N]}^\mu \tilde{w}^0$, so that (C.2) is equal to

$$\frac{1}{N^2} \dagger X D_{[N]}^\mu (\sigma^2 \text{Id}_T + D_{[N]}^\mu)^{-1} X + \frac{2}{N} \sum_{t=1}^T \langle \dagger O_{[N],t}^\mu, c^\mu \rangle X_t, \quad (\text{C.3})$$

where $O_{[N],t}^\mu$ is the t -th column vector of $O_{[N]}^\mu$. In order that (C.3) is bounded below, we require that the coefficient of X converges to zero when $D_{[N]}^\mu$ does. The following lemma is sufficient.

Lemma C.1. *For each $1 \leq t \leq T$,*

$$\langle c^\mu, O_{[N],t}^\mu \rangle^2 \leq \frac{\bar{J}^2}{\tilde{\Lambda}_{\min}} D_{[N],tt}^\mu,$$

where $\tilde{\Lambda}^{\min}$ is given in proposition 3.2.

Proof. If $\bar{J} = 0$ the conclusion is evident, thus we assume throughout this proof that $\bar{J} \neq 0$. Since $D_{[N],tt}^\mu = \dagger \bar{O}_{[N],t}^\mu \tilde{K}_{[N]}^{\mu,0} O_{[N],t}^\mu$, we find from the definition that

$$D_{[N],tt}^\mu = \sum_{k,m=-n}^n \Lambda^N(k,m) \dagger O_{[N],t}^\mu M^{\mu,m} O_{[N],t}^\mu.$$

We introduce the matrixes $(L^{\mu,k})$, $k \in \mathbb{Z}$, where for $1 \leq s, t \leq T$,

$$L_{st}^{\mu,k} = M_{st}^{\mu,k} - \bar{c}_s^\mu \bar{c}_t^\mu = \int_{\mathcal{T}^N} (f(u_{s-1}^0) - \bar{c}_{s-1}^\mu)(f(u_{t-1}^k) - \bar{c}_{t-1}^\mu) \mu(du)$$

where $\bar{c}^\mu = \frac{1}{\bar{J}} c^\mu$.

These matrices have the same properties as the matrixes $M^{\mu,k}$, in particular the discrete Fourier Transform $(\tilde{L}^{\mu^N,l})_{l=-n,\dots,n}$ is Hermitian positive. Using this spectral representation we write

$$D_{[N],tt}^\mu = \tilde{\Lambda}^N(0,0) \langle \bar{c}^\mu, O_{[N],t}^\mu \rangle^2 + \frac{1}{N} \sum_{l=-n}^n \tilde{\Lambda}^N(0,-l) \dagger O_{[N],t}^\mu \tilde{L}^{\mu,l} O_{[N],t}^\mu,$$

and since $\tilde{\Lambda}^N(0,-l)$ is positive for all $l = -n, \dots, n$ and $\dagger O_{[N],t}^\mu \tilde{L}^{\mu^N,l} O_{[N],t}^\mu$ is positive for all $t = 1, \dots, T$, we have

$$D_{[N],tt}^\mu \geq \frac{\tilde{\Lambda}^N(0,0)}{\bar{J}^2} \langle c^\mu, O_{[N],t}^\mu \rangle^2,$$

and the conclusion follows from assumption (3.5). □

We may use the previous lemma to obtain a lower-bound for the quadratic form (C.3). We recall the easily-proved identity from the calculus of quadratics that, for all $x \in \mathbb{R}$,

$$ax^2 + 2bx \geq -\frac{b^2}{a}.$$

We therefore find, through lemma C.1, that (C.3) is greater than or equal to

$$-\frac{\bar{J}^2}{\tilde{\Lambda}_{\min}} \left(T\sigma^2 + \sum_{t=1}^T D_{[N],tt}^\mu \right) = -\frac{\bar{J}^2}{\tilde{\Lambda}_{\min}} \left(T\sigma^2 + \text{Trace}(\tilde{K}_{[N]}^{\mu,0}) \right). \quad (\text{C.4})$$

We have already noted in the proof of proposition 4.8 that $\text{Trace}(\tilde{K}^{\mu^N,0}) \leq T\Lambda^{sum}$. Thus, pulling these results together, we find that $\phi^N(\mu, v)$ is greater than $-\beta_2$, where

$$\beta_2 = \frac{T\bar{J}^2}{2\sigma^2 \tilde{\Lambda}_{\min}} (\sigma^2 + \Lambda^{sum}). \quad (\text{C.5})$$

D Proof of lemmas 5.4 and 5.14

For technical reasons, we need the following definition which is also used in appendix E. The motivation is that when we analyse the function $\Phi^N(\mu, v)$ defined in (4.26) we are led to use spectral representations and to introduce the Fourier transform \tilde{v} of v . Since $\tilde{v} \in (\mathbb{C}^T)^N$ the correspondence $v \rightarrow \tilde{v}$ from $\mathcal{T}_{1,T}^N$ to $(\mathbb{C}^T)^N$ is not one to one. We need to take into account the symmetries of \tilde{v} , hence the following definition.

Definition 8. For $v = (v^j)_{j=-n \dots n} \in \mathcal{T}_{1,T}^N$, we note $\mathcal{H}^N(v) = v_\diamond = (v_\diamond^{-n}, \dots, v_\diamond^n) \in \mathcal{T}_{1,T}^N$, where v_\diamond is defined from the Discrete Fourier Transform $\tilde{v} = (\tilde{v}^{-n}, \dots, \tilde{v}^n)$ of v as follows

$$\tilde{v}^k = \sum_{j=-n}^n v^j \exp\left(-\frac{2\pi i j k}{N}\right). \quad (\text{D.1})$$

The inverse transform is given by $v^j = \frac{1}{N} \sum_{k=-n}^n \tilde{v}^k \exp\left(\frac{2\pi i j k}{N}\right)$.

Because v is in $\mathcal{T}_{1,T}^N$ the real part of its DFT is even ($\text{Re}(\tilde{v}^{-k}) = \text{Re}(\tilde{v}^k)$, $k = -n, \dots, n$) and similarly its imaginary part is odd. As a consequence we define

$$v_\diamond^k = \begin{cases} \tilde{v}^0 & k = 0 \\ \sqrt{2}\text{Im}(\tilde{v}^k) & k = -n, \dots, -1 \\ \sqrt{2}\text{Re}(\tilde{v}^k) & k = 1, \dots, n \end{cases} \quad (\text{D.2})$$

It is easily verified that the mapping $v \rightarrow v_\diamond = \mathcal{H}^N(v)$ is a bijection from $\mathcal{T}_{1,T}^N$ to itself the inverse being given by

$$v^j = \frac{1}{N} \left[v_\diamond^0 + \sqrt{2}\text{Re} \left(\sum_{k=1}^n (v_\diamond^k + i v_\diamond^{-k}) e^{\frac{2\pi i j k}{N}} \right) \right]$$

and that

$$\sum_{k=-n}^n \|v_\diamond^k\|^2 = \sum_{k=-n}^n \dagger \tilde{v}^k * \tilde{v}^k = N \sum_{k=-n}^n \|v^k\|^2,$$

For a probability measure $\mu^N \in \mathcal{M}_1^+(\mathcal{T}^N)$, we define $\mu_\diamond^N = \mu_{1,T}^N \circ (\mathcal{H}^N)^{-1}$ to be the image law.

We also note $\underline{\mu}_\diamond^N$ the measure $\underline{\mu}_{1,T}^N \circ (\mathcal{H}^N)^{-1}$ (where $\underline{\mu}^N$ is given in definition 1). We note that

$$\underline{P}_\diamond \simeq \mathcal{N}_T(0_T, N\sigma^2 \text{Id}_T). \quad (\text{D.3})$$

We notice that $\Gamma_{[N],2}(\mu) = \int_{\mathcal{T}_{1,T}^N} \phi_\diamond^N(\mu, v_\diamond) \underline{\mu}_\diamond^N(dv_\diamond)$, where

$$\begin{aligned} \phi_\diamond^N(\mu, v_\diamond) &= \frac{1}{2N^2\sigma^2} \sum_{l=-n}^n \dagger \tilde{v}^l * \tilde{A}_{[N]}^{\mu, -l} \tilde{v}^l \\ &+ \frac{1}{N\sigma^2} \dagger \tilde{v}^0 \left(\text{Id}_T - \tilde{A}_{[N]}^\mu(0) \right) c^\mu - \frac{1}{2\sigma^2} \dagger c^\mu \left(\text{Id}_T - \tilde{A}_{[N]}^\mu(0) \right) c^\mu, \end{aligned} \quad (\text{D.4})$$

and \tilde{v} is implicitly given by (D.2) as a function of v_\diamond . We have used definition 8 and the DFT diagonalisation of Lemma A.2. We note that, since $\tilde{A}_{[N]}^{\mu, l}$ is Hermitian positive, $\dagger \tilde{v}^l * \tilde{A}_{[N]}^{\mu, l} \tilde{v}^l$ is real and positive. We recall lemma 5.4 and give a proof.

Lemma

There exist positive constants $c > 0$ and $a > 1$ such that, for all N ,

$$\int_{\mathcal{T}^N} \exp(aN\phi^N(\hat{\mu}_N(u), \Psi_{1,T}(u))) P^{\otimes N}(du) \leq \exp(Nc),$$

where ϕ^N is defined in (4.26).

Proof. We have from (C.1) that $\phi^N(\mu, v) = \underline{\phi}_\diamond^N(\mu, w_\diamond)$, where $w_\diamond^j = v_\diamond^j$ for all j , except that $w_\diamond^0 = v_\diamond^0 - Nc^\mu$. Since (by (D.3)) the distribution of the variables v_\diamond under $\underline{P}_\diamond^{\otimes N}$ is $\mathcal{N}_T(0_T, N\sigma^2 \text{Id}_T)^{\otimes N}$, the distribution of w_\diamond under $\underline{P}_\diamond^{\otimes N}$ is $\mathcal{N}_T(Nc^\mu, N\sigma^2 \text{Id}_T)^{\otimes N}$. By lemma 4.10, the eigenvalues of $\tilde{A}_{[N]}^{\mu, j}$ are upperbounded by $0 < \alpha < 1$, for all j . Thus

$$N\underline{\phi}_\diamond^N(\mu, w_\diamond) \leq \frac{\alpha}{2N\sigma^2} \sum_{l=-n}^n \|w_\diamond^l\|^2 + \frac{1}{\sigma^2} \langle c^\mu, w_\diamond^0 \rangle + \frac{N}{2\sigma^2} \|c^\mu\|^2. \quad (\text{D.5})$$

Hence we find that

$$\int_{\mathcal{T}^N} \exp(aN\phi^N(\hat{\mu}_N(u), \Psi(u))) P^{\otimes N}(du) \leq \left(\sqrt{2\pi N\sigma^2}\right)^{-NT} \times \int_{\mathcal{T}_{1,T}^{N-1}} \mathcal{G}_1 \exp\left(\frac{1}{2N\sigma^2} \left[\sum_{|j|=1}^n a\alpha \|y^j\|^2 - \|y^j\|^2 \right]\right) \prod_{|j|=1}^n \prod_{t=1}^T dy_t^j,$$

where

$$\mathcal{G}_1 = \int_{\mathcal{T}_{1,T}} \exp\left[\frac{1}{2N\sigma^2} \times [a\alpha \|y^0\|^2 + 2aN\langle c^{\hat{\mu}_N}, y^0 \rangle + aN^2 \|c^{\hat{\mu}_N}\|^2 - \|y^0 + Nc^{\hat{\mu}_N}\|^2]\right] \prod_{t=1}^T dy_t^0.$$

We note the dependency of \mathcal{G}_1 on (y^j) (for all $|j| \neq n$) via $c^{\hat{\mu}_N}$. After diagonalisation, we find that

$$\mathcal{G}_1 = \int_{\mathcal{T}_{1,T}} \exp\left[\frac{N\|c^{\hat{\mu}_N}\|^2 a(a-1)(1-\alpha)}{2(1-\alpha)\sigma^2} - \frac{(1-\alpha\alpha)}{2N\sigma^2} \sum_{s=1}^T \left(y_s^0 - \frac{Nc_s^{\hat{\mu}_N}(a-1)}{1-\alpha}\right)^2\right] \prod_{s=1}^T dy_s^0.$$

We assume that $a > 1$ is such that $1 - \alpha\alpha > 0$. To bound this expression, we note the identity that if $\mathcal{A} : \mathbb{R} \rightarrow \mathbb{R}$ satisfies $|\mathcal{A}| \leq \mathcal{B} > 0$ and $\gamma_c > 0$, then

$$\int_{\mathbb{R}} \exp\left(-\frac{1}{2\gamma_c} (t - \mathcal{A}(t))^2\right) dt \leq 2\mathcal{B} + \sqrt{2\pi\gamma_c}.$$

Since $|c_s^{\hat{\mu}_N}| \leq |\bar{J}|$, $s = 1, \dots, T$, and hence $\|c^{\hat{\mu}_N}\|^2 \leq T\bar{J}^2$, we therefore find that $\mathcal{G}_1 \leq \mathcal{G}_1^c$, where

$$\mathcal{G}_1^c = \exp\left[\frac{NT\bar{J}^2 a(a-1)(1-\alpha)}{2\sigma^2(1-\alpha\alpha)}\right] \left(\frac{2N|\bar{J}|(a-1)}{1-\alpha\alpha} + \sqrt{\frac{2\pi N\sigma^2}{1-\alpha\alpha}}\right)^T.$$

Thus

$$\int_{\mathcal{T}^N} \exp(aN\phi^N(\hat{\mu}_N(u), \Psi_{1,T}(u))) P^{\otimes N}(du) \leq \mathcal{G}_1^c (1-\alpha\alpha)^{-\frac{T(N-1)}{2}} (2\pi N\sigma^2)^{-\frac{T}{2}},$$

which yields the lemma. \square

We include the proof of lemma 5.14 which is used in the proof of the upper bound on compact sets in section 5.3.2.

Lemma

There exist constants $a > 1$ and $c > 0$ such that for all $\mu \in \mathcal{M}_{1,S}^+(\mathcal{T}^Z) \cap \mathcal{E}_2$,

$$\Gamma(\mu) \leq \frac{(I^{(3)}(\mu, P^Z) + c)}{a}.$$

Proof. $a > 1$ is chosen as in the proof of lemma 5.4. We have (from (5.3) that

$$I^{(3)}(\mu, P^Z) = \lim_{N \rightarrow \infty} N^{-1} I^{(2)}(\mu^N, P^{\otimes N}).$$

We recall that $I^{(2)}$ may be expressed using the variational expression (5.2) as

$$I^{(2)}(\mu^N, P^{\otimes N}) = \sup_{\zeta^N \in C_b(\mathcal{T}^N)} \left(\int_{\mathcal{T}^N} \zeta^N(u) \mu^N(du) - \log \int_{\mathcal{T}^N} \exp(\zeta^N(u)) P^{\otimes N}(du) \right), \quad (\text{D.6})$$

where ζ^N is a continuous, bounded function on \mathcal{T}^N . We let $\zeta_M^N = a 1_{B_M} \zeta_*^N$, where $\zeta_*^N(u) = N(\phi^N(\mu^N, \Psi_{1,T}(u)) + \Gamma_{[N],1}(\mu))$, and $u \in B_M$ only if either $\|\Psi(u)\| \leq NM$ or $(\phi^N(\mu^N, \Psi_{1,T}(u)) + \Gamma_{[N],1}(\mu)) \leq 0$. We proved in section 4.3.2 that $\phi^N(\mu^N, \Psi_{1,T}(u))$ possesses a lower bound, which means that ζ_M^N is continuous and bounded. Furthermore ζ_M^N grows to ζ_*^N , so that after substituting ζ_M^N into (D.6) and taking $M \rightarrow \infty$ (i.e. applying the dominated convergence theorem), we obtain

$$a \int_{\mathcal{T}^N} \zeta_*^N(u) \mu^N(du) \leq \log \int_{\mathcal{T}^N} \exp(a \zeta_*^N(u)) P^{\otimes N}(du) + I^{(2)}(\mu^N, P^{\otimes N}). \quad (\text{D.7})$$

It can be easily shown, similarly to lemma 5.4, that

$\log \int_{\mathcal{T}^N} \exp(a \zeta_*^N(u)) P^{\otimes N}(du) \leq Nc$. We may thus divide both sides by aN and let $N \rightarrow \infty$ to obtain the required result. \square

E Proof of lemma 5.11

We prove lemma 5.11.

Lemma

There exists a constant C_0 such that for all ν in $\mathcal{M}_{1,S}^+(\mathcal{T}^Z)$, all $\varepsilon > 0$ and all $\mu \in V_\varepsilon(\nu) \cap \mathcal{E}_2$,

$$\left| \Gamma_{[N]}(\mu) - \Gamma_{[N]}^\nu(\mu) \right| \leq C_0(C_N^\nu + \varepsilon)(1 + \mathbb{E}^{\underline{\mu}, T}[\|v^0\|^2]).$$

Here $V_\varepsilon(\nu)$ is the open neighbourhood defined in proposition 4.6, and $\underline{\mu}$ is given in definition 1.

Proof. We firstly bound Γ_1 . From (5.13) and (5.14) and lemma A.2 we have

$$\begin{aligned} \left| \Gamma_{[N],1}(\mu) - \Gamma_{[N],1}^\nu(\mu) \right| &\leq \frac{1}{2N} \sum_{l=-n}^n \left| \log \det \left(\text{Id}_T + \sigma^{-2} \tilde{K}_{[N]}^{\mu,l} \right) - \log \det \left(\text{Id}_T + \sigma^{-2} \tilde{K}_{[N]}^{\nu,l} \right) \right| \\ &\quad + \frac{1}{2N} \sum_{l=-n}^n \left| \log \det \left(\text{Id}_T + \sigma^{-2} \tilde{K}_{[N]}^{\nu,l} \right) - \log \det \left(\text{Id}_T + \sigma^{-2} \tilde{K}_{[N]}^{\nu,N,l} \right) \right|. \end{aligned}$$

It thus follows from proposition 4.6 and lemma 5.10 that

$$\left| \Gamma_{[N],1}(\mu) - \Gamma_{[N],1}^\nu(\mu) \right| \leq C_0^*(C_N^\nu + \varepsilon),$$

for some constant C_0^* which is independent of ν and N .

We define $\phi_{\infty,\diamond}^N(\nu, v_\diamond) = \phi_\infty^N(\nu, (\mathcal{H}^N)^{-1}(v))$, where \mathcal{H}^N is given in definition 8 and ϕ_∞^N is given in (5.12), and find that

$$\begin{aligned} \phi_{\infty,\diamond}^N(\mu, v_\diamond) &= \frac{1}{2N^2\sigma^2} \sum_{l=-n}^n \dagger \tilde{v}^{l,*} \tilde{A}^{\nu,-l} \tilde{v}^l \\ &\quad + \frac{1}{N\sigma^2} \dagger \tilde{v}^0 \left(\text{Id}_T - \tilde{A}^\nu(0) \right) c^\nu + \frac{1}{2\sigma^2} \dagger c^\nu \left(\text{Id}_T - \tilde{A}^\nu(0) \right) c^\nu. \quad (\text{E.1}) \end{aligned}$$

This means that

$$\Gamma_{[N],2}^\nu(\mu) - \Gamma_{[N],2}(\mu) = \int_{\mathcal{T}_{1,T}^N} \phi_{\infty,\diamond}^N(\nu, v_\diamond) - \phi_\diamond^N(\mu^N, v_\diamond) \underline{\mu}_\diamond^N(dv_\diamond). \quad (\text{E.2})$$

Upon expansion of the above expression, we find that

$$|\phi_{\infty,\diamond}^N(\nu, v_\diamond) - \phi_\diamond^N(\mu^N, v_\diamond)| \leq \frac{1}{2\sigma^2} \left(\frac{1}{N^2} \sum_{l=-n}^n \|\tilde{A}_{[N]}^{\mu,-l} - \tilde{A}^{\nu,N,-l}\| \|\tilde{v}^l\|^2 + \frac{2}{N} \|d_{\nu,\mu}\| \|\tilde{v}^0\| + |e_{\nu,\mu}| \right),$$

where $d_{\nu,\mu} = c^\mu - c^\nu + \tilde{A}^{\nu,N,0} c^\nu - \tilde{A}_{[N]}^{\mu,0} c^\mu$ and $e_{\nu,\mu} = \dagger c^\mu \tilde{A}_{[N]}^{\mu,0} c^\mu - \|c^\mu\|^2 - \dagger c^\nu \tilde{A}^{\nu,N,0} c^\nu + \|c^\nu\|^2$. It follows from proposition 4.5 and lemma 5.10 that the (Euclidean) norm each of the above terms is bounded by $C^*(C_N^\nu + \varepsilon)$ for some constant C^* .

The lemma now follows after consideration of the fact that $\int_{\mathcal{T}_T^Z} \|v^k\|^2 \underline{\mu}_{1,T}(dv) = \mathbb{E}^{\underline{\mu}_{1,T}}[\|v^0\|^2]$, $\|\tilde{v}^0\|^2 \leq N \sum_{k=-n}^n \|v^k\|^2$ (Cauchy-Schwarz) and, because of the properties of the DFT, $\sum_{l=-n}^n \|v^l\|^2 = N \sum_{k=-n}^n \|v^k\|^2$. \square

References

- [1] J. BALADRON, D. FASOLI, O. FAUGERAS, AND J. TOUBOUL, *Mean-field description and propagation of chaos in networks of Hodgkin-Huxley and Fitzhugh-Nagumo neurons*, The Journal of Mathematical Neuroscience, 2 (2012).
- [2] J. R. BAXTER AND N. C. JAIN, *An approximation condition for large deviations and some applications*, in Convergence in Ergodic Theory and Probability, V. Bergulson, ed., Ohio State University Mathematical Research Institute Publications, 1993.
- [3] G. BEN-AROUS AND A. GUIONNET, *Large deviations for langevin spin glass dynamics*, Probability Theory and Related Fields, 102 (1995), pp. 455–509.
- [4] ———, *Symmetric Langevin Spin Glass Dynamics*, The Annals of Probability, 25 (1997), pp. 1367–1422.
- [5] P. BRESSLOFF, *Stochastic neural field theory and the system-size expansion*, SIAM J. Appl. Math, 70 (2009), pp. 1488–1521.
- [6] A. BUDHIRAJA, P. DUPUIS, AND F. M., *Large deviation properties of weakly interacting processes via weak convergence methods*, Annals of Probability, 40 (2012), pp. 74–102.
- [7] M. BUICE AND J. COWAN, *Field-theoretic approach to fluctuation effects in neural networks*, Physical Review E, 75 (2007).
- [8] M. BUICE, J. COWAN, AND C. CHOW, *Systematic fluctuation expansion for neural network activity equations*, Neural computation, 22 (2010), pp. 377–426.
- [9] M. J. CÁCERES, J. A. CARILLO, AND B. PERHAME, *Analysis of nonlinear noisy integrate and fire neuron models: blow-up and steady states*, The Journal of Mathematical Neuroscience, 1 (2011).
- [10] B. CESSAC, *Increase in complexity in random neural networks*, Journal de Physique I (France), 5 (1995), pp. 409–432.

-
- [11] B. CESSAC AND M. SAMUELIDES, *From neuron to neural networks dynamics.*, EPJ Special topics: Topics in Dynamical Neural Networks, 142 (2007), pp. 7–88.
- [12] T. CHIYONOBU AND S. KUSUOKA, *The large deviation principle for hypermixing processes*, Probability Theory and Related Fields, 78 (1988), pp. 627–649.
- [13] A. CRISANTI AND H. SOMPOLINSKY, *Dynamics of spin systems with randomly asymmetric bonds: Langevin dynamics and a spherical model*, Physical Review A, 36 (1987), pp. 4922–4939.
- [14] ———, *Dynamics of spin systems with randomly asymmetric bounds: Ising spins and Glauber dynamics*, Phys. Review A, 37 (1987), p. 4865.
- [15] L. F. CUGLIANDOLO, J. KURCHAN, P. LE DOUSSAL, AND L. PELITI, *Glassy behaviour in disordered systems with nonrelaxational dynamics*, Physical review letters, 78 (1997), pp. 350–353.
- [16] D. DALEY AND D. VERE-JONES, *An introduction to the theory of point processes: volume II: general theory and structure*, vol. 2, Springer, 2007.
- [17] D. DAWSON AND J. GARTNER, *Large deviations from the mckean-ulasov limit for weakly interacting diffusions*, Stochastics, 20 (1987).
- [18] ———, *Multilevel large deviations and interacting diffusions*, Probability Theory and Related Fields, 98 (1994), pp. 423–487.
- [19] P. DAYAN AND L. ABBOTT, *Theoretical Neuroscience : Computational and Mathematical Modeling of Neural Systems*, MIT Press, 2001.
- [20] A. DEMBO AND O. ZEITOUNI, *Large deviations techniques*, Springer, 1997. 2nd Edition.
- [21] J.-D. DEUSCHEL AND D. W. STROOCK, *Large Deviations*, vol. 137 of Pure and Applied Mathematics, Academic Press, 1989.
- [22] M. DONSKER AND S. VARADHAN, *Asymptotic evaluation of certain markov process expectations for large time, iv*, Communications on Pure and Applied Mathematics, XXXVI (1983), pp. 183–212.
- [23] M. DONSKER AND S. VARADHAN, *Large deviations for stationary Gaussian processes*, Commun. Math. Phys., 97 (1985), pp. 187–210.
- [24] S. ELBOUSTANI AND A. DESTEXHE, *A master equation formalism for macroscopic modeling of asynchronous irregular activity states*, Neural computation, 21 (2009), pp. 46–100.
- [25] R. ELLIS, *Entropy, large deviations and statistical mechanics*, Springer, 1985.
- [26] G. B. ERMENTROUT AND D. TERMAN, *Foundations of Mathematical Neuroscience*, Interdisciplinary Applied Mathematics, Springer, 2010.
- [27] O. FAUGERAS AND J. MACLAURIN, *A large deviation principle and an analytical expression of the rate function for a discrete stationary gaussian process*, tech. rep., ArXiv: <http://arxiv.org/abs/1311.4400>, Nov. 2013.
- [28] M. FISCHER, *On the form of the large deviation rate function for the empirical measures of weakly interacting systems*, Archiv, (2012).

-
- [29] W. GERSTNER, *Time structure of the activity in neural network models*, Physical Review E, 51 (1995), pp. 738–758.
- [30] W. GERSTNER AND W. KISTLER, *Spiking Neuron Models*, Cambridge University Press, 2002.
- [31] W. GERSTNER AND J. VAN HEMMEN, *Coherence and incoherence in a globally coupled ensemble of pulse-emitting units*, Physical review letters, 71 (1993), pp. 312–315.
- [32] I. GINZBURG AND H. SOMPOLINSKY, *Theory of correlations in stochastic neural networks*, Physical Review E, 50 (1994).
- [33] A. GUIONNET, *Dynamique de Langevin d'un verre de spins*, PhD thesis, Université de Paris Sud, 1995.
- [34] ———, *Averaged and quenched propagation of chaos for spin glass dynamics*, Probability Theory and Related Fields, 109 (1997), pp. 183–215.
- [35] E. IZHIKEVICH, *Dynamical Systems in Neuroscience: The Geometry of Excitability And Bursting*, MIT Press, 2007.
- [36] E. KANDEL, J. SCHWARTZ, AND T. JESSEL, *Principles of Neural Science*, McGraw-Hill, 4th ed., 2000.
- [37] L. LAPICQUE, *Recherches quantitatives sur l'excitation des nerfs traitée comme une polarisation*, J. Physiol. Paris, 9 (1907), pp. 620–635.
- [38] T. M. LIGGETT, *Interacting Particle Systems*, Springer Berlin Heidelberg, 2005.
- [39] O. MOYNOT, *Etude mathématique de la dynamique des réseaux neuronaux aléatoires récurrents*, PhD thesis, Université Paul Sabatier, Toulouse, 1999.
- [40] O. MOYNOT AND M. SAMUELIDES, *Large deviations and mean-field theory for asymmetric random recurrent neural networks*, Probability Theory and Related Fields, 123 (2002), pp. 41–75.
- [41] J. NEVEU, *Processus aléatoires gaussiens*, vol. 34, Presses de l'Université de Montréal, 1968.
- [42] M. SAMUELIDES AND B. CESSAC, *Random recurrent neural networks*, European Physical Journal - Special Topics, 142 (2007), pp. 7–88.
- [43] H. SOMPOLINSKY, A. CRISANTI, AND H. SOMMERS, *Chaos in Random Neural Networks*, Physical Review Letters, 61 (1988), pp. 259–262.
- [44] H. SOMPOLINSKY AND A. ZIPPELIUS, *Dynamic theory of the spin-glass phase*, Physical Review Letters, 47 (1981), pp. 359–362.
- [45] ———, *Relaxational dynamics of the Edwards-Anderson model and the mean-field theory of spin-glasses*, Physical Review B, 25 (1982), pp. 6860–6875.



**RESEARCH CENTRE
SOPHIA ANTIPOLIS – MÉDITERRANÉE**

2004 route des Lucioles - BP 93
06902 Sophia Antipolis Cedex

Publisher
Inria
Domaine de Voluceau - Rocquencourt
BP 105 - 78153 Le Chesnay Cedex
inria.fr

ISSN 0249-6399