

On randomly colouring locally sparse graphs

Alan Frieze, Juan Vera

► **To cite this version:**

Alan Frieze, Juan Vera. On randomly colouring locally sparse graphs. Discrete Mathematics and Theoretical Computer Science, DMTCS, 2006, 8, pp.121-128. hal-00961105

HAL Id: hal-00961105

<https://hal.inria.fr/hal-00961105>

Submitted on 19 Mar 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

On randomly colouring locally sparse graphs

Alan Frieze and Juan Vera[†]

Department of Mathematics
Carnegie Mellon University
Pittsburgh PA15213, USA

alan@random.math.cmu.edu
jvera@andrew.cmu.edu

received Aug 30, 2004, accepted Apr 13, 2006.

We consider the problem of generating a random q -colouring of a graph $G = (V, E)$. We consider the simple Glauber Dynamics chain. We show that if for all $v \in V$ the average degree of the subgraph H_v induced by the neighbours of $v \in V$ is $\ll \Delta$ where Δ is the maximum degree and $\Delta > c_1 \ln n$ then for sufficiently large c_1 , this chain mixes rapidly provided $q/\Delta > \alpha$, where $\alpha \approx 1.763$ is the root of $\alpha = e^{1/\alpha}$. For this class of graphs, which includes planar graphs, triangle free graphs and random graphs $G_{n,p}$ with $p \ll 1$, this beats the $11\Delta/6$ bound of Vigoda [20] for general graphs.

Keywords: Counting Colourings, Sampling, Markov Chains

1 Introduction

Markov Chain Monte Carlo (MCMC) is an important tool in sampling from complex distributions. It has been successfully applied in several areas of Computer Science, most notably volume computation [3], [15], [16] and estimating the permanent of a non-negative matrix [12]. It was used by Jerrum [10] to generate a random q -colouring of a graph G , provided $q > 2\Delta$. This has led to the challenging problem of determining the smallest value of q for which it is possible to generate a (near)-uniform sample from the set \mathcal{Q} of proper q -colourings of G in polynomial time. We cannot expect the chain to mix for $q \leq \Delta + 1$ and at present it is unknown as to whether or not it mixes rapidly for say $q = \Delta + 2$. Vigoda [20] improved Jerrum's result by reducing the lower bound on q to $11\Delta/6$. This is still the best lower bound on q for general graphs.

The lack of complete success on the general problem has led to the analysis of restricted classes of graphs. Suppose that we consider *Glauber dynamics* on the set \mathcal{Q} . Specifically we will consider the *heat bath* dynamics, which may be described as follows. We start from an arbitrary proper q -colouring $X_0 \in \mathcal{Q}$. At step $t > 0$ of the process, in state $X_{t-1} \in \mathcal{Q}$, we choose a vertex $v_t \in V$ uniformly

[†]Research supported by NSF grant CCR-0200945.

at random. Then we choose j_t uniformly at random from the colours with which v_t may be properly coloured, given $X_{t-1}(V \setminus v_t)$. We recolour v_t with j_t to give $X_t \in \mathcal{Q}$.

Dyer and Frieze [2] considered this process restricted to the class of graphs $\mathcal{G}(c_1, c_2)$: the set of graphs with n vertices, maximum degree $\Delta \geq c_1 \log n$ and girth $g \geq c_2 \log \Delta$. They showed using the idea of “burn-in” that for c_1, c_2 sufficiently large, Glauber Dynamics mixed in $O(n \log n)$ time, provided $q > \alpha \Delta$ where $\alpha \approx 1.763$ is the root of $\alpha = e^{1/\alpha}$. Molloy [17] improved this result by reducing the lower bound on q to being more than $\beta \Delta$ where $\beta \approx 1.489$ is the root of $(1 - e^{-1/\beta})^2 + \beta e^{-1/\beta} = 1$. The girth assumptions were then relaxed by Hayes [7] to $g \geq 5$ for $k/\Delta > \alpha$ and $g \geq 6$ for $k/\Delta > \beta$. Subsequently, Hayes and Vigoda [8] made considerable progress, using a non-Markovian coupling, and reduced the lower bound on k/Δ to $(1 + \epsilon)$ for all $\epsilon > 0$, which is nearly optimal. Their result requires girth $g \geq 9$. However, the large maximum degree restriction remained. This was replaced by $\Delta \geq \Delta_0$ in Dyer, Frieze, Hayes and Vigoda [5], with the same restrictions on girth as in [7]. Dyer, Flaxman, Frieze and Vigoda [4] show that for sparse random graphs, the number of colours required for rapid mixing is of order the average rather than maximum degree **whp**. Goldberg, Martin and Paterson [6] prove results on the related notion of strong spatial mixing.

In this paper we avoid girth restrictions and consider locally sparse graphs instead. We say that a graph $G = (V, E)$ is γ -locally sparse if for all $v \in V$, the average degree of the graph induced by the neighbourhood $N(v)$ is at most γ . Thus planar graphs are always 6-locally-sparse and triangle free graphs are 0-locally-sparse.

Theorem 1.1 *Suppose that $q \geq (\alpha + \epsilon)\Delta$ where ϵ is a small positive constant. Let G be an n -vertex γ -locally sparse graph with $\gamma \leq \epsilon^2 \Delta / 10$ and $\Delta \geq c_1 \log n$. If $c_1 = c_1(\epsilon)$ is sufficiently large then the Glauber dynamics converges to within variation distance e^{-1} from uniform over \mathcal{Q} in at most $O(n \ln n)$.*

Notice that if $G = G_{n,p}$ and $\frac{c_1 \log n}{n} \leq p \leq \epsilon^2 / 11$ then **whp** G satisfies the conditions of the theorem. Note also that the chromatic number of a triangle-free graph is $O(\Delta / \log \Delta)$ – see Johansson [14] or Molloy and Reed [18] or Alon, Krivelevich and Sudakov [1] or Vu [21].

Our proof uses coupling and relies on a recent idea from Hayes and Vigoda [9] that utilises the fact that we can couple against the steady state distribution of the chain. Note that the theorem generalises Theorem 4 of [9].

In what follows we will assume that n is sufficiently large and ϵ is sufficiently small to satisfy our inequalities.

2 Preliminaries

We will consider two copies of Glauber Dynamics, $(X_t, t \geq 0)$ and $(Y_t, t \geq 0)$. Here X_0 is an arbitrary colouring and Y_0 is chosen from the uniform (*stationary*) distribution over \mathcal{Q} . At time t , the Hamming distance between X_t, Y_t is defined by

$$H(X_t, Y_t) = \sum_{v \in V} 1_{X_t(v) \neq Y_t(v)}.$$

We will couple the two processes as in Jerrum [10]. Here v_t is the same in both processes and then the choice of colours is maximally coupled. For vertex w let

$$A(X_t, w) = \{c \in [q] : c \notin X_t(N(w))\}$$

be the set of colours available to colour w in X_t if $v_t = w$.

Let $a(X_t, w) = |A(X_t, w)|$ and define the terms $A(Y_t, w)$, $a(Y_t, w)$ analogously.

It is shown in [9] that

$$\mathbf{E}(H(X_{t+1}, Y_{t+1}) - H(X_t, Y_t)) \leq -\frac{1}{n}H(X_t, Y_t) + \frac{1}{n} \sum_{w \in V} \frac{|\{u \in N(w) : X_t(u) \neq Y_t(u)\}|}{\max\{a(X_t, w), a(Y_t, w)\}}. \quad (1)$$

We will show that for $w \in V$ and $\delta = \epsilon/10$,

$$\Pr(a(Y_t, w) \leq \Delta/(1 - \delta)) \leq n^{-4}. \quad (2)$$

Assuming that $a(Y_t, w) \geq \Delta/(1 - \delta)$ in (1) we get

$$\begin{aligned} \mathbf{E}(H(X_{t+1}, Y_{t+1}) - H(X_t, Y_t)) &\leq -\frac{1}{n}H(X_t, Y_t) + \frac{1}{n} \frac{H(X_t, Y_t)\Delta}{\Delta/(1 - \delta)} \\ &\leq -\frac{\delta}{n}H(X_t, Y_t). \end{aligned}$$

So conditional on an event of probability $1 - O(n^{-3})$, we have

$$\mathbf{E}(H(X_{t+1}, Y_{t+1}) \mid X_t, Y_t) \leq \left(1 - \frac{\delta}{n}\right) H(X_t, Y_t).$$

Thus if $T = n(1 + \ln n)\delta^{-1}$ then conditional on an event of probability $1 - O(n^{-2} \log n)$, we have

$$\mathbf{E}(H(X_T, Y_T)) \leq e^{-1}$$

and so unconditionally

$$\mathbf{E}(H(X_T, Y_T)) \leq e^{-1} + o(1).$$

Hence the mixing time of the Glauber Dynamics is $O(n \ln n)$ as claimed.

3 Bounding the number of available colours

Fix $v \in W$ and let H_v be the subgraph of G induced by $N(v)$. Let $B(v)$ be the vertices of $N(v)$ that have degree at least $\gamma\delta^{-1}$ in H_v . Note that $\gamma\delta^{-1} \leq \epsilon\Delta$ and

$$|B(v)| \leq \delta|N(v)|, \quad (3)$$

since G is γ -locally-sparse.

Let

$$N^*(v) = N(v) \setminus B(v) = \{w_1, w_2, \dots, w_d\}.$$

Now let us fix the colours $\kappa(v)$ used at

$$v \in W_v = V \setminus N^*(v).$$

Let us use the term *allowable* for colorings of $N^*(v)$ which respect this conditioning. Let Ω be the set of allowable colourings of $N^*(v)$.

Let $a^*(\sigma, v)$ be the number of colours not used on $N^*(v)$. Note that (3) implies

$$a(\sigma, v) \geq a^*(\sigma, v) - \delta|N(v)|. \quad (4)$$

Now consider the following process \mathcal{P}_σ for producing an allowable colouring of H_v . Here $\sigma \in \Omega$. We let $\sigma_0 = \sigma$ and for $j = 1, 2, \dots, d$ let σ_j be obtained from σ_{j-1} as follows: Keep $\sigma_j(w_k) = \sigma_{j-1}(w_k)$ for $k \neq j$ and choose $\sigma_j(w_j)$ randomly from what is available to it.

Let Z_σ be the number of colours not appearing on a vertex in $N^*(v)$ if we start with $\sigma_0 = \sigma$.

Lemma 3.1 *If σ is chosen uniformly from Ω then for any $c > 0$,*

$$\Pr(a^*(\sigma, v) \geq c) = \Pr(Z_\sigma \geq c).$$

Proof We first prove that

$$\text{If } \sigma_0 \text{ is chosen uniformly from } \Omega \text{ then } \sigma_d \text{ is also uniform over } \Omega. \quad (5)$$

We do this by induction on j , with base case $j = 0$.

$$\begin{aligned} \Pr(\sigma_j = \sigma) &= \sum_{\sigma' \in \Omega} \Pr(\sigma_j = \sigma \mid \sigma_{j-1} = \sigma') \Pr(\sigma_{j-1} = \sigma') \\ &= \frac{1}{|\Omega|} \sum_{\sigma' \sim \sigma} \Pr(\sigma_j = \sigma \mid \sigma_{j-1} = \sigma') \end{aligned}$$

Here $\sigma' \sim \sigma$ if σ, σ' differ only at w_j .

$$\begin{aligned} &= \frac{1}{|\Omega|} \sum_{\sigma' \sim \sigma} \frac{1}{|\{\sigma' : \sigma' \sim \sigma\}|} \\ &= \frac{1}{|\Omega|}. \end{aligned}$$

Now $a^*(\sigma_d, v) = Z_{\sigma_0}$ and so

$$\Pr(a^*(\sigma_d, v) \geq c) = \Pr(Z_{\sigma_0} \geq c)$$

and the lemma follows from (5) □

For $w \in N^*(v)$ let

$$L(w) = [q] \setminus \{\kappa(u) : u \in N(w) \setminus N^*(v)\}$$

be the colours not specifically barred from w by the current conditioning. Then let

$$L^*(w_j) = [q] \setminus \{\sigma_{j-1}(u) : u \neq w_j\} \quad \text{for } j = 1, 2, \dots, d$$

be the colours available to w_j when it is re-coloured by σ_j .

We will first estimate the (conditional) expectation of Z_σ for arbitrary σ . Suppose that $x \in [q]$. Let $\theta_{x,j} = 1_{x \in L(w_j)}$ and let $\theta_{x,j}^* = 1_{x \in L^*(w_j)}$. Then we have

$$\begin{aligned} \Pr(x \notin \sigma_d(N^*(v))) &= \prod_{j=1}^d \Pr(\sigma_d(w_j) \neq x \mid \sigma_d(w_i) \neq x, 1 \leq i < j) \\ &= \prod_{j=1}^d \mathbf{E} \left(\left(1 - \frac{1}{|L^*(w_j)|} \right)^{\theta_{x,j}^*} \right) \\ &\geq \prod_{j=1}^d \left(1 - \frac{1}{|L(w_j)| - \gamma\delta^{-1}} \right)^{\theta_{x,j}} \end{aligned}$$

since $|L^*(w_j)| \geq |L(w_j)| - \gamma\delta^{-1}$ and $L^*(w_j) \subseteq L(w_j)$ implying $\theta_{x,j}^* \leq \theta_{x,j}$.

Then, following [2],

$$\begin{aligned} \mathbf{E}(Z_\sigma) &\geq \sum_{x \in [q]} \prod_{j=1}^d \left(1 - \frac{1}{|L(w_j)| - \gamma\delta^{-1}} \right)^{\theta_{x,j}} \\ &\geq q \left(\prod_{x \in [q]} \prod_{j=1}^d \left(1 - \frac{1}{|L(w_j)| - \gamma\delta^{-1}} \right)^{\theta_{x,j}} \right)^{1/q} \\ &= q \left(\prod_{j=1}^d \left(1 - \frac{1}{|L(w_j)| - \gamma\delta^{-1}} \right)^{|L(w_j)|} \right)^{1/q} \\ &\geq q \exp \left\{ -\frac{1}{q} \sum_{j=1}^d \frac{|L(w_j)|}{|L(w_j)| - 1 - \gamma\delta^{-1}} \right\}, \quad \text{using } 1 - x \geq e^{-x/(1-x)} \text{ for } 0 < x < 1, \\ &\geq q \exp \left\{ -\frac{\Delta}{q} \cdot \frac{q - \Delta}{q - \Delta - 1 - \gamma\delta^{-1}} \right\} \\ &\geq \left(1 + \frac{\epsilon}{2} \right) \Delta. \end{aligned} \tag{6}$$

(If $f(x) = xe^{-1/x}$ then $f(\alpha) = 1$ and $f'(\alpha) \sim .891$.)

We will now prove that for all $\sigma \in \Omega$, Z_σ is concentrated around its mean via the use of the Azuma-Hoeffding martingale inequality. To this end, let x_1, x_2, \dots, x_d be the colours assigned to w_1, w_2, \dots, w_d . Thus we can write $Z_\sigma = Z_\sigma(x_1, x_2, \dots, x_d)$. Now let

$$Z_{\sigma,i} = Z_{\sigma,i}(x_1, x_2, \dots, x_i) = \mathbf{E}(Z \mid x_1, x_2, \dots, x_i).$$

We will show next that for all feasible colours $x_1, x_2, \dots, x_i, x'_i$ that

$$|Z_{\sigma,i}(x_1, x_2, \dots, x_{i-1}, x_i) - Z_{\sigma,i}(x_1, x_2, \dots, x_{i-1}, x'_i)| \leq 2. \tag{7}$$

The aforementioned inequality will then imply that for any $t \geq 0$,

$$\Pr(Z_\sigma - \mathbf{E}(Z_\sigma) \leq -t) \leq e^{-t^2/(2d)}$$

and then taking $t = \epsilon\Delta/4$ and using (6) we get

$$\Pr\left(Z_\sigma \leq \left(1 + \frac{\epsilon}{4}\right)\Delta\right) \leq e^{-\epsilon^2\Delta/32}.$$

This together with Lemma 3.1 and (4) implies (2).

To prove (7), fix $i, x_1, x_2, \dots, x_i, x_i^*$. In one instance of \mathcal{P}_σ we start by colouring w_1, w_2, \dots, w_i with x_1, x_2, \dots, x_i to produce colouring τ . In another instance we start by colouring w_1, w_2, \dots, w_i with x_1, x_2, \dots, x_i^* to produce colouring τ^* .

We couple these two constructions in order to minimise the expected difference in the number of vertices U with a different colour. A *paths of disagreement* argument gives that

$$\mathbf{E}(U) \leq 1 + \sum_{j=i+1}^d \left(\frac{\gamma\delta^{-1}}{|L(w_j)| - \gamma\delta^{-1}} \right)^{j-i} \leq 2 \quad (8)$$

and (7) follows. \square

Explanation of (8): We claim that if c_j, c_j^* is the colour of v_j in σ_d, σ_d^* respectively, then

$$\Pr(c_j \neq c_j^*) \leq \left(\frac{\gamma\delta^{-1}}{|L(w_j)| - \gamma\delta^{-1}} \right)^{j-i}.$$

This is because if $c_j \neq c_j^*$ then there is a path of disagreements $v_{i_1}, v_{i_2}, \dots, v_{i_s}$ where $i = i_1 < i_2 < \dots < i_s = j$ such that $c_{i_r} \neq c_{i_r}^*$ for $1 \leq r \leq s$. There are at most $(\lambda\delta^{-1})^{j-i}$ such paths and each has probability at most $(|L(w_j)| - \gamma\delta^{-1})^{i-j}$ of all vertices being coloured differently. \square

References

- [1] N. Alon, M. Krivelevich and B. Sudakov, Coloring graphs with sparse neighborhoods, *Journal of Combinatorial Theory, Ser. B* 77 (1999) 73-82.
- [2] M. Dyer, A. Frieze, Randomly colouring graphs with lower bounds on girth and maximum degree, *Random Structures and Algorithms*, 23 167-179, 2003.
- [3] M.E. Dyer, A.M. Frieze and R. Kannan, A random polynomial time algorithm for approximating the volume of convex bodies, *Journal of the Association for Computing Machinery* 38(1):1-17, 1991.
- [4] M.E. Dyer, A. Flaxman, A.M. Frieze and E. Vigoda, Randomly colouring sparse random graphs.
- [5] M.E. Dyer, A.M. Frieze, T. P. Hayes and E. Vigoda, Randomly coloring constant degree graphs.
- [6] L. Goldberg, R. Martin and M. Paterson, Strong spatial mixing for graphs with fewer colours.
- [7] T. P. Hayes, Randomly coloring graphs of girth at least five, *Proceedings of the 35th Annual ACM Symposium on Theory of Computing*, (2003) 269-278
- [8] T. P. Hayes and E. Vigoda, A Non-Markovian Coupling for Randomly Sampling Colorings. *Proceedings of the 44th Annual IEEE Symposium on Foundations of Computer Science*, (2003) 618-627.
- [9] T. P. Hayes and E. Vigoda, Coupling with the stationarity distribution and improved sampling for colorings and independent sets.
- [10] M.R. Jerrum, A very simple algorithm for estimating the number of k -colourings of a low-degree graph, *Random Structures and Algorithms* 7 (1995), 157-165.
- [11] M.R. Jerrum, *Counting, sampling and integrating: algorithms and complexity*, Lectures in Mathematics-ETH Zürich, Birkhäuser, to appear January 2003.
- [12] M.R. Jerrum, A. Sinclair and E. Vigoda, A polynomial-time approximation algorithm for the permanent of a matrix with non-negative entries, *Proceedings of the 12th Annual ACM-SIAM Symposium on Discrete Algorithms* (2001), 712-721.
- [13] M.R. Jerrum, L.G. Valiant and V.V. Vazirani, Random generation of combinatorial structures from a uniform distribution, *Theoretical Computer Science* 43 (1986), 169-188.
- [14] A. Johansson, Asymptotic choice number for triangle-free graphs, DIMACS technical report 1996.
- [15] R. Kannan, L. Lovász and M. Simonovits, Random walks and an $O^*(n^5)$ volume algorithm for convex bodies, *Random Structures and Algorithms* 11 (1997), 1-50.
- [16] L. Lovász and S. Vempala, Simulated annealing in convex bodies and an $O^*(n^4)$ volume algorithm, to appear in JCSS.
- [17] M. Molloy, The Glauber dynamics on colorings of a graph with high girth and maximum degree. To appear in *SIAM Journal on Computing*.
- [18] M. Molloy and B. Reed, *Graph Colouring and the Probabilistic Method*, Springer, 2002.
- [19] J. van den Berg and J.E. Steif, Percolation and the hard-core lattice gas model, *Stochastic Processes and their Applications* 49 (1994), 179-197.
- [20] E. Vigoda, Improved bounds for sampling colorings, *Proceedings of the 40th Annual IEEE Symposium on Foundations of Computer Science* (1999), 51-59.
- [21] V. Vu, An upper bound on the list chromatic number of locally sparse graphs, *Combinatorics, Probability and Computing* 11 (2002), 103-111

