

Some new optimal and suboptimal infinite families of undirected double-loop networks

Bao-Xing Chen, Ji-Xiang Meng, Wen-Jun Xiao

► **To cite this version:**

Bao-Xing Chen, Ji-Xiang Meng, Wen-Jun Xiao. Some new optimal and suboptimal infinite families of undirected double-loop networks. *Discrete Mathematics and Theoretical Computer Science, DMTCS*, 2006, 8, pp.299–311. <hal-00961120>

HAL Id: hal-00961120

<https://hal.inria.fr/hal-00961120>

Submitted on 19 Mar 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Some new optimal and suboptimal infinite families of undirected double-loop networks

Bao-Xing Chen^{1,2} †, Ji-Xiang Meng² and Wen-Jun Xiao³

¹Dept. of Computer Science, Zhangzhou Teacher's College, Zhangzhou, P.R. of China
Email: cbaoxing@hotmail.com

²College of Mathematics & System Science, Xinjiang University, Wulumuqi, P.R. of China

³Dept. of Computer Science, South China University of Technology, Guangzhou, P.R. of China

received Apr 3, 2006, revised Jul 16, 2006, accepted Oct 24, 2006.

Let n, s be positive integers such that $2 \leq s < n$ and $s \neq \frac{n}{2}$. An undirected double-loop network $G(n; 1, s)$ is an undirected graph (V, E) , where $V = \mathbb{Z}_n = \{0, 1, 2, \dots, n-1\}$ and $E = \{(i, i+1 \pmod{n}), (i, i+s \pmod{n}) \mid i \in \mathbb{Z}\}$. It is a circulant graph with n nodes and degree 4. In this paper, the sufficient and necessary conditions for a class of undirected double-loop networks to be optimal are presented. By these conditions, 6 new optimal and 5 new suboptimal infinite families of undirected double-loop networks are given.

AMS Subject Classifications: 05C12

Keywords: undirected double-loop networks, diameter, optimal, suboptimal

1 Introduction

Double-loop networks are popular in the design and implementation of metropolitan networks and parallel processing computer systems. They have many attractive properties like vertex symmetry, incremental extensibility, low valency, ease of implementation, etc. Some researchers are interested in the study of double-loop networks [1-22]. They mainly focus on designs of optimal double-loop networks [1], [2], [4], [9], [12]–[15], [18], [19], diameters [1]–[4], [9]–[15], [18]–[22] and routing [5], [7], [8], [17]. For more details we refer readers to [3], [16] and the references therein.

Let n, s be positive integers such that $2 \leq s < n$ and $s \neq \frac{n}{2}$. The undirected double-loop network $G(n; 1, s)$ is an undirected graph (V, E) , where $V = \mathbb{Z}_n = \{0, 1, 2, \dots, n-1\}$ and $E = \{(i, i+1 \pmod{n}), (i, i+s \pmod{n}) \mid i \in \mathbb{Z}_n\}$. It is a circulant graph with n nodes and degree 4. Let $d(i, j)$ be the length of a shortest path from node i to node j . Let $d(n; 1, s)$ denote the diameter of $G(n; 1, s)$. Since $G(n; 1, s)$ is vertex symmetric, $d(n; 1, s) = \max\{d(i, j) \mid 0 \leq i, j < n\} = \max\{d(0, i) \mid 0 \leq i < n\}$. Let $D(n) = \min\{d(n; 1, s) \mid 1 < s < n\}$. Wong and Coppersmith [20] gave the lower bound $\frac{1}{2}(\sqrt{2n} - 3)$

†This work was supported by the Natural Science Foundation of Fujian Province(No. Z0511035).

for $D(n)$. Boesch and Wang [4] sharpened the bound by giving $lb(n) = \lceil \frac{\sqrt{2n-1}-1}{2} \rceil$, where $\lceil x \rceil$ denotes the minimum integer $\geq x$.

If s exists such that $D(n) = d(n; 1, s) = lb(n)$, then n , s , and $G(n; 1, s)$ will be called optimal. If s exists such that $D(n) = d(n; 1, s) = lb(n) + 1$, then n , s , and $G(n; 1, s)$ will be called suboptimal.

A set Θ of natural numbers will be called an optimal (suboptimal) family if each $n \in \Theta$ is optimal (suboptimal).

Some optimal infinite families of undirected double-loop networks were given in [2], [11], [12], [19]. And as we know, only one suboptimal infinite family of undirected double-loop networks $\{2t^2 + 2t \mid t \geq 2\}$ has been given so far [12]. In this paper, the sufficient and necessary conditions for a class of undirected double-loop networks to be optimal are presented. And by using these conditions, we obtain 6 new optimal and 5 new suboptimal infinite families of undirected double-loop networks.

2 Definitions and some lemmas

Let \mathbb{Z} and \mathbb{Z}^+ be the set of integers and nonnegative integers respectively. Let $\lfloor x \rfloor$ denote the maximum integer $\leq x$.

Definition 2.1 (a_1, a_2) is said to be a non-negative solution of the equation

$$x + ys \equiv 0 \pmod{n} \quad (1)$$

if $a_1 + a_2s \equiv 0 \pmod{n}$, $a_1 \geq 0$, $a_2 \geq 0$ and $(a_1, a_2) \neq (0, 0)$.

(u, v) is said to be the smallest non-negative solution of the equation (1) if (u, v) is a non-negative solution of the equation (1) and the following conditions hold:

- (1) if (a_1, a_2) is a non-negative solution of the equation (1), then $u + v \leq a_1 + a_2$.
- (2) if (a_1, a_2) is a non-negative solution of the equation (1), where $(a_1, a_2) \neq (u, v)$ and $u + v = a_1 + a_2$, then $u > a_1$.

Clearly, the smallest non-negative solution of equation (1) is unique. For example, it is easy to see that $(4, 1)$, $(2, 3)$, $(0, 5)$, $(8, 2)$, $(4, 6)$, \dots are non-negative solutions of the equation $x + 6y \equiv 0 \pmod{10}$. Thus $(4, 1)$ is the smallest non-negative solution of the equation $x + 6y \equiv 0 \pmod{10}$.

Definition 2.2 Let (u, v) be the smallest non-negative solution of the equation (1). $(-a_1, a_2)$ is said to be a cross solution of the equation (1) if $-a_1 + a_2s \equiv 0 \pmod{n}$, $a_1 \geq 0$, $a_2 \geq 0$, $(-a_1, a_2) \neq (0, 0)$, and the three points $(-a_1, a_2)$, $(0, 0)$, (u, v) are not on the same line. $(-a, b)$ is said to be the smallest cross solution of the equation (1) if $(-a, b)$ is a cross solution of the equation (1) and the following conditions hold:

- (1) if $(-a_1, a_2)$ is a cross solution of the equation (1), then $a + b \leq a_1 + a_2$.
- (2) if $(-a_1, a_2)$ is a cross solution of the equation (1), where $(-a_1, a_2) \neq (-a, b)$ and $a + b = a_1 + a_2$, then $b > a_2$.

Clearly, the smallest cross solution of equation (1) is unique. For example, it is easy to see that $(2, 2)$ is the smallest non-negative solution of the equation $x + 5y \equiv 0 \pmod{12}$, and $(-5, 1), (-3, 3), (-1, 5), (-12, 0), (-10, 2), (-8, 4), (-6, 6), (-4, 8), (-2, 10), \dots$ are cross solutions of the equation $x + 5y \equiv 0 \pmod{12}$. Thus $(-1, 5)$ is the smallest cross solution of the equation $x + 5y \equiv 0 \pmod{12}$.

From [11] we have the following two lemmas.

Lemma 2.3 *Let (u, v) be the smallest non-negative solution of the equation (1) and $(-a, b)$ be the smallest cross solution of the equation (1) If $u < v$, then $a > u, a > b, b < v$ and $n=av + bu$.*

Lemma 2.4 *Let (u, v) be the smallest non-negative solution of the equation (1) and $(-a, b)$ be the smallest cross solution of the equation (1). If $u \geq v$, then $a < u, a \leq b, v < b$ and $n=av + bu$.*

Given $G(n; 1, s)$, we construct an infinite grid $G_{n,1,s}$ in \mathbb{Z}^2 as D. Tzvieli did in [19], labelling each lattice point (i, j) by $i + js \pmod{n}$. Every label $m, 0 \leq m < n$, is repeated in $G_{n,1,s}$ infinitely many times. We refer to a lattice point with label i as an i -point. If $i + js \equiv 0 \pmod{n}$, then we call (i, j) a 0-point.

The smallest non-negative solution (u, v) of the equation (1) can be seen as a 0-point in the first quadrant with $u + v$ is minimum (in case of tie, take the maximum u). The smallest cross solution $(-a, b)$ of the equation (1) can be seen as a 0-point in the second quadrant with $a + b$ is minimum (in case of tie, take the maximum b).

For $G(12; 1, 5)$, one can see that $(2, 2)$ is a 0-point in the first quadrant, and $(-1, 5)$ is a 0-point in the second quadrant(see Fig. 1).




Fig. 1 0-points in the $G_{12;1,5}$.

From Lemma 4 [10] and Lemma 5 [10] we have the following lemma:

Lemma 2.5 *Given $G(n; 1, s)$ with $1 < s < n$ and $s \neq \frac{n}{2}$. Suppose that (u, v) and $(-a, b)$ are two 0-points and u, v, a, b are all non-negative integers. Let $r_1 = \lfloor (u + v)/2 \rfloor, r_2 = \lfloor (a + b)/2 \rfloor, r_3 = \lfloor (|u - a| + v + b)/2 \rfloor$ and $r_4 = \lfloor (u + a + |v - b|)/2 \rfloor$. If $n = av + bu$ and $b \geq a, u \geq v, u > a, v < b$ (or $a > b, v > u, a > u, b < v$), then*

$$d(n; 1, s) = \begin{cases} r_3 - 1, & \text{if } r_3 = r_4 \text{ and } (u + a)(v + b) \equiv 1 \pmod{2} \\ \max\{r_1, r_2, \min\{r_3, r_4\}\}, & \text{otherwise.} \end{cases}$$

By Lemma 2.5, it is easy to prove the following corollary.

Corollary 2.6 Given $G(n; 1, s)$ with $1 < s < n$ and $s \neq \frac{n}{2}$. Suppose that (u, v) and $(-a, b)$ are two 0-points and u, v, a, b are all non-negative integers. Let d denote the diameter of $G(n; 1, s)$. If $n = av + bu$ and $b \geq a, u \geq v, u > a, v < b$ (or $a > b, v > u, a > u, b < v$), then $u + v \leq 2d + 1$ and $a + b \leq 2d + 1$.

Lemma 2.7 Suppose that u, v, a, b are all integers and there exist two integers λ, χ such that $\lambda b + \chi v = 1$. Let $s_0 \equiv \lambda a - \chi u$ and $s = s_0 \pmod{n}$. If $n = av + bu$, then $u + vs \equiv 0 \pmod{n}$ and $-a + bs \equiv 0 \pmod{n}$, i.e., (u, v) and $(-a, b)$ are two 0-points.

Proof: Since $\begin{pmatrix} b & v \\ a & -u \end{pmatrix} \begin{pmatrix} \lambda \\ \chi \end{pmatrix} = \begin{pmatrix} 1 \\ s_0 \end{pmatrix}$, we have $n \begin{pmatrix} \lambda \\ \chi \end{pmatrix} = \begin{pmatrix} u & v \\ a & -b \end{pmatrix} \begin{pmatrix} 1 \\ s_0 \end{pmatrix}$. That is, $u + vs \equiv u + vs_0 \pmod{n} = n\lambda \pmod{n} = 0$ and $-a + bs \equiv -a + bs_0 \pmod{n} = -n\chi \pmod{n} = 0$. \square

We use $\gcd(a_1, a_2)$ to denote the greatest common divisor of two integers a_1, a_2 . As the following lemma is well-known and easy to prove, its proof is omitted.

Lemma 2.8 Suppose that a_1, b_1 and c_1 are integers.

- (1) $\gcd(a_1, b_1) = 1$ if and only if there exist two integers λ, χ such that $\lambda a_1 + \chi b_1 = 1$.
- (2) If $\gcd(a_1, b_1) = 1$ and $\gcd(a_1, c_1) = 1$, then $\gcd(a_1, b_1 c_1) = 1$.

3 Main results

The sufficient and necessary conditions for a class of undirected double-loop networks to be optimal are given in the following theorem. By using this theorem, we can give some new optimal or suboptimal infinite families of undirected double-loop networks in this section.

Theorem 3.1 Suppose that $n = 2t^2 + 2t - B$, where $t > B > 0$, and $t, B \in \mathbb{Z}^+$. Then there exists a positive integer s such that $d(n; 1, s) = t = lb(n)$ if and only if there exist four non-negative integers a, b, u, v satisfying the following five conditions:

- 1 $n = av + bu$;
- 2 $a + b \leq 2t + 1$ and $u + v \leq 2t + 1$;
- 3 $b \geq a, u \geq v, u > a, v < b$;
- 4 One of the following two conditions holds:
 - (a) $x = 0, b - a = y, y$ is odd, $y \mid 2B + 1$ and $1 \leq y \leq \sqrt{2B + 1}$.
 - (b) $y = 0, u - v = x, x$ is odd, $x \mid 2B + 1$ and $1 \leq x \leq \sqrt{2B + 1}$. where $x = 2t + 1 - (a + b)$, $y = 2t + 1 - (u + v)$.
- 5 $\gcd(b, v) = 1$.

Proof: Suppose that there exists a positive integer s such that $d(n; 1, s) = t$. Let (u, v) be the smallest non-negative solution of the equation (1) and $(-a, b)$ be the smallest cross solution of the equation (1).

In the following we consider two cases: $u \geq v$ and $u < v$.

Case: $u \geq v$ Then by Lemma 2.4 and Corollary 2.6 , we see that (1), (2) and (3) hold.

Now we will prove that condition (4) holds.

Let $r_1 = \lfloor (u+v)/2 \rfloor, r_2 = \lfloor (a+b)/2 \rfloor, r_3 = \lfloor (u-a+v+b)/2 \rfloor$ and $r_4 = \lfloor (u+a+b-v)/2 \rfloor$.

If $r_3 = r_4$ and $(u+a)(v+b) \equiv 1 \pmod{2}$, then as $u-a+v+b \equiv 0 \pmod{2}$ and $u+a-v+b \equiv 0 \pmod{2}$, we have $u-a+v+b = u+a-v+b$, i.e., $a = v$. Thus the diameter of $G(n; 1, s)$ is $t = (u+b)/2 - 1$. So $b-a = b-v = (u+b) - (u+v) = 2t+2 - (2t+1-y) = 1+y$ and $u-v = u-a = (u+b) - (a+b) = 2t+2 - (2t+1-x) = 1+x$. As $(u+a)(v+b) \equiv 1 \pmod{2}$ and $a = v$, we have $x = 2t+1 - (a+b) = 2t+1 - (v+b) \equiv 0 \pmod{2}$ and $y = 2t+1 - (u+v) = 2t+1 - (u+a) \equiv 0 \pmod{2}$.

As $0 \leq (b-a)(u-v) = (1+y)(u-v) \leq (1+y)(u+v)$ and $(b-a)(u-v) = 2(av+bu) - (a+b)(u+v) = 2n - (2t+1-x)(2t+1-y)$, we have $2n - (2t+1-x)(2t+1-y) \leq (1+y)(u+v) = (1+y)(2t+1-y)$, i.e., $(2t+2-x+y)(2t+1-y) \geq 2n$. Thus we have $(y + \frac{1-x}{2})^2 + (x-2)(2t+1 - \frac{x}{4}) \leq \frac{1}{4} + 2B - 2t$. As $\frac{1}{4} + 2B - 2t < 0$ and $2t+1 - \frac{x}{4} > 0$, we have $x < 2$. Similarly we can prove that $y < 2$. Since $x \equiv 0 \pmod{2}, y \equiv 0 \pmod{2}, x \geq 0$ and $y \geq 0$, we have $x = y = 0$. As $b-a = 1+y = 1, u-v = 1+x = 1, a+b = 2t+1-x = 2t+1$ and $u+v = 2t+1-y = 2t+1$, we have $a = v = t, b = u = t+1$. Since $av+bu = t^2 + (t+1)^2 = 2t^2 + 2t+1$ and $n = av+bu$, we get a contradiction. Thus $r_3 = r_4$ and $(u+a)(v+b) \equiv 1 \pmod{2}$ can not hold simultaneously.

In other cases, the diameter of $G(n; 1, s)$ is $\max\{r_1, r_2, \min\{r_3, r_4\}\} = t$. Thus $r_3 \leq t$ or $r_4 \leq t$.

If $r_3 \leq t$, we have $b-a \leq y$. As $0 \leq (b-a)(u-v) \leq y(u+v)$ and $(b-a)(u-v) = 2(av+bu) - (a+b)(u+v) = 2n - (2t+1-x)(2t+1-y)$, we have $(2t+1-x+y)(2t+1-y) \geq 2n$, i.e., $4t^2 + (4-2x)t + (1-x+y)(1-y) \geq 2n$. Thus we have $(y - \frac{x}{2})^2 + (x-1)(2t+1 - \frac{x+1}{4}) \leq \frac{1}{4} + 2B - 2t$. As $\frac{1}{4} + 2B - 2t < 0$ and $2t+1 - \frac{x+1}{4} > 0$, we have $x < 1$. Thus $x = 0$ and $y \leq \sqrt{2B+1}$. As $x = 0$ and $0 \leq (b-a)(u-v) = -2B - 1 - xy + (x+y)(2t+1) = -2B - 1 + y(2t+1)$, we have $y \geq 1$. As $u-v = \frac{-2B-1+y(2t+1)}{b-a} = \frac{2t-2B+(y-1)(2t+1)}{b-a} \leq u+v = 2t+1-y$, we have $b-a > y-1$. Since $b-a \leq y$, we have $b-a = y$. As $b+a = 2t+1, b-a = b+a-2a$, we see that $y = b-a$ is odd. As $b-a = y$ and $(b-a)(u-v) = -2B - 1 + y(2t+1)$, we have $2B+1 = y(2t+1-u+v)$. Thus y divides $2B+1$, i.e., $y \mid 2B+1$.

If $r_4 \leq t$, we have $u-v \leq x$. As in the case of $r_3 \leq t$, we can prove that $y = 0, u-v = x, x$ is odd, $x \mid 2B+1$ and $1 \leq x \leq \sqrt{2B+1}$.

From above we see that condition (4) holds.

Since $u+vs \equiv 0 \pmod{n}$ and $-a+bs \equiv 0 \pmod{n}$, there exist two integers λ and χ such that $\begin{pmatrix} u & v \\ -a & b \end{pmatrix} \begin{pmatrix} 1 \\ s \end{pmatrix} = n \begin{pmatrix} \lambda \\ \chi \end{pmatrix}$. Thus $\begin{pmatrix} 1 \\ s \end{pmatrix} = \begin{pmatrix} b & -v \\ a & u \end{pmatrix} \begin{pmatrix} \lambda \\ \chi \end{pmatrix}$. So $\lambda b + (-\chi)v = 1$. By Lemma 2.8, we have $\gcd(v, b) = 1$. Thus condition (5) holds.

Case: $u < v$ By a similar argument we can deduce the following:

- (1) $n = av + bu$;
- (2) $a + b \leq 2t + 1$ and $u + v \leq 2t + 1$;
- (3) $a > b, v > u, a > u, b < v$;
- (4) One of the following two conditions holds:
 - (a) $x = 0, a - b = y, y$ is odd, $y \mid 2B + 1$ and $1 \leq y \leq \sqrt{2B + 1}$.

(b) $y = 0, v - u = x, x$ is odd, $x \mid 2B + 1$ and $1 \leq x \leq \sqrt{2B + 1}$. where $x = 2t + 1 - (a + b)$,
 $y = 2t + 1 - (u + v)$.

(5) $\gcd(b, v) = 1$.

By letting $a' = u, b' = v, u' = a, v' = b, x' = 2t + 1 - (a' + b') = y$ and $y' = 2t + 1 - (u' + v') = x$, it is routine to verify that a', b', u' and v' satisfy the five conditions of the theorem.

On the other hand, if there exist four non-negative integers a, b, u, v satisfying the five conditions of the theorem, then by condition 5 there exist two integers λ and χ such that $\lambda b + \chi v = 1$. Let $s \equiv \lambda a - \chi u \pmod{n}$. By Lemma 2.7, we see that (u, v) and $(-a, b)$ are two 0-points.

As $u \geq v, b \geq a, u > a$, and $v < b$, by Lemma 2.5 we have $d(n; 1, s) \leq t$. Since $d(n; 1, s) \geq lb(n) = t$, we have $d(n; 1, s) = t$. \square

Corollary 3.2 Suppose that $n = 2t^2 + 2t - B$, where $t > B > 0, t, B \in \mathbb{Z}^+$ and $2B + 1$ is a prime number. Then n is optimal if and only if $\gcd(t, B) = 1$ or $\gcd(t + 1, B) = 1$.

Proof: Suppose that n is optimal, by Theorem 3.1 there exist four non-negative integers a, b, u, v satisfying the condition 4 of Theorem 3.1. Let $x = 2t + 1 - (a + b)$ and $y = 2t + 1 - (u + v)$. Since $2B + 1$ is a prime number, we have (a) $x = 0, b - a = y = 1$ or (b) $y = 0, u - v = x = 1$.

When $x = 0, b - a = y = 1$, as $(b - a)(u - v) = 2(av + bu) - (a + b)(u + v) = 2n - (2t + 1) * 2t = 2t - 2B$, we have $a = t, b = t + 1, u = 2t - B$ and $v = B$. Since a, b, u, v satisfy the condition 5 of Theorem 3.1, we have $1 = \gcd(b, v) = \gcd(t + 1, B)$.

When $y = 0, u - v = x = 1$, as $(b - a)(u - v) = 2(av + bu) - (a + b)(u + v) = 2n - 2t * (2t + 1) = 2t - 2B$, we have $a = B, b = 2t - B, u = t + 1$ and $v = t$. Since a, b, u, v satisfy the condition 5 of Theorem 3.1, we have $1 = \gcd(b, v) = \gcd(2t - B, t) = \gcd(-B, t) = \gcd(t, B)$.

Thus we have proved that if n is optimal, then $\gcd(t, B) = 1$ or $\gcd(t + 1, B) = 1$.

If $\gcd(t, B) = 1$ or $\gcd(t + 1, B) = 1$, we will prove that n is optimal.

When $\gcd(t, B) = 1$, let $a = B, b = 2t - B, u = t + 1$ and $v = t$. It is routine to verify that four non-negative integers a, b, u, v satisfy the five conditions of Theorem 3.1.

When $\gcd(t + 1, B) = 1$, let $a = t, b = t + 1, u = 2t - B$ and $v = B$. It is easy to see that four non-negative integers a, b, u, v satisfy the five conditions of Theorem 3.1.

From above we see that Corollary 3.2 holds. \square

By Theorem 3.1 and Corollary 3.2, we can give some new optimal and suboptimal infinite families of undirected double-loop networks in the following theorems.

Theorem 3.3 (1) Let $\Theta = \{2t^2 + 2t - 6 \mid t > 6, t \neq 6e + 2 \text{ and } t \neq 6e + 3, e \in \mathbb{Z}^+\}$. Then Θ is an optimal infinite family, and when $t = 6e, 6e + 1, 6e + 4, 6e + 5$ respectively, the optimal step s is $12e^2, 12e^2 + 4e - 2, 60e^2 + 88e + 28, 60e^2 + 108e + 42$ correspondingly.

(2) $\{G(2t^2 + 2t - 6; 1, 2t^2 - 3) \mid t = 6e + 2 \text{ or } t = 6e + 3, 1 \leq e \in \mathbb{Z}^+\}$ is a suboptimal infinite family.

Proof:

(1) Let n be $2t^2 + 2t - 6$. In the following we consider two cases: $t = 6e + 1$ or $t = 6e + 5$ and $t = 6e$ or $t = 6e + 4$.

$t = 6e + 1$ or $t = 6e + 5$ When $t = 6e + 1$ or $t = 6e + 5$, we have $\gcd(t, 6) = 1$.

$t = 6e$ or $t = 6e + 4$ When $t = 6e$ or $t = 6e + 4$, we have $\gcd(t + 1, 6) = 1$.

As $2 \times 6 + 1 = 13$ and 13 is a prime number, by Corollary 3.2, we see that Θ is an optimal infinite family.

When $t = 6e + 1$, let $a = 6, b = 2t - 6, u = t + 1$, and $v = t$. As $e \times b + (-2e + 1) \times v = 1$, the optimal step s is $e \times a - (-2e + 1) \times u \pmod{n} = 12e^2 + 4e - 2$.

When $t = 6e + 5$, let $a = 6, b = 2t - 6, u = t + 1$, and $v = t$. As $-(e + 1) \times b + (2e + 1) \times v = 1$, $s = -(e + 1) \times a - (2e + 1) \times u \pmod{n} = 60e^2 + 108e + 42$.

When $t = 6e$, let $a = t, b = t + 1, u = 2t - 6$, and $v = 6$. As $1 \times b - e \times v = 1$, $s = 1 \times a - (-e) \times u \pmod{n} = 12e^2$.

When $t = 6e + 4$, let $a = t, b = t + 1, u = 2t - 6$, and $v = 6$. As $-1 \times b + (e + 1) \times v = 1$, $s = -1 \times a - (e + 1) \times u \pmod{n} = 60e^2 + 88e + 28$.

(2) When $t = 6e + 2$, we have $\gcd(t + 1, 6) = \gcd(6e + 3, 6) = \gcd(3, 6) = 3$ and $\gcd(t, 6) = \gcd(6e + 2, 6) = \gcd(2, 6) = 2$.

When $t = 6e + 3$, we have $\gcd(t + 1, 6) = \gcd(6e + 4, 6) = \gcd(4, 6) = 2$ and $\gcd(t, 6) = \gcd(6e + 3, 6) = \gcd(3, 6) = 3$.

As $2 \times 6 + 1 = 13$ and 13 is a prime number, by Corollary 3.2, we see that when $t = 6e + 2$ or $t = 6e + 3$, $1 \leq e \in \mathbb{Z}^+$, $n = 2t^2 + 2t - 6$ can not be optimal.

Let $a = t, b = t + 2, u = 2t - 3$ and $v = 1$. As $0 \times b + 1 \times v = 1$, let $s = 0 \times a - 1 \times u \equiv 2t^2 - 3 \pmod{n}$. By Lemma 2.7, we see that $(2t - 3, 1)$ and $(-t, t + 2)$ are two 0-points. By Lemma 2.5 we have $d(n; 1, s) = t + 1$.

From above we see that when $t = 6e + 2$ or $t = 6e + 3$, $1 \leq e \in \mathbb{Z}^+$, $G(2t^2 + 2t - 6; 1, 2t^2 - 3)$ is suboptimal. \square

Theorem 3.4 1. Let $n = 2t^2 + 2t - 12$ and $\Theta = \{2t^2 + 2t - 12 \mid t > 12, t \neq 6e + 2 \text{ and } t \neq 6e + 3, e \in \mathbb{Z}^+\}$. Then Θ is an optimal infinite family, and when $t = 6e, 6e + 1, 6e + 4, 6e + 5$ respectively, the optimal step s is $6e - 36e^3 \pmod{n}, 36e^3 + 24e^2 - 2e - 2 \pmod{n}, -36e^3 - 96e^2 - 78e - 16 \pmod{n}, 36e^3 + 96e^2 + 78e + 18 \pmod{n}$ correspondingly.

2. Let $\Psi = \{2t^2 + 2t - 12 \mid t = 6e + 2 \text{ or } t = 6e + 3, \text{ where } e \in \mathbb{Z}^+ \text{ and } e \geq 2\}$. Then Ψ is a suboptimal infinite family, and when $t = 6e + 2$ or $6e + 3$, the suboptimal step s is $t^2 + 2t - 3, t^2 - 4$ correspondingly.

Proof:

(1) By Theorem 3.1, we only need to find four non-negative integers a, b, u, v which satisfy the five conditions of Theorem 3.1. In the following we consider two cases: $t = 6e$ or $t = 6e + 4$, and $t = 6e + 1$ or $t = 6e + 5$.

$t = 6e$ or $t = 6e + 4$ When $t = 6e$ or $t = 6e + 4$, let $a = t, b = t + 1, u = 2t - 12$ and $v = 12$.

When $t = 6e$ or $t = 6e + 4$, As $\gcd(2, t + 1) = 1$ and $\gcd(3, t + 1) = 1$, by Lemma 2.8, we have that $\gcd(4, t + 1) = 1$ and $\gcd(12, t + 1) = 1$. That is, $\gcd(v, b) = 1$. Thus it is easy to verify that four integers a, b, u, v satisfy the five conditions of Theorem 3.1.

$t = 6e + 1$ or $t = 6e + 5$ When $t = 6e + 1$ or $t = 6e + 5$, let $a = 12, b = 2t - 12, u = t + 1$ and $v = t$.

When $t = 6e + 1$ or $t = 6e + 5$, As $\gcd(2, t) = 1$ and $\gcd(3, t) = 1$, by Lemma 2.8, we have that $\gcd(4, t) = 1$ and $\gcd(12, t) = 1$. Thus $\gcd(v, b) = \gcd(t, 2t - 12) = \gcd(t, -12) = \gcd(t, 12) = 1$. So it is easy to verify that four integers a, b, u, v satisfy the five conditions of Theorem 3.1.

From above we see that Θ is an optimal infinite family.

When $t = 6e$, let $a = t, b = t + 1, u = 2t - 12$, and $v = 12$. As $(1 - 6e) \times b + 3e^2 \times v = 1$, $s = (1 - 6e) \times a - 3e^2 \times u \pmod{n} = 6e - 36e^3 \pmod{n}$.

When $t = 6e + 1$ or $6e + 4$ or $6e + 5$, the corresponding optimal step s can be computed similarly. They are $36e^3 + 24e^2 - 2e - 2 \pmod{n}$, $-36e^3 - 96e^2 - 78e - 16 \pmod{n}$, $36e^3 + 96e^2 + 78e + 18 \pmod{n}$ respectively.

(2) For $n = 2t^2 + 2t - 12$, where $t = 6e + 2$ or $t = 6e + 3, 2 \leq e \in \mathbb{Z}^+$, in the following we will prove that n can not be optimal.

If n is optimal, then there exist four non-negative integers a, b, u, v which satisfy the five conditions of Theorem 3.1. Let $x = 2t + 1 - (a + b)$ and $y = 2t + 1 - (u + v)$.

In the following we consider two cases:

(A) $x = 0, b - a = y, y$ is odd, $y \mid 2B + 1$ and $1 \leq y \leq \sqrt{2B + 1}$.

(B) $y = 0, u - v = x, x$ is odd, $x \mid 2B + 1$ and $1 \leq x \leq \sqrt{2B + 1}$, where $B = 12$.

Case A If $x = 0, b - a = y, y$ is odd, $y \mid 25$ and $1 \leq y \leq \sqrt{25}$, then $y = 1$ or $y = 5$.

Subcase: $x = 0, y = 1$: As $b - a = y$ and $(b - a)(u - v) = 2n - (a + b)(u + v) = 2n - (2t + 1 - x)(2t + 1 - y) = 2t - 24$, we have that $b - a = 1$ and $u - v = 2t - 24$. Thus $a = t, b = t + 1, u = 2t - 12$ and $v = 12$. When $t = 6e + 2$ or $t = 6e + 3, \gcd(b, v) \neq 1$.

Subcase: $x = 0, y = 5$: As $b - a = y$ and $(b - a)(u - v) = 2n - (a + b)(u + v) = 2n - (2t + 1 - x)(2t + 1 - y) = 10t - 20$, we have that $b - a = 5$ and $u - v = 2t - 4$. Thus $a = t - 2, b = t + 3, u = 2t - 4$ and $v = 0$. So $\gcd(b, v) = t + 3 \neq 1$.

Case B If $y = 0, u - v = x, x$ is odd, $x \mid 25$ and $1 \leq x \leq \sqrt{25}$, then $x = 1$ or $x = 5$.

Subcase: $y = 0, x = 1$: As $u - v = x$ and $(b - a)(u - v) = 2n - (a + b)(u + v) = 2n - (2t + 1 - x)(2t + 1 - y) = 2t - 24$, we have $u - v = 1$ and $b - a = 2t - 24$. Thus $a = 12, b = 2t - 12, u = t + 1$ and $v = t$. When $t = 6e + 2$ or $t = 6e + 3, \gcd(b, v) \neq 1$.

Subcase: $y = 0, x = 5$: As $u - v = x$ and $(b - a)(u - v) = 2n - (a + b)(u + v) = 2n - (2t + 1 - x)(2t + 1 - y) = 10t - 20$, we have that $u - v = 5$ and $b - a = 2t - 4$. Thus $a = 0, b = 2t - 4, u = t + 3$ and $v = t - 2$. So $\gcd(b, v) = \gcd(2t - 4, t - 2) = t - 2 \neq 1$.

From above we see that in any cases, two non-negative integers b, v can not satisfy condition 5 of Theorem 3.1. So n can not be optimal.

When $t = 6e + 2$, let $a = 0, b = 2t - 4, u = t + 3$ and $v = t - 3$. As $\frac{4-t}{2} \times b + (t - 3) \times v = 1$, let $s = \frac{4-t}{2} \times a - (t - 3) \times u \equiv 9 - t^2 \pmod{n} = t^2 + 2t - 3$. By Lemma 2.7, we see that $(t + 3, t - 3)$ and $(0, 2t - 4)$ are two 0-points. By Lemma 2.5 we have $d(n; 1, s) = t + 1$.

When $t = 6e + 3$, let $a = t - 2, b = t + 2, u = 2t - 4$ and $v = 2$. As $1 \times b - \frac{t+1}{2} \times v = 1$, let $s = 1 \times a + \frac{t+1}{2} \times u \pmod{n} = t^2 - 4$. By Lemma 2.7, we see that $(2t - 4, 2)$ and $(-t + 2, t + 2)$ are two 0-points. By Lemma 2.5 we have $d(n; 1, s) = t + 1$.

Thus when $t = 6e + 2$ or $t = 6e + 3, 2 \leq e \in \mathbb{Z}^+, 2t^2 + 2t - 12$ is suboptimal. \square

Theorem 3.5 Let $\Theta = \{2t^2 + 2t - (2A^2 + 2A - 2) \mid A > 0, A \in \mathbb{Z}^+, t \geq \frac{A^2 + A}{2}\}$. Then Θ is an optimal infinite family. When $t + A$ is odd, the optimal step s is $t^2 - A^2 + 1$. When $t + A$ is even, the optimal step s is $t^2 - A^2 - 2A$.

Proof: Let $n = 2t^2 + 2t - (2A^2 + 2A - 2)$. When $A > 0, A \in \mathbb{Z}^+, t \geq \frac{A^2 + A}{2}, lb(n) = t$.

In the following we consider two cases:

$t + A$ is an odd number When $t + A$ is odd, suppose that $t + A = 2j + 1$. Let $a = t - A + 1, b = t + A, u = 2t - 2A$ and $v = 2$. As $1 \times b - jv = 1$, let $s \equiv 1 \times (t - A + 1) + j(2t - 2A) \pmod{n} = t^2 - A^2 + 1$. By Lemma 2.7, we see that (u, v) and $(-a, b)$ are two 0-points. By Lemma 2.5 we have $d(n; 1, s) = t$.

$t + A$ is an even number When $t + A$ is even, let $a = 2, b = 2t - 2A, u = t + A$ and $v = t - A + 1$. Since $\frac{t-A}{2}b + (1-t+A)v = 1$, let $s \equiv \frac{t-A}{2}a - (1-t+A)u \pmod{n} = t^2 - A^2 - 2A$. By Lemma 2.7, we see that (u, v) and $(-a, b)$ are two 0-points. By Lemma 2.5 we have $d(n; 1, s) = t$.

Thus when $t \geq \frac{A^2 + A}{2}, 0 < A \in \mathbb{Z}^+, 2t^2 + 2t - (2A^2 + 2A - 2)$ is optimal. \square

By Theorem 3.5, it is easy to prove the following corollary.

Corollary 3.6 Let $\Gamma = \{2t^2 - 2 \mid t = A^2 + A - 2, A \geq 2, A \in \mathbb{Z}^+\}$. Then Γ is an optimal infinite family. When A is odd, the optimal step s is $t^2 - A^2 + 1$. When A is even, the optimal step s is $t^2 - A^2 - 2A$.

By APPENDIX B [19], we see that $390 = 2 \times 14^2 - 2$ is suboptimal. In other words, there are some suboptimal integers in the set $\Sigma = \{2t^2 - 2 \mid t > 2, t \in \mathbb{Z}^+\}$. From Corollary 3.6, we see that the set Γ , which is a subset of the set Σ , is optimal. It would be interesting to find out a subset of the set Σ which is a suboptimal infinite family.

Theorem 3.7 (1) Let $n = 2t^2 + 2t - 14$ and $\Theta = \{2t^2 + 2t - 14 \mid t > 14, t \neq 14e + 6 \text{ and } t \neq 14e + 7, e \in \mathbb{Z}^+\}$. Then Θ is an optimal infinite family, and when $t = 14e, 14e + 1, 14e + 2, 14e + 3, 14e + 4, 14e + 5, 14e + 8, 14e + 9, 14e + 10, 14e + 11, 14e + 12, 14e + 13$ respectively, the optimal step s is $28e^2, 28e^2 + 4e - 2, 140e^2 + 48e, 140e^2 + 68e + 2, 84e^2 + 52e + 6, 84e^2 + 64e + 8, -84e^2 - 104e - 28 \pmod{n}, -84e^2 - 116e - 38 \pmod{n}, -140e^2 - 212e - 74 \pmod{n}, -140e^2 - 232e - 92 \pmod{n}, -28e^2 - 52e - 22 \pmod{n}, -28e^2 - 56e - 28 \pmod{n}$ correspondingly.

(2) Let $\Psi = \{2t^2 + 2t - 14 \mid t = 14e + 6 \text{ or } t = 14e + 7, \text{ where } e \in \mathbb{Z}^+ \text{ and } e \geq 1\}$. Then Ψ is a suboptimal infinite family, and when $t = 14e + 6$ or $14e + 7$, the suboptimal step s is $t^2 + 2t - 4, t^2 - 5$ correspondingly.

Proof:

(1) In the following we consider two cases:

(A) $t = 14e$, or $t = 14e + 2$, or $t = 14e + 4$, or $t = 14e + 8$, or $t = 14e + 10$, or $t = 14e + 12$;

(B) $t = 14e + 1$, or $t = 14e + 3$, or $t = 14e + 5$, or $t = 14e + 9$, or $t = 14e + 11$, or $t = 14e + 13$.

Case A When $t = 14e$, or $t = 14e + 2$, or $t = 14e + 4$, or $t = 14e + 8$, or $t = 14e + 10$, or $t = 14e + 12$, we have $\gcd(t + 1, 14) = 1$.

Case B When $t = 14e + 1$, or $t = 14e + 3$, or $t = 14e + 5$, or $t = 14e + 9$, or $t = 14e + 11$, or $t = 14e + 13$, we have $\gcd(t, 14) = 1$.

As $2 \times 14 + 1 = 29$ and 29 is a prime number, by Corollary 3.2, we see that Θ is an optimal infinite family.

When $t = 14e$, let $a = t, b = t + 1, u = 2t - 14$, and $v = 14$. As $1 \times b - e \times v = 1, s = 1 \times a + e \times u \pmod{n} = 28e^2$.

When $t = 14e + 1$ or $14e + 2$ or $14e + 3$ or $14e + 4$ or $14e + 5$ or $14e + 8$ or $14e + 9$ or $14e + 10$ or $14e + 11$ or $14e + 12$ or $14e + 13$, the corresponding optimal step s can be computed similarly. They are $28e^2 + 4e - 2, 140e^2 + 48e, 140e^2 + 68e + 2, 84e^2 + 52e + 6, 84e^2 + 64e + 8, -84e^2 - 104e - 28 \pmod{n}, -84e^2 - 116e - 38 \pmod{n}, -140e^2 - 212e - 74 \pmod{n}, -140e^2 - 232e - 92 \pmod{n}, -28e^2 - 52e - 22 \pmod{n}, -28e^2 - 56e - 28 \pmod{n}$ respectively.

(2) When $t = 14e + 6$, we have $\gcd(t, 14) = \gcd(14e + 6, 14) = \gcd(6, 14) = 2$ and $\gcd(t + 1, 14) = \gcd(14e + 7, 14) = \gcd(7, 14) = 7$.

When $t = 14e + 7$, we have $\gcd(t, 14) = \gcd(14e + 7, 14) = 7$ and $\gcd(t + 1, 14) = \gcd(14e + 8, 14) = \gcd(8, 14) = 2$.

As $2 \times 14 + 1 = 29$ and 29 is a prime number, by Corollary 3.2, we see that when $t = 14e + 6$ or $t = 14e + 7, 1 \leq e \in \mathbb{Z}^+, n = 2t^2 + 2t - 14$ can not be optimal.

When $t = 14e + 6$, let $a = 2, b = 2t - 4, u = t + 2$ and $v = t - 3$. As $\frac{4-t}{2} \times b + (t - 3) \times v = 1$, let $s = \frac{4-t}{2} \times a - (t - 3) \times u \equiv 10 - t^2 \pmod{n} = t^2 + 2t - 4$. By Lemma 2.7, we see that $(t + 2, t - 3)$ and $(-2, 2t - 4)$ are two 0-points. By Lemma 2.5, we have $d(n; 1, s) = t + 1$.

When $t = 14e + 7$, let $a = t - 3, b = t + 2, u = 2t - 4$ and $v = 2$. As $1 \times b - \frac{t+1}{2} \times v = 1$, let $s \equiv 1 \times a + \frac{t+1}{2} \times u \pmod{n} = t^2 - 5$. By Lemma 2.7, we see that $(2t - 4, 2)$ and $(-t + 3, t + 2)$ are two 0-points. Thus by Lemma 2.5 we have $d(n; 1, s) = t + 1$.

From above we see that when $t = 14e + 6$ or $t = 14e + 7$, where $1 \leq e \in \mathbb{Z}^+, 2t^2 + 2t - 14$ is suboptimal. \square

Theorem 3.8 1. Let $\Theta = \{2t^2 + 2t - 15 \mid t > 15, t \neq 15e + 5 \text{ and } t \neq 15e + 9, e \in \mathbb{Z}^+\}$. Then Θ is an optimal infinite family, and when $t = 15e, 15e + 1, 15e + 2, 15e + 3, 15e + 4, 15e + 6, 15e + 7, 15e + 8, 15e + 10, 15e + 11, 15e + 12, 15e + 13, 15e + 14$ respectively, the optimal step s is $30e^2, 240e^2 + 46e - 5, 240e^2 + 78e - 3, 120e^2 + 54e + 3, 120e^2 + 70e + 5, -60e^2 - 54e - 9 \pmod{n}, 60e^2 + 58e + 13, 60e^2 + 66e + 15, 330e^2 + 460e + 150, 330e^2 + 504e + 180, 210e^2 + 348e + 138, -30e^2 - 56e - 24 \pmod{n}, -30e^2 - 60e - 30 \pmod{n}$ correspondingly.

2. $\{G(2t^2 + 2t - 15; 1, 2t^2 - 10) \mid t = 15e + 5 \text{ or } t = 15e + 9, \text{ where } 1 \leq e \in \mathbb{Z}^+\}$ is a suboptimal infinite family.

Proof:

(1) In the following we consider two cases:

(A) $t = 15e$, or $t = 15e + 1$, or $t = 15e + 3$, or $t = 15e + 6$, or $t = 15e + 7$, or $t = 15e + 10$, or $t = 15e + 12$, or $t = 15e + 13$;

(B) $t = 15e + 2$, or $t = 15e + 4$, or $t = 15e + 8$, or $t = 15e + 11$, or $t = 15e + 14$.

Case A When $t = 15e$, or $t = 15e + 1$, or $t = 15e + 3$, or $t = 15e + 6$, or $t = 15e + 7$, or $t = 15e + 10$, or $t = 15e + 12$, or $t = 15e + 13$, we have $\gcd(t + 1, 15) = 1$.

Case B When $t = 15e + 2$, or $t = 15e + 4$, or $t = 15e + 8$, or $t = 15e + 11$, or $t = 15e + 14$, we have $\gcd(t, 15) = 1$.

As $2 \times 15 + 1 = 31$ and 31 is a prime number, by Corollary 3.2, we see that Θ is an optimal infinite family.

When $t = 15e$, let $a = t, b = t + 1, u = 2t - 15$, and $v = 15$. As $1 \times b - e \times v = 1, s = 1 \times a + e \times u \pmod{n} = 30e^2$.

When $t = 15e + 1$ or $15e + 2$ or $15e + 3$ or $15e + 4$ or $15e + 6$ or $15e + 7$ or $15e + 8$ or $15e + 10$ or $15e + 11$ or $15e + 12$ or $15e + 13$ or $15e + 14$, the corresponding optimal step s can be computed similarly. They are $240e^2 + 46e - 5, 240e^2 + 78e - 3, 120e^2 + 54e + 3, 120e^2 + 70e + 5, -60e^2 - 54e - 9 \pmod{n}, 60e^2 + 58e + 13, 60e^2 + 66e + 15, 330e^2 + 460e + 150, 330e^2 + 504e + 180, 210e^2 + 348e + 138, -30e^2 - 56e - 24 \pmod{n}, -30e^2 - 60e - 30 \pmod{n}$ respectively.

(2) When $t = 15e + 5$, we have $\gcd(t, 15) = \gcd(15e + 5, 15) = \gcd(5, 15) = 5$ and $\gcd(t + 1, 15) = \gcd(15e + 6, 15) = \gcd(6, 15) = 3$.

When $t = 15e + 9$, we have $\gcd(t, 15) = \gcd(15e + 9, 15) = \gcd(9, 15) = 3$ and $\gcd(t + 1, 15) = \gcd(15e + 10, 15) = \gcd(10, 15) = 5$.

As $2 \times 15 + 1 = 31$ and 31 is a prime number, by Corollary 3.2, we see that when $t = 15e + 5$ or $t = 15e + 9, 1 \leq e \in \mathbb{Z}^+, n = 2t^2 + 2t - 15$ can not be optimal.

When $t = 15e + 5$ or $t = 15e + 9$, let $a = t, b = t + 3, u = 2t - 5$ and $v = 1$. As $0 \times b + 1 \times v = 1$, let $s = 0 \times a - 1 \times u \equiv 5 - 2t \pmod{n} = 2t^2 - 10$. By Lemma 2.7, we see that $(2t - 5, 1)$ and $(-t, t + 3)$ are two 0-points. By Lemma 2.5, we have $d(n; 1, s) = t + 1$.

From above we see that when $t = 15e + 5$ or $t = 15e + 9$, where $1 \leq e \in \mathbb{Z}^+, G(2t^2 + 2t - 15; 1, 2t^2 - 10)$ is suboptimal. \square

Acknowledgements

The authors thank the referees for their helpful comments that improved the accuracy and clarity of our presentation.

References

- [1] F. Aguiló and M. A. Fiol, An efficient algorithm to find optimal double loop networks, *Discrete Mathematics* 138(1995), 15-29.
- [2] J. -C. Bermond and D. Tzvieli, Minimal diameter double-loop networks: dense optimal families, *Networks* 21(1991), 1-9.
- [3] J. -C. Bermond, F. Comellas and D. F. Hsu, Distributed loop computer networks: a survey, *Journal of Parallel and Distributed Computing* 24(1995), 2-10.
- [4] F. T. Boesch and J. F. Wang, Reliable circulant networks with minimum transmission delay, *IEEE Trans. Circuits Syst. CAS-32*(1985), 1286-1291.

- [5] N. Chalamaiah and B. Ramamurty, Finding shortest paths in distributed loop networks, *Information Processing Letters* 67(1998), 157-161.
- [6] R. C.-F. Chan, C.-Y. Chen, Z.-X. Hong, A simple algorithm to find the steps of double-loop networks, *Discrete Applied Mathematics* 121(2002), 61-72.
- [7] B. X. Chen, J. X. Meng and W. J. Xiao, A constant time optimal routing algorithm for undirected double-loop networks, *International Conference on Mobile Ad-hoc and Sensor Networks, Wuhan, China, Lecture Notes in Computer Science (LNCS), Springer Verlag , 3794(2005), 308-316.*
- [8] B. X. Chen and W. J. Xiao, A constant time optimal routing algorithm for directed double loop networks $G(n; s_1, s_2)$, In the proceeding of 5th International Conference on Software Engineering, Artificial Intelligence, Networking, and Parallel/Distributed Computing(SNPD 2004), 1-5.
- [9] B. X. Chen and W. J. Xiao, Optimal designs of directed double-loop networks, *International Symposium on Computational and Information Sciences (CIS'04), Lecture Notes in Computer Science (LNCS), Springer Verlag, 3314(2004), 19-24.*
- [10] B. X. Chen and W. J. Xiao, B. Parhami, Diameter Formulas for a Class of Undirected Double-loop networks, *Journal of Interconnection Networks* 6(2005), 1: 1-15.
- [11] B. X. Chen and W. J. Xiao, A diameter formula for an undirected double-loop network, Accepted by *Ars Combinatoria*.
- [12] D. Z. Du, D. F. Hsu, Li Qiao and Xu Jun-ming, A combinatorial problem related to distributed loop networks, *Networks* 20(1990), 173-180.
- [13] P. Esqué, F. Aguiló and M. A. Fiol, Double commutative-step digraphs with minimum diameters, *Discrete Mathematics* 114(1993), 147-157.
- [14] D. F. Hsu and J. Shapiro, Bounds for the minimal number of transmission delays in double loop networks, *Journal of Combinatorics, Information & System Sciences* 16(1991), 55-62.
- [15] F. K. Hwang and Y. H. Xu, Double loop networks with minimum delay, *Discrete Mathematics* 66(1987), 109-118.
- [16] F. K. Hwang, A complementary survey on double-loop networks, *Theoretical Computer Science* 263(2001), 211-229.
- [17] K. Mukhopadhyaya and B. P. Sinha, Fault-tolerant routing in distributed loop networks, *IEEE Transactions on Computers* 44(1995), 12:1452-1456.
- [18] Q. Li, J. M. Xu and Z. L. Zhang, Infinite families of optimal double loop networks, *Science in China, Ser A* 23(1993), 979-992.
- [19] D. Tzvieli, Minimal diameter double-loop networks I. Large infinite Optimal families, *Networks* 21(1991), 387-415.
- [20] C. K. Wong and D. Coppersmith, A combinatorial problem related to multimodule memory organizations, *Journal of the Association for Computing Machinery* 21(1974), 3:392-402.

- [21] J. A. L. Yebra, M. A. Fiol, P. Morillo and I. Alegre, The diameter of undirected graphs associated to plane tessellations, *Ars Combinatoria* 20-B(1985), 151-171.
- [22] J. Zerovnik and T. Pisanski, Computing the diameter in multi-loop networks, *Journal of Algorithm* 14(1993), 226-243.

