

Long cycles in hypercubes with distant faulty vertices

Petr Gregor, Riste Škrekovski

► **To cite this version:**

Petr Gregor, Riste Škrekovski. Long cycles in hypercubes with distant faulty vertices. *Discrete Mathematics and Theoretical Computer Science, DMTCS*, 2009, 11 (1), pp.185–198. <hal-00988176>

HAL Id: hal-00988176

<https://hal.inria.fr/hal-00988176>

Submitted on 7 May 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Long cycles in hypercubes with distant faulty vertices

Petr Gregor^{1,†} and Riste Škrekovski^{2,‡}

¹*Department of Theoretical Computer Science and Mathematical Logic, Charles University, Malostranské nám. 25, 118 00 Prague, Czech Republic.*

²*Department of Mathematics, University of Ljubljana, Jadranska 21, 1000 Ljubljana, Slovenia.*

received 21st July 2008, revised 17th March 2009, accepted 25th March 2009.

In this paper, we study long cycles in induced subgraphs of hypercubes obtained by removing a given set of faulty vertices such that every two faults are distant. First, we show that every induced subgraph of Q_n with minimum degree $n - 1$ contains a cycle of length at least $2^n - 2f$ where f is the number of removed vertices. This length is the best possible when all removed vertices are from the same bipartite class of Q_n . Next, we prove that every induced subgraph of Q_n obtained by removing vertices of some given set M of edges of Q_n contains a Hamiltonian cycle if every two edges of M are at distance at least 3. The last result shows that the shell of every linear code with odd minimum distance at least 3 contains a Hamiltonian cycle. In all these results we obtain significantly more tolerable faulty vertices than in the previously known results. We also conjecture that every induced subgraph of Q_n obtained by removing a balanced set of vertices with minimum distance at least 3 contains a Hamiltonian cycle.

Keywords: hypercube, faulty vertex, Hamiltonian cycle, long cycle, linear code

1 Introduction

Given a set U of binary strings of length n , is there a (cyclic) ordering of U such that every two consecutive strings differ in a single bit? This can be viewed as a problem of Hamiltonian paths (cycles) in the induced subgraph of the n -dimensional hypercube Q_n obtained by removing a given set $F = V(Q_n) \setminus U$ of faulty vertices. Clearly, since Q_n is bipartite and balanced in the sense that it contains the same number of vertices in each bipartite class (called parity), it follows that F must be balanced as well (almost-balanced for Hamiltonian paths between vertices of the same parity).

If there are no restrictions on U , this problem was recently proved to be NP-complete [4]. On the other hand, if there are only few faulty vertices, the answer to this problem is positive. In particular, it is well-known that Q_n for any $n \geq 1$ has a Hamiltonian path between every two vertices of opposite parity. The corresponding (cyclic) ordering of $V(Q_n) = \{0, 1\}^n$ is called a (cyclic) Gray code, named after Frank

[†]Supported in part by the Czech Science Foundation Grant 201/08/P298 and by ARRS Research Program P-0297.

[‡]Supported in part by the Czech-Slovenian bilateral grant MEB 080905 and by KAM, Charles University in Prague.

Gray who patented a scheme to generate such code [7]. These codes have applications in many areas such as data compression, image processing, information retrieval, signal encoding, processor allocation in hypercubic networks. More on Gray codes can be found in a survey by Savage [11].

As for other previous results, Lewinter and Widulski [8] showed that for every vertex u , the graph $Q_n - \{u\}$ contains a Hamiltonian path between every two vertices of the parity other than the parity of u . Then, Locke and Stong [9] showed that $Q_n - \{u, v\}$ has a Hamiltonian cycle for every two vertices u and v of opposite parity. Furthermore, by result due to Dvořák and Gregor [4], $Q_n - F$ has a Hamiltonian cycle for every balanced set $F \subseteq V(Q_n)$ with $|F| \leq (n - 5)/3$. It is also claimed (without a proof) in [9] that this bound can be improved to $|F| + 3 \log_2 |F| + 1 \leq n$. Note that the best possible upper bound would be $|F| \leq 2n - 3$ since there is a balanced set F of $2n - 2$ vertices of Q_n such that $Q_n - F$ contains a vertex of degree 1.

In this paper we try to significantly improve the number of tolerable faulty vertices under some additional condition on their structure. A similar approach has been successfully applied for the analogous problem with faulty edges in [3] where it has been characterized for every set M of independent edges (matching) of Q_n when $Q_n - M$ contains a Hamiltonian cycle.

The considered condition on the faulty vertices is that they are not close to each other. For $F \subseteq V(Q_n)$ let $d(F)$ denote the minimum (Hamming) distance between every two vertices of F . This parameter plays an important role in the design of error-correcting codes. If we encode a message into a sequence of blocks from $F \subseteq V(Q_n)$, then we can correct any up to $(d(F) - 1)/2$ errors occurring in each block. For more information on error-correcting codes, we refer to the monograph by MacWilliams and Sloane [10].

Note that the above mentioned balanced set F of $2n - 2$ vertices of Q_n has $d(F) \leq 2$. Thus, it is natural to require, apart from the balance of F , that $d(F) \geq 3$. We conjecture that these necessary conditions are also sufficient.

Conjecture 1 *Let F be a balanced set of vertices of Q_n with $d(F) \geq 3$. Then $Q_n - F$ contains a Hamiltonian cycle.*

A stronger form of the above conjecture requires only vertices of a same parity to be distant. Note that the minimum distance 3 is now replaced by 4 since every two vertices of the same parity are at even distance.

Conjecture 2 *Let A and B be equal-sized sets of even and odd (respectively) vertices of Q_n with $d(A) \geq 4$ and $d(B) \geq 4$. Then $Q_n - (A \cup B)$ contains a Hamiltonian cycle.*

The above conjectures are stated in the strongest possible form regarding the prescribed minimum distance between vertices. Although weaker, it is still interesting to establish variants of the above conjectures when the set of vertices have higher distance, i.e. replacing the minimum distances 3 and 4 by bigger integers.

To support Conjecture 1, we have three partial results indicating that it should hold. In Section 3 we study long cycles in induced subgraphs G of Q_n with minimum degree $\delta(G) \geq n - 1$.

Theorem 1 *Let G be an induced subgraph of Q_n ($n \geq 1$) with $\delta(G) \geq n - 1$. Then G contains a cycle of length at least $2|V(G)| - 2^n$.*

Note that $d(F) \geq 3$ where $F = V(Q_n) \setminus V(G)$ implies that $\delta(G) \geq n - 1$ but not vice-versa. Thus, Theorem 1 considers more general sets than Conjecture 1. Moreover, the set F can be unbalanced so

Theorem 1 can only find shorter cycles than Hamiltonian cycles. But in the particular case when F contains only vertices of the same parity, the length of the cycle obtained by Theorem 1 is the best possible.

Cycles of at least this length have been previously found by Fu [6] in Q_n with any set F of faulty vertices such that $|F| \leq 2n - 5$. Then, this bound was improved by Castañeda and Gotchev [1] to $|F| \leq 3n - 7$ for $n \geq 5$. Recently, it was further increased by Fink and Gregor [5] to $|F| \leq n^2/10 + n/2 + 1$ for $n \geq 15$ which is also asymptotically optimal (if there are no restrictions on F). However, in Theorem 1 we may have a nontrivial set F of up to 2^{n-1} vertices.

In Section 4 we study induced subgraphs of Q_n obtained by removing the vertices of a given set of edges with minimum distance $d(M) \geq 3$. Note that we state the following theorem for the dimension $n + 2$ instead of n only for technical reasons.

Theorem 2 *Let M be a set of edges of Q_{n+2} ($n \geq 1$) with $d(M) \geq 3$. Then $Q_{n+2} - V(M)$ contains a Hamiltonian cycle.*

Clearly, for every (nontrivial) set M of edges of Q_n , the condition $d(M) \geq 3$ is equivalent to the condition that the graph $G = Q_n - V(M)$ has minimum degree $\delta(G) \geq n - 1$ and its complement in Q_n has regular degree 1. In this sense, Theorem 2 studies a subclass of induced subgraphs of Q_n that are considered in Theorem 1. On the other hand, Theorem 2 gives Hamiltonian cycles instead of long cycles in Theorem 1.

Moreover, the bound $d(M) \geq 3$ in Theorem 2 is tight. To see this, consider a matching M consisting of $n - 1$ edges of the same direction $i \in [n]$ that cover all neighbors of some vertex u except the neighbor in the direction i . Then $d(M) = 2$ but $Q_n - V(M)$ is disconnected.

In Section 5 we study induced subgraphs of Q_n obtained by removing some linear code.

Theorem 3 *Let F be a linear code in Q_n with odd $d(F) \geq 3$. Then $Q_n - F$ is Hamiltonian.*

Note that linear codes with odd minimum distance are balanced. In a particular case when F is the well-known Hamming code [10], which exists for $n = 2^r - 1$ where $r > 1$ is integer, we obtain an optimal number of $2^n/(n+1)$ faulty vertices with minimum distance $d(F) \geq 3$. It could be noted that the induced subgraph of Q_n obtained by removing the Hamming code is called the Hamming shell and it has been previously studied by Dejter [2].

2 Preliminaries

For a positive integer n , we denote by $[n]$ the set $\{1, 2, \dots, n\}$. As usual, the vertex and edge sets of a graph G are denoted by $V(G)$ and $E(G)$, respectively. For sets $V \subseteq V(G)$ and a set $E \subseteq E(G)$, let $G - V$ denote the subgraph of G induced by $V(G) \setminus V$, and let $G - E$ denote the graph with vertices $V(G)$ and edges $E(G) \setminus E$.

The n -dimensional hypercube Q_n is a (bipartite) graph with all binary vectors of length n as vertices and with edges joining every two vertices that differ in exactly one coordinate, i.e.

$$V(Q_n) = \{0, 1\}^n \text{ and } E(Q_n) = \{uv; |\Delta(u, v)| = 1\},$$

where $\Delta(u, v) = \{i \in [n]; u_i \neq v_i\}$. Thus the distance of vertices u and v is $d(u, v) = |\Delta(u, v)|$. The distance of two edges uv and xy is the minimum distance between a vertex of uv to a vertex of xy . A vertex v is said to be *even* (*odd*) if it has even (odd) weight. The *weight* of v is the number of 1's in v . Note that vertices of each parity form bipartite classes of Q_n . Consequently, u and v have the same parity if and only if $d(u, v)$ is even.

Now we introduce notations to describe subcubes of Q_n . Assume that $D \subseteq [n]$ is a (nontrivial) set of $d = |D|$ coordinates of Q_n . We can represent every vertex x of Q_n as a pair $x = (u, v)_D$, where $u \in \{0, 1\}^d$ and $v \in \{0, 1\}^{n-d}$ are the vectors obtained by projection of x on the coordinates from D and $[n] \setminus D$, respectively. The index D in $(u, v)_D$ may be omitted whenever it is clear from the context. For $v \in \{0, 1\}^{n-d}$ we define the set $V_D(v) = \{(u, v)_D \in \{0, 1\}^n \mid u \in \{0, 1\}^d\}$. In other words, the set $V_D(v)$ contains the vertices of Q_n with the coordinates from $[n] \setminus D$ fixed by v . Furthermore, let $Q_D(v)$ be the subgraph of Q_n induced by $V_D(v)$. Since $Q_D(v)$ is isomorphic to Q_d , we say that $Q_D(v)$ is a d -dimensional *subcube* of Q_n . We usually replace the index D in $V_D(u)$ and $Q_D(u)$ with d to indicate just the size of D when the set D itself is clear or irrelevant.

In a particular case when $D = [n] \setminus \{i\}$, let Q_L^i and Q_R^i denote the $(n-1)$ -dimensional subcubes $Q_D(0)$ and $Q_D(1)$, respectively. That is, Q_L^i and Q_R^i are induced by vertices with 0 and 1 (respectively) in the i -th coordinate. The set of all edges between Q_L^i and Q_R^i is called a *layer*. Thus $E(Q_n)$ can be partitioned into n layers for each $i \in [n]$. An edge xy of Q_n such that $x \in V(Q_L^i)$ and $y \in V(Q_R^i)$ is *even* (*odd*) if the vertex x is even (odd). The set of all edges between Q_L^i and Q_R^i of the same parity is called a *half-layer* of Q_n . Hence every layer consists of two half-layers.

It is easy to show that every Hamiltonian cycle of Q_n contains at least one edge from each half-layer. In Section 4 we will need the following result on Hamiltonian cycles in hypercubes with a faulty matching.

Proposition 1 (Dimitrov et al. [3]) *Let M be a matching in Q_n and $n \geq 4$. Then $Q_n - M$ has a Hamiltonian cycle if and only if M does not contain a half-layer.*

3 Long paths in induced subgraphs of large degree

In this section we study the hypercube Q_n with a set F of faulty vertices such that every non-faulty vertex of Q_n has at most one faulty neighbor in Q_n . That is, we have an induced subgraph G of Q_n with a minimum degree $\delta(G) \geq n-1$. For our convenience, we allow $\delta(G) = n$ (i.e. $F = \emptyset$), but we assume that G is nonempty (i.e. $F \neq V(Q_n)$). Notice that a faulty vertex may have many faulty neighbors.

It is easy to see that G is always connected. Indeed, the statement holds for $n=1$, and for an arbitrary $i \in [n]$, $n \geq 2$, let $G_L^i = G \cap Q_L^i$ and $G_R^i = G \cap Q_R^i$. If one of them is empty, say G_L^i , then $G = G_R^i = Q_R^i$, which is connected. Otherwise there exist vertices $x \in V(G_L^i)$ and $y \in V(G_R^i)$, such that x and y are adjacent in G . Since $\delta(G_L^i), \delta(G_R^i) \geq n-2$, by induction it follows that G_L^i, G_R^i , and hence also G are connected.

Furthermore, F contains at most 2^{n-1} vertices with the extremal case when F forms a subcube of dimension $n-1$. Since F can be unbalanced, instead of Hamiltonian cycles we find so called long cycles. A cycle in $G = Q_n - F$ is *long* if it has length at least $2^n - 2|F| = 2|V(G)| - 2^n$. In a particular case when F contains only vertices of the same parity, such length is the best possible. Similarly, a path in G is *long* if it has length at least $2^n - 2|F| - 2 = 2|V(G)| - 2^n - 2$. Note that every long path between vertices u and v has length at least $2^n - 2|F| - 1 = 2|V(G)| - 2^n - 1$ if u and v are of different parity.

For long cycles and paths in G we will need the following lemma on interconnecting two parts of G . Recall that the parity of an edge $xy \in E(Q_n)$ where $x \in V(Q_L^i)$ and $y \in V(Q_R^i)$ is defined as the parity of the vertex x .

Lemma 1 *Let G be an induced subgraph of Q_n ($n \geq 4$) with $\delta(G) \geq n-1$. Then, for every $i \in [n]$ such that G_L^i and G_R^i are nonempty, there are at least 2 edges between G_L^i and G_R^i of each parity.*

Proof: If $\delta(G) = n$, then $G = Q_n$ and the statement trivially holds. Now we have $\delta(G) = n-1$.




Fig. 1: All nontrivial induced subgraphs G of Q_3 (up to isomorphism) with $\delta(G) = 2$.

For $n = 4$ we assume that $|V(G_L^i)| \leq |V(G_R^i)|$. We consider all possibilities up to isomorphism for the graph G_L^i on Figure 1. Note that the crossed red vertices on Figure 1 represent the removed vertices of Q_3 . If G_L^i is as on Figure 1(a)-(c), then it has at least two vertices of each parity with degree exactly 2 in G_L^i . Since $\delta(G) = 3$, it follows that these vertices are connected with G_R^i so the statement holds. If G_L^i is as on Figure 1(d), then G_R^i is also as on Figure 1(d) or $G_R^i = Q_3$. In both cases, G_L^i is connected with G_R^i by at least 6 edges. Since at most 4 of these edges have the same parity, the statement holds also in this case.

For $n \geq 5$ we split G into G_L^j and G_R^j for an arbitrary $j \in [n] \setminus \{i\}$. By the assumptions that both G_L^i and G_R^i are nonempty and $\delta(G) = n - 1$, it follows that $G_{LL} = G_L^i \cap G_L^j$ and $G_{RL} = G_R^i \cap G_L^j$ are nonempty, or $G_{LR} = G_L^i \cap G_R^j$ and $G_{RR} = G_R^i \cap G_R^j$ are nonempty. By symmetry, assume that the former case occurs. Then, applying induction for G_L^j we obtain at least 2 edges between G_{LL} and G_{RL} of each parity. Since G_{LL} and G_{RL} are subgraphs of G_L^i and G_R^i , the claim is established. \square

Now we show that induced subgraphs of Q_n with large minimum degree have long paths between any two vertices.

Theorem 4 *Let G be an induced subgraph of Q_n ($n \geq 1$) with $\delta(G) \geq n - 1$. Then G contains a long path between every two distinct vertices u and v .*

Proof: We proceed by induction. For $n \leq 2$, the statement trivially holds. For $n = 3$ we have either $G = Q_3$ or G is as on Figure 1(a)-(d). If $G = Q_3$, then the statement holds since G contains a Hamiltonian path between every two vertices of opposite parity, and an almost (up to one vertex) Hamiltonian path between every two vertices of the same parity [8]. If G is as on Figure 1(a), then it contains only half of the vertices of Q_3 , so it suffices that G is connected. If G is as on Figure 1(b)-(c), then every two vertices of G are connected by some path of length at least 3, which is even more than it is required for a long path. It remains to consider the case when G is as on Figure 1(d). First observe that there are 3 pairs of vertices u and v in G such that $d(u, v) = 3$. For all these pairs, G contains a path of length 5 between them. Furthermore, every two vertices u and v lie on a common 6-cycle of G . Thus G contains a path of length at least 4 between u and v also when $d(u, v) \leq 2$. So we have a long path in all cases.

For $n \geq 4$ we split G into G_L^i and G_R^i for some $i \in [n]$ such that u and v are in different parts, say $u \in V(G_L^i)$ and $v \in V(G_R^i)$. By Lemma 1, there are at least 2 edges between G_L^i and G_R^i of each parity. Hence there is an edge $xy \in E(G)$ such that $x \in V(G_L^i)$ is of opposite parity than u and $y \in V(G_R^i)$ is not v . Applying induction we obtain a path P_L in G_L^i between u and x of length at least

$2|V(G_L^i)| - 2^{n-1} - 2$, and a path P_R in G_R^i between y and v of length at least $2|V(G_R^i)| - 2^{n-1} - 2$. Observe that the length of P_L is actually at least $2|V(G_L^i)| - 2^{n-1} - 1$ since u and x are at odd distance. Therefore, by concatenating these paths with the edge xy we obtain a long path between u and v in G . \square

Now, the result on long cycles follows directly.

Proof of Theorem 1: Applying Theorem 4 for some two neighbors u and v in G , we obtain a path P of length at least $2|V(G)| - 2^n - 2$ between u and v . Observe that the length of P is actually at least $2|V(G)| - 2^n - 1$ since u and v are at odd distance. Closing P with the edge uv we obtain the desired long cycle in G . \square

4 Hamiltonicity of induced subgraphs of large degree given by faulty matchings

In this section we show that every induced subgraph of Q_{n+2} obtained by removing the vertices of a given set M of edges (matching) of Q_{n+2} with $d(M) \geq 3$ has a Hamiltonian cycle as stated by Theorem 2. Recall that $d(M)$ denotes the minimum distance between every two edges of M .

The idea of the proof of Theorem 2 is as follows. We split Q_{n+2} into Q_2 's and start with a 2-factor C of $Q_{n+2} - V(M)$ consisting of cycles inside these Q_2 's or inside two adjacent Q_2 's. By contracting the Q_2 's we obtain the hypercube Q_n . Next, we find some special spanning tree T of this Q_n that determines which cycles of C will be interconnected together. Finally, we assign some values to the edges of T to describe how to interconnect the cycles of C together so that we obtain a Hamiltonian cycle of $Q_{n+2} - V(M)$.

To formalize the proof, we need some additional definitions. Assume that M is a matching of Q_{n+2} where $n \geq 1$ such that $d(M) \geq 3$. The hypercube Q_{n+2} can be viewed as a Cartesian product $Q_{n+2} = Q_2 \square Q_n$. Thus for the set of coordinates $D = \{1, 2\} \subseteq [n+2]$, we consider Q_{n+2} as the (contracted) Q_n in which every vertex u represents the subcube $Q_D(u)$. For the rest of this section, the set D is fixed, so let us replace the index D with 2, which is the size of D . Thus, for example $Q_2(u)$ denotes $Q_D(u)$.

Some edges of M are inside these Q_2 's, the other edges of M interconnect distinct Q_2 's. However, each Q_2 may contain (one or two) vertices of at most one edge of M because of the minimum distance 3 between the edges of M . Thus, we may assign the red, black or green color to each vertex u of Q_n as follows:

- if $Q_2(u)$ contains an edge of M , then u receives the red color;
- if $Q_2(u)$ contains only one vertex of some edge of M , then u receives the black color;
- if $Q_2(u)$ contains no vertex covered by M , then u receives the green color.

Moreover, if some neighbors u and v of Q_n receive the black color because of the same edge of M , then we also assign the black color to the edge uv of Q_n . Notice that the edge uv corresponds to a copy of Q_3 in Q_{n+2} . Let $Q_n(M)$ denote the contracted Q_n with assigned colors as described above. See Figures 2 and 4(a) for an illustration. On Figure 2 the dashed squares represent the copies of Q_2 's in Q_{n+2} , the dotted edges represent edges between two adjacent Q_2 's, and the bold black edges represent the matching M .

For our convenience in dealing with small dimensions, let us make one additional assumption on $Q_n(M)$ based on the following observation.




Fig. 2: The example for $n = 4$ of the graph $Q_{n+2} = Q_2 \square Q_n$ containing (black) matching M with $d(M) \geq 3$.

Proposition 2 For $n \in \{1, 2, 3\}$, every matching M of Q_{n+2} such that $d(M) \geq 3$ has at most two edges.

Proof: Clearly, it suffices to consider only the case $n = 3$. Assume that uv and xy are two edges of Q_5 with $d(uv, xy) \geq 3$. There are three possible configurations (up to relabeling the vertices): $d(uv, xy) = d(u, x) = d(v, y) = 4$, $d(uv, xy) = d(u, x) = d(v, y) = 3$, and $d(uv, xy) = d(u, x) = 3$ and $d(v, y) = 5$ as depicted on Figure 3. Note that on Figure 3 the edges between the four copies of Q_3 in Q_5 are omitted except the edges between the red vertices. Observe that in all three cases, every vertex of Q_5 is at distance at most 2 from some vertex of $\{u, v, x, y\}$. Thus, there is no other edge in Q_5 at distance at least 3 from both uv and xy . \square

Hence for $n \in \{1, 2, 3\}$, by the symmetry of Q_{n+2} we may assume that the edges of M are in the




Fig. 3: All configurations (up to isomorphism) of two edges uv and xy in Q_5 with $d(uv, xy) \geq 3$.




Fig. 4: (a) The (contracted) colored hypercube $Q_n(M)$ corresponding to Figure 2, (b) a spanning tree T of $Q_n(M)$ obtained by Lemma 3.

directions 1 and 2. It follows that the graph $Q_n(M)$ has no black vertices for $n \in \{1, 2, 3\}$ as every edge of M is inside some Q_2 . The following lemma summarizes basic properties of $Q_n(M)$.

Lemma 2 *Let M be a matching of Q_{n+2} ($n \geq 1$) such that $d(M) \geq 3$. Then the graph $Q_n(M)$ has the following properties:*

- (i) every red vertex has only green neighbors;
- (ii) every black vertex has at most 2 black neighbors;
- (iii) no two black edges are incident.

Proof: Let u and v be neighbors in $Q_n(M)$. If u is red, that is, $Q_2(u)$ contains some edge $xy \in M$, then all vertices in $Q_2(v)$ are at distance at most 2 from xy . Since $d(M) \geq 3$, it follows that v is green, so (i) holds.

If u is black, then it has some black neighbor joined with its black edge. Suppose on the contrary that u has another two black neighbors v and w . That is, $Q_2(u)$, $Q_2(v)$, and $Q_2(w)$ contain vertices x , y , and z , respectively, that are covered by M . Moreover, we have $d(x, y) = d(x, z) = 3$ since $d(M) \geq 3$. In both $Q_2(v)$ and $Q_2(w)$ there is only one vertex at distance 3 from x , so the position of y and z is fixed. But it follows that $d(y, z) = 2$, a contradiction.

Finally, suppose on the contrary that two black edges are incident at some vertex u . Then $Q_2(u)$ contains two vertices covered by distinct edges of M . But it follows that every two vertices of $Q_2(u)$ are at distance at most 2, a contradiction. \square

Now we find a spanning tree of $Q_n(M)$ that will determine which Q'_2 s will be consecutive on the desired Hamiltonian cycle of $Q_{n+2} - V(M)$.

Lemma 3 *Let M be a matching of Q_{n+2} ($n \geq 1$) such that $d(M) \geq 3$. Let A be the set of black edges of $Q_n(M)$, and let B be the set of other edges between black vertices in $Q_n(M)$. Then $Q_n(M)$ has a spanning tree T with the following properties:*

- (i) T contains A ;
- (ii) T avoids B ;
- (iii) every black vertex has degree 2 or 3 in T ;
- (iv) every red or green vertex has degree at most 2 in T .

Proof: Recall that for $n \in \{1, 2, 3\}$ we may assume that $Q_n(M)$ contains no black vertices, so the statement trivially holds; just take any Hamiltonian path of $Q_n(M)$. Now assume that $n \geq 4$. First observe that B forms a matching in Q_n . Indeed, suppose that two edges of B are incident at some vertex u . Then u is black, and it has also one edge from A . But this means that u has at least 3 black neighbors, contrary to Lemma 2(ii).

Moreover, we claim that B does not contain a half-layer of $Q_n(M)$. Suppose on the contrary that B contains a half-layer C . Then all 2^{n-1} vertices of C are black, so they are covered by the matching A . Since A connects no two vertices of C and C contains exactly a half of the vertices of $Q_n(M)$, it follows that actually all 2^n vertices of $Q_n(M)$ are black. But this contradicts Lemma 2(ii) so the claim holds. Hence, by Proposition 1, there is a Hamiltonian cycle of $Q_n(M)$ that avoids B . By removing some edge between non-black vertices on this cycle, we obtain a spanning path T' of $Q_n(M)$ satisfying (ii)-(iv).

In addition to satisfy (i), we iteratively add the edges of A into T' and at the same time, we remove some edges to keep the tree structure. In particular, assume that we want to add the black edge uv into T' . Since the black vertices u and v have only one black edge by Lemma 2(iii), it follows that the degree of both u and v in T' is 2. Thus, this procedure may increase the degree of a black vertex only to 3. Let w be the neighbor of the vertex v on the cycle of $T' \cup \{uv\}$. Note that w is neither red by Lemma 2(i), nor black since T' avoids B . Hence w is green. Then, by removing the edge vw from $T' \cup \{uv\}$ we obtain a new spanning tree T' of $Q_n(M)$ containing the edge $uv \in A$ and satisfying conditions (ii)-(iv). After we consider the whole set A , we have the required spanning tree. \square

Figure 4 shows an example of a spanning tree of $Q_n(M)$ obtained by Lemma 3. Notice that $A = \{uv, xy\}$ and $B = \{ux, vy\}$ in this example.

In the hypercube Q_{n+2} , each subcube $Q_2(u)$ represented by a vertex u of $Q_n(M)$ contains up to 4 edges that are not incident with the matching M . Let $S(u) \subseteq E(Q_2)$ be the set of these edges projected to D ; that is, for $a, b \in \{0, 1\}^2$,

$$ab \in S(u) \text{ if and only if the edge } (a, u)(b, u) \in E(Q_2(u)) \text{ is not incident with } M.$$

See Figure 5 for some examples. Note that $|S(u)| = 1$ if u is red, $|S(u)| = 2$ if u is black, and $|S(u)| = 4$ if u is green. The following lemma describes some properties of S with respect to our spanning tree T .

Lemma 4 *Let M be a matching of Q_{n+2} ($n \geq 1$) such that $d(M) \geq 3$, and let T be the spanning tree of $Q_n(M)$ obtained by Lemma 3. For every two vertices u and v of $Q_n(M)$ with distance 2 in T , it holds that*

- (i) if u and v are red with $S(u) = \{ab\}$ and $S(v) = \{cd\}$, then $\{a, b, c, d\} = \{0, 1\}^2$;




Fig. 5: $S(u) = S(v) = \{\{00, 01\}, \{01, 11\}\}$, $S(x) = E(Q_2)$, $S(y) = \{\{00, 01\}\}$.




Fig. 6: An illustration for Lemma 4.

- (ii) if u is red and v is black, then $S(u) \cap S(v) = \emptyset$;
- (iii) if u and v are black, then $S(u) \neq S(v)$.

Proof: Note that in all three cases the common neighbor of u and v is green by Lemmas 2(i) and 3(ii). See Figure 6 for an illustration. First, assume that u and v are red. Let $a, b, c, d \in \{0, 1\}^2$ be such that $S(u) = \{ab\}$ and $S(v) = \{cd\}$. That is, $(a, u)(b, u)$ and $(c, v)(d, v)$ are the edges of $Q_2(u)$ and $Q_2(v)$, respectively, that are not incident with M . Thus, the edges $(c, u)(d, u)$ and $(a, v)(b, v)$ are in M . Since $d(u, v) = 2$ and $d(M) \geq 3$, it follows that the vectors a, b, c, d of $\{0, 1\}^2$ are distinct, so (i) holds.

Next, assume that u is red and v is black. Let $a, b, c \in \{0, 1\}^2$ be such that $(a, u)(b, u)$ is the edge of M in $Q_2(u)$ and (c, v) is the vertex of $Q_2(v)$ covered by M . Again, since $d(u, v) = 2$ and $d(M) \geq 3$, it follows that the vectors a, b, c of $\{0, 1\}^2$ are distinct. Let $d \in \{0, 1\}^2$ be the fourth vector of $\{0, 1\}^2$, so $S(u) = \{cd\}$. As (c, v) is covered by M , we have $cd \notin S(v)$. Hence also (ii) holds.

Finally, assume that u and v are black. Let $a, b \in \{0, 1\}^2$ be such that (a, u) and (b, v) are covered by M . Similarly as above, we obtain that $a \neq b$, and therefore $S(u) \neq S(v)$. \square




Fig. 7: The 2-factor C of $Q_{n+2} - V(M)$.

Now we assign to the non-black edges uv of T values $c(uv)$ from $E(Q_2)$ that describe in the main proof how to interconnect cycles of some special 2-factor C of $Q_{n+2} - V(M)$ into a Hamiltonian cycle. A value $c(uv) = ab$ where $a, b \in \{0, 1\}^2$ means that the cycles of C in $Q_2(u)$ and $Q_2(v)$ will be interconnected by replacing the edges $(a, u)(b, u)$ and $(a, v)(b, v)$ of C with the edges $(a, u)(a, v)$ and $(b, u)(b, v)$. Since the sets $S(u)$ and $S(v)$ contain the list of available edges for this operation, we need this assignment to satisfy the conditions described by the following lemma.

Lemma 5 *Let M be a matching of Q_{n+2} ($n \geq 1$) such that $d(M) \geq 3$, let A be the set of black edges of $Q_n(M)$, and let T be the spanning tree of $Q_n(M)$ obtained by Lemma 3. Then, there is an assignment $c : E(T) \setminus A \rightarrow E(Q_2)$ such that*

- (i) $c(uv) \in S(u) \cap S(v)$, for every edge $uv \in E(T) \setminus A$;
- (ii) $c(uv) = c(vw)$ if and only if v is red, for every two incident edges $uv, vw \in E(T) \setminus A$.

Proof: From Lemma 3 it follows that by removing the edges of A , the spanning tree T splits into spanning paths. Recall that by Lemmas 2 and 3, every red or black vertex has only green neighbors on these paths. First we assign the single value of $S(u)$ to the edges from each red vertex u . By Lemma 4(i), no conflict may arise in a green vertex that has two red neighbors. Clearly, two edges incident at some red vertex receive the same value of c . Hence the conditions in the statement hold so far.

Next, we traverse each path of $T - A$ starting in an arbitrary end, and along the way we assign the value of c to all unassigned edges of this path so that the statement of the lemma is satisfied. Recall that there are no two consecutive black vertices on any path of $T - A$ by Lemma 3(ii). Assume that we are in an unassigned edge uv . Hence the vertices u and v are not red. Let xu and vy be the previous and the next edge on the same path, respectively, if they exist. If y is red, then the edge vy has already some assigned value $c(vy)$. We consider the colors of u and v .

If both u and v are green, then we choose $c(uv) \in S(u) \cap S(v) = E(Q_2)$ such that $c(xu) \neq c(uv) \neq c(vy)$. Note that such value $c(uv)$ exists, since there are four possibilities in $S(u) \cap S(v)$ and at most two of them are prohibited. If u is green and v is black, then $|S(u) \cap S(v)| = 2$ and y is not red by Lemma 2(i). We choose $c(uv) \in S(u) \cap S(v)$ such that $c(xu) \neq c(uv)$. Finally, if u is black and v is green, then $|S(u) \cap S(v)| = 2$ and y can be red. Similarly as above, we choose $c(uv) \in S(u) \cap S(v)$ such that $c(xu) \neq c(uv)$. By Lemma 4(ii), it follows that $c(uv) \neq c(vy)$ if y is red.

Therefore, after we traverse all paths, the statement of the lemma holds. \square




Fig. 8: An example of a Hamiltonian cycle of $Q_{n+2} - V(M)$ obtained by Theorem 2.

Now we prove the main theorem of this section.

Proof of Theorem 2: Let $Q_n(M)$ be the contracted hypercube with colors assigned as defined at the beginning of this section, and let A be the set of black edges in $Q_n(M)$. Recall that A forms a matching in $Q_n(M)$ by Lemma 2. First, we define the following 2-factor C of $Q_{n+2} - V(M)$. For every green vertex u , we take the 4-cycle of $Q_2(u)$ into C . For every red vertex u , we take twice into C the only edge in $Q_2(u)$ that is not incident with M . Because of technical reasons, let us consider this multi-edge to be a 2-cycle in Q_{n+2} . For every black edge uv we take into C the only possible 6-cycle in the subgraph of $Q_{n+2} - V(M)$ induced on vertices of $Q_2(u)$ and $Q_2(v)$ as illustrated by Figure 7(a). Hence, C is a 2-factor of $Q_{n+2} - V(M)$. Moreover, it is easy to see that for every $u \in \{0, 1\}^n$ and $a, b \in \{0, 1\}^2$,

$$(a, u)(b, u) \in E(C) \text{ if and only if } ab \in S(u). \quad (1)$$

See Figure 7 for an illustration.

Let T be the spanning tree of $Q_n(M)$ obtained by Lemma 3. We use T to interconnect C into a Hamiltonian cycle. To interconnect cycles of C in the adjacent Q_2 's we apply the assignment $c : E(T) \setminus A \rightarrow E(Q_2)$ from Lemma 5. Assume that $c(uv) = ab$ for an edge $uv \in E(T) \setminus A$. By Lemma 5(i) and by (1), we have that $(a, u)(b, u) \in E(C)$ and $(a, v)(b, v) \in E(C)$. Hence, by replacing these two edges with the edges $(a, u)(a, v)$ and $(b, u)(b, v)$ we still obtain a 2-factor of $Q_{n+2} - V(M)$.

By Lemma 5(ii), no edge can be replaced more than once this way, except the multi-edges, which may be replaced twice. Since T is a spanning tree, after we proceed with all edges of $E(T) \setminus A$, we obtain a Hamiltonian cycle of $Q_{n+2} - V(M)$. See Figure 8 for an illustration. \square

5 Hamiltonicity of shells of linear codes

In this section we study induced subgraphs of Q_n obtained by removing some linear code. A (binary) linear code \mathcal{C} given by (binary) parity-check matrix H is the set of (binary) vectors x such that $Hx^T = 0$. The length of \mathcal{C} is the number n of columns of H . The dimension of \mathcal{C} is the smallest k such that H has $n - k$ linearly independent rows, that is $k = n - rk(H)$ where $rk(H)$ denotes the rank of H . The minimum distance in \mathcal{C} is the smallest d such that H has d linearly dependent columns. A linear code of length n , dimension k , and minimum distance d is shortly called $[n, k, d]$ -code. The vectors in \mathcal{C} are called codewords. The shell of \mathcal{C} is the graph $Q_n - \mathcal{C}$.

If \mathcal{C} is $[n, k, d]$ -code, then it forms k -dimensional linear subspace of $\{0, 1\}^n$, so it has 2^k codewords. Moreover, d is the smallest weight of a nonzero codeword, or equivalently, d is the smallest distance between every two codewords. If d is odd, then exactly half of codewords are even, so \mathcal{C} is balanced. For more properties of linear codes see for example the monograph by MacWilliams and Sloane [10].

We will need the following result on hypercubes with two faulty vertices.

Proposition 3 (Locke and Stong [9]) *Let x and y be two vertices of Q_n ($n \geq 3$) at odd distance. Then $Q_n - \{x, y\}$ is Hamiltonian.*

Let us restate Theorem 3 in a slightly different form than it was mentioned in Section 1.

Theorem 3 *The shell of every $[n, k, d]$ -code \mathcal{C} with odd $d \geq 3$ is Hamiltonian.*

Proof: Let z be a codeword of weight d , and let $D \subseteq [n]$ be the set of d coordinates in z containing 1. If $d = n$, then $\mathcal{C} = \{\mathbf{0}, \mathbf{1}\}$ where $\mathbf{0}$ and $\mathbf{1}$ are vectors of all 0's and of all 1's, respectively. So the statement of the theorem follows by Proposition 3. Now we assume that $d < n$.

First, consider the subcube $Q_D(u)$ for some $u \in \{0, 1\}^{n-d}$ containing some codeword x . By the linearity of \mathcal{C} , $Q_D(u)$ contains also the codeword $y = x \oplus z$ and no other codewords because of the minimum distance d . Moreover, $Q_D(v)$ contains no codeword for every neighbor v of u in Q_{n-d} since $d(x, w) \leq (d+1)/2$ or $d(y, w) \leq (d+1)/2$ for every $w \in V_D(v)$ and $(d+1)/2 < d$ for $d \geq 3$.

The construction of a Hamiltonian cycle of the shell of \mathcal{C} proceeds as follows. We start by choosing an arbitrary Hamiltonian cycle $u_1, u_2, \dots, u_{2^{n-d}}, u_1$ of Q_{n-d} . For sake of simplicity, for $d = n - 1$ let us consider the sequence $0, 1, 0$ of vertices of Q_1 to be a Hamiltonian cycle of Q_1 . Then, we choose in each subcube $Q_D(u_i)$ two vertices $a_i, b_i \in V_D(u_i) \setminus \mathcal{C}$ such that $Q_D(u_i) - \mathcal{C}$ contains a Hamiltonian path between a_i and b_i , and $b_i a_{i+1} \in E(Q_n)$ (where $a_{2^{n-d}+1} = a_0$) for every $i \in [2^{n-d}]$. Therefore, by gluing these paths we obtain a Hamiltonian cycle of $Q_n - \mathcal{C}$.

It remains to determine precisely the vertices a_i and b_i . We may assume that $Q_D(u_1)$ contains no codeword and we choose $b_1 \in V_D(u_1)$ and $a_2 \in V_D(u_2)$ such that a_2 is the neighbor of b_1 in Q_n and a_2 is not in \mathcal{C} . Now, for each $i \in \{2, \dots, 2^{n-d}\}$ we choose the vertices $b_i \in V_D(u_i)$ and $a_{i+1} \in V_D(u_{i+1})$ in the following way.

If $Q_D(u_i)$ contains no codeword, we choose $b_i \in V_D(u_i)$ and $a_{i+1} \in V_D(u_{i+1})$ such that $d(a_i, b_i)$ is odd, a_{i+1} is the neighbor of b_i in $Q_D(u_{i+1})$ and a_{i+1} is not in \mathcal{C} . Since $d(a_i, b_i)$ is odd, $Q_D(u_i)$ contains a Hamiltonian path between a_i and b_i .

If $Q_D(u_i)$ contains some codeword x , then $Q_D(u_i) - \{x, y\}$ where $y = x \oplus z$ contains a Hamiltonian cycle C by Lemma 3. Thus we choose $b_i \in V_D(u_i)$ and $a_{i+1} \in V_D(u_{i+1})$ such that b_i is some neighbor of a_i on C and a_{i+1} is the neighbor of b_i in $Q_D(u_{i+1})$. Hence, $d(a_i, b_i) = 1$ and $Q_D(u_i) - \mathcal{C}$ contains a Hamiltonian path between a_i and b_i .

Observe that in the last subcube when $i = 2^{n-d}$ we obtain for $a_{i+1} = a_1$ that $d(a_1, b_1)$ is odd, so also $Q_D(u_1)$ contains a Hamiltonian path between a_1 and b_1 . Therefore, for every $i \in [2^{n-d}]$ the vertices a_i and b_i have the required properties. \square

Acknowledgements

The authors would like to thank the anonymous referees for their valuable comments.

References

- [1] N. CASTAÑEDA, I. GOTCHEV, *Embedded Paths and Cycles in Faulty Hypercubes*, J. Comb. Optim., (2009), doi:10.1007/s10878-008-9205-6.
- [2] I. J. DEJTER, *Equitable factorizations of Hamming shells*, Discrete Math., 261 (2003), 177–187.
- [3] D. DIMITROV, T. DVOŘÁK, P. GREGOR, R. ŠKREKOVSKI, *Gray Codes Faulting Matchings*, IMFM Preprints, 45 (2007).
- [4] T. DVOŘÁK, P. GREGOR, *Partitions of faulty hypercubes into paths with prescribed endvertices*, SIAM J. Discrete Math., 22 (2008), 1448–1461.
- [5] J. FINK, P. GREGOR, *Long paths and cycles in hypercubes with faulty vertices*, submitted.
- [6] J.-S. FU, *Fault-tolerant cycle embedding in the hypercube*, Parallel Comput., 29 (2003), 821–832.
- [7] F. GRAY, *Pulse Code Communication*, U.S. Patent 2,632,058, filed 13 November 1947, issued 17 March 1953.
- [8] M. LEWINTER, W. WIDULSKI, *Hyper-Hamilton laceable and caterpillar-spannable product graphs*, Comput. Math. Appl., 34 (1997), 99–104.
- [9] S. C. LOCKE, R. STONG, *Spanning Cycles in Hypercubes: 10892*, The Amer. Math. Monthly, 110 (2003), 440–441.
- [10] F. J. MACWILLIAMS, N. J. A. SLOANE, *The Theory of Error-Correcting Codes*. North-Holland, New York (1981).
- [11] C. SAVAGE, *A survey of combinatorial Gray codes*, SIAM Rev., 39 (1997), 605–629.