

Independent sets in $(P, \text{diamond})$ -free graphs

Raffaele Mosca

► **To cite this version:**

Raffaele Mosca. Independent sets in $(P, \text{diamond})$ -free graphs. Discrete Mathematics and Theoretical Computer Science, DMTCS, 2009, 11 (1), pp.125–140. <hal-00988179>

HAL Id: hal-00988179

<https://hal.inria.fr/hal-00988179>

Submitted on 7 May 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Independent sets in $(P_6, \text{diamond})$ -free graphs

Raffaele Mosca

Dipartimento di Scienze, Università degli Studi "G. d'Annunzio", Pescara 65127, Italy.
r.mosca@unich.it

received June 6, 2006, revised June 6, 2007, March 17, 2008, January 23, 2009, accepted January 27, 2009.

We prove that on the class of $(P_6, \text{diamond})$ -free graphs the Maximum-Weight Independent Set problem and the Minimum-Weight Independent Dominating Set problem can be solved in polynomial time.

Keywords: Maximum-Weight Independent Set; Minimum-Weight Independent Dominating Set; polynomial-time algorithm.

1 Introduction

An *independent set* (or a *stable set*) in a graph G is a subset of pairwise nonadjacent vertices of G . An independent set of G is *maximal* if it is not properly contained in any other independent set of G .

The Maximum-Weight Independent Set (*WIS*) problem is the following: Given a graph $G = (V, E)$ and a weight function w on V , determine an independent set of G of maximum weight. Let $\alpha_w(G)$ denote the maximum weight of an independent set of G . The WIS problem reduces to the *IS* problem if all vertices v have the same weight $w(v) = 1$.

The WIS problem is NP-hard [34] and remains difficult for cubic graphs [27] and for planar graphs [26], while it can be efficiently solved for various graph classes which include perfect graphs [33] (and the class of perfect graphs includes the chordal graphs), $K_{1,3}$ -free graphs [2, 37, 40, 42, 45], and $2K_2$ -free graphs [21, 22, 38].

The Minimum-Weight Independent Dominating Set (*WID*) problem is the following: Given a graph $G = (V, E)$ and a weight function w on V , determine a maximal independent set of G of minimum weight. Let $\iota_w(G)$ denote the minimum weight of a maximal independent set of G . The WID problem reduces to the *ID* problem if all vertices v have the same weight $w(v) = 1$.

The WID problem is NP-hard [28] and remains difficult for chordal graphs [18] and for $2P_3$ -free perfect graphs [46], while it can be efficiently solved for various graph classes which include permutation graphs [15], locally independent well-dominated graphs [47], and $2K_2$ -free graphs [21, 22, 38].

Both WIS and WID remain difficult for triangle-free graphs [43]. Also, for both IS and ID, the class of P_5 -free graphs is the unique minimal class, defined by forbidding a single connected subgraph, for which the computational complexity is an open question (see [1, 3, 7]).

On the other hand, several papers introduced structural properties on graphs containing no long induced paths (see e.g. [5, 6, 19, 39]), often applied to design efficient algorithms for solving various NP-hard

problems, including WIS or WID, in subclasses of such graphs: concerning subclasses of P_5 -free graphs, see e.g. [4, 9, 10, 11, 13, 16, 17, 23, 29, 31, 36]; concerning subclasses of P_6 -free graphs, see e.g. [9, 24, 30, 35, 41, 44].

Let us focus on two such graph subclasses which involve triangle-free graphs as well.

The class of $(P_5, \text{diamond})$ -free graphs: a recent paper [8] shows that such graphs have bounded clique-width and that a corresponding clique-width expression can be constructed in $\mathcal{O}(n + m)$ time, which implies that a large class of NP-hard problems including WIS and WID can be solved for such graphs in $\mathcal{O}(n + m)$ time.

The class of $(P_6, \text{triangle})$ -free graphs: a recent paper [14] shows that such graphs have bounded clique-width and that a corresponding clique-width expression can be constructed in $\mathcal{O}(n^2)$ time, which implies that a large class of NP-hard problems including WIS and WID can be solved for such graphs in $\mathcal{O}(n^2)$ time.

This paper introduces a proof that WIS and WID can be solved for $(P_6, \text{diamond})$ -free graphs in $\mathcal{O}(n^7)$ time.

2 Notation and preliminaries

For any missing notation or reference, let us refer to [12].

Throughout this paper let $G = (V, E)$ be a finite undirected graph without self-loops and multiple edges and let $|V| = n, |E| = m$. For every $u \in V$, let $N(u) = \{v \in V : uv \in E\}$ be the set of *neighbors* of u . Let U, W be two subsets of V . Let $N(U) = \{v \in V \setminus U : \text{there exists } u \in U \text{ such that } uv \in E\}$. Let $N_W(U) = N(U) \cap W$. Let us say that U has a *join* (a *co-join*, respectively) with W , if each vertex in U is adjacent (is nonadjacent) to each vertex in W . Let $v \in V$. Let us say that: v *contacts* U if v is adjacent to some vertex of U ; v is *universal* for U if v is adjacent to each vertex of U ; v is *partial* to U if v contacts U but is not universal for U . Then let us say that U , with $\emptyset \subset U \subset V$, is a *module* of G – often called *homogeneous set* in the literature – if no vertex of $V \setminus U$ is partial to U .

Let $G[U]$ denote the subgraph of G induced by the vertex subset U . For any graph F , G is *F-free* if G contains no induced subgraph isomorphic to F .

A *component* of G is the vertex set of a maximal connected subgraph of G . A component of G is *trivial* if it is a singleton, and *nontrivial* otherwise.

Concerning WIS and WID, algorithmically an easy reduction works if the graph is disconnected: that is, if G has components V_1, \dots, V_k , then $\alpha_w(G) = \alpha_w(G[V_1]) + \dots + \alpha_w(G[V_k])$ and $\iota_w(G) = \iota_w(G[V_1]) + \dots + \iota_w(G[V_k])$.

A path P_k has vertices v_1, v_2, \dots, v_k and edges $v_j v_{j+1}$ for $1 \leq j < k$. A cycle C_k has vertices v_1, v_2, \dots, v_k and edges $v_j v_{j+1}$ for $1 \leq j \leq k$ (subscript addition taken modulo k). A *triangle* is a graph of three vertices a, b, c and edges ab, ac, bc . A *diamond* is a graph of four vertices a, b, c, d and edges ab, ac, ad, bc, bd .

A *clique* of G is a set of pairwise adjacent vertices of G . Notice that each component of G is a clique if and only if G is P_3 -free.

A graph is *chordal* if it contains no induced C_k , $k \geq 4$.

For chordal graphs, WIS and ID can be efficiently solved (see [25] and [20], respectively), while WID remains NP-hard on them [18].

In [32] the authors proved that distance-hereditary graphs have bounded clique-width, and that a corresponding clique-width expression can be constructed in $\mathcal{O}(n + m)$ time. Since chordal diamond-free

graphs are distance-hereditary, a direct consequence is:

Theorem 1 ([32]) *Both the WIS and the WID problems can be solved for chordal diamond-free graphs in $\mathcal{O}(n + m)$ time.* \square

In [14] the authors proved that $(P_6, \text{triangle})$ -free graphs have bounded clique-width, and that a corresponding clique-width expression can be constructed in $\mathcal{O}(n^2)$ time. A direct consequence is:

Theorem 2 ([14]) *Both the WIS and the WID problems can be solved for $(P_6, \text{triangle})$ -free graphs in $\mathcal{O}(n^2)$ time.* \square

Obviously, the WIS (or WID) problem on a graph G with vertex weight function w can be reduced to the same problem on subgraphs $G[V \setminus N(v)]$ for every $v \in V$ in the following way:

$$\alpha_w(G) = \max\{\alpha_w(G[V \setminus N(v)]) \mid v \in V\}$$

$$\iota_w(G) = \min\{\iota_w(G[V \setminus N(v)]) \mid v \in V\}$$

Thus, whenever WIS (or WID) is solvable in time T for every subgraph $G[V \setminus N(v)]$ of G with $v \in V$, then it is solvable for G in time nT , plus $\mathcal{O}(n^3)$ additional steps to generate those subgraphs.

Let us conclude with an observation which will be often used later.

Observation 1 *Let $G = (V, E)$ be a graph, and $U \subseteq V$ with $|U| = k$. If one can solve WIS (or WID) for each induced subgraph of $G[V \setminus U]$ in time T , then one can solve WIS (or WID) for G in time $2^k(T + n^2)$.*

Proof: Let $I(U)$ be the family of independent sets of $G[U]$. Then to solve WIS (or WID) for G one can consider WIS (or WID) for $|I(U)|$ subgraphs of G , i.e., for $G[V \setminus U]$ and for $G[I \cup (V \setminus (N(I) \cup U))]$ for every $I \in I(U)$. Since $|I(U)| \leq 2^k$, the assertion follows. \square

Remark: The results of the next section are introduced only for WIS, by meaning that they hold for WID as well (by interchanging WIS with WID, and α with ι).

3 Independent sets in $(P_6, \text{diamond})$ -free graphs

Let us introduce a preliminary result.

Definition 1 *A graph $G = (V, E)$ is green if there exists a partition $\{X, Y\}$ of V (with X or Y possibly empty) such that:*

- (i) $G[X]$ and $G[Y]$ are P_3 -free;
- (ii) each component of $G[Y]$ is a module of G ;
- (iii) each vertex in Y is adjacent to at most one vertex in each component of $G[X]$.

Notice that every P_3 -free graph is green.

Lemma 1 *One can solve WIS for every green P_6 -free graph in $\mathcal{O}(n^3)$ time.*

Proof: Let $G = (V, E)$ be a green P_6 -free graph. Assume without loss of generality that G is connected. Let $\{X, Y\}$ be a partition of V according to Definition 1. In particular, to our aim, by ii one can assume that Y is an independent set. In fact, one can contract each component K of $G[Y]$ into a singleton u with $w(u) = \alpha_w(G[K])$: that can be done in polynomial time since each component of $G[Y]$ is a clique.

Let W be the family of nontrivial components of $G[X]$. In particular, to our aim (similarly to above), one can assume that in each component $Q \in W$ at most one vertex is nonadjacent to any vertex in Y .

Claim 3.1 *There exists $y^* \in Y$ such that y^* contacts every element of W .*

Proof: For any $y \in Y$, let $J(y) = \{Q \in W : y \text{ contacts } Q\}$. Let $y^* \in Y$ be such that $|J(y^*)| \geq |J(y)|$ for every $y \in Y$. We state that this vertex y^* is a proper choice for Claim 3.1. Assume for a contradiction that there exists a component $Q \in W$ such that y^* does not contact Q . Since G is connected, one can select $y \in Y$ belonging to a shortest path from y^* to Q , such that y contacts Q . By definition of W and by iii, Q contains two (adjacent) vertices q_1 and q_2 such that y is adjacent to q_1 and nonadjacent to q_2 . Then, since G is P_6 -free, y contacts all the elements of W which are contacted by y^* . This implies $|J(y^*)| < |J(y)|$, a contradiction. \square

Claim 3.2 *There exists at most one element of W of cardinality greater than 2.*

Proof: Assume for a contradiction that there exist two elements of W , say \tilde{Q} and Q , with $|\tilde{Q}| \geq 3$ and $|Q| \geq 3$. Let $y^* \in Y$ according to Claim 3.1. Let $\tilde{q} \in \tilde{Q}$ and $q \in Q$ be adjacent to y^* . Since $|\tilde{Q}| \geq 3$, there exists $a \in Y$ adjacent to $q_a \in \tilde{Q}$, with $q_a \neq \tilde{q}$. Then a is adjacent to \tilde{q} , otherwise $a, q_a, \tilde{q}, y^*, q$ and any vertex in Q nonadjacent to a induce a P_6 . Then, since $|Q| \geq 3$, there exists $b \in Y$ adjacent to $q_b \in Q$, with $q_b \neq q$. By symmetry, one has that b is adjacent to \tilde{q} . Since $|Q| \geq 3$, there exists $q' \in Q$ such that $q' \neq q$ and q' is nonadjacent to b . Then $q', q_b, b, \tilde{q}, q_a, a$ induce a P_6 , a contradiction. \square

If $|Q| \leq 2$ for every $Q \in W$, then by iii G is triangle-free. Otherwise, by Claim 3.2 there exists at most one element, say \tilde{Q} , of W of cardinality greater than 2. One can solve WIS in G by solving WIS in $G[V \setminus \tilde{Q}]$ and in $G[V \setminus N(\tilde{q})]$ for every $\tilde{q} \in \tilde{Q}$. Since such graphs are triangle-free, the lemma follows by Theorem 2. \square

3.1 Deleting C_6 's in $(P_6, \text{diamond})$ -free graphs

Throughout this subsection assume that $G = (V, E)$ is a $(P_6, \text{diamond})$ -free graph containing a 6-cycle C , say with vertices v_i and edges $v_i v_{i+1}$, $i \in \{1, \dots, 6\}$ (subscript addition taken modulo 6). Let $N(C)$ be the set of vertices from $V \setminus C$ which are adjacent to some vertex in C . For any subset S of C , let M_S be the set formed by those vertices in $N(C)$ which are adjacent to each vertex in S and are nonadjacent to each vertex in $C \setminus S$. In particular, let us write M_1 for $S = \{v_1\}$, $M_{1,2}$ for $S = \{v_1, v_2\}$, and so on. Then let $Z(k)$ denote the set of vertices of $N(C)$ having exactly k neighbors in C .

Since G is $(P_6, \text{diamond})$ -free: $Z(1) = Z(5) = Z(6) = \emptyset$; each vertex in $Z(2)$ belongs to some of the sets $M_{i,i+2}$ or $M_{i,i+3}$, $i \in \{1, \dots, 6\}$ (subscript addition taken modulo 6); each vertex in $Z(3)$ belongs to some of the sets $M_{i,i+2,i+4}$ or $M_{i,i+2,i+3}$ or $M_{i,i+3,i+4}$, $i \in \{1, \dots, 6\}$ (subscript addition taken modulo 6).

Lemma 2 *Every component of $G[Z(0)]$ is green.*

Proof: Let K be a component of $G[Z(0)]$. Since G is connected, there exists $v \in V \setminus K$ which contacts K . If v is universal for K , then $G[K]$ is P_3 -free (since G is diamond-free). Then let us assume that v is partial to K , and prove that $G[K]$ is green. Let us write $X = K \cap N(v)$, and $Y = K \setminus N(v)$. Since G is diamond-free, $G[X]$ is P_3 -free. Let T be a component of $G[Y]$. Then T is a module of $G[K]$: otherwise, for any $x \in X$ partial to T , one has that two adjacent vertices in T together with x , v and two vertices in C would induce a P_6 (since G is $(P_6, \text{diamond})$ -free, v is the endpoint of an induced P_3 involving two vertices in C). Then T is a clique (since G is diamond-free), i.e., $G[Y]$ is P_3 -free. Furthermore, each vertex in Y is adjacent to at most one vertex in each component of X , otherwise a diamond arises involving v . Then the lemma follows. \square

Let us fix any vertex of C , say v_2 .

Let us prove that WIS can be solved for $G[V \setminus N(v_2)]$ in $\mathcal{O}(n^6)$ time.

A partition of $V \setminus N(v_2)$ is given by $\{\{v_2, v_4, v_5, v_6\}, M_{1,3,4,6}, M_{1,3,5}, M_{1,3,4}, M_{1,4,5}, M_{3,5,6}, M_{3,6,1}, M_{4,6,1}, M_{6,3,4}, M_{1,3}, M_{1,4}, M_{1,5}, M_{3,5}, M_{3,6}, M_{4,6}, Z(0)\}$.

Since G is diamond-free, the sets $M_{1,3,4,6}, M_{1,3,4}, M_{1,4,5}, M_{3,5,6}, M_{3,6,1}, M_{4,6,1}, M_{6,3,4}$ have cardinality at most one. Then, by Observation 1, to our aim it is sufficient to prove that WIS can be solved for each induced subgraph of $G[U]$ in polynomial time, where a partition of U is given by

$\{M_{1,3,5}, M_{1,3}, M_{1,4}, M_{1,5}, M_{3,5}, M_{3,6}, M_{4,6}, Z(0)\}$.

Since G is diamond-free: $M_{1,3,5}, M_{1,3}, M_{1,4}, M_{1,5}, M_{3,5}, M_{3,6}, M_{4,6}$ are independent sets. Since G is $(P_6, \text{diamond})$ -free: $M_{1,3,5} \cup M_{1,3} \cup M_{1,5} \cup M_{3,5}$ is an independent set; $M_{1,3} \cup M_{1,5} \cup M_{3,5} \cup M_{4,6}$ has a co-join with $Z(0)$.

For any $W \subseteq U$, let us write $W^* = W \cap Z(0)$.

For any $W \subseteq U$, let us say that a component K of $G[W^*]$ is of:

type 1 if K is not a clique and there exists a vertex in $W \setminus Z(0)$ partial to K ;

type 2 if K is a clique and there exists a vertex in $W \setminus Z(0)$ partial to K ;

type 3 otherwise.

Let $T_1(W), T_2(W), T_3(W)$ respectively denote the union of components of $G[W^*]$ of type 1, 2, 3.

Let us fix a subset $W \subseteq U$.

Notice that $M_{1,3,5}$ is an independent set, since G is diamond-free.

Let us consider the following binary relation ' \leq_1 ' on $M_{1,3,5}$: for any $a, b \in M_{1,3,5}$, $a \leq_1 b$ if $N_{T_1(W)}(a) \subseteq N_{T_1(W)}(b)$. It is immediate to verify that $(M_{1,3,5}, \leq_1)$ is a partially ordered set.

Lemma 3 *Let $y \in M_{1,3,5}$ be maximal for $(M_{1,3,5}, \leq_1)$. Then $G[T_1(W) \setminus N(y)]$ is P_3 -free.*

Proof: First let us prove that y contacts every component Q of type 1 of $G[W^*]$. Assume for a contradiction that there exists a component Q_1 of type 1 of $G[W^*]$ such that y does not contact Q_1 . By definition of component of type 1, there exists $y_1 \in W \setminus Z(0)$ partial to Q_1 . Since G is P_6 -free (also recall that $W \subseteq U$), $y_1 \in M_{1,3,5}$. By the maximality of y there exists a vertex \tilde{q} in some component \tilde{Q} of type 1 of $G[W^*]$ such that \tilde{q} is adjacent to y and nonadjacent to y_1 . Then \tilde{q}, y, v_1, y_1 and two adjacent vertices of Q_1 induce a P_6 , a contradiction.

To conclude the proof of the lemma one has to show that if y contacts a component Q of type 1 of $G[W^*]$, then $G[Q \setminus N(y)]$ is P_3 -free. This can be shown by applying the last part of the argument of Lemma 2. \square

Let us consider the following binary relation ‘ \leq_2 ’ on $M_{1,3,5}$: for any $a, b \in M_{1,3,5}$, $a \leq_2 b$ if $N_{T_2(W)}(a) \subseteq N_{T_2(W)}(b)$. It is immediate to verify that $(M_{1,3,5}, \leq_2)$ is a partially ordered set.

Lemma 4 *Let $y \in M_{1,3,5}$ be maximal for $(M_{1,3,5}, \leq_2)$. If $T_1(W) = \emptyset$, then $G[(W \setminus N(y))^*]$ admits at most one component of type 2.*

Proof: Let Q_1 be a component of type 2 of $G[(W \setminus N(y))^*]$. By definition of component of type 2 there exists $y_1 \in (W \setminus N(y)) \setminus Z(0)$ partial to Q_1 . Since G is P_6 -free (also recall that $W \subseteq U$), $y_1 \in M_{1,3,5}$. Let $q_1 \in Q_1$ be adjacent to y_1 ; let $q' \in Q_1$ be nonadjacent to y_1 . Since $T_1(W) = \emptyset$, there exists a component of type 2 of $G[W^*]$, say \tilde{Q}_1 , such that $Q_1 \subseteq \tilde{Q}_1$. In particular, there exists $\tilde{q}_1 \in \tilde{Q}_1 \setminus Q_1$ adjacent to y and nonadjacent to y_1 , otherwise, by the maximality of y there would exist a vertex $t \in W \setminus Q_1$ adjacent to y and nonadjacent to y_1 , i.e., vertices q', q_1, y_1, v_1, y, t would induce a P_6 .

Let us prove that Q_1 is the unique component of type 2 of $G[(W \setminus N(y))^*]$. Assume for a contradiction that there exists another component of type 2, say Q_2 , of $G[(W \setminus N(y))^*]$. Notice that y_1 is nonadjacent to any vertex q_2 of Q_2 , otherwise $q', \tilde{q}_1, y, v_1, y_1$ and q_2 would induce a P_6 . Then let $y_2 \in M_{1,3,5}$ be partial to Q_2 . By symmetry, y_2 is nonadjacent to any vertex of Q_1 . Then a vertex of Q_2 , y_2, v_1, y_1, q_1, q' induce a P_6 , a contradiction. \square

Now, let us consider the following cases:

1. $T_1(U) = T_2(U) = \emptyset$.

Then each component K of $G[Z(0)]$ is a module of $G[U]$. Then, to our aim, one can assume that K is a singleton. In fact, one can contract K into a singleton u with $w(u) = \alpha_w(G[K])$: that can be done in polynomial time by Lemmas 2 and 1. So in general, one can assume that $Z(0)$ is an independent set. One can solve WIS in $G[U]$ by solving WIS in $G[U \setminus M_{1,4}]$ and in $G[U \setminus N(y)]$, for every $y \in M_{1,4}$.

That can be done in $\mathcal{O}(n^3)$ time. In fact, by the assumptions and by the above properties, one can verify that $G[U \setminus M_{1,4}]$ is triangle-free, and that $G[U \setminus N(y)]$ is triangle-free for every vertex $y \in M_{1,4}$ (in particular, no vertex of $M_{3,6} \setminus N(y)$ is adjacent to a vertex of $Z(0) \setminus N(y)$, otherwise a P_6 arises). Then the assertion follows by Theorem 2.

For the other two cases we note that the existence of a component Q of type 1 or type 2 in $U \cap Z(0)$ implies the existence of a vertex $a \in M_{1,3,5}$ which is partial to Q , similarly to the proof above. So, $M_{1,3,5}$ is nonempty.

2. $T_1(U) = \emptyset, T_2(U) \neq \emptyset$.

Based on $(M_{1,3,5}, \leq_2)$, the vertices y_1, \dots, y_h of $M_{1,3,5}$ can be totally ordered so that y_i is maximal for $(\{y_i, \dots, y_h\}, \leq_2)$ for $i = 1, \dots, h$. Then one can solve WIS in $G[U]$ by sequentially solving WIS in $G[U \setminus N(y_1)]$, in $G[(U \setminus \{y_1, \dots, y_{i-1}\}) \setminus N(y_i)]$ for $i = 2, \dots, h$, and in $G[U \setminus M_{1,3,5}]$.

That can be done in $\mathcal{O}(n^5)$ time. In fact, let us first consider $G[U \setminus N(y_1)]$. If $G[(U \setminus N(y_1))^*]$ admits no component of type 2, then one can refer to Case 1. Otherwise, by Lemma 4, $G[(U \setminus N(y_1))^*]$ admits a unique component of type 2, say Q . Then one can solve WIS in $G[U \setminus N(y_1)]$ by solving WIS in $G[(U \setminus N(y_1)) \setminus Q]$ and in $G[(U \setminus N(y_1)) \setminus N(q)]$, for every $q \in Q$: since for each of such graphs $G[H]$ one has that $G[H^*]$ has no component of type 2, one can refer to Case 1 and to Lemmas 1 and 2. Now, let us consider $G[(U \setminus \{y_1, \dots, y_{i-1}\}) \setminus N(y_i)]$ for $i = 2, \dots, h$: by the mentioned total order, one can apply the argument applied for $G[U \setminus N(y_1)]$ in order to show that WIS can be

solved for such graphs in polynomial time. Finally, let us consider $G[U \setminus M_{1,3,5}]$: since no vertex in $U \setminus M_{1,3,5}$ is partial to any component of $G[Z(0)]$ (otherwise a P_6 arises), one can refer to Case 1.

3. $T_1(U) \neq \emptyset$.

Based on $(M_{1,3,5}, \leq_1)$, the vertices y_1, \dots, y_h of $M_{1,3,5}$ can be totally ordered so that y_i is maximal for $(\{y_i, \dots, y_h\}, \leq_1)$ for $i = 1, \dots, h$. Then one can solve WIS in $G[U]$ by sequentially solving WIS in $G[U \setminus N(y_1)]$, in $G[(U \setminus \{y_1, \dots, y_{i-1}\}) \setminus N(y_i)]$ for $i = 2, \dots, h$, and in $G[U \setminus M_{1,3,5}]$.

That can be done in $\mathcal{O}(n^6)$ time. In fact, let us first consider $G[U \setminus N(y_1)]$. By Lemma 3, $G[T_1(W) \setminus N(y_1)]$ is P_3 -free. Then $G[(U \setminus N(y_1))^*]$ admits no component of type 1. Then one can refer to Case 2 and to Lemmas 1 and 2. Now, let us consider $G[(U \setminus \{y_1, \dots, y_{i-1}\}) \setminus N(y_i)]$ for $i = 2, \dots, h$: by the mentioned total order, one can apply the argument applied for $G[U \setminus N(y_1)]$ in order to show that WIS can be solved for such graphs in polynomial time. Finally, let us consider $G[U \setminus M_{1,3,5}]$: since no vertex in $U \setminus M_{1,3,5}$ is partial to any component of $G[Z(0)]$ (otherwise a P_6 arises), one can refer to Case 1.

Let us summarize the above argument as follows.

Theorem 3 *Let $G = (V, E)$ be a $(P_6, \text{diamond})$ -free graph containing a 6-cycle C . Then one can solve WIS for $G[V \setminus N(c)]$ in $\mathcal{O}(n^6)$ time, for any vertex c of C . \square*

3.2 Deleting C_5 's in $(P_6, \text{diamond}, C_6)$ -free graphs

Throughout this subsection assume that $G = (V, E)$ is a $(P_6, \text{diamond}, C_6)$ -free graph containing a 5-cycle C , say with vertices v_i and edges $v_i v_{i+1}$, $i \in \{1, \dots, 5\}$ (subscript addition taken modulo 5). Let $N(C)$ be the set of vertices from $V \setminus C$ which are adjacent to some vertex in C . For any subset S of C , let M_S be the set formed by those vertices in $N(C)$ which are adjacent to each vertex in S and are nonadjacent to each vertex in $C \setminus S$. In particular, let us write M_1 for $S = \{v_1\}$, $M_{1,2}$ for $S = \{v_1, v_2\}$, and so on. Then let $Z(k)$ denote the set of vertices of $N(C)$ having exactly k neighbors in C .

Since G is $(P_6, \text{diamond})$ -free: $Z(4) = Z(5) = \emptyset$; each element of $Z(3)$ belongs to some of the sets $M_{i, i+2, i+3}$, $i \in \{1, \dots, 5\}$ (subscript addition taken modulo 5).

Similarly to the previous subsection, one has the following fact.

Lemma 5 *Every component of $G[Z(0)]$ is green. \square*

Lemma 6 *There exists a vertex c of C such that one of the following statements holds:*

- (i) $M_i \setminus N(c) = \emptyset$ for all $i \in \{2, \dots, 5\}$, and $M_{i, i+1} \setminus N(c) = \emptyset$ for all $i \in \{1, \dots, 5\}$ (subscript addition taken modulo 5);
- (ii) $M_i \setminus N(c) = \emptyset$ for all $i \in \{1, \dots, 5\}$, and $M_{i, i+1} \setminus N(c) = \emptyset$ for all $i \in \{1, \dots, 4\}$ (subscript addition taken modulo 5).

Proof: Since G is (P_6, C_6) -free, for all $i \in \{1, \dots, 5\}$ (subscript addition taken modulo 5) one has that: if $M_i \neq \emptyset$, then $M_{i+2} = M_{i+3} = M_{i+1, i+2} = M_{i+3, i+4} = \emptyset$; if $M_{i, i+1} \neq \emptyset$, then $M_{i-1, i} = M_{i+1, i+2} = \emptyset$. This implies the lemma. \square

Let us fix any vertex of C , say v_2 , guaranteed by Lemma 6.

Let us prove that one can solve WIS for $G[V \setminus N(v_2)]$ in $\mathcal{O}(n^4)$ time.

A partition of $V \setminus N(v_2)$ is given by $\{\{v_2, v_4, v_5\}, M_{1,3,4}, M_{3,4,5}, M_{1,3}, M_{1,4}, M_{3,5}, M_1, Z(0)\}$.

Since G is diamond-free, the sets $M_{1,3,4}, M_{3,4,5}$ have cardinality at most one. Then, by Observation 1, to our aim it is sufficient to prove that WIS can be solved for each induced subgraph of $G[U]$ in polynomial time, where a partition of U is given by $\{M_{1,3}, M_{1,4}, M_{3,5}, M_1, Z(0)\}$.

Since G is diamond-free, $M_{1,3}, M_{1,4}, M_{3,5}$ are independent sets. Since G is P_6 -free: M_1 has a co-join with $Z(0)$; each vertex in $M_{1,3} \cup M_{1,4} \cup M_{3,5}$ is not partial to any component of $G[Z(0)]$, i.e., each component of $G[Z(0)]$ is a module of $G[U]$. So by assertions similar to Lemmas 5 and 1, one can assume that $Z(0)$ is an independent set.

Now, let us consider the following cases, which are exhaustive by symmetry.

Case 1 statement i of Lemma 6 holds.

Case 1.1 $M_1 = \emptyset$. One can solve WIS in $G[U]$ by solving WIS in $G[U \setminus M_{3,5}]$ and in $G[U \setminus N(y)]$ for every $y \in M_{3,5}$. Since G is diamond-free, one can verify that such graphs are triangle-free. Then in this case one can solve WIS for $G[U]$ in $\mathcal{O}(n^3)$ time by Theorem 2.

Case 1.2 $M_1 \neq \emptyset$.

First assume that M_1 is a clique. One can solve WIS in $G[U]$ by solving WIS in $G[U \setminus M_1]$ and in $G[U \setminus N(y)]$ for every $y \in M_1$. Then, by referring to Case 1.1, in this case one can solve WIS for $G[U]$ in $\mathcal{O}(n^4)$ time.

Then assume that M_1 is not a clique, i.e., $G[M_1]$ is disconnected (since $G[M_1]$ is P_3 -free). Then $M_{3,5}$ is partitioned into $\{M_{3,5}^0, M'_{3,5}\}$, where $M'_{3,5} = \{x \in M_{3,5} : x \text{ is universal for } M_1\}$, and $M_{3,5}^0 = \{x \in M_{3,5} : x \text{ does not contact } M_1\}$ (in fact if $y, z \in M_1$ are nonadjacent and a vertex $x \in M_{3,5}$ is adjacent to y and nonadjacent to z , then v_4, v_3, x, y, v_1, z induce a P_6). One can solve WIS in $G[U]$ by solving WIS in $G[U \setminus M'_{3,5}]$ and in $G[U \setminus N(y)]$ for every $y \in M'_{3,5}$.

That can be done in $\mathcal{O}(n^4)$ time. Concerning graphs $G[U \setminus N(y)]$ for every $y \in M'_{3,5}$, one can refer to Case 1.1. Then let us consider $G[U \setminus M'_{3,5}]$. Notice that $M_{3,5}^0$ has a co-join with $Z(0)$, otherwise a P_6 arises involving a vertex of M_1 . Then $U \setminus M'_{3,5}$ is partitioned into $\{X, Y\}$, where $X = M_1 \cup M_{1,3} \cup M_{1,4}$ (i.e., $G[X]$ is P_3 -free) and $Y = M_{3,5}^0 \cup Z(0)$ (i.e., Y is an independent set). Then each component of $G[U \setminus M'_{3,5}]$ is either P_3 -free or green. Then the assertion follows by Lemma 1.

Case 2 statement ii of Lemma 6 holds.

One can solve WIS in $G[U]$ by solving WIS in $G[U \setminus M_{4,5}]$ and in $G[U \setminus N(y)]$ for every $y \in M_{4,5}$. Since G is diamond-free, $M_{4,5}$ is a clique. Then, by referring to Case 1.1 (3.2), in this case one can solve WIS for $G[U]$ in $\mathcal{O}(n^4)$ time.

Let us summarize the above argument as follows.

Theorem 4 Let $G = (V, E)$ be a $(P_6, \text{diamond}, C_6)$ -free graph containing a 5-cycle C .

Then there exists a vertex c of C (which can be easily found) such that one can solve WIS for $G[V \setminus N(c)]$ in $\mathcal{O}(n^4)$ time. \square

3.3 Deleting C_4 's in $(P_6, \text{diamond}, C_6, C_5)$ -free graphs

Throughout this subsection assume that $G = (V, E)$ is a $(P_6, \text{diamond}, C_6, C_5)$ -free graph containing a 4-cycle C , say with vertices v_i and edges $v_i v_{i+1}$, $i \in \{1, \dots, 4\}$ (subscript addition taken modulo 4). Let $N(C)$ be the set of vertices from $V \setminus C$ which are adjacent to some vertex in C . For any subset S of C , let M_S be the set formed by those vertices in $N(C)$ which are adjacent to each vertex in S and are nonadjacent to each vertex in $C \setminus S$. In particular, let us write M_1 for $S = \{v_1\}$, $M_{1,2}$ for $S = \{v_1, v_2\}$, and so on. Then let $Z(k)$ denote the set of vertices of $N(C)$ having exactly k neighbors in C .

Since G is $(P_6, \text{diamond})$ -free: $Z(3) = Z(4) = \emptyset$.

Similarly to the previous subsection, one has the following fact.

Lemma 7 *Every component of $G[Z(0)]$ is green.* □

Let us fix any vertex of C , say v_2 .

Let us prove that WIS can be solved for $G[V \setminus N(v_2)]$ in $\mathcal{O}(n^6)$ time.

A partition of $V \setminus N(v_2)$ is given by $\{\{v_2, v_4\}, M_{1,3}, M_{3,4}, M_{4,1}, M_1, M_3, M_4, Z(0)\}$. Then, by Observation 1, to our aim it is sufficient to prove that WIS can be solved for each induced subgraph of $G[U]$ in polynomial time, where a partition of U is given by $\{M_{1,3}, M_{3,4}, M_{4,1}, M_1, M_3, M_4, Z(0)\}$,

Let us introduce some preliminary definitions and lemmas.

Let us write:

$$\begin{aligned} M_1^0 &= \{x \in M_1 : x \text{ does not contact } M_4\} \\ M_3^0 &= \{x \in M_3 : x \text{ does not contact } M_4\} \\ M_4^0 &= \{x \in M_4 : x \text{ does not contact } M_1 \cup M_3\} \\ X &= \{x \in M_1 : x \text{ contacts } M_4\} \cup \{x \in M_4 : x \text{ contacts } M_1\} \\ Y &= \{x \in M_3 : x \text{ contacts } M_4\} \cup \{x \in M_4 : x \text{ contacts } M_3\} \end{aligned}$$

Let us write:

$$\begin{aligned} Z_1 &= M_1^0 \cup \{z \in Z(0) : z \text{ contacts } M_1^0\} \\ Z_3 &= M_3^0 \cup \{z \in Z(0) : z \text{ contacts } M_3^0\} \\ Z_4 &= M_4^0 \cup \{z \in Z(0) : z \text{ contacts } M_4^0\} \\ Z_X &= X \cup \{z \in Z(0) : z \text{ contacts } X\} \\ Z_Y &= Y \cup \{z \in Z(0) : z \text{ contacts } Y\} \\ \tilde{Z} &= \{z \in Z(0) : z \text{ does not contact } M_1 \cup M_3 \cup M_4\}. \end{aligned}$$

Lemma 8 *The following facts hold:*

- (i) *each pair of the sets $Z_1, Z_3, Z_4, Z_X, Z_Y, \tilde{Z}$ has a co-join;*
- (ii) *each component of $G[Z_1 \cup Z_3 \cup Z_4 \cup \tilde{Z}]$ is green;*
- (iii) *each component of $G[Z_X \cup Z_Y]$ is either P_3 -free or bipartite.*

Proof:

Proof of i. Since G is C_5 -free, M_1 has a co-join with M_3 . Then, since G is C_6 -free, no vertex in M_4 can be adjacent to a vertex in M_1 and to a vertex in M_3 at the same time. It follows that each pair of the sets $M_1^0, M_3^0, M_4^0, X, Y$ has a co-join. Furthermore, since G is (P_6, C_6) -free, one can verify that two vertices chosen in two different sets – among the mentioned sets – cannot contact a component of $Z(0)$ at the same time. Finally, since G is P_6 -free, if a vertex in $y \in M_1 \cup M_3 \cup M_4$ contacts a component K of $G[Z(0)]$, then y is universal for K . Then i follows.

Proof of ii. It is enough to deal with $G[Z_1]$, as the other subsets can be treated similarly. If $Z_1 = M_1^0$, then $G[Z_1]$ is P_3 -free. Otherwise, since G is $(P_6, \text{diamond})$ -free, one can verify (similarly to the argument of Lemma 2) that each component of $G[Z_1]$ is green.

Proof of iii. It is enough to deal with $G[Z_1]$, as the other subsets can be treated similarly. If $Z_X = X$, then since G is $(\text{diamond}, C_5)$ -free, each component of $G[Z_X]$ is bipartite. Otherwise, since G is P_6 -free, each vertex in $Z(0)$ contacting a component K of $G[X]$ dominates K . Then, since G is diamond-free, each component of $G[Z_X]$ is a clique. \square

Lemma 9 *If $M_1 \neq \emptyset$ and $M_3 \neq \emptyset$, then:*

- (i) $Z(0)$ has a co-join with $M_1 \cup M_3$;
- (ii) $X = Y = \emptyset$.

Proof:

Proof of i. It follows since G is (P_6, C_6, C_5) -free.

Proof of ii. By symmetry, let us only prove that $X = \emptyset$. Assume for a contradiction that $X \neq \emptyset$. Then let $x_1 \in M_1$ be adjacent to $x_4 \in M_4$. Let $x_3 \in M_3$. By i of Lemma 8, x_1 and x_4 are nonadjacent to x_3 . Then $x_4, x_1, v_1, v_2, v_3, x_3$ induce a P_6 . \square

Lemma 10 *If a vertex $y \in M_{1,3}$ contacts a component K of $G[Z_X \cup Z_Y]$, then $K \setminus N(y)$ is either a clique or an independent set.*

Proof: By symmetry, let us consider only $G[Z_X]$. Let K be a component of $G[X]$. If $G[K]$ is P_3 -free, then the assertion trivially follows. Then, by iii of Lemma 8, assume that $G[K]$ is bipartite. Let $y \in M_{1,3}$ contact K . Notice that y cannot be adjacent to two adjacent vertices of K , otherwise a diamond arises involving v_1 . Then, to avoid a P_6 , y is adjacent to all the vertices of a side of the bipartite graph, i.e., $K \setminus N(y)$ is an independent set. \square

Let us write $Z = Z_1 \cup Z_3 \cup Z_4 \cup Z_X \cup Z_Y \cup \tilde{Z}$. Then $\{M_{1,3}, Z\}$ is a partition of U .

For any $W \subseteq U$, let us write $W^* = W \cap Z$.

For any $W \subseteq U$, let us say that a component K of $G[W^*]$ is of:

type 1 if K is not a clique and there exists a vertex in $W \setminus Z$ partial to K ;

type 2 if K is a clique and there exists a vertex in $W \setminus Z$ partial to K ;

type 3 otherwise.

Let $T_1(W), T_2(W), T_3(W)$ respectively denote the union of components of $G[W^*]$ of type 1, 2, 3.

Let us fix a subset $W \subseteq U$.

Notice that $M_{1,3}$ is an independent set, since G is diamond-free.

Let us consider the following binary relation ' \leq_1 ' on $M_{1,3}$: for any $a, b \in M_{1,3}$, $a \leq_1 b$ if $N_{T_1(W)}(a) \subseteq N_{T_1(W)}(b)$. It is immediate to verify that $(M_{1,3}, \leq_1)$ is a partially ordered set.

Lemma 11 *Let $y \in M_{1,3}$ be maximal for $(M_{1,3}, \leq_1)$. Then $G[T_1(W) \setminus N(y)]$ is P_3 -free.*

Proof: If either $M_1 = \emptyset$ or $M_3 = \emptyset$, then one can apply an argument similar to that of Lemma 3, by considering also (i) of Lemma 8 and Lemma 10 – in detail, if $M_1 = \emptyset$ (if $M_3 = \emptyset$), then vertex v_1 (vertex v_3) is universal for $M_{1,3}$ and does not contact Z .

If $M_1 \neq \emptyset$ and $M_3 \neq \emptyset$, then one can apply an argument similar to that of Lemma 3, by considering Lemma 9 and the fact that no element of $M_{1,3}$ can be partial to a component of $G[M_1 \cup M_3]$ (since G is diamond-free). \square

Let us consider the following binary relation ' \leq_2 ' on $M_{1,3}$: for any $a, b \in M_{1,3}$, $a \leq_2 b$ if $N_{T_2(W)}(a) \subseteq N_{T_2(W)}(b)$. It is immediate to verify that $(M_{1,3}, \leq_2)$ is a partially ordered set.

Lemma 12 *Let $y \in M_{1,3}$ be maximal for $(M_{1,3}, \leq_2)$. If $T_1(W) = \emptyset$, then $G[(W \setminus N(y))^*]$ admits at most one component of type 2.*

Proof: One can apply the argument in the proof of Lemma 11, by considering Lemma 4 instead of Lemma 3. \square

Now, let us consider the following cases.

Case 1 $M_{3,4} = M_{4,1} = \emptyset$.

Case 1.1 $T_1(U) = T_2(U) = \emptyset$.

Then each component K of $G[Z]$ is a module of $G[U]$. Then, to our aim, one can assume that K is a singleton. In fact, one can contract K into a singleton k with $w(k) = \alpha_w(G[K])$: that can be done in $\mathcal{O}(n^3)$ time by ii–iii of Lemma 8 and Lemma 1.

So in general, one can assume that Z is an independent set. Then $G[U]$ is bipartite. In this case one can solve WIS for $G[U]$ in time $\mathcal{O}(n^3)$ by Theorem 2.

Case 1.2 $T_1(U) = \emptyset, T_2(U) \neq \emptyset$.

Based on $(M_{1,3}, \leq_2)$, the vertices y_1, \dots, y_h of $M_{1,3}$ can be totally ordered so that y_i is maximal for $(\{y_i, \dots, y_h\}, \leq_2)$ for $i = 1, \dots, h$. Then one can solve WIS in $G[U]$ by sequentially solving WIS in $G[U \setminus N(y_1)]$, in $G[(U \setminus \{y_1, \dots, y_{i-1}\}) \setminus N(y_i)]$ for $i = 2, \dots, h$, and in $G[U \setminus M_{1,3}]$.

That can be done in $\mathcal{O}(n^5)$ time. In fact, let us first consider $G[U \setminus N(y_1)]$. If $G[(U \setminus N(y_1))^*]$ admits no component of type 2, then one can refer to Case 1.1. Otherwise, by Lemma 12, $G[(U \setminus N(y_1))^*]$ admits a unique component of type 2, say Q . Then one can solve WIS in $G[U \setminus N(y_1)]$ by solving WIS in $G[(U \setminus N(y_1)) \setminus Q]$ and in $G[(U \setminus N(y_1)) \setminus N(q)]$, for every $q \in Q$: since for each of such graphs $G[H]$ one has that $G[H^*]$ has no component of type 2, one can refer to Case 1.1, to ii–iii of Lemma 8 and to Lemmas 7 and 1. Now, let us consider $G[(U \setminus \{y_1, \dots, y_{i-1}\}) \setminus N(y_i)]$ for $i = 2, \dots, h$: by the mentioned total order, one can apply the argument applied for $G[U \setminus N(y_1)]$ in

order to show that WIS can be solved for such graphs in polynomial time. Finally, let us consider $G[U \setminus M_{1,3}] = G[Z]$: then one can refer to Case 1.1.

Case 1.3 $T_1(U) \neq \emptyset$.

Based on $(M_{1,3}, \leq_2)$, the vertices y_1, \dots, y_h of $M_{1,3}$ can be totally ordered so that y_i is maximal for $(\{y_i, \dots, y_h\}, \leq_2)$ for $i = 1, \dots, h$. Then one can solve WIS in $G[U]$ by sequentially solving WIS in $G[U \setminus N(y_1)]$, in $G[(U \setminus \{y_1, \dots, y_{i-1}\}) \setminus N(y_i)]$ for $i = 2, \dots, h$, and in $G[U \setminus M_{1,3}]$.

That can be done in $\mathcal{O}(n^6)$ time. In fact, let us first consider $G[U \setminus N(y_1)]$. By Lemma 11, $G[T_1(W) \setminus N(y_1)]$ is P_3 -free. Then $G[(U \setminus N(y_1))^*]$ admits no component of type 1. Then one can refer to Case 1.2 and to Lemmas 7 and 1. Now, let us consider $G[(U \setminus \{y_1, \dots, y_{i-1}\}) \setminus N(y_i)]$ for $i = 2, \dots, h$: by the total order, one can apply the argument applied for $G[U \setminus N(y_1)]$ in order to show that WIS can be solved for such graphs in polynomial time. Finally, let us consider $G[U \setminus M_{1,3}] = G[Z]$: then one can refer to Case 1.1.

Case 2 $M_{3,4} \cup M_{4,1} \neq \emptyset$.

Since G is diamond-free, $M_{3,4}$ and $M_{4,1}$ are cliques. Since G is $(P_6, \text{diamond}, C_6, C_5)$ -free, the following facts hold: $M_{3,4} \cup M_{4,1}$ has a co-join with $N(C) \setminus (N(v_2) \cup M_{3,4} \cup M_{4,1})$; a vertex in $M_{3,4} \cup M_{4,1}$ and a vertex in $N(C) \setminus (N(v_2) \cup M_{3,4} \cup M_{4,1})$ cannot contact a component of $G[Z(0)]$ at the same time; $M_{3,4}$ has a co-join with $M_{4,1}$.

First assume that $M_{3,4} \neq \emptyset$ and $M_{4,1} \neq \emptyset$. Then $M_{3,4} \cup M_{4,1}$ has a co-join with $Z(0)$ (otherwise a P_6 or a C_6 arises). In general, by the above facts, $M_{3,4} \cup M_{4,1}$ has a co-join with $U \setminus (M_{3,4} \cup M_{4,1})$. Then, since $M_{3,4}$ and $M_{4,1}$ are cliques, one can directly refer to Case 1.

Then assume that $M_{3,4} = \emptyset$. One can solve WIS for $G[U]$ by solving WIS in $G[U \setminus M_{4,1}]$ and in $G[U \setminus N(y)]$, for every $y \in M_{4,1}$. Since $M_{4,1}$ is a clique, one can directly refer to Case 1.

The case in which $M_{4,1} = \emptyset$ can be similarly treated, by symmetry.

Let us summarize the above argument as follows.

Theorem 5 *Let $G = (V, E)$ be a $(P_6, \text{diamond}, C_6, C_5)$ -free graph containing a 4-cycle C . Then one can solve WIS for $G[V \setminus N(c)]$ in $\mathcal{O}(n^6)$ time, for any vertex c of C . \square*

3.4 A solution for WIS and WID in $(P_6, \text{diamond})$ -free graphs

In this subsection we formalize an efficient method for solving WIS (or WID) in $(P_6, \text{diamond})$ -free graphs. To this end, let us first summarize the results of the previous subsections in the following theorem.

Theorem 6 *Let $G = (V, E)$ be a connected $(P_6, \text{diamond})$ -free graph containing a C_6 or a C_5 or a C_4 . Then there exists a vertex c (which can be easily found) such that one can solve WIS (or WID) for $G[V \setminus N(c)]$ in $\mathcal{O}(n^6)$ time.*

Proof: If G contains a C_6 , then the assertion follows by Theorem 3. If G is C_6 -free and contains a C_5 , then the assertion follows by Theorem 4. If G is (C_6, C_5) -free and contains a C_4 , then the assertion follows by Theorem 5. \square

To prove that WIS (or WID) is solvable in polynomial time on the class of $(P_6, \text{diamond})$ -free graphs, it suffices to find a polynomial upper bound $p(n) = \mathcal{O}(n^7)$ on the number of steps sufficient for any

allowed input of order n . If G is chordal, then we are done by Theorem 1. Otherwise, there exists a sixth-degree polynomial $q(n)$ with the property that in any $(P_6, \text{diamond})$ -free non-chordal graph one can determine a vertex x such that WIS (or WID) can be solved on $G_1 = G[V \setminus N(x)]$ in $q(n)$ time. If $G'_1 = G - x$ is not chordal, then again one can find a vertex x' such that the problem can be solved on $G_2 = G[V(G'_1) \setminus N(x')]$ in $q(n)$ time, and so on. In this way we obtain some graphs G_1, G_2, \dots, G_k with $k < n$, such that each G_i is either chordal or admits an efficient WIS (or WID) algorithm. Thus, the total running time is $O(n(q(n) + r(n)))$ where $r(n)$ is the time needed to check whether the current graph is chordal and if it is not, then to find a suitable vertex under the conditions of Theorem 6.

Now, by Theorems 1 and 6 one obtains:

Theorem 7 *Both the WIS and the WID problems can be solved for $(P_6, \text{diamond})$ -free graphs in $\mathcal{O}(n^7)$ time. \square*

Acknowledgements

I would like to thank the referees for their valuable remarks and suggestions, and the handling editor for his helpful and resolving comments.

References

- [1] V.E. Alekseev, On the local restriction effect on the complexity of finding the graph independence number, *Combinatorial-algebraic Methods in Applied Mathematics*, Gorkiy University Press, Gorkiy (1983) 3-13 (in Russian)
- [2] V.E. Alekseev, A polynomial algorithm for finding largest independent sets in fork-free graphs, *Discrete Analysis and Operations Research Ser. 1*, 6 (1999) 3-19 (in Russian), *Discrete Applied Mathematics* 135 (2004) 3-16
- [3] V.E. Alekseev, On easy and hard hereditary classes of graphs with respect to the independent set problem, *Discrete Applied Mathematics* 132 (2004) 17-26
- [4] C. Arbib, R. Mosca, On $(P_5, \text{diamond})$ -free graphs, *Discrete Mathematics* 250 (2002) 1-22
- [5] G. Bacsó, Zs. Tuza, Dominating cliques in P_5 -free graphs, *Periodica Mathematica Hungarica* 21, 4 (1990) 303-308
- [6] G. Bacsó, Zs. Tuza, A characterization of graphs without long induced paths, *J. Graph Theory* 14, 4 (1990) 455-464
- [7] R. Boliac, V.V. Lozin, Independent domination in finitely defined classes of graphs, *Theoretical Computer Science* 301 (2003) 271-284
- [8] A. Brandstädt, $(P_5, \text{diamond})$ -free graphs revisited: structure and linear time optimization, *Discrete Applied Mathematics* 138 (2004) 13-27
- [9] A. Brandstädt, Chính T. Hoàng, On clique separators, nearly chordal graphs and the Maximum Weight Stable Set problem, M. Jünger and V. Kaibel (Eds.): IPCO 2005, LNCS 3509, pp. 265-275, 2005
- [10] A. Brandstädt, H.-O. Le, V.B. Le, On α -redundant vertices in P_5 -free graphs, *Information Processing Letters* 82 (2002) 119-122
- [11] A. Brandstädt, V.B. Le, S. Mahfud, New applications of clique separators decomposition for the Maximum Weight Stable Set problem, *Theoretical Computer Science* 370 (2007) 229-239
- [12] A. Brandstädt, V.B. Le, J.P. Spinrad, Graph Classes: A Survey, *SIAM Monographs on Discrete Math. Appl.*, Vol. 3, SIAM, Philadelphia (1999)
- [13] A. Brandstädt, V.V. Lozin, A note on α -redundant vertices in graphs, *Discrete Applied Mathematics* 108 (2001) 301-308
- [14] A. Brandstädt, T. Klemmt, S. Mahfud, P_6 - and Triangle-Free Graphs Revisited: Structure and Bounded Clique-Width, *Discrete Mathematics and Theoretical Computer Science* 8 (2006) 173-188
- [15] A. Brandstädt, D. Kratsch, On domination problems on permutation and other graphs, *Theoretical Computer Science* 54 (1987) 181-198

- [16] A. Brandstädt, D. Kratsch, On the structure of (P_5, gem) -free graphs, *Discrete Applied Mathematics* 145 (2005) 155-166
- [17] A. Brandstädt, R. Mosca, On the structure and stability number of P_5 - and co-chair-free graphs, *Discrete Applied Mathematics* 132 (2004) 47-65
- [18] G. Chang, The weighted independent domination problem is NP-complete for chordal graphs, *Discrete Applied Mathematics* 143 (2004) 351-352
- [19] J. Dong, On the i -diameter of i -center in a graph without long induced paths, *J. Graph Theory* 30 (1999) 235-241
- [20] M. Farber, Independent domination in chordal graphs, *Operations Research Letters* 1 (1982) 134-138
- [21] M. Farber, On diameters and radii of bridged graphs, *Discrete Mathematics* 73 (1989) 249-260
- [22] M. Farber, M. Hujter, Zs. Tuza, An upper bound on the number of cliques in a graph, *Networks* 23 (1993) 75-83
- [23] J.-L. Fouquet, V. Giakoumakis, F. Maire, H. Thuillier, On graphs without P_5 and \overline{P}_5 , *Discrete Mathematics* 146 (1995) 33-44
- [24] J.-L. Fouquet, V. Giakoumakis, J.-M. Vanherpe, Bipartite graphs totally decomposable by canonical decomposition, *International Journal of Foundations of Computer Science* 10 (1999) 513-533
- [25] A. Frank, Some polynomial algorithms for certain graphs and hypergraphs, *Proceedings of the Fifth British Combinatorial Conference* (Univ. Aberdeen, Aberdeen 1975) 211-226; *Congressus Numerantium* No. XV, Utilitas Math., Winnipeg, Man. (1976)
- [26] M.R. Garey, D.S. Johnson, L. Stockmeyer, Some simplified NP-complete graph problems, *Theoretical Computer Science* 1 (1976) 237-267
- [27] M.R. Garey, D.S. Johnson, The rectilinear Steiner tree problem is NP-complete, *SIAM J. Applied Mathematics* 32 (1977) 826-834
- [28] M.R. Garey, D.S. Johnson, *Computers and Intractability: A Guide to the Theory of NP-completeness*, Freeman, San Francisco, CA (1979)
- [29] M. Gerber, V.V. Lozin, On the stable set problem in special P_5 -free graphs, *Discrete Applied Mathematics* 125 (2003) 215-224
- [30] V. Giakoumakis, J.-M. Vanherpe, Linear time recognition and optimization for weak-bisplit graphs, bi-cographs and bipartite P_6 -free graphs, *International Journal of Foundations of Computer Science* 14 (2003) 107-136
- [31] V. Giakoumakis, I. Rusu, Weighted parameters in (P_5, \overline{P}_5) -free graphs, *Discrete Applied Mathematics* 80 (1997) 255-261
- [32] M.C. Golumbic, U. Rotics, On the clique-width of some perfect graph classes, *International Journal of Foundations of Computer Science* 11 (2000) 423-443

- [33] M. Grötschel, L. Lovász, A. Schrijver, Polynomial algorithms for perfect graphs, *Annals of Discrete Mathematics* 21 (1984) 325-356
- [34] R.M. Karp, Reducibility among combinatorial problems, *Complexity of Computer Computations*, R.E. Miller, J.W. Thatcher (eds.), Plenum Press, New York (1972) 85-103
- [35] J. Liu, H. Zhou, Dominating subgraphs in graphs with some forbidden structure, *Discrete Mathematics* 135 (1994) 163-168
- [36] V.V. Lozin, Stability in P_5 - and banner-free graphs, *European J. Operational Research* 125 (2000) 292-297
- [37] V.V. Lozin, M. Milanič, A polynomial algorithm to find an independent set of maximum weight in a fork-free graph, RUTCOR Research Report, Rutgers University, 30-2005
- [38] V.V. Lozin, R. Mosca, Independent sets in extensions of $2K_2$ -free graphs, *Discrete Applied Mathematics* 146 (2005) 74-80
- [39] V.V. Lozin, D. Rautenbach, Some results on graphs without long induced paths, *Information Processing Letters* 88 (2003) 167-171
- [40] G.J. Minty, On maximal independent sets of vertices in claw-free graphs, *J. Combinatorial Theory, Series B* 28 (1980) 284-304
- [41] R. Mosca, Stable sets in certain P_6 -free graphs, *Discrete Applied Mathematics* 92 (1999) 177-191
- [42] D. Nakamura, A. Tamura, A revision of Minty's algorithm for finding a maximum weight stable set in a claw-free graph, *J. Operations Research Society of Japan* 44 (2001) 194-204
- [43] S. Poljak, A note on stable sets and colorings of graphs, *Commun. Math. Univ. Carolinae* 15 (1974) 307-309
- [44] B. Randerath, I. Schiermeyer, M. Tewes, Three-colourability and forbidden subgraphs. II: polynomial algorithms, *Discrete Mathematics* 251 (2002) 137-153
- [45] N. Sbihi, Algorithme de recherche d'un stable de cardinalité maximum dans un graphe sans étoile, *Discrete Mathematics* 29 (1980) 53-76
- [46] I.E. Zverovich, Independent domination on $2P_3$ -free perfect graphs, DIMACS Technical Report 2003-22, Rutgers University, 2003
- [47] I.E. Zverovich, V.E. Zverovich, Locally well-dominated and locally independent well-dominated graphs, *Graphs and Combinatorics* 19 (2) (2003) 279-288