

Self-complementing permutations of k -uniform hypergraphs

Artur Szymański, Adam Pawel Wojda

► **To cite this version:**

Artur Szymański, Adam Pawel Wojda. Self-complementing permutations of k -uniform hypergraphs. Discrete Mathematics and Theoretical Computer Science, DMTCS, 2009, 11 (1), pp.117–123. <hal-00988180>

HAL Id: hal-00988180

<https://hal.inria.fr/hal-00988180>

Submitted on 7 May 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Self-complementing permutations of k -uniform hypergraphs

Artur Szymański and A. Paweł Wojda

Faculty of Applied Mathematics, AGH University of Science and Technology, Al. Mickiewicza 30, 30-053 Kraków, Poland

received January 20, 2008, revised January 16, 2009, accepted January 23, 2009.

A k -uniform hypergraph $H = (V; E)$ is said to be *self-complementary* whenever it is isomorphic with its complement $\overline{H} = (V; \binom{V}{k} - E)$. Every permutation σ of the set V such that $\sigma(e)$ is an edge of \overline{H} if and only if $e \in E$ is called *self-complementing*. 2-self-complementary hypergraphs are exactly self-complementary graphs introduced independently by Ringel (1963) and Sachs (1962).

For any positive integer n we denote by $\lambda(n)$ the unique integer such that $n = 2^{\lambda(n)}c$, where c is odd.

In the paper we prove that a permutation σ of $[1, n]$ with orbits O_1, \dots, O_m is a self-complementing permutation of a k -uniform hypergraph of order n if and only if there is an integer $l \geq 0$ such that $k = a2^l + s$, a is odd, $0 \leq s < 2^l$ and the following two conditions hold:

- (i) $n = b2^{l+1} + r$, $r \in \{0, \dots, 2^l - 1 + s\}$, and
- (ii) $\sum_{i: \lambda(|O_i|) \leq l} |O_i| \leq r$.

For $k = 2$ this result is the very well known characterization of self-complementing permutation of graphs given by Ringel and Sachs.

Keywords: Self-complementing permutations, k -uniform hypergraphs

1 Introduction

Let V be a set of n elements. The set of all k -subsets of V is denoted by $\binom{V}{k}$. A k -uniform hypergraph H consists of a *vertex-set* $V(H)$ and an *edge-set* $E(H) \subseteq \binom{V(H)}{k}$. Two k -uniform hypergraphs G and H are *isomorphic*, if there is a bijection $\sigma : V(G) \rightarrow V(H)$ such that $e \in E(G)$ if and only if $\{\sigma(x) | x \in e\} \in E(H)$. The complement of a k -uniform hypergraph H is the hypergraph \overline{H} such that $V(\overline{H}) = V(H)$ and the edge set of which consists of all k -subsets of $V(H)$ not in $E(H)$ (in other words $E(\overline{H}) = \binom{V(H)}{k} - E$). A k -uniform hypergraph H is called *self-complementary* (*s-c* for short) if it is isomorphic with its complement \overline{H} . Isomorphism of a k -uniform self-complementary hypergraph onto its complement is called *self-complementing permutation* (or *s-c permutation*).

The 2-uniform self-complementary hypergraphs are exactly self-complementary graphs. This class of graphs has been independently discovered by Ringel and Sachs who proved the following.

Theorem 1 (Ringel (Rin63) and Sachs (Sac62)) *Let n be a positive integer. A permutation σ of $[1, n]$ is a self-complementing permutation of a self-complementary graph of order n if and only if all the orbits of σ have their cardinalities congruent to 0 (mod 4) except, possibly, one orbit of cardinality 1.*

Observe that by Theorem 1 an s-c graph of order n exists if and only if $n \equiv 0$ or $n \equiv 1 \pmod{4}$ or, equivalently, whenever $\binom{n}{2}$ is even. In (SW) we prove that a similar result is true for k -uniform hypergraphs.

Theorem 2 ((SW)) *Let k and n be positive integers, $k \leq n$. A k -uniform self-complementary hypergraph of order n exists if and only if $\binom{n}{k}$ is even.*

A simple criterion for evenness of $\binom{n}{k}$ has been given in (Gla99) (and then rediscovered in (KHRM58)).

Theorem 3 ((Gla99; KHRM58)) *Let k and n be positive integers, $k = \sum_{i=0}^{+\infty} c_i 2^i$ and $n = \sum_{i=0}^{+\infty} d_i 2^i$, where $c_i, d_i \in \{0, 1\}$ for every i . $\binom{n}{k}$ is even if and only if there is i_0 such that $c_{i_0} = 1$ and $d_{i_0} = 0$.*

Theorem 3 asserts that $\binom{n}{k}$ is even if and only if k has 1 in a certain binary place while n has 0 in the corresponding binary place. For example, $\binom{27}{13}$ is even since $13 = 1 \cdot 2^3 + 1 \cdot 2^2 + 0 \cdot 2^1 + 1 \cdot 2^0$ and $27 = 1 \cdot 2^4 + 1 \cdot 2^3 + 0 \cdot 2^2 + 1 \cdot 2^1 + 1 \cdot 2^0$ (so we have $c_2 = 1$ and $d_2 = 0$).

Except for Theorem 1 which is a characterization of the self-complementing permutations for graphs, there are already two published results characterizing the permutations of k -uniform s-c hypergraphs for $k > 2$. Namely, Kocay in (Koc92) (see also (Pal73)) and Szymański in (Szy05) have characterized the s-c permutations of s-c k -uniform hypergraphs for, respectively, $k = 3$ and $k = 4$. This work is a continuation of the work of (SW) and (Woj06). We generalize all the results mentioned above by giving a characterization of the s-c permutations of k -uniform hypergraphs for any integers k and n .

2 Result

Any positive integer n may be written in the form $n = 2^l c$, where c is an odd integer. Moreover, l and c are uniquely determined. We write then $\lambda(n) = l$. Note that in the binary expansion of n , $\lambda(n)$ is the index of the first 1-bit. For any set A we shall write $\lambda(A)$ in place of $\lambda(|A|)$, for short.

In the proof of our main result we shall need the following lemma proved in (Woj06).

Lemma 1 *Let k, m and n be positive integers, and let $\sigma : V \rightarrow V$ be a permutation of a set V , $|V| = n$, with orbits O_1, \dots, O_m . σ is a self-complementing permutation of a self-complementary k -uniform hypergraph, if and only if, for every $p \in \{1, \dots, k\}$ and for every decomposition*

$$k = k_1 + \dots + k_p$$

of k ($k_j > 0$ for $j = 1, \dots, p$), and for every subsequence of orbits

$$O_{i_1}, \dots, O_{i_p}$$

such that $k_j \leq |O_{i_j}|$ for $j = 1, \dots, p$, there is a subscript $j_0 \in \{1, \dots, p\}$ such that

$$\lambda(k_{j_0}) < \lambda(O_{i_{j_0}})$$

Given any integer $l \geq 0$. If the binary expansion of k is 1-bit in position l , then k can be written in the form $k = a_l 2^l + s_l$, where a_l is odd and $0 \leq s_l < 2^l$.

Theorem 4 *Let k and n be integers, $k \leq n$. A permutation σ of $[1, n]$ with orbits O_1, \dots, O_m is a self-complementing permutation of a k -uniform hypergraph of order n if and only if there is a nonnegative integer l such that $k = a_l 2^l + s_l$, where a_l is odd and $0 \leq s_l < 2^l$, and the following two conditions hold:*

$$(i) \quad n = b_l 2^{l+1} + r_l, \quad r_l \in \{0, \dots, 2^l - 1 + s_l\}, \text{ and}$$

$$(ii) \quad \sum_{i: \lambda(O_i) \leq l} |O_i| \leq r_l.$$

Proof:

Sufficiency. By contradiction. Let n, k, l, a_l, b_l, s_l and r_l be integers verifying the conditions of the theorem, let σ be a permutation of $[1, n]$ with orbits O_1, \dots, O_m verifying (ii), and let us suppose that σ is not a s-c permutation of any k -uniform s-c hypergraph of order n . Then, by Lemma 1, there is a decomposition of $k = k_1 + \dots + k_t$ and a subsequence of orbits O_{i_1}, \dots, O_{i_t} such that

$$0 < k_j \leq |O_{i_j}| \tag{1}$$

and

$$\lambda(k_j) \geq \lambda(O_{i_j}) \tag{2}$$

for $j = 1, \dots, t$.

Since a_l is odd, we have $k \equiv 2^l + s_l \pmod{2^{l+1}}$. By (2), $\sum_{j: \lambda(O_{i_j}) > l} k_j \equiv 0 \pmod{2^{l+1}}$. Therefore

$$k = \sum_{j=1}^t k_j = \sum_{j: \lambda(O_{i_j}) > l} k_j + \sum_{j: \lambda(O_{i_j}) \leq l} k_j \equiv \sum_{j: \lambda(O_{i_j}) \leq l} k_j \pmod{2^{l+1}}$$

Hence, and by (1), (i) and (ii) we have $\sum_{j: \lambda(O_{i_j}) \leq l} k_j \leq \sum_{j: \lambda(O_{i_j}) \leq l} |O_{i_j}| < 2^{l+1}$, and therefore

$$2^l + s_l = \sum_{j: \lambda(O_{i_j}) \leq l} k_j \leq \sum_{j: \lambda(O_{i_j}) \leq l} |O_{i_j}| \leq r_l < 2^l + s_l$$

a contradiction.

Necessity. Let $1 \leq k \leq n$ and let σ be a permutation of the set $[1, n]$ with orbits O_1, \dots, O_m . Let us suppose that for every integer l such that $k = a_l 2^l + s_l$, where a_l is odd positive integer, $0 \leq s_l < 2^l$, and $n = b_l 2^{l+1} + r_l$, $0 \leq r_l < 2^{l+1}$ we have either

$$r_l \in \{2^l + s_l, \dots, 2^{l+1} - 1\}$$

or

$$r_l \in \{0, \dots, 2^l - 1 + s_l\} \quad \text{and} \quad \sum_{i: \lambda(O_i) \leq l} |O_i| > r_l$$

We shall prove that σ is not a s-c permutation of any s-c k -uniform hypergraph of order n . For this purpose we shall give two claims.

Claim 1 For every nonnegative integer l such that $k = a_l 2^l + s_l$, where a_l is odd and $0 \leq s_l < 2^l$, we have

$$\sum_{i: \lambda(O_i) \leq l} |O_i| \geq 2^l + s_l$$

Proof of Claim 1. Let us write $\sum_{i: \lambda(O_i) \leq l} |O_i|$ and $\sum_{i: \lambda(O_i) > l} |O_i|$ in their binary forms:

$$\sum_{i: \lambda(O_i) \leq l} |O_i| = \sum_{j=0}^{\infty} e_j 2^j$$

$$\sum_{i: \lambda(O_i) > l} |O_i| = \sum_{j=0}^{\infty} f_j 2^j$$

where $e_j, f_j \in \{0, 1\}$ for every j . Observe that $f_j = 0$ for $j = 0, \dots, l$ and therefore

$$\sum_{j=0}^l e_j 2^j = r_l \tag{3}$$

We shall consider two cases.

Case 1. $r_l \in \{0, \dots, 2^l + s_l - 1\}$ and $\sum_{i: \lambda(O_i) \leq l} |O_i| > r_l$.

We have $n \geq 2^{l+1}$ (otherwise $r_l = n = \sum_{i: \lambda(O_i) \leq l} |O_i|$).

Since $\sum_{j=0}^{\infty} e_j 2^j > r_l$, and by (3), we obtain $\sum_{j=0}^{\infty} e_j 2^j \geq 2^{l+1} > 2^l + s_l$.

Case 2. $r_l \in \{2^l + s_l, \dots, 2^{l+1} - 1\}$.

We have $\sum_{i: \lambda(O_i) \leq l} |O_i| = \sum_{j=0}^{\infty} e_j 2^j \geq \sum_{j=0}^l e_j 2^j = r_l \geq 2^l + s_l$, and the claim is proved. \square

Claim 2 Let $\alpha_1, \dots, \alpha_q$ and $\lambda_1, \dots, \lambda_q$ be integers such that $0 < \alpha_i, 0 \leq \lambda_i \leq \lambda(\alpha_i)$ and $\lambda_i \leq l$ for $i = 1, \dots, q$ and $\sum_{i=1}^q \alpha_i \geq 2^l$. Then there are β_1, \dots, β_q such that for every $i = 1, \dots, q$

$$0 \leq \beta_i \leq \alpha_i \tag{4}$$

and

$$\text{either } \beta_i = 0 \text{ or } \lambda(\beta_i) \geq \lambda_i \tag{5}$$

and

$$\sum_{i=1}^q \beta_i = 2^l \tag{6}$$

Proof of Claim 2. The existence of β_1, \dots, β_q verifying (4)-(5) and $\sum_{i=1}^q \beta_i \leq 2^l$ is very easy. Indeed, it is immediate that $\beta_1 = 2^{\lambda_1}, \beta_2 = \dots, \beta_q = 0$ is a sequence with the desired properties.

So let us suppose that β_1, \dots, β_q is a sequence verifying (4)-(5) and $\sum_{i=1}^q \beta_i \leq 2^l$ such that $\sum_{i=1}^q \beta_i$ is maximal. If $\sum_{i=1}^q \beta_i = 2^l$ then the proof is complete. So let us suppose that $\sum_{i=1}^q \beta_i < 2^l$. Then there is $i_0 \in \{1, \dots, q\}$ such that $\beta_{i_0} < \alpha_{i_0}$. Observe that $\beta_{i_0} + 2^{\lambda_{i_0}} \leq \alpha_{i_0}$. The sequence $\bar{\beta}_1, \dots, \bar{\beta}_q$ defined by $\bar{\beta}_{i_0} = \beta_{i_0} + 2^{\lambda_{i_0}}$ and $\bar{\beta}_i = \beta_i$ for $i \neq i_0$ also verifies (4)-(5) and $\sum_{i=1}^q \bar{\beta}_i \leq 2^l$, which contradicts the maximality of the sum $\sum_{i=1}^q \beta_i$, and the claim is proved. \square

We shall use our claims to construct a decomposition of k in the form $k = k_1 + \dots + k_m$ such that

- (1) k_1, \dots, k_m are nonnegative integers,
- (2) $k_i \leq |O_i|$ for $i = 1, \dots, m$, and
- (3) $\lambda(k_i) \geq \lambda(O_i)$ whenever $k_i > 0$

By Lemma 1, this will imply that σ is not a s-c permutation of any k -uniform s-c hypergraph. Let us write k in its binary form:

$$k = 2^{l_t} + 2^{l_{t-1}} + \dots + 2^{l_1} + 2^{l_0}$$

where $l_0 < l_1 < \dots < l_t$.

By Claim 1, $\sum_{i:\lambda(O_i) \leq l_0} |O_i| \geq 2^{l_0}$. Hence, and by Claim 2, there are nonnegative integers $k_1^{(0)}, k_2^{(0)}, \dots, k_m^{(0)}$ such that $k_i^{(0)} = 0$ for i such that $\lambda(O_i) > l_0$ and

$$k_i^{(0)} \leq |O_i| \text{ for } i = 1, \dots, m$$

$$\lambda(k_i^{(0)}) \geq \lambda(O_i) \text{ whenever } k_i^{(0)} > 0$$

and

$$\sum_{i=1}^m k_i^{(0)} = 2^{l_0}$$

Note that, for $i = 1, \dots, m$, we have $\lambda(|O_i| - k_i^{(0)}) \geq \lambda(O_i)$.

Let us suppose that we have already constructed $k_1^{(j)}, \dots, k_m^{(j)}$, ($j \leq t$), such that $k_i^{(j)} = 0$ for i such that $\lambda(O_i) > l_j$ and

$$k_i^{(j)} \leq |O_i| \text{ for } i = 1, \dots, m$$

$$\lambda(k_i^{(j)}) \geq \lambda(O_i) \text{ whenever } k_i^{(j)} > 0$$

$$\sum_{i=0}^m k_i^{(j)} = 2^{l_j} + 2^{l_{j-1}} + \dots + 2^{l_0}$$

and

$$\lambda(|O_i| - k_i^{(j)}) \geq \lambda(O_i)$$

If $j = t$, then we have already found a desired decomposition of k . If $j < t$, then, by Claim 1, we have

$$\sum_{i:\lambda(O_i) \leq l_{j+1}} (|O_i| - k_i^{(j)}) \geq 2^{l_{j+1}}.$$

$\lambda(|O_i| - k_i^{(j)}) \geq \lambda(O_i)$ for every $i \in \{1, \dots, m\}$ such that $|O_i| - k_i^{(j)} > 0$. Hence, and by Claim 2, there are β_1, \dots, β_m such that $\beta_i = 0$ for i such that $\lambda(O_i) > l_{j+1}$ and

$$0 \leq \beta_i \leq |O_i| - k_i^{(j)} \text{ for } i = 1, \dots, m$$

$$\lambda(O_i) \leq \lambda(\beta_i) \text{ for } i = 1, \dots, m \text{ whenever } \beta_i \neq 0$$

$$\sum_{i=1}^m \beta_i = 2^{l_{j+1}}$$

Thus we may define for every $i = 1, \dots, m$

$$k_i^{(j+1)} = k_i^{(j)} + \beta_i$$

to obtain the sequence $(k_1^{(j+1)}, \dots, k_m^{(j+1)})$ verifying for every $i \in \{1, \dots, m\}$

$$k_i^{(j+1)} = 0 \text{ for } i \text{ such that } \lambda(O_i) > l_{j+1}$$

$$k_i^{(j+1)} \leq |O_i|$$

$$\lambda(k_i^{(j+1)}) \geq \lambda(O_i) \text{ whenever } k_i^{(j+1)} > 0$$

and

$$\sum_{i=1}^m k_i^{(j+1)} = 2^{l_{j+1}} + 2^{l_j} + \dots + 2^{l_0}$$

It is clear that $k = \sum_{i=1}^m k_i^{(t)}$ and the proof of Theorem 4 is complete. \square

Theorem 4 implies very easily the following theorem first proved by Kocay.

Corollary 1 (Kocay (Koc92)) *σ is a self-complementing permutation of a self-complementary 3-uniform hypergraph if and only if either all the orbits of σ have even cardinalities, or else, it has 1 or 2 fixed points and the all remaining orbits of σ have their cardinalities being multiples of 4.*

For $k = 2^l$ Theorem 4 may be written as follows.

Corollary 2 *Let l and n be nonnegative integers, $2^l < n$, and let $0 \leq r < 2^{l+1}$ be such that $n \equiv r \pmod{2^{l+1}}$. A permutation σ of $[1, n]$ with orbits O_1, \dots, O_m is a self-complementing permutation of a 2^l -uniform self-complementary hypergraph if and only if*

$$(i) \ r \in \{0, \dots, 2^l - 1\} \text{ and}$$

$$(ii) \ \sum_{i: \lambda(O_i) \leq l} |O_i| \leq r.$$

Theorem 2 for $l = 1$ (i.e. for graphs) is exactly Theorem 1, and for $l = 2$ the following theorem proved by Szymański in (Szy05).

Corollary 3 *A permutation σ is self-complementing permutation of a 4-uniform hypergraph of order n if and only if $n \equiv r \pmod{8}$ with $r = 0, 1, 2$ or 3 , and the sum of the cardinalities of orbits which are not multiples of 8 is at most 3.*

Acknowledgements

The research was partially supported by AGH local grant No 11 420 04.

References

- [Gla99] J.W.L. Glaisher. On the residue of a binomial coefficient with respect to a prime modulus. *Quarterly Journal of Mathematics*, 30:150–156, 1899.
- [KHRM58] S.H. Kimball, T.R. Hatcher, J.A. Riley, and L. Moser. Solution to problem e1288: Odd binomial coefficients. *Amer. Math. Monthly*, 65:368–369, 1958.
- [Koc92] W. Kocay. Reconstructing graphs as subsumed graphs of hypergraphs, and some self-complementary triple systems. *Graphs and Combinatorics*, 8:259–276, 1992.
- [Pal73] E.M. Palmer. On the number of n -plexes. *Discrete Math.*, 6:377–390, 1973.
- [Rin63] G. Ringel. Selbstkomplementare graphen. *Arch. Math.*, 14:354–358, 1963.
- [Sac62] H. Sachs. Über selbstkomplementare graphen. *Publ. Math. Debrecen*, 9:270–288, 1962.
- [SW] A. Szymański and A.P. Wojda. A note on k -uniform self-complementary hypergraphs of given order. *submitted*.
- [Szy05] A. Szymański. A note on self-complementary 4-uniform hypergraphs. *Opuscula Mathematica*, 25/2:319–323, 2005.
- [Woj06] A.P. Wojda. Self-complementary hypergraphs. *Discussiones Mathematicae – Graph Theory*, 26:217–224, 2006.

