

On the length of shortest 2-collapsing words

Alessandra Cherubini, Andrzej Kisielewicz, Brunetto Piochi

► **To cite this version:**

Alessandra Cherubini, Andrzej Kisielewicz, Brunetto Piochi. On the length of shortest 2-collapsing words. *Discrete Mathematics and Theoretical Computer Science, DMTCS*, 2009, 11 (1), pp.33–44. <hal-00988185>

HAL Id: hal-00988185

<https://hal.inria.fr/hal-00988185>

Submitted on 7 May 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

On the Length of Shortest 2-Collapsing Words

Alessandra Cherubini¹ and Andrzej Kisielewicz² and Brunetto Piochi³ †

¹ Politecnico di Milano, Dipart. di Matematica “F. Brioschi”, Piazza L. da Vinci 32 - 20133 Milano, Italy
alessandra.cherubini@polimi.it

² University of Wrocław, Institute of Mathematics, pl. Grunwaldzki 2 - 50-384 Wrocław, Poland
kisiel@math.uni.wroc.pl

³ Università di Firenze, Dipart. di Matematica “U. Dini”, viale Morgagni 67/a - 50134 Firenze, Italy;
piochi@math.unifi.it

received May 12, 2008, revised October 24, 2008, accepted December 10, 2008.

Given a word w over a finite alphabet Σ and a finite deterministic automaton $\mathcal{A} = \langle Q, \Sigma, \delta \rangle$, the inequality $|\delta(Q, w)| \leq |Q| - k$ means that under the natural action of the word w the image of the state set Q is reduced by at least k states. The word w is k -collapsing (k -synchronizing) if this inequality holds for any deterministic finite automaton (with $k + 1$ states) that satisfies such an inequality for at least one word. We prove that for each alphabet Σ there is a 2-collapsing word whose length is $\frac{|\Sigma|^3 + 6|\Sigma|^2 + 5|\Sigma|}{2}$. Then we produce shorter 2-collapsing and 2-synchronizing words over alphabets of 4 and 5 letters.

Keywords: property Δ_k , finite automata, synchronizing finite automata

1 Introduction and preliminaries.

In this paper by an automaton $\mathcal{A} = \langle Q, \Sigma, \delta \rangle$ we mean a finite deterministic automaton with state set Q , input alphabet Σ , and transition function $\delta : Q \times \Sigma \rightarrow Q$. The action of Σ on Q given by δ will be denoted by concatenation: $qa = \delta(q, a)$. This action extends naturally, by composition, to the action of the words of Σ^* on Q . Given a word $w \in \Sigma^*$, we will be interested in the difference of the cardinalities $|Q| - |Qw|$. This difference is called the *deficiency* of the word w with respect to \mathcal{A} and will be denoted $df_{\mathcal{A}}(w)$.

For a fixed $k \geq 1$, a word $w \in \Sigma^*$ is called k -compressing for \mathcal{A} if $df_{\mathcal{A}}(w) \geq k$. An automaton \mathcal{A} is k -compressible if there exists a k -compressing word for \mathcal{A} . A word $w \in \Sigma^*$ is k -collapsing (over Σ) if it is k -compressing for every k -compressible automaton with the input alphabet Σ . A word v is called

†This research was partially supported by the ESF programme Automatha

k -synchronizing if it is k -compressing for all k -compressible automata with $k + 1$ states. Of course a k -collapsing word is also k -synchronizing.

The concept of a k -collapsing word is a natural automata-theoretic interpretation of the notion of words with the property Δ_k introduced for algebraic motivations in [12]. Actually k -collapsing words can be seen as universal testers for checking whether an automaton is k -compressible. They are a black-box versions of words considered in the so called *generalized Černý conjecture* that extends the celebrated *Černý conjecture* [5]. Recall that the former conjecture deals with the length of the shortest k -compressing word for a given k -compressible automaton. In [12] it was proved that k -collapsing words always exist, for each Σ and each $k \geq 1$, by means of a recursive construction which gives a word whose length is doubly exponential as function of k . Then better lower and upper bounds for the length $c(k, t)$ of the shortest k -collapsing words on an alphabet of cardinality t were given in [9] and the lower bound for the case $k = 2$ was slightly improved in [10]. So far for the case $k = 2$ the best known bounds are the following: $2t^2 \leq c(2, t) \leq t^3 + 3t^2 + 2t$.

In [1, 2], two different algorithms for deciding whether a word is 2-collapsing are given and another more combinatorial approach to the same problem is presented in [8]. Here we get a better upper bound for $c(2, t)$, namely $c(2, t) \leq \frac{t^3 + 6t^2 + 5t}{2}$, by means of the latter approach. Moreover we prove that this bound can be improved at least for small values of t : we build, again by using this approach, shorter 2-collapsing words on alphabets of cardinality 4 and 5. The reader is referred to [9] and to [4] for some further references and connections to some topics in Theoretical Computer Science, Language Theory and Combinatorics. Here for sake of completeness we shortly recall notation and main properties.

We view an automaton $\mathcal{A} = \langle Q, \Sigma, \delta \rangle$ as a set of transformations on Q labelled by letters of Σ rather than as a standard triple. By *transformations of \mathcal{A}* we mean those transformations on Q that are induced via δ by letters of Σ . In order to define an automaton it is enough just to assign to every letter of Σ a transformation on Q . Now, for $a \in \Sigma$, $\text{df}_{\mathcal{A}}(a) = 0$ if and only if the corresponding transformation is a permutation on Q . If $\text{df}_{\mathcal{A}}(a) = 1$, then there is a state $z \in Q$ which does not belong to the image Qa and two different states $x, y \in Q$ satisfying $xa = ya$; in such a case the corresponding transformation will be referred to as a *transformation of type $\{x, y\} \setminus z$* (x, y have the same image under the transformation, z is missed). The semigroup generated by the transformations of \mathcal{A} consists precisely of the transformations corresponding to words in Σ^* . It contains a group generated by those transformations that are permutations; this group is called the *group of permutations of \mathcal{A}* .

It is well known ([9]) that each 2-collapsing word over a fixed alphabet Σ is 2-full, i.e. it contains any word of length 2 on Σ among its factors, hence to characterize 2-collapsing words it is enough to consider 2-full words compressing each 2-compressible automata such that no word of length 2 is 2-compressing for them (*proper 2-compressible automata*, for short). Proper 2-compressible automata must have both non-permutation and permutation transformations and they are characterized by the following:

Proposition 1 ([8]) *An automaton \mathcal{A} is proper 2-compressible if and only if \mathcal{A} fulfils one of the following conditions:*

1. *there are two states x, y such that all non-permutation transformations are of the same type $\{x, y\} \setminus x$, and the group of permutations fixes neither the element x nor the set $\{x, y\}$;*
2. *there is a state x such that each non-permutation transformation is of type $\{x, z\} \setminus x$ for some state z , at least two different types occur, and the group of permutations does not fix x ;*

3. there are two states x, y such that each non-permutation transformation is of type $\{x, y\} \setminus x$ or $\{x, y\} \setminus y$, both types occur, and the group of permutations does not fix the set $\{x, y\}$.

Automata of the first two types are called MONO-automata, of the latter one are called STEREO-automata.

In [8] it is proved that a word $w \in \Sigma^*$ is 2-collapsing if and only if certain systems of conditions on permutations have only trivial solutions. We recall some notations from [6]. In the sequel, as no confusion can raise, we view the set Q of the states of \mathcal{A} as a set of natural numbers: $Q = \{1, 2, \dots, n\}$, so that permutations on Q can be viewed as elements of the symmetric group Sym_n . The two states x, y occurring in Proposition 1 will be then denoted by 1 and 2. By a *role assignment* we mean an arbitrary partition of the alphabet Σ with a distinguished nonempty block $\Pi \subset \Sigma$. Roughly speaking letters in Π play the role of permutation letters of Σ and the remaining letters play the role of non-permutation letters. Let $\Upsilon = \Sigma \setminus \Pi$. Since non-permutation letters in proper 2-compressible automata can be of different types, a role assignment induces on Υ a further partition in non empty blocks. We denote this partition either as $\{D_2, \dots, D_h\}$, $h \geq 2$, where letters in D_i will play the role of the non-permutation letters of the form $\{1, i\} \setminus 1$ or as $\{E_1, E_2\}$ where letters in E_1, E_2 will play the role of non-permutation letters of the form $\{1, 2\} \setminus 1$ and $\{1, 2\} \setminus 2$ respectively. In the first case the partition of Σ will be called a *MONO-role assignment* (DB-partition in [8]) and in the latter a *STEREO-role assignment* (3DB-partition in [8]).

Let (Π, Υ) be an arbitrary role assignment, then each word $w \in \Sigma^+$ can be uniquely represented in the following form:

$$w = u_0 p_1 u_1 \dots u_{m-1} p_m u_m \quad (1)$$

where $p_1, \dots, p_m \in \Pi^+$, $u_1, \dots, u_{m-1} \in \Upsilon^+$, $u_0, u_m \in \Upsilon^*$ and m is a positive integer. We say that a factor p_i of the decomposition (1) is an *inner factor* if both u_{i-1} and u_i are non-empty. Then for a MONO-role assignment (Π, Υ) for each $a \in \Upsilon$ we denote by S_a the set of the inner factors p_i of w such that a is the first letter in u_i . Otherwise if (Π, Υ) is a STEREO-role assignment $(\Pi, \{E_1, E_2\})$ we denote by S_k , $k = 1, 2$, the sets of the inner factors p_i of w such that the last letter of u_{i-1} belongs to E_k .

We assign to the word $w \in \Sigma^+$, for each role assignment and for each inner factor p_i , a permutation condition of the form:

- $1p_i \in \{1, j\}$ if the role assignment is MONO and $p_i \in S_a$ with $a \in D_j$,
- $kp_i \in \{1, 2\}$, $k = 1, 2$, if the role assignment is STEREO and $p_i \in S_k$.

Then each role assignment associates to w a system of permutation conditions formed by all the permutation conditions corresponding to the inner factors of w . In case of a MONO-role assignment (Π, Υ) such system will be denoted $\Gamma_w(\Pi, \Upsilon)$ otherwise it will be denoted $\Gamma'_w(\Pi, \Upsilon)$.

Since letters of Π are regarded as permutations acting on a finite set of positive integers $\{1, 2, \dots, n\}$ (representing Q), conditions of the form $jv \in A$ with j positive integer, $v \in \Pi^+$, $A \subseteq \{1, 2, \dots, n\}$ mean that the image of j under the product v of permutations belongs to the set A . We say that a system of permutation conditions *has a solution* if there exists an assignment of permutations on a finite set $\{1, 2, \dots, n\}$ to letters in Π such that all the conditions in the system are satisfied. Obviously the systems associated to w by each role assignment have always the solution where all permutations are the identity. A solution of the system $\Gamma_w(\Pi, \Upsilon)$, associated to w by a MONO-role assignment, is called *trivial* if all permutations fix 1. Also, in the special case when Υ consists of the unique block D_2 (and as a consequence, all j 's on the right hand side of the conditions are equal to 2), a solution with all permutations fixing the set $\{1, 2\}$ is considered trivial. The remaining solutions are *nontrivial*. Similarly a solution of the system

$\Gamma'_w(\Pi, \Upsilon)$, associated to w by a STEREO-role assignment (Π, Υ) is called trivial if all the permutations fix $\{1, 2\}$.

In [8] 2-collapsing words were characterized in terms of permutation conditions in the following way.

Theorem 2 ([8]) *A word $w \in \Sigma^+$ is 2-collapsing if and only if it is 2-full and the following conditions hold:*

1. *For each MONO-role assignment (Π, Υ) of Σ all the solutions of the system $\Gamma_w(\Pi, \Upsilon)$ are trivial.*
2. *For each STEREO-role assignment (Π, Υ) of Σ all the solutions of the system $\Gamma'_w(\Pi, \Upsilon)$ are trivial.*

By the above Theorem 2 one can readily derive the following characterization of 2-synchronizing words.

Corollary 3 *A word $w \in \Sigma^+$ is 2-synchronizing if and only if it is 2-full and for each role assignment (Π, Υ) of Σ all the solutions of the systems $\Gamma_w(\Pi, \Upsilon)$ (in case of MONO-role assignments) and $\Gamma'_w(\Pi, \Upsilon)$ (in case of STEREO-role assignments) in the semigroup \mathbf{T}_3 of all transformations on 3 letters are trivial.*

2 Main result.

We introduce some useful notations.

Definition 1 . *A permutation p on $\{1, 2, \dots, n\}$ is special with respect to the ordered pair (i, j) , $i, j \in \{1, 2, \dots, n\}$, whether it fixes either $\{i\}$ or $\{i, j\}$. Two permutations p, z are special of the same type with respect to the ordered pair (i, j) if both p and z fix $\{i\}$ or both of them fix $\{i, j\}$.*

In the sequel by $(i)\dots, (ij)\dots, (ijh\dots)\dots$ we will denote respectively a permutation fixing i , containing a cycle (i, j) , containing a cycle $(ijh\dots)$ of arbitrary length.

Note that the property of being special of the same type with respect to (i, j) is not transitive; e.g. both the permutations $p = (1)(2\dots n)$ and $z = (12)(3\dots n)$ are special of the same type as $y = (1)(2)(3\dots n)$ with respect to $(1, 2)$, but they are not of the same type. Note however that two special permutations p and z can be of the same type of y with respect to (i, j) without being of the same type only if one of them fixes $\{i\}$ and does not fix j , the other has a cycle (ij) while y fixes both $\{i\}$ and $\{j\}$. Trivial solutions of systems introduced before Theorem 2 are special of the same type with respect to $(1, 2)$, so we are interested in sets of equations whose occurrence in a system guarantees that solutions are special of the same type with respect to $(1, 2)$.

Lemma 4 *Let p, z be two permutations on $\{1, 2, \dots, n\}$ and let $i, j \in \{1, 2, \dots, n\}$. The permutations p, z fulfil the conditions:*

1. $ip \in \{i, j\}$
2. $ip^2 \in \{i, j\}$
3. $iz \in \{i, j\}$
4. $iz^2 \in \{i, j\}$
5. $ipzp \in \{i, j\}$ (or $izpz \in \{i, j\}$)

if and only if they are special of the same type with respect to (i, j) .

Proof: The "if part" is trivial. Conversely from condition 1. we immediately get that either p fixes i or it is of the form $p = (ij\dots)$ and condition 2. gives that in such case $p = (ij)\dots$. Similarly conditions 3. and 4. give that z either fixes i or has the form $z = (ij)\dots$. Now if $z = (ij)\dots$ and p fixes i then p fixes also j otherwise condition 5. is not fulfilled. If $p = (ij)\dots$ then $jpz = ipz \in \{i, j\}$ by condition 5. whence z fixes $\{i, j\}$. \square

Similarly, the following can be proved:

Lemma 5 *Let p, z be two permutations on $\{1, 2, \dots, n\}$ and let $i, j \in \{1, 2, \dots, n\}$. The permutations p, z fulfil the conditions:*

1. $ip \in \{i, j\}$
2. $ip^2 \in \{i, j\}$
3. $iz^2 \in \{i, j\}$
4. $ipz \in \{i, j\}$
5. $izp \in \{i, j\}$

if and only if they are special of the same type with respect to (i, j) .

Theorem 6 *The minimal 2-collapsing word on t letters has length less than or equal to*

$$\frac{t^3 + 6t^2 + 5t}{2}.$$

Proof: To show our bound we want to exhibit a 2-collapsing word, similar to the one introduced in [9]. Consider an alphabet $\Sigma = \{a_1, a_2, \dots, a_t\}$ on t letters in the lexicographic order and let $u = a_1a_2\dots a_t$ and

$$w = u \left(\prod_{q \in X} qu \right)$$

where

$$X = \{x, x^2, xyx : x, y \in \Sigma, y > x\}.$$

We want to prove that the word w is 2-collapsing. Let us consider different role assignments (Π, Υ) of Σ .

Assume that (Π, Υ) is a STEREO-role assignment with $\Upsilon = \{E_1, E_2\}$; let h, k be respectively the least and the greatest index of an element in Υ and let a_j be an element in Υ belonging to a different block than a_k (eventually $h = j$). Each $a_i, i > k$ (if any) fixes $\{1, 2\}$, since the factors $a_k a_i a_k, a_j a_i a_j$ occur in w whence both the equations $1a_i \in \{1, 2\}$ and $2a_i \in \{1, 2\}$ occur in $\Gamma'_w(\Pi, \Upsilon)$. More the product $a_1 a_2 \dots a_{h-1}$ fixes $\{1, 2\}$. In fact $a_j(a_1 a_2 \dots a_{h-1})a_h$ and $a_k(a_1 a_2 \dots a_{h-1})a_h$ are factors of w and all a_1, \dots, a_{h-1} are in Π hence both the equations $1(a_1 a_2 \dots a_{h-1}) \in \{1, 2\}$ and $2(a_1 a_2 \dots a_{h-1}) \in \{1, 2\}$ occur in $\Gamma'_w(\Pi, \Upsilon)$. This yields that also any $a_i, i < h$ (if any) fixes $\{1, 2\}$, since also $a_i a_j a_i(a_1 a_2 \dots a_{h-1})a_h$ and $a_i a_k a_i(a_1 a_2 \dots a_{h-1})a_h$ are factors of w whence both the equations $1a_i(a_1 a_2 \dots a_{h-1}) \in \{1, 2\}$ and $2a_i(a_1 a_2 \dots a_{h-1}) \in \{1, 2\}$ occur in $\Gamma'_w(\Pi, \Upsilon)$ with $a_1 a_2 \dots a_{h-1}$ fixing $\{1, 2\}$. Similarly, at last, each

$a_i : h < i < k$ (if any) fixes $\{1, 2\}$. In fact either a_h and a_k are in different blocks or there is an $a_j \in \Upsilon$ belonging to a different block with respect to a_h and a_k , so for each i , $h < i < k$ there are two indices s, r , $h \leq s < i < r \leq k$ such that a_s and a_r are in different blocks of Υ , hence $a_i a_r a_i (a_1 a_2 \dots a_{h-1}) a_h$ and $a_s a_i a_s$ are factors of w .

Then let (Π, Υ) be a MONO-role assignment with $\Upsilon = \{D_2, D_3, \dots, D_g\}$ with $g > 2$. Let k be the greatest index such that $a_k \in \Upsilon$ then $a_k \in D_{k'}$ for some k' . Let $D_{j'}$ be a block of Υ different from $D_{k'}$ and let $a_j \in D_{j'}$. We will prove that every $a_i \in \Pi$ fixes $\{1\}$. In fact if $i > k$ then $a_k a_i a_k, a_j a_i a_j$ are factors of w hence both the equations $1a_i \in \{1, k'\}, 1a_i \in \{1, j'\}$ occur in the system $\Gamma_w(\Pi, \Upsilon)$, whence a_i fixes $\{1\}$ for all $a_i \in \Pi$ with $i > k$, if any. Now let $i < k$. If $j < i < k$ then $a_j a_i a_j, a_k a_{k+1} \dots a_t a_i a_k a_i$ are factors of w , if $i < j$ then $a_k a_{k+1} \dots a_t a_i a_j a_i, a_k a_{k+1} \dots a_t a_i a_k a_i$ are factors of w and all elements a_{k+1}, \dots, a_t are in Π , hence they fix $\{1\}$. Then in the system $\Gamma_w(\Pi, \Upsilon)$ there are either the equations $1a_i \in \{1, j'\}$ and $1a_{k+1} \dots a_t a_i \in \{1, k'\}$ or $1a_{k+1} \dots a_t a_i \in \{1, j'\}$ and $1a_{k+1} \dots a_t a_i \in \{1, k'\}$ from which in both cases we get that $1a_i \in \{1, k'\}$ and $1a_i \in \{1, j'\}$, thus all $a_i \in \Pi$ fix $\{1\}$.

At last suppose that Υ consists of only one block, i.e. $g = 2$, and let h, k respectively be the least and the greatest index of an element in Υ . Denote $v_1 = a_1 \dots a_{h-1}$ and $v_2 = a_{k+1} \dots a_t$, then $a_h v_1 a_h, a_k v_2 a_k$ are factors of w whence $1v_1, 1v_2 \in \{1, 2\}$ are equations occurring in $\Gamma_w(\Pi, \Upsilon)$.

In this case, suppose first that $1v_2 = 1$. Since for all $a_i \in \Sigma$, $a_k v_2 a_i v_1 a_h$ and $a_k v_2 a_i^2 v_1 a_h$ are factors of w , equations in $\Gamma_w(\Pi, \Upsilon)$ give $1a_i v_1 \in \{1, 2\}$ and $1a_i^2 v_1 \in \{1, 2\}$. Moreover since w contains $a_k a_i a_k$ or $a_k v_2 a_i a_k a_i$ we have $1a_i \in \{1, 2\}$ for every $a_i \in \Sigma$. If all the letters a_i fix 1, then we are done. Otherwise suppose there is a letter a_i such that $1a_i = 2$. Then we get $2v_1 \in \{1, 2\}$ and since $1v_1 \in \{1, 2\}$ we obtain that v_1 fixes $\{1, 2\}$. Thus from $1a_i^2 v_1 \in \{1, 2\}$ we get $2a_i = 1$ and $a_i = (12) \dots$. To prove that in this case all permutations are of the same type, note that whenever a factor $a_k v_2 a_i a_j a_i v_1 a_h$ appears in w we have $1a_i a_j a_i \in \{1, 2\}$ and then apply Lemma 4.

Suppose now $1v_2 = 2$. Then $2a_i v_1, 2a_i^2 v_1 \in \{1, 2\}$ for every $a_i \in \Pi$. Let us prove that v_1 must fix $\{1, 2\}$. This is trivial if $h = 1$ and v_1 is empty, else let $\{1, 2\} = \{1v_1, zv_1\}$ for some z and prove that $z = 2$. For all $i : 1 \leq i < h$ $a_k v_2 a_i a_h a_i$ and $a_i a_h a_i v_1 a_h$ are factors of w , hence $1v_2 a_i \in \{1, 2\}$ and $1a_i v_1 \in \{1, 2\}$ are equations in $\Gamma_w(\Pi, \Upsilon)$, so that $2a_i \in \{1, 2\}$ and $a_i v_1$ fixes $\{1, 2\}$. If for some $i : 1 \leq i < h$: $2a_i = 2a_i^2$ then $a_i = (2) \dots$ and $2v_1 \in \{1, 2\}$ whence $z = 2$; else $\{2a_i, 2a_i^2\} = \{1, z\}$ which immediately yields $a_i = (21z \dots) \dots$ since $2 \neq 2a_i \in \{1, 2\}$. If there exist two such elements a_i, a_j , $1 \leq i < j < h$ then the factor $a_k v_2 a_i a_j a_i v_1 a_h$ occurs in w so that $1v_2 a_i a_j a_i v_1 = (2a_i a_j) a_i v_1 \in \{1, 2\}$ is an equation in $\Gamma_w(\Pi, \Upsilon)$ whence $2a_i a_j = z \in \{1, 2\}$ and $z = 2$; but if only one element a_i of this type exists then $z = 1a_i = 1v_1 \in \{1, 2\}$. Thus v_1 fixes $\{1, 2\}$, which immediately yields $2a_i, 1a_i \in \{1, 2\}$ for every $a_i \in \Pi$, $i < k$ by the occurrence of the factor $a_k v_2 a_i a_k a_i v_1 a_h$, and for every $a_i \in \Pi$, $i > k$ by the occurrence of the factors $a_k a_i a_k$ and $a_k v_2 a_i v_1 a_h$; hence all elements in Π fix $\{1, 2\}$ so that they are special of the same type.

Thus w is 2-collapsing because the systems associated to w for each role assignment (Π, Υ) have only trivial solutions and the length of the word w is:

$$t + t(t+1) + t(t+2) + \frac{t(t-1)}{2}(t+3) = \frac{t^3 + 6t^2 + 5t}{2}.$$

□

3 Short 2-collapsing words.

The bound found in Theorem 6 is lower than the one given in [9], but still for small values of n it reveals to be not efficient: indeed for $n = 3, 4, 5$ it respectively gives $|w| \leq 48, 90, 150$. It was proved in [3] that the shortest 2-collapsing words on 3 letters have length 21 and the shortest 2-synchronizing words on 3 letters have length 20, while a 2-collapsing word on 4 letters is known whose length equals 58 ([4]). But we can use our lemmas to get shorter 2-collapsing and 2-synchronizing words on 4 and 5 letters.

We will make a systematic use of the following

Lemma 7 *Let w be a word on an alphabet Σ with $|\Sigma| \geq 2$, let $\Sigma' \subseteq \Sigma$ and $w' \in \Sigma'^+$ be a factor of w , which is a 2-collapsing word on Σ' . For a role assignment (Π, Υ) of Σ let $\Pi' = \Pi \cap \Sigma'$ and $\Upsilon' = \Upsilon \cap \Sigma'$ and for each block $E_i (D_i)$ in Υ let $E'_i = E_i \cap \Sigma'$ ($D'_i = D_i \cap \Sigma'$). Then for each role assignment (Π, Υ) such that $\emptyset \neq \Pi'$ and $\emptyset \neq \Upsilon'$ the following properties hold:*

- *If (Π, Υ) is a MONO-role assignment with $\Upsilon = D_2$, then in each solution of $\Gamma_w(\Pi, \Upsilon)$ the elements in Π' are special of the same type with respect to $(1, 2)$.*
- *If (Π, Υ) is a MONO-role assignment with $\Upsilon = \{D_2, D_3, \dots, D_g\}$, $g > 2$, and there are at least two non empty blocks D'_{i_1}, D'_{i_2} , then in each solution of $\Gamma_w(\Pi, \Upsilon)$ all the elements in Π' fix $\{1\}$.*
- *If (Π, Υ) is a STEREO-role assignment and both E'_1, E'_2 are nonempty, then in each solution of $\Gamma'_w(\Pi, \Upsilon)$ all the elements in Π' fix the set $\{1, 2\}$.*

In all the above cases, if $\Pi' = \Pi$ then the system associated to w for (Π, Υ) has only trivial solutions.

Proof: If (Π, Υ) is a MONO-role assignment, then (Π', Υ') is a MONO-role assignment of Σ' and the equations of $\Gamma_{w'}(\Pi', \Upsilon')$ occur in $\Gamma_w(\Pi, \Upsilon)$, hence all solutions of the latter system are also solutions of the former. Hence if Υ and consequently Υ' has only one block, then all elements in Π' are special permutations relative to $(1, 2)$ which all fix $\{1\}$ or all fix $\{1, 2\}$, because w' is 2-collapsing. Similarly if Υ' is formed by at least two blocks then all elements in Π' fix $\{1\}$. If (Π, Υ) is a STEREO-role assignment of Σ , and Υ' has two non trivial blocks, then (Π', Υ') is a STEREO-role assignment of Σ' hence the equations of $\Gamma'_{w'}(\Pi', \Upsilon')$ occur in $\Gamma'_w(\Pi, \Upsilon)$ and so all elements in Π' are special permutations relative to $(1, 2)$ which all fix $\{1, 2\}$. The last statement is trivial. \square

It is important to remark, using the same notation and the same arguments of the above Lemma 7, that if (Π, Υ) is a MONO-role assignment with more than one block different from Π , while Υ' is formed by a unique block D'_i , then in each solution of $\Gamma_w(\Pi, \Upsilon)$ all the elements in Π' are special of the same type with respect to $\{1, i\}$. Similarly if (Π, Υ) is a STEREO-role assignment of Σ and $E'_1 = \emptyset$ then in each solution of $\Gamma'_w(\Pi, \Upsilon)$ all the elements in Π' are special of the same type with respect to $(2, 1)$ while if $E'_2 = \emptyset$ then all the elements in Π' are special of the same type with respect to $(1, 2)$.

Proposition 8

$$c(2, 4) \leq 56$$

Proof: Let $\Sigma = \{a, b, c, d\}$ and let

$$\begin{aligned} r &= cabac^2bacbcacb^2a^2c, \\ s &= dadbda^2b^2dabd^2ab, \\ u &= adacdbcadac^2d^2cbdca. \end{aligned}$$

Consider the word $w = r(bab)su$. Of course $|w| = 56$; we want to prove that w is 2-collapsing. In order to exhibit w we made use of two of the minimal 2-collapsing words on 3 letters. Namely $u_1 = r(bab)$ is the 55-th word in the list in [3] on the alphabet $\{c, a, b\}$, while $u_2 = (bab)s(ad)$ is the 11-th one in that list on the alphabet $\{b, a, d\}$. Both the words u_1 and u_2 are 2-collapsing, as shown in [3]. Put $\Sigma_1 = \{a, b, c\}$ and $\Sigma_2 = \{a, b, d\}$.

Let us consider different role assignments (Π, Υ) of Σ and put $\Pi_i = \Pi \cap \Sigma_i$, $\Upsilon_i = \Upsilon \cap \Sigma_i$, for $i = 1, 2$.

Let $|\Pi| = 1$ and $\Pi \subset \{a, b\}$ then $\Pi_1 = \Pi_2 = \Pi$. Moreover if (Π, Υ) induces on Υ a partition in more than one block then there is at least an $i = 1, 2$ such that Υ_i contains two non empty blocks, so the system associated to w for (Π, Υ) has only trivial solutions by Lemma 7. Then let $\Pi = \{c\}$. If (Π, Υ) is a MONO-role assignment and Υ consists of only one block, then all the solutions of $\Gamma_w(\Pi, \Upsilon)$ are trivial by Lemma 7. If Υ consists of more than one block then one has to consider the factors acb, bca of w if the letters a and b belong to different blocks, else consider the factors acd and dca . In both cases, if (Π, Υ) is a MONO-role assignment then the equations $1c \in \{1, 2\}, 1c \in \{1, 3\}$ occur in $\Gamma_w(\Pi, \Upsilon)$ hence c fixes $\{1\}$. If (Π, Υ) is a STEREO-role assignment then the equations $1c \in \{1, 2\}, 2c \in \{1, 2\}$ occur in $\Gamma'_w(\Pi, \Upsilon)$, hence c fixes the set $\{1, 2\}$. The same argument applies when $\Pi = \{d\}$, considering the factors adb, bda or bdc, cdb .

Let $|\Pi| = 2$. Assume $\Upsilon = \{a, b\}$. Then for all $i = 1, 2$, $\emptyset \neq \Pi_i \neq \Pi$ and $\Upsilon = \Upsilon_i$. Then if Υ is formed by two blocks then all the solutions of the system associated to w for (Π, Υ) are trivial: by Lemma 7 if (Π, Υ) is a MONO-role assignment then both c, d fix $\{1\}$, if (Π, Υ) is a STEREO-role assignment then both c, d fix $\{1, 2\}$. Otherwise if Υ is formed by a unique block we only get that c and d are special permutations relative to $(1, 2)$. But since w contains the factors $acdb, bdca$, the equations $1cd \in \{1, 2\}, 1dc \in \{1, 2\}$ occur in $\Gamma_w(\Pi, \{a, b\})$ hence c, d are special permutations of the same type with respect to $(1, 2)$, i.e. the system $\Gamma_w(\Pi, \{a, b\})$ has only trivial solutions. Then let $\Upsilon \neq \{a, b\}$, whence $\Pi = \Pi_i$ for some $i = 1, 2$. Hence if Υ is formed by a unique block then $\Gamma_w(\Pi, \Upsilon)$ has only trivial solutions by Lemma 7. Otherwise we only get that the solutions of the systems associated to w for (Π, Υ) are either special permutations relative to $(1, 2)$ or fix $\{2\}$. Denote by β_1, β_2 the elements in Π and by α_1, α_2 the elements in Υ ; for every choice of (Π, Υ) where $\Upsilon \neq \{a, b\}$ is formed by two blocks, there is an $i = 1, 2$ such that $\alpha_h \beta_i \alpha_h$ and either $\alpha_h \beta_i^\epsilon \beta_{3-i} \alpha_h$ or $\alpha_h \beta_i^\epsilon \beta_{3-i} \alpha_{3-h}$ (where $\epsilon \in \{0, 1\}$ and β_i^0 is the empty word) are factors of w for all $h = 1, 2$. Then if (Π, Υ) is a STEREO-role assignment then the equations $1\beta_i \in \{1, 2\}, 2\beta_i \in \{1, 2\}, 1\beta_i^\epsilon \beta_{3-i} \in \{1, 2\}, 2\beta_i^\epsilon \beta_{3-i} \in \{1, 2\}$ occur in $\Gamma'_w(\Pi, \Upsilon)$, and this yields that the solutions of the systems are pairs of special permutations which both fix the set $\{1, 2\}$, i.e. the system has only trivial solutions. Otherwise if (Π, Υ) is a MONO-role assignment with three blocks the equations $1\beta_i \in \{1, 2\}, 1\beta_i \in \{1, 3\}, 1\beta_i^\epsilon \beta_{3-i} \in \{1, 2\}, 1\beta_i^\epsilon \beta_{3-i} \in \{1, 3\}$ occur in $\Gamma_w(\Pi, \Upsilon)$, and this yields that the solutions of the systems are pair of trivial permutations which both fix $\{1\}$, hence again the system has only trivial solutions.

Let $|\Pi| = 3$. If $\Upsilon = \{\alpha\} \subset \{a, b\}$ then for all $i = 1, 2$ $\Upsilon_i = \Upsilon$, hence the solutions of $\Gamma_w(\Pi, \Upsilon)$ are triples of special permutations of the same type of $\gamma \in \{a, b\} - \Upsilon$, since they are also solutions of $\Gamma_{u_i}(\Pi_i, \Upsilon_i)$. If these permutations are not all of the same type then it must be $\gamma = (1)(2)\dots$. Then if $\gamma = b$ consider the factor $acdbca$ of w , otherwise consider the factor $badacdb$ of w . In the first case the equation $1cdbc \in \{1, 2\}$ occurs in $\Gamma_w(\Pi, \Upsilon)$, in the second case $1adacd \in \{1, 2\}$ occurs in $\Gamma_w(\Pi, \Upsilon)$, and both the equations together with $\gamma = (1)(2)\dots$ yield that c, d are of the same type. Hence the system $\Gamma_w(\Pi, \Upsilon)$ has only trivial solutions.

If $\Upsilon = \{c\}$ then a, b are special permutations of the same type with respect to $(1, 2)$ since they are solutions of $\Gamma_{u_1}(\{a, b\}, \{c\})$, whence $1a, 1a^2, 1b, 1b^2 \in \{1, 2\}$ and b, d are special solutions of the same

type with respect to $(1, 2)$ by applying Lemma 5 to $\Gamma_u(\{a, b\}, \{c\}) \cup \{1b, 1b^2 \in \{1, 2\}\}$. Whence $1d, 1d^2 \in \{1, 2\}$ and again a, d are special solutions of the same type with respect to $(1, 2)$ by applying Lemma 4 to

$$\Gamma_u(\{a, b\}, \{c\}) \cup \{1a, 1a^2, 1d, 1d^2 \in \{1, 2\}\}.$$

If $\Upsilon = \{d\}$, then a, b are special solutions of the same type with respect to $(1, 2)$ since they are solutions of $\Gamma_{u_2}(\{a, b\}, \{d\})$, whence $1a, 1a^2, 1b, 1b^2 \in \{1, 2\}$. More $1ac, 1bca, 1ac^2, 1cb \in \{1, 2\}$ by $\Gamma_u(\{a, b\}, \{d\})$. If a fixes $\{1\}$ but does not fix $\{2\}$, then b too fixes $\{1\}$ and we immediately get $1ac, 1ca, 1c^2, 1c \in \{1, 2\}$ whence a and c are special solutions of the same type with respect to $(1, 2)$ by applying Lemma 5 so that also c fixes $\{1\}$. If $a = (1)(2)\dots$ then $1c, 1bc, 1c^2, 1cb \in \{1, 2\}$ whence again b and c are trivial solutions of the same type by Lemma 5. Finally, let $a = (12)\dots$; then b fixes $\{1, 2\}$ whence $2c, 1c \in \{1, 2\}$. \square

Proposition 9

$$c(2, 5) \leq 119$$

Proof: Let $\Sigma = \{a, b, c, d, e\}$ and let

$$\begin{aligned} r &= cabac^2bacbcacb^2a^2c, \\ s &= dadbda^2b^2dabd^2ab, \\ u &= adacdbcadac^2d^2cbdca, \\ s' &= abeaebea^2b^2eabe^2ab, \\ u' &= aeacebcaea^2e^2cbe. \end{aligned}$$

Consider the words $w_1 = r(bab)su \in \{a, b, c, d\}^+$ and $w_2 = bs'u'r(bab) \in \{a, b, c, e\}^+$.

Like in the proof of Proposition 8 the word w_1 is 2-collapsing on the letters $\{a, b, c, d\}$ and symmetrically the word w_2 is 2-collapsing on the letters $\{a, b, c, e\}$: we only changed the order of some parts, but still we involved as factors some of the minimal 2-collapsing words on three letters. The factors in u and $u'(ca)$ strictly correspond to the one we considered in the proof of Proposition 8.

More, let $v = cedbeacdebdedbedade^2d^2ecedce$ and finally $w = s'u'r(bab)sv$.

(Note that w contains all the factors of w_2). Of course $|w| = 119$; we want to prove that w is 2-collapsing. Let us consider different role assignments (Π, Υ) of Σ and put $\Pi_i = \Pi \cap \Sigma_i$, $\Upsilon_i = \Upsilon \cap \Sigma_i$, for $i = 1, 2$.

If $|\Pi| = 1$ arguments like the ones applied in the proof of Proposition 8 can be used to prove that the system associated to w for (Π, Υ) has only trivial solutions.

Let $|\Pi| = 2, 3$. Assume that, for some $i = 1, 2$, $\Pi \subseteq \Sigma_i$. If Υ is formed by a unique block, then the system associated to w for (Π, Υ) has only trivial solutions by Lemma 7; the same happens when both Υ_i and Υ are decomposed in more than one block. So, for all i such that $\Pi \subseteq \Sigma_i$, Υ_i is formed by a unique block while Υ is formed by more than one block. It is easy to check that if $\Pi \subset \{a, b, c\}$ then either Υ_1 or Υ_2 are formed by two blocks. Then assume $\Pi = \{a, b, c\}$, whence $\Upsilon = (\{d\}, \{e\})$. Hence all the solutions of the system associated to w for (Π, Υ) fix $\{1, 2\}$ by the remark after Lemma 7. Assume that Π is not a subset of $\{a, b, c\}$ and $\Pi \subseteq \Sigma_1$, then $\Pi_1 = \Pi$ and $\Upsilon_2 = \Upsilon = (\Upsilon_1, \{e\})$. In all the solutions of the system associated to w for (Π, Υ) by Lemma 7 the elements in Π_2 fix $\{1, 2\}$ if (Π, Υ) is a STEREO-role assignment and fix $\{1\}$ if it is a MONO-role assignment. Moreover each solution of the system associated with w for (Π, Υ) is formed by permutations which either fix $\{1, 2\}$ or fix $\{1\}$ or fix $\{2\}$ (by remark after

Lemma 7). So let (Π, Υ) be a STEREO-role assignment. Then the solutions of the system $\Gamma'_w(\Pi, \Upsilon)$ either fix all $\{1, 2\}$, or all fix $\{1\}$ ($\{2\}$) and the elements in Π_2 are of the form $(1)(2)\dots$. In the latter case, since $edbe, edce$ are factors of w , when either b or c are in Π_2 we get that d too has the form $(1)(2)\dots$, otherwise we get again that d has the form $(1)(2)\dots$ because w has the factors edb, edc . Hence $\Gamma'_w(\Pi, \Upsilon)$ has only trivial solutions. Then let (Π, Υ) be a MONO-role assignment (with more than one block in Υ). Since w contains all the factors ade, bde, cde we get that d fixes $\{1\}$, whence $\Gamma_w(\Pi, \Upsilon)$ has only trivial solutions. Similar arguments apply when Π is not a subset of $\{a, b, c\}$ and $\Pi \subseteq \Sigma_2$. So let $\{d, e\} \subseteq \Pi$. Obviously $\Upsilon_i = \Upsilon$ for all $i = 1, 2$. Then if Υ contains at least two blocks then the solutions of the system associated to w for (Π, Υ) are trivial by Lemma 7. So let (Π, Υ) be a MONO-role assignment with a unique block in Υ . Then all the solutions of $\Gamma_w(\Pi, \Upsilon)$ are formed by special permutations with respect to $(1, 2)$ and if $|\Pi_i| > 1$ then the permutations in Π_i are of the same type. Assume that the permutations in Π_1 and in Π_2 are not of the same type then if $\Pi \subseteq \{a, d, e\}$, the factors $cdeb, bdec, cedb, bedc$ give that both d, e fix $\{1, 2\}$. Moreover $beaeb$ is a factor of w so again both a, e fix $\{1, 2\}$ provided that $a \in \Pi$. Now let $\Pi = \{b, d, e\}$. If there is a solution of $\Gamma_w(\Pi, \Upsilon)$ which is not trivial then b has the form $(1)(2)\dots$, and the factor $cedbea$ gives that both d, e fix $\{1, 2\}$. If $\Pi = \{c, d, e\}$ then if there is a solution of $\Gamma_w(\Pi, \Upsilon)$ which is not trivial then c has the form $(1)(2)\dots$, and again the factor $bdedb$ gives that both d and e fix $\{1, 2\}$. Hence the systems associated to w for each partition (Π, Υ) with $|\Pi| = 2, 3$ have only trivial solutions.

So, let $|\Pi| = 4$ and let $\Upsilon = \{\alpha\}$. If $\alpha \in \{a, b, c\}$ then in each solution of $\Gamma_w(\Pi, \Upsilon)$ all the elements in Π_1 are special permutations of the same type with respect to $(1, 2)$ and the same holds for all the elements in Π_2 . So if in Π there exist elements of different form then the elements in $\{a, b, c\} - \{\alpha\}$ must have the form $(1)(2)\dots$. But, in such case, considering the equations generated by some of the factors $acedbea, bdcacedb, cedbeac$ of w it is easy to verify that also d and e fix $\{1, 2\}$. If $\alpha = d$ then in each solution of $\Gamma_w(\Pi, \Upsilon)$ a, b, c are special of the same type with respect to $(1, 2)$ by Lemma 7 and $1cace, 1beac, 1eb, 1e, 1be, 1e^2, 1ece \in \{1, 2\}$ occur in $\Gamma_{(bdca)v}(\Pi, \Upsilon)$ whence they occur also in $\Gamma_w(\Pi, \Upsilon)$; hence b, e and c, e are special of the same type with respect to $(1, 2)$. If one among b, c is different from $(1)(2)\dots$ then all of them are of the same type; but if both of them fix both 1 and 2 then one gets $1ae, 1ea \in \{1, 2\}$ and so all the solutions of $\Gamma_w(\Pi, \Upsilon)$ are trivial. If $\alpha = e$ then again in each solution of $\Gamma_w(\Pi, \Upsilon)$ a, b, c are special of the same type with respect $(1, 2)$. Moreover $1db, 1acd, 1bd, 1dad, 1d^2, 1dc \in \{1, 2\}$ occur in $\Gamma_{(bdca)v}(\Pi, \Upsilon)$, whence d, b and d, a are special of the same type by Lemmas 5 and 4. Again, assume $a = b = (1)(2)\dots$: this yields at last $1cd, 1dc \in \{1, 2\}$ and again all the solutions of $\Gamma_w(\Pi, \Upsilon)$ are trivial. \square

In Proposition 8 of [7] we considered a slightly longer word, so proving that $c(2, 5) \leq 120$. But in [11] an algorithm is presented to find shorter 2-collapsing words from a given 2-collapsing word. The author run her algorithm on our word and got four 2-collapsing words of length 119, one of which is the word considered in the above Proposition 9. She also did the same on the word in our Proposition 8: she could not find shorter 2-collapsing words, but found three more 2-collapsing words of the same length.

Of course Propositions 8 and 9 give upper bounds to the length of the minimal 2-synchronizing words respectively on 4 and 5 letters. But this bounds can be further improved: denote by $s(2, t)$ the length of the least 2-synchronizing word on t letters.

Proposition 10

$$s(2, 4) \leq 52$$

Proof: Let $\Sigma = \{a, b, c, d\}$ and let

$$\begin{aligned} r &= cabab^2cbcaca^2b^2c^2a \in \{a, b, c\}^+, \\ s &= bd^2a^2b^2dbdada^2babd \in \{a, b, d\}^+. \end{aligned}$$

The word $w = (cd^2cbcd)r(ab)s(cdadc^2d)$ is 2-synchronizing and $|w| = 52$. In order to exhibit w we used two of the minimal 2-synchronizing words on 3 letters listed in [3]. Namely $u_1 = r(ab)$ is the 11-th word in the list in [3], on the alphabet $\Sigma_1 = \{c, a, b\}$, while $u_2 = (ab)s$ is the 15-st one in that list, on the alphabet $\Sigma_2 = \{a, b, d\}$. Both the words u_1 and u_2 are 2-collapsing ([3]).

Let us consider different role assignments (Π, Υ) of Σ and put $\Pi_i = \Pi \cap \Sigma_i$, $\Upsilon_i = \Upsilon \cap \Sigma_i$, for $i = 1, 2$. The cases when $|\Pi| = 1, 2$ can be proved with similar arguments as in the proof of Proposition 8.

Let $|\Pi| = 3$. We are looking for solutions of our systems of equations on 3 permutations t, y, z in the symmetric group Sym_3 . If a system has only special solutions with respect to $\{1, 2\}$ and t, y are of different types while both of them are of the same of z , then without loss of generality $t = (12)(3)$ and $y = (1)(23)$; in such a case we immediately get that $z = (1)(2)(3) = t^2 = y^2$.

Now let $\alpha \in \Upsilon$. If $\alpha = a$ (resp. $\alpha = b$) the permutations b, c (resp. a, c) are trivial solutions of $\Gamma_{u_1}(\Pi_1, \{\alpha\})$ and b, d (resp. a, d) are trivial solutions of $\Gamma_{u_2}(\Pi_2, \{\alpha\})$, whence both c and d are of the same type as b (resp. of a). If b (resp. a) is different from the identity, then both c and d must be of the same type by the previous remark, but this is true also on the other case as the equation $1bcd \in \{1, 2\}$ belongs to $\Gamma_w(\Pi, \{a\})$ (resp. $1cdca \in \{1, 2\}$ belongs to $\Gamma_w(\Pi, \{b\})$). If $\alpha = c$ then a, b are trivial solutions of $\Sigma_{u_1}(\Pi_1, \{\alpha\})$. More $cdc, cd^2c, cdadc$ are factors of w . Since $1a, 1a^2 \in \{1, 2\}$ then only special permutations of the same type are solutions of $\Gamma_w(\Pi, \{c\})$ by Lemma 4. Thus b and d are special permutations of the same type as a . But the equation $1au_2 \in \{1, 2\}$ appears in $\Gamma_w(\Pi, \{c\})$: if a is the identity it immediately yields that $1dbd \in \{1, 2\}$ by previous remark. By Lemma 4 b and d are of the same type. A similar argument runs when $\alpha = d$ since $dcd, dc^2d, dcbed, du_1bd$ are factors of w . \square

Proposition 11

$$s(2, 5) \leq 111$$

Proof: Let $\Sigma = \{a, b, c, d, e\}$. We want to exhibit a 2-synchronizing word of length 111 on Σ , like in the proof of Proposition 10. We shall consider two slightly longer 2-synchronizing word on 4 letters with a better overlapping:

$$\begin{aligned} r &= ab^2c^2abacab^2cbca^2c ; s = bd^2abadaba^2bdb^2ad ; u = acd^2c^2dabcadcdbd ; \\ w_1 &= ur(ab)s ; s' = s ; r' = dedea^2eadaead^2e^2a ; u' = dbaebdeabe^2b^2edb ; \\ w_2 &= (ab)sr'u' ; v = cebeacdaece ; v' = cec^2e^2cbcedec. \end{aligned}$$

The word w_1 (resp. w_2) is 2-synchronizing on the alphabet $\{a, b, c, d\}$ (resp. $\{a, b, d, e\}$), by similar arguments as in the proof of Proposition 10. In fact it is built from two words in the list in [3] for suitable alphabets: $r(ab)$ is the 6-th and $(ab)s$ is the 16-th (resp. $(ab)s'$ is the 16-th while $(dad)r'$ is the 1-st) and u (resp. u') gives us the other necessary factors.

The word $w = vur(ab)sr'u'v'$ is 2-synchronizing and its length equals 111. \square

Acknowledgements

The authors thank Elena Pribavkina for informing them about the results of applying her algorithm on the short 2-collapsing words presented in [7] and for her careful reading of the present paper.

References

- [1] D. S. Ananichev, A. Cherubini, M. V. Volkov, *Image reducing words and subgroups of free groups*, Theor. Comput. Sci. **307**, no.1, (2003), 77–92.
- [2] D. S. Ananichev, A. Cherubini, M. V. Volkov, *An inverse automata algorithm for recognizing 2-collapsing words*, in: M. Ito, M. Toyama (eds.), *Developments in Language Theory*, Lect. Notes Comp. Sci. 2450, Springer, Berlin 2003, 270–282.
- [3] D. S. Ananichev, I. V. Petrov, *Quest for short synchronizing words and short collapsing words*, WORDS 2003. Proc. 4th Int. Conf., Univ. of Turku, Finland, 411–418.
- [4] D. S. Ananichev, I. V. Petrov, M. V. Volkov, *Collapsing words: a progress report*, Internat. J. Foundations Comp. Sci. **17**, no.3, (2006), 507–518.
- [5] J. Černý, *Poznámka k homogénnym experimentom s konečnými automatami*, Mat.-Fyz. Cas. Slovensk. Akad. Vied. **14**, (1964), 208–216 [in Slovak].
- [6] A. Cherubini, *Synchronizing and collapsing words*, Milan J. Math. **75**, (2007), 305–321.
- [7] A. Cherubini, A. Kisielewicz, B. Piochi, *A bound for the length of shortest 2-collapsing words*, in: P. Arnoux, N. Bedaride, J. Cassaigne (eds.), WORDS 2007. Proc. 6th Int. Conf. on Words, 90–99.
- [8] A. Cherubini, P. Gawrychowski, A. Kisielewicz, B. Piochi, *A combinatorial approach to collapsing words*, in: R. Krýlovič, P. Urzyczyn (eds.), *Mathematical Foundations of Computer Science 2006*, Lect. Notes Comp. Sci. 4162, Springer, Berlin 2006, 256–266.
- [9] S. W. Margolis, J.-E. Pin, M. V. Volkov, *Words guaranteeing minimum image*, Internat. J. Foundations Comp. Sci. **15**, no.2, (2004), 259–276.
- [10] E. V. Pribavkina, *On some properties of the language of 2-collapsing words*, Internat. J. Foundations Comp. Sci. **17**, no.3, (2006), 665–676.
- [11] E. V. Pribavkina, *2-collapsing words and a sequence reconstruction problem*, CD Proceedings of AutoMathA Conference 2007, Mondello, Italy.
- [12] N. Sauer, M. G. Stone, *Composing functions to reduce image size*, Ars Combinatoria **31**, (1991), 171–176.