

On a 1, 2 Conjecture

Jakub Przybylo, Mariusz Woźniak

► **To cite this version:**

Jakub Przybylo, Mariusz Woźniak. On a 1, 2 Conjecture. Discrete Mathematics and Theoretical Computer Science, DMTCS, 2010, 12 (1), pp.101-108. <hal-00990444>

HAL Id: hal-00990444

<https://hal.inria.fr/hal-00990444>

Submitted on 13 May 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

On a 1, 2 Conjecture

Jakub Przybyło[†] and Mariusz Woźniak[‡]

AGH University of Science and Technology, Al. Mickiewicza 30, 30-059 Kraków, Poland

received February 12, 2008, accepted February 3, 2010.

Let us assign positive integers to the edges and vertices of a simple graph G . As a result we obtain a vertex-colouring of G with integers, where a vertex colour is simply a sum of the weight assigned to the vertex itself and the weights of its incident edges. Can we obtain a proper colouring using only weights 1 and 2 for an arbitrary G ?

We give a positive answer when G is a 3-colourable, complete or 4-regular graph. We also show that it is enough to use weights from 1 to 11, as well as from 1 to $\lfloor \frac{\chi(G)}{2} \rfloor + 1$, for an arbitrary graph G .

Keywords: neighbour-distinguishing total-weighting, irregularity strength

1 Introduction

A k -total-weighting of a simple graph G is an assignment of an integer weight, $w(e), w(v) \in \{1, \dots, k\}$ to each edge e and each vertex v of G . The k -total-weighting is *neighbour-distinguishing* (or *vertex colouring*, see (1)) if for every edge uv , $w(u) + \sum_{e \ni u} w(e) \neq w(v) + \sum_{e \ni v} w(e)$. If it exists, we say that G *permits* a neighbour-distinguishing k -total-weighting. The smallest k for which G permits a neighbour-distinguishing k -total-weighting we denote by $\tau(G)$.

A similar parameter, but in the case of an *edge* (not total) weighting was introduced and studied in (8) by Karoński, Łuczak and Thomason. They conjectured that every simple connected graph that is not K_2 permits a *neighbour-distinguishing 3-edge-weighting*, and showed that this statement holds *e.g.* for the 3-colourable graphs. It is also known, see (1), that each *nice graph* (*i.e.*, a graph without a K_2 -component) permits a neighbour-distinguishing 16-edge-weighting, hence this parameter is finite.

Note that if a graph permits a neighbour-distinguishing k -edge-weighting, then it also permits a neighbour-distinguishing k -total-weighting (it is enough to put ones at all vertices), hence we obtain an upper bound $\tau(G) \leq 16$ for all graphs and $\tau(G) \leq 3$ for the 3-colourable graphs (for all graphs if the conjecture by Karoński, Łuczak and Thomason holds). Therefore, we ask if, maybe, the weights 1 and 2 are enough in the case of a total-weighting.

Conjecture 1 *Every simple graph permits a neighbour-distinguishing 2-total-weighting.*

[†]The research of the first author was partially supported by MNiSW grant N N201 389134.

[‡]The research of the second author was partially supported by polish ministry MNiSW

It might seem quite plausible in the face of the result by Addario-Berry, Dalal and Reed from (1), which says that for any fixed $p \in (0, 1)$ the random graph $G_{n,p}$ (asymptotically) almost surely permits neighbour-distinguishing 2-edge-weighting.

In the following section we discuss the correspondence between $\chi(G)$ and $\tau(G)$. The obtained upper bound for $\tau(G)$ yields a positive verification of Conjecture 1 for the 3-colourable graphs, discussed, together with the results for the complete and 4-regular graphs, in Section 3. The last section, Sec. 4, is devoted to decreasing the general upper bound for $\tau(G)$ from 16 down to 11.

It is also worth mentioning here that our reasonings correspond with the recent study of Bača, Jendroľ, Miller and Ryan. In (3) they introduced and studied a parameter called a *total vertex irregularity strength*, which is the smallest k for which there exists a k -total-weighting such that each vertex of a graph receive a different colour, i.e. $w(u) + \sum_{e \ni u} w(e) \neq w(v) + \sum_{e \ni v} w(e)$ for each pair of vertices u, v (which do not have to be adjacent). This parameter, as well as the other mentioned in this section, may be viewed as descendants of the well known *irregularity strength* of a graph, see (4) or (9).

Due to a considerable amount of time consumed by the reviewing process, we conclude the introduction with a short list of up-to-date results in the field. All of these were however predated and (most of them) inspired by a manuscript of this paper. First Przybyło showed in (10) that $\tau(G) \leq 7$ for all regular graphs. Then a huge break-through was a result of Kalkowski (5), who proved that $\tau(G) \leq 3$ for all graphs. Following and (considerably) modifying his algorithm led Kalkowski, Karoński and Pfender (7) to decreasing the general bound of 16 to 5 in the case of edge-weightings. Also the list version of the “1,2 Conjecture” has been introduced by Wong and Zhu (15), and independently by Przybyło and Woźniak (13). Among others one can read in either of these, that Proposition 9 of this paper can be extended to its list version. See also (2; 6; 11; 12; 14) for more related results.

2 Bounds for graphs with low chromatic number

Our aim is to show that $\tau(G) \leq 2$ for a simple graph G . Note that $\tau(G) = 1$ iff there are no neighbours with the same degree in G . Since we wish to distinguish only neighbours, we may assume that G is a connected graph. Given a total-weighting w of G , let $c_w(v) := w(v) + \sum_{e \ni v} w(e)$ (or $c(v)$ for short if the weighting w is obvious) define the resulting colouring for each $v \in V(G)$. We shall call $c(v)$ a *colour* or a *total weight* of v . Surprisingly easily one may prove the following statement.

Proposition 2 $\tau(G) \leq 2$ for bipartite graphs.

Proof: Let us first arbitrarily assign weights 1 or 2 to the edges of G . Then put 1 or 2 at vertices so that the resulting total weights of the vertices in one colour class are even and the total weights of the vertices in the other colour class are odd. \square

Consequently, $\tau(G) = 2$ if G is a single edge, hence our parameter makes sense for all graphs (while considering edge-weightings we must restrict ourselves to graphs without K_2 -components).

Actually, similar reasoning provides us with a more general observation.

Proposition 3 Given a graph G and a set of target colours t_v for all $v \in V(G)$, there exists a 2-total-weighting w of G such that $c_w(v) \equiv t_v \pmod{2}$ for each $v \in V(G)$. \square

In this section we discuss a generalization of this statement. Consequently, we obtain an upper bound for $\tau(G)$ based on the chromatic number of G .

Lemma 4 Given a cycle C , a set of target colours t_v for all $v \in V(C)$, and a positive integer $p \geq 3$, there exists a $(\lfloor \frac{p}{2} \rfloor + 1)$ -total-weighting w of C such that $c_w(v) \equiv t_v \pmod{p}$ for each $v \in V(C)$.

Proof: Let v_1, \dots, v_n be the consecutive vertices on the cycle C , $|C| = n$. We may assume $t_{v_i} \in [3, p+2]$ for each i . Let $V(C) = S \cup L$, where $v_i \in S$ if $t_{v_i} \in [3, \lfloor \frac{p}{2} \rfloor + 2]$ or $v_i \in L$ if $t_{v_i} \in [\lfloor \frac{p}{2} \rfloor + 3, p+2]$. Denote t_{v_i} by s_i if $v_i \in S$ or by l_i if $v_i \in L$, and set $h = \lfloor \frac{p}{2} \rfloor + 1$. We will use only weights $1, \dots, h$.

Note first, that if $|L|$ is even, then we can easily find the required weighting. To do this, it is sufficient to use only numbers 1 and h as the weights of the edges. We start by setting $w(v_n v_1) = 1$, and then assign 1 or h to the consecutive edges of C one after another, so that the two edges adjacent to v_i obtain the same weights if $v_i \in S$ or different ones otherwise. This way, the arities (mod p) of the current total weights of the vertices in S belong to the set $\{1, 2\}$ (depending on the evenness of p) and are equal to $\lfloor \frac{p}{2} \rfloor + 2$ for the ones in L . Hence, it is now easy to complete the weighting using numbers from the set $\{1, \dots, h\}$ as the weights of the vertices of C .

Let then $|L|$ be odd and assume first that a vertex in L , say v_2 , has the two neighbours (v_1 and v_3) in S . If

$$s_1 - 2 + s_3 - 2 \geq l_2 - h, \quad (1)$$

then there exist $s'_1 \in [1, s_1 - 2]$, $s'_3 \in [1, s_3 - 2]$ such that $s'_1 + s'_3 = l_2 - h$. Set $w(v_1 v_2) = s'_1$, $w(v_2) = h$ and $w(v_2 v_3) = s'_3$ to obtain $c_w(v_2) = l_2$. Then set $w(v_n v_1) = w(v_3 v_4) = 1$ (we have $v_n v_1 = v_3 v_4$ if $C = C_3$). Since there remain an even number of weightless vertices in L , we apply the same reasoning as in the above second paragraph of the proof (by assigning weights 1 or h to the remaining edges of C first, and dealing with the vertices then) to obtain the desired weighting. On the other hand, if

$$p + s_1 - 2h + p + s_3 - 2h \leq l_2 - 1, \quad (2)$$

then there exist $s'_1 \in [p + s_1 - 2h, h]$, $s'_3 \in [p + s_3 - 2h, h]$ such that $s'_1 + s'_3 = l_2 - 1$. Analogously as above, set $w(v_1 v_2) = s'_1$, $w(v_2) = 1$ and $w(v_2 v_3) = s'_3$ to obtain $c_w(v_2) = l_2$. Then put $w(v_n v_1) = w(v_3 v_4) = h$ and apply the technique from the second paragraph to finish the weighting. It is easy to verify that all the weights used were in the range $[1, h]$, and that one of the inequalities (1) or (2) holds if only $h \geq 2$.

Finally, we may assume that L contains two, say v_2, v_3 , consecutive vertices of the cycle (and $|L|$ is odd). Let $l_2 \leq l_3$. Then $h + l_2 - l_3, l_3 - h - 1 \in \{1, \dots, h\}$, hence it is enough to set $w(v_1 v_2) = 1$, $w(v_2) = h + l_2 - l_3$, $w(v_2 v_3) = l_3 - h - 1$, $w(v_3) = 1$ and $w(v_3 v_4) = h$ to obtain $c_w(v_2) = l_2$ and $c_w(v_3) = l_3$. This time we are left with an odd number of weightless vertices in L . However, thanks to the weights assigned to $v_1 v_2$ and $v_3 v_4$, we can again complete the weighting of C analogously as in the second paragraph. \square

Lemma 5 Given a connected graph G , $u \in V(G)$, a set of target colours t_v for all $v \in V(G)$, and a positive integer p , there exists a $(\lfloor \frac{p}{2} \rfloor + 1)$ -total-weighting w of G such that $c_w(v) \equiv t_v \pmod{p}$ for each $v \in V(G) \setminus \{u\}$.

Proof: Given G and $u \in V(G)$, we use weights $1, \dots, \lfloor \frac{p}{2} \rfloor + 1$ to obtain the desired weighting as follows. First assign weight 1 to all the edges and vertices of G . If the obtained colouring does not comply with our requirements, i.e. there exists $v \in V(G) \setminus \{u\}$ with improper colour arity (not congruent to $t_v \pmod{p}$), choose a path from v to u in G . By changing the weights of v and an edge from this path adjacent to

v , we obtain $c_w(v) \equiv t_v \pmod{p}$ for the resulting weighting w . Then continue traversing the path, at each step changing, if necessary, the weights of one consecutive vertex and one edge, so that at the end the colours of all the vertices from the path different from u have the proper arities mod p . Note, we do not change the total weights of the vertices that do not belong to the path. This way we have reduced the number of the vertices in $V(G) \setminus \{u\}$ with improper colour arities. Therefore, applying the procedure described repeatedly yields the desired result. \square

Theorem 6 *Given a connected graph G which is not a tree, a set of target colours t_v for all $v \in V(G)$, and a positive integer $p \geq 3$, there exists a $(\lfloor \frac{p}{2} \rfloor + 1)$ -total-weighting w of G such that $c_w(v) \equiv t_v \pmod{p}$ for each $v \in V(G)$.*

Proof: Since G is not a tree, it contains a subgraph C which is a cycle. Let u be an arbitrary vertex of this cycle. By Lemma 5 we may find such a weighting of G that the resulting total weights of all vertices $v \in V(G) \setminus \{u\}$ are as required. Then put a temporary weight 0 at all the vertices and on each edge of C . Denote the resulting weighting of G by w' , and let $s_v = c_{w'}(v)$ for $v \in V(C)$. To obtain the desired weighting, it is enough to apply Lemma 4 to C with the target arities $t_v - s_v$ for each vertex v of C . \square

Corollary 7 $\tau(G) \leq \lfloor \frac{\chi(G)}{2} \rfloor + 1$ for an arbitrary graph G .

Proof: By Theorem 6 (or Proposition 3) there is a $(\lfloor \frac{\chi(G)}{2} \rfloor + 1)$ -total-weighting of G such that the resulting total weights have distinct arities mod $\chi(G)$ for vertices in different colour classes. \square

In the last section of this paper we show that $\tau(G) \leq 11$, see Theorem 12, hence the above corollary provides us with a reasonable upper bound for graphs with $\chi(G) \leq 19$.

3 Exact values of $\tau(G)$

By Corollary 7, we obtain the following one.

Corollary 8 *Each 3-colourable graph permits a neighbour-distinguishing 2-total-weighting.* \square

The next two results exemplify the fact that $\tau(G)$ not necessarily depends on $\chi(G)$. Though the following proposition is the consequence of (3), we present here our proof for the cohesion of the article.

Proposition 9 $\tau(G) = 2$ for complete graphs.

Proof: We apply induction to show that it is enough to use 1 and 2 to obtain such a weighting of K_n , that the colours of its vertices are n consecutive integers, either from n to $2n - 1$ or from $n + 1$ to $2n$. The assertion is trivial for $n = 2$.

Assume then that $n \geq 3$ and that we have found such a weighting for K_{n-1} . Let us add a new vertex v joining it by a single edge with each vertex of K_{n-1} . Note that the vertices of K_{n-1} carry colours from the interval $[n - 1, 2n - 2]$ which are consecutive integers. If the greatest of them equals $2n - 3$, we put 2 at v and on all the edges incident with it. This way, the vertices of K_n obtain n different colours from the interval $[n + 1, 2n]$. Analogously, if the greatest total weight at a vertex of K_{n-1} equals $2n - 2$, we put 1 at v and on all the edges incident with it. \square

Theorem 10 $\tau(G) = 2$ for all 4-regular graphs.

Proof: Let G be a connected 4-regular graph. If $G = K_5$ or $\chi(G) \leq 3$, then we are done by Proposition 9 or Corollary 8, respectively. Hence, by Brooks' Theorem, we may assume $\chi(G) = 4$. Choose the colour classes, A, B, C and D for G so that A is as large as possible, and, subject to the choice of A , B is also as large as possible, and finally, subject to the choice of A and B , C is as large as possible as well. Consequently, each vertex in $B \cup C \cup D$ has at least one neighbour in A , each vertex in $C \cup D$ has at least one neighbour in B , and each vertex in D has at least one neighbour in C . Let $D = D_1 \cup D_2$, where $v \in D_i$ if v has exactly i neighbours in A , and let for $X, Y \subset V(G)$, $E(X, Y)$ denote the set of all edges between X and Y in G . We define a weighting w in the following way.

Assign 2 to the edges in $E(A, B \cup C \cup D)$, 1 to the edges in $E(D, B \cup C)$, 2 to the vertices in $A \cup D_2$, and 1 to the vertices in D_1 . This way the induced subgraph $G[B \cup C]$ remains weightless, while $c(v) = 10$ if $v \in A$, $c(v) = 6$ if $v \in D_1$, or $c(v) = 8$ if $v \in D_2$. For each edge $xy \in E(B, C)$ where $y \in C$, we assign 2 to it if y has a neighbour in D_1 or 1 otherwise. Then choose the weights for each vertex u in B so that their total weights are odd. Since each of them is incident with at least one edge (in $E(B, A)$) with weight 2, we have $c(u) \in \{7, 9\}$. Now all adjacent vertices in $A \cup B \cup D$ are distinguished. We will put 1 or 2 at the vertices in C and adjust the weights of some of the edges (in $E(C, D_2)$) and vertices (in D_2), so that the colours of the vertices in $A \cup B \cup D$ stay unchanged, while the vertices in C are distinguished from the others. Given an arbitrary vertex v in C , note that it is incident with at least one edge (in $E(C, A)$) with weight 2 and at least one with weight 1. Consider the four edges incident with v .

If three of them have weight 1, hence $N(v) \cap D_1 = \emptyset$, we set $w(v) = 1$, and thus $c(v) = 6$.

If at least two of them have weight 2 and $N(v) \cap D_2 = \emptyset$, we choose $w(v)$ so that $c(v) = 8$.

Finally, if at least two of them have weight 2 and there is a vertex $y \in N(v) \cap D_2$, then (by our construction) $w(y) = 2$, $w(yv) = 1$ and v has exactly one neighbour, say x , in B . Moreover, exactly two edges incident with v have weight 1. If $c(x) = 9$, we put 1 at v and get $c(v) = 7$. On the other hand, if $c(x) = 7$, set $w(y) = 1$, $w(yv) = 2$ and $w(v) = 2$. This way the colour of y remains unchanged, while $c(v) = 9$.

After processing all the vertices v in C we obtain a weighting that is neighbour-distinguishing. \square

4 General bound

In (1), the following theorem was proved.

Theorem 11 Let $G = (V, E)$ be a bipartite graph with bipartition $V = X \cup Y$. For each $v \in X$, let $a_v^- = \lfloor \frac{d(v)}{2} \rfloor$ and set $a_v^+ = a_v^- + 1$. For each $v \in Y$, fix any a_v^-, a_v^+ such that $a_v^- \leq \lfloor \frac{d(v)}{2} \rfloor \leq a_v^+$ and

$$a_v^+ \leq \min \left(\frac{d(v) + a_v^-}{2} + 1, 2a_v^- + 1 \right). \quad (3)$$

Then there is a spanning subgraph H of G such that $d_H(v) \in \{a_v^-, a_v^+\}$ for all $v \in V$.

We use it, together with the construction by Addario-Berry, Dalal and Reed from (1), to prove the following statement.

Theorem 12 Every graph permits a neighbour-distinguishing 11-total-weighting.

Proof: Let G be a connected graph. For any ordering of a set of its vertices, let $F(v_i) = \{v_j \mid v_j \in N(v_i) \text{ and } j > i\}$ and $B(v_i) = \{v_j \mid v_j \in N(v_i) \text{ and } j < i\}$. Call these sets the *forward* and the *backward* neighbours of v_i , respectively. Choose an ordering of $V(G)$ that maximizes $k = \max\{j : |F(v_i)| > |B(v_i)| \text{ for } i \leq j\}$. Put the first k vertices in V_1 and the remaining ones into a temporary set T . Note that k defined above does not decrease regardless of any reordering of the vertices within T . Moreover, $d_T(v) \leq d_{V_1}(v)$ for all $v \in T$. (Otherwise, we could move v to the $(k+1)$ 'st position of the ordering and thereby create an ordering with a larger value of k .)

Next, apply the above described procedure to the induced graph $G[T]$, to find a set $V_2 \subseteq T$ and a new ordering of the vertices in T . Remove the vertices of V_2 from the temporary set T . Then repeat this procedure twice more to find V_3, V_4 and the final ordering of the vertices of G . Let $V_5 = V(G) \setminus (V_1 \cup V_2 \cup V_3 \cup V_4)$. Note that each $v \in V_i, i = 1, 2, 3, 4$, has strictly fewer backward neighbours in V_i than forward neighbours. By the observation at the end of the previous paragraph, for each $v \in V_5 \subset T$, $d_{V_2}(v) + d_{V_3}(v) + d_{V_4}(v) + d_{V_5}(v) = d_{V_2 \cup V_3 \cup V_4 \cup V_5}(v) \leq d_{V_1}(v)$. Analogously we have $d_{V_3}(v) + d_{V_4}(v) + d_{V_5}(v) \leq d_{V_2}(v)$, $d_{V_4}(v) + d_{V_5}(v) \leq d_{V_3}(v)$, $d_{V_5}(v) \leq d_{V_4}(v)$, and hence $8d_{V_5}(v) \leq d_{V_1}(v)$ for each $v \in V_5$.

Consider the edges from V_5 to V_1 . Since every vertex $v \in V_5$ has at least $8d_{V_5}(v)$ edges to V_1 , we can choose a subset (of these edges) where each $v \in V_5$ has exactly $8d_{V_5}(v)$ edges to V_1 . Let B be the bipartite graph spanned by this reduced set of edges.

Now set $w(e) = 2$ for each edge of G . Then assign weights between 3 and 10 to the vertices so that the resulting colour of each vertex has an arity mod 8 as specified in Table 1. We will modify this weighting

V_1	V_2	V_3	V_4	V_5
1	3	5	7	0, 2, 4 or 6

Tab. 1: Target arity for partition elements

maintaining the specified colour arities mod 8 and eliminating possible internal conflicts within the sets V_1, \dots, V_5 .

Process the vertices of $V_1 \cup V_2 \cup V_3 \cup V_4$ one after another according to the specified order. Given a vertex $v \in V_i$, we add 8 to the weights of an arbitrary subset of the forward edges of v in order to distinguish its colour from the colours of its backward neighbours in V_i . This is always feasible, since v has at least one more forward neighbour (not necessarily in V_i) than backward neighbours in V_i .

After processing all the vertices in $V_1 \cup V_2 \cup V_3 \cup V_4$, $w(e) \in \{2, 10\}$, $3 \leq w(v) \leq 10$ for each edge e and each vertex v of G , while the arities (mod 8) of colours are as specified in Table 1. Moreover, every two neighbours from $V_1 \cup V_2 \cup V_3 \cup V_4$ have distinct colours. In our final step, we adjust the weights of edges in B to distinguish adjacent vertices in V_5 . First we use Theorem 11 where $X = V_1 \cap V(B)$ and $Y = V_5 \cap V(B)$ to determine a subgraph H of B . Then for each edge e of B we will add 1 to its weight if $e \in E(H)$ or subtract 1 otherwise (we may do so, since $w(e) \in \{2, 10\}$). At the end we modify the weights of the vertices in V_1 so that their colours are as they were before this final step.

Define $\{a_v^-, a_v^+\}$ for each $v \in X$ by setting $a_v^- = \lfloor \frac{d_B(v)}{2} \rfloor$ and $a_v^+ = a_v^- + 1$. Then choose $\{a_v^-, a_v^+\}$ for each vertex in Y as follows. Process the vertices of Y in any order. For each $v \in Y$ in turn, we choose $a_v^- \in [\frac{d_B(v)}{4}, \frac{d_B(v)}{2}]$ (this range has integer endpoints, because 8 divides $d_B(v)$), and set $a_v^+ = a_v^- + \frac{d_B(v)}{4} + 1$. We make our choice to ensure that for any previously processed neighbour $u \in V_5$ of v , for any

$a_v \in \{a_v^-, a_v^+\}$, and for any $a_u \in \{a_u^-, a_u^+\}$, we have $c(v) + a_v - (d_B(v) - a_v) \neq c(u) + a_u - (d_B(u) - a_u)$, where by $c(v)$, $c(u)$ we mean the current colours of v , u (and $c(v) + a_v - (d_B(v) - a_v)$ will be the colour of v after the final step.) This is possible since each previously processed neighbour of v can prevent at most two choices for a_v^- and there are precisely $2d_{V_5}(v) + 1$ choices available for a_v^- .

This set of degree choices satisfies the conditions of Theorem 11. Indeed, it is clear for the vertices in X . Also obviously $a_v^- \leq \lfloor \frac{d_B(v)}{2} \rfloor \leq a_v^+$ for each $v \in Y$, so it remains to show that for all $v \in Y$, (3) holds. Since $a_v^- \leq \frac{d_B(v)}{2}$, then $a_v^+ = a_v^- + \frac{d_B(v)}{4} + 1 = \frac{d_B(v)}{4} + \frac{a_v^-}{2} + \frac{a_v^-}{2} + 1 \leq \frac{d_B(v)}{2} + \frac{a_v^-}{2} + 1$. On the other hand, since $a_v^- \geq \frac{d_B(v)}{4}$, then $a_v^+ = a_v^- + \frac{d_B(v)}{4} + 1 \leq 2a_v^- + 1$. Thus, by Theorem 11, a subgraph H of B exists such that after performing the previously described additions and subtractions on the set of edges of B , all adjacent vertices in V_5 have distinct colours. Note that such changes may influence the total weights of the vertices in V_1 , adding 1 or 2 to some of them, or decreasing them by 1 (depending on the evenness of $d_B(v)$ and the choice of a_v^- or a_v^+). Then however, for each such $v \in V_1$ (since $3 \leq w(v) \leq 10$) we balance this change by subtracting 1 or 2, or by adding 1, respectively, to its weight $w(v)$.

Note that the arities (mod 8) of colours of the vertices in V_5 remain as they are specified in Table 1 (they are even numbers), and that $w(v), w(e) \in \{1, \dots, 11\}$ for all $v \in V(G)$ and all $e \in E(G)$, hence the described construction provides us with a neighbour-distinguishing 11-total-weighting. \square

References

- [1] L. Addario-Berry, K. Dalal, B. A. Reed, *Degree constrained subgraphs*, Proceedings of GRACO2005, volume 19 of Electron. Notes Discrete Math., Amsterdam (2005), 257-263 (electronic), Elsevier.
- [2] M. Anholcer, M. Kalkowski, J. Przybyło, *A new upper bound for the total vertex irregularity strength of graphs*, Discrete Math. 309 (21) (2009) 6316-6317.
- [3] M. Bača, S. Jendroř, M. Miller, J. Ryan, *On Irregular Total Labelings*, Discrete Math. 307 (2007) 1378-1388.
- [4] G. Chartrand, M. S. Jacobson, J. Lehel, O. R. Oellermann, S. Ruiz, F. Saba, *Irregular networks*, Proc. of the 250th Anniversary Conf. on Graph Theory, Fort Wayne, Indiana (1986).
- [5] M. Kalkowski, *On 1,2-conjecture*, submitted.
- [6] M. Kalkowski, M. Karoński, F. Pfender, *A new upper bound for the irregularity strength of graphs*, manuscript.
- [7] M. Kalkowski, M. Karoński, F. Pfender, *Vertex-coloring edge-weightings: Towards the 1-2-3-conjecture*, to appear in J. Combin. Theory Ser. B.
- [8] M. Karoński, T. Łuczak, A. Thomason, *Edge weights and vertex colours*, J. Combin. Theory, Ser. B 91 (2004) 151-157.
- [9] J. Lehel, *Facts and quests on degree irregular assignments*, Graph Theory, Combinatorics and Applications, Willey, New York, (1991) 765-782.

- [10] J. Przybyło, *A note on a neighbour-distinguishing regular graphs total-weighting*, Electron. J. Combin. 15 (1) (2008) #N35.
- [11] J. Przybyło, *Irregularity strength of regular graphs*, Electron. J. Combin. 15 (1) (2008) #R82.
- [12] J. Przybyło, *Linear bound on the irregularity strength and the total vertex irregularity strength of graphs*, SIAM J. Discrete Math. 23 (1) (2009) 511-516.
- [13] J. Przybyło, M. Woźniak, *Total weight choosability of graphs*, submitted.
- [14] T. Wong, D. Yang, X. Zhu, *Total weighting of graphs by max-min method*, submitted.
- [15] T. Wong, X. Zhu, *Total weight choosability of graphs*, submitted.