

Asymptotics of the Stirling numbers of the first kind revisited: A saddle point approach

Guy Louchard

► **To cite this version:**

Guy Louchard. Asymptotics of the Stirling numbers of the first kind revisited: A saddle point approach. Discrete Mathematics and Theoretical Computer Science, DMTCS, 2010, 12 (2), pp.167-184. hal-00990462

HAL Id: hal-00990462

<https://hal.inria.fr/hal-00990462>

Submitted on 13 May 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Asymptotics of the Stirling numbers of the first kind revisited: A saddle point approach

Guy Louchard

Université Libre de Bruxelles, Dép. d'Informatique, CP 212, B-1050 Bruxelles, Belgium. louchard@ulb.ac.be

received April 11, 2009, revised Dec. 21, 2009, accepted Feb. 18, 2010.

Using the saddle point method, we obtain from the generating function of the Stirling numbers of the first kind $\left[\begin{smallmatrix} n \\ j \end{smallmatrix} \right]$ and Cauchy's integral formula, asymptotic results in central and non-central regions. In the central region, we revisit the celebrated Goncharov theorem with more precision. In the region $j = n - n^\alpha$, $\alpha > 1/2$, we analyze the dependence of $\left[\begin{smallmatrix} n \\ j \end{smallmatrix} \right]$ on α .

Keywords: Stirling numbers, saddle point method.

To Philippe

1 Introduction

Let $\left[\begin{smallmatrix} n \\ j \end{smallmatrix} \right]$ be the Stirling number of the first kind (unsigned version). Their generating function is given by

$$\phi_n(z) = \prod_0^{n-1} (z+i) = \frac{\Gamma(z+n)}{\Gamma(z)}, \quad \phi_n(1) = n!.$$

In the sequel all asymptotics are meant for $n \rightarrow \infty$.

An asymptotic expansion for $j = \mathcal{O}(1)$ is given in Wilf [14], which has been extended to the range $j = \mathcal{O}(\ln n)$ by Hwang [6]. The generalized Stirling numbers have been considered by Tsylova [13] and Chelluri et al. [2]. The q -Stirling numbers are studied in Kyriakoussis and Vamvakari [9].

In Sec.2, we revisit the asymptotic expansions in the central region and in Sec.3, we analyse the non-central region $j = n - n^\alpha$, $\alpha > 1/2$. We use Cauchy's integral formula and the saddle point method.

2 Central region

Consider the random variable J_n , with probability distribution

$$\mathbb{P}[J_n = j] = Z_n(j),$$

$$Z_n(j) := \frac{\binom{n}{j}}{n!}.$$

The mean and variance are given by

$$M := \mathbb{E}(J_n) = \sum_0^{n-1} \frac{1}{1+i} = H_n = \psi(n+1) + \gamma,$$

$$\sigma^2 := \mathbb{V}(J_n) = \sum_0^{n-1} \frac{i}{(1+i)^2} = \psi(1, n+1) + \psi(n+1) - \frac{\pi^2}{6} + \gamma,$$

where $\psi(x)$ is the digamma function, $\psi(k, x)$ is the k th polygamma function, and

$$M \sim \ln(n) + \gamma + \frac{1}{2n} + \mathcal{O}\left(\frac{1}{n^2}\right),$$

$$\sigma^2 \sim \ln(n) - \frac{\pi^2}{6} + \gamma + \frac{3}{2n} + \mathcal{O}\left(\frac{1}{n^2}\right).$$

It is convenient to set

$$A_n := \ln(n) - \frac{\pi^2}{6} + \gamma = \ln\left(ne^{\gamma - \pi^2/6}\right),$$

and to consider all our next asymptotics ($n \rightarrow \infty$) as functions of A_n . Of course, all asymptotics can be reformulated in terms of $\ln(n)$.

We have

$$M \sim A_n + \frac{\pi^2}{6} + \mathcal{O}\left(\frac{1}{n}\right),$$

$$\sigma^2 \sim A_n + \mathcal{O}\left(\frac{1}{n}\right).$$

A celebrated central limit theorem of Goncharov says that

$$J_n \sim \mathcal{N}(M, \sigma),$$

where \mathcal{N} is the Gaussian distribution, with a rate of convergence $\mathcal{O}(1/\sqrt{\ln(n)})$. This can also be deduced from the Quasi-Power theorem of Hwang [7],[8].

In this Section, we want to obtain a more precise local limit theorem for J_n in terms of $x := \frac{J_n - M}{\sigma}$ and A_n . Actually, we obtain the following theorem, where we use $B_n := \sqrt{A_n}$ to simplify the expressions.

Theorem 2.1

$$Z_n(j) \sim \frac{1}{\sqrt{2\pi}B_n} e^{-x^2/2} \cdot \left[1 + \frac{x^3/6 - x/2}{B_n} + \frac{3x^2/8 - x^4/6 - 1/12 + x^6/72}{B_n^2} + \frac{-\pi^2 x^3/18 + 37x^5/240 - 355x^3/144 + x/8 - x^7/48 + x^9/1296 + \pi^2 x/6 - \zeta(3)x + \zeta(3)x^3/3}{B_n^3} + \dots \right].$$

Proof: By Cauchy's theorem,

$$\begin{aligned} Z_n(j) &= \frac{1}{2\pi i} \int_{\Omega} \frac{\phi_n(z)}{z^{j+1}n!} dz \\ &= \frac{1}{2\pi i} \int_{\Omega} e^{S(z)} dz, \end{aligned}$$

where Ω is inside the analyticity domain of the integrand and encircles the origin and

$$S(z) = S_1(z) + S_2(z), \quad S_1(z) = \sum_{i=0}^{n-1} \ln(z+i) - \ln(n!), \quad S_2(z) = -(j+1)\ln(z).$$

We will use the Saddle point method (for a good introduction to this method, see Flajolet and Sedgewick [3], ch.VIII). Set

$$S^{(i)} := \frac{d^i S}{dz^i}.$$

These derivatives can be expressed in terms of $\psi(k, z+n)$ and $\psi(k, z)$.

First we must find the solution of

$$S^{(1)}(\tilde{z}) = 0 \tag{1}$$

with smallest module.

Set $\tilde{z} := z^* - \varepsilon$, where $z^* = \lim_{n \rightarrow \infty} \tilde{z}$. Here, it is easy to check that $z^* = 1$. Set $j = M + x\sigma$, x fixed and $B_n := \sqrt{A_n}$.

This leads, to first order (keeping only the ε term in (1)), to

$$\varepsilon := \frac{-x}{B_n} + \frac{x^2 - 1}{B_n^2} + \dots + \frac{1}{n} \left(\frac{3x}{4B_n^3} + \dots \right) + \mathcal{O}\left(\frac{1}{n^2 B_n^4}\right).$$

This shows that, asymptotically, ε is given by a series of powers of n^{-1} , where each coefficient is given by a series of powers of B_n^{-1} . To obtain more precision, we set again $j = M + x\sigma$, expand in powers of n^{-1} , and equate each coefficient to 0. . This leads to (here and in the following, we provide only a few terms but Maple knows more)

$$\varepsilon = \frac{-x}{B_n} - \frac{1}{B_n^2} + \frac{0}{B_n^3} + \dots + \frac{1}{n} \left(\frac{3x}{4B_n^3} + \frac{x^2 + 3/2}{B_n^4} + \dots \right) + \mathcal{O}\left(\frac{1}{n^2 B_n^4}\right).$$

We have, with $\tilde{z} := z^* - \varepsilon = 1 - \varepsilon$,

$$Z_n(j) = \frac{1}{2\pi\mathbf{i}} \int_{\Omega} \exp \left[S(\tilde{z}) + S^{(2)}(\tilde{z})(z - \tilde{z})^2/2! + \sum_{l=3}^{\infty} S^{(l)}(\tilde{z})(z - \tilde{z})^l/l! \right] dz.$$

Note that the linear term vanishes. Set $z = \tilde{z} + \mathbf{i}\tau$. This gives

$$Z_n(j) = \frac{1}{2\pi} \exp[S(\tilde{z})] \int_{-\infty}^{\infty} \exp \left[S^{(2)}(\tilde{z})(\mathbf{i}\tau)^2/2! + \sum_{l=3}^{\infty} S^{(l)}(\tilde{z})(\mathbf{i}\tau)^l/l! \right] d\tau. \quad (2)$$

The justification of the integration procedure is given in the Appendix. Let us first analyze $S(\tilde{z})$. We obtain

$$\begin{aligned} S(\tilde{z}) &= -x^2/2 + \frac{x^3/6 - x}{B_n} + \frac{-x^4/12 + x^2/2 - 1/2}{B_n^2} \\ &+ \frac{-x^3/3 + x^5/20 + x/2 - \pi^2 x^3/18 + \zeta(3)x^3/3}{B_n^3} + \dots + \mathcal{O}\left(\frac{1}{nB_n^3}\right). \end{aligned}$$

Also,

$$\begin{aligned} S^{(2)}(\tilde{z}) &= B_n^2 - B_n x - 1 + x^2 + \dots, \\ S^{(3)}(\tilde{z}) &= -2B_n^2 + 4B_n x - \pi^2/3 + 2\zeta(3) - 6x^2 + 4 + \dots, \\ S^{(4)}(\tilde{z}) &= 6B_n^2 - 18B_n x + 36x^2 - 18 + \pi^2 - \pi^4/15 + \dots, \\ S^{(l)}(\tilde{z}) &= \mathcal{O}(B_n^2), \quad l \geq 5. \end{aligned}$$

We need these many terms in the following. Note that, with $z = \tilde{z}e^{i\theta}$, this leads to

$$S^{(2)}(\tilde{z}) \frac{(z - \tilde{z})^2}{2} \sim -\frac{1}{2} \ln(n)\theta^2. \quad (3)$$

We can now compute (2), for instance by using the classical trick of setting

$$S^{(2)}(\tilde{z})(i\tau)^2/2! + \sum_{l=3}^{\infty} S^{(l)}(\tilde{z})(i\tau)^l/l! = -u^2/2.$$

Computing τ as a truncated series in u , this gives, by inversion,

$$\tau = \left[u(1 + x/(2B_n) + \dots) + u^2(\mathbf{i}/(3B_n) + \dots) + u^3(-1/(36B_n^2) + \dots) \right] / B_n + \dots$$

Setting $d\tau = \frac{d\tau}{du} du$, expanding w.r.t. B_n and integrating on $[u = -\infty.. \infty]$, this gives

$$\frac{1}{\sqrt{2\pi}B_n} \left[1 + \frac{x}{2B_n} + \frac{5/12 - x^2/8}{B_n^2} + \frac{x(8\pi^2 - 10 - 93x^2 - 48\zeta(3))}{48B_n^3} + \dots \right].$$




Fig. 1: Comparison between $Z_n(j)$ and $\frac{1}{\sqrt{2\pi\sigma}} \exp\left[-\left(\frac{j-M}{\sigma}\right)^2/2\right]$, $n = 3000$

Finally (2) leads to

$$\begin{aligned}
 Z_n(j) &\sim \frac{1}{\sqrt{2\pi}B_n} e^{-x^2/2} \\
 &\cdot \exp\left[\frac{x^3/6 - x}{B_n} + \frac{-x^4/12 + x^2/2 - 1/2}{B_n^2} + \frac{-x^3/3 + x^5/20 + x/2 - \pi^2 x^3/18 + \zeta(3)x^3/3}{B_n^3} + \dots\right] \\
 &\cdot \left[1 + \frac{x}{2B_n} + \frac{5/12 - x^2/8}{B_n^2} + \frac{x(8\pi^2 - 10 - 93x^2 - 48\zeta(3))}{48B_n^3} + \dots\right],
 \end{aligned}$$

or

$$Z_n(j) \sim R_1,$$

$$\begin{aligned}
 R_1 &= \frac{1}{\sqrt{2\pi}B_n} e^{-x^2/2} \\
 &\cdot \left[1 + \frac{x^3/6 - x/2}{B_n} + \frac{3x^2/8 - x^4/6 - 1/12 + x^6/72}{B_n^2} \right. \\
 &\quad \left. + \frac{-\pi^2 x^3/18 + 37x^5/240 - 355x^3/144 + x/8 - x^7/48 + x^9/1296 + \pi^2 x/6 - \zeta(3)x + \zeta(3)x^3/3}{B_n^3} + \dots\right].
 \end{aligned}$$

□

For $n = 3000$, a comparison between $Z_n(j)$ and $\frac{1}{\sqrt{2\pi\sigma}} \exp\left[-\left(\frac{j-M}{\sigma}\right)^2/2\right]$ is given in Figure 1.




Fig. 2: $Z_n(j) / \left[\frac{1}{\sqrt{2\pi\sigma}} \exp \left[- \left(\frac{j-M}{\sigma} \right)^2 / 2 \right] \right]$, color=red, $Z_n(j)/R_1$, color=blue, $n = 3000$

Of course, only few values of j are significant and also the quality of the Gaussian is low, all asymptotic expressions depend actually on powers of A_n^{-1} , but A_n is not large.

A comparison of $Z_n(j) / \left[\frac{1}{\sqrt{2\pi\sigma}} \exp \left[- \left(\frac{j-M}{\sigma} \right)^2 / 2 \right] \right]$ with $Z_n(j)/R_1$, with 2 terms in R_1 , is given in Figure 2.

The precision of R_1 is of order 10^{-2} . Using 3 terms in R_1 leads to a less good result: A_n is not large enough to take advantage of the $A_n^{-3/2}$ term: $A_n = 6.94$ here, we deal with asymptotic series, not necessarily convergent ones. More terms can be computed in R_1 (which is almost automatic with Maple).

3 Large deviation, $j = n - n^\alpha$, $1 > \alpha > 1/2$

The case $j = \mathcal{O}(n)$ is analyzed in Timashev [12]. But he obtains a series of powers of n^{-1} , determined by a power series of a certain function that depends on the solution of a given non-linear differential equation of the first order. The coefficients obey some linear recurrence relations in the complex plane. The case $j = n - c$, c constant, is considered in Grünberg [5]. As previous work for the case $j = n - n^\alpha$, let us mention Bender [1], Temme [11], Moser and Wyman [10] (see also the comments by Odlyzko in [4], p.1182). They all use, explicitly or not, the Saddle point method. For $\alpha < 1/2$, Moser and Wyman (6.9) give an explicit asymptotic expression. For $\alpha > 1/2$, they first compute in (4.52) the numerical solution zn of $S'(zn) = 0$ and give in (4.51) an asymptotic expression. This is rather precise: for $n = 50$, this gives a precision of order 10^{-4} . [1] and [11] also compute numerically zn .

However, all these results do not shed light on the dependence of $[z^j]\phi(z)$ on n^α . This is what we want

to explicit in this Section. It appears that the range $\alpha > 1/2$ is more delicate than the other range.

Recall that

$$\phi_n(z) = \prod_0^{n-1} (z+i) = \frac{\Gamma(z+n)}{\Gamma(z)}.$$

We have

$$G_n(z) := \frac{\Gamma(z+n)}{\Gamma(z)z^{j+1}} = \exp[S(z)],$$

with

$$S(z) = S_1(z) + S_2(z), S_1(z) = \sum_0^{n-1} \ln(z+i), S_2(z) = -(j+1) \ln(z).$$

We first compute \tilde{z} such that

$$S'(\tilde{z}) = 0. \quad (4)$$

We have

$$S'(z) = \psi(z+n) - \psi(z) - \frac{j+1}{z}.$$

Similarly (we need these expressions later on)

$$S^{(2)}(z) = \psi(1, z+n) - \psi(1, z) + \frac{j+1}{z^2},$$

$$S^{(k)}(z) = \psi(k-1, z+n) - \psi(k-1, z) + (-1)^k (k-1)! \frac{j+1}{z^k}.$$

Some experiments with some values for α ($\alpha = 5/8$ is a good choice) show that \tilde{z} must be a combination of $x = n^\alpha$ and $y = n^{1-\alpha}$ and $x \gg y \gg 1$. Note that both x and y are large. We will derive series of powers of x^{-1} , where each coefficient is a series of powers of y^{-1} . We obtain the following theorem

Theorem 3.1

$$[z^j]\phi_n(z) \sim \frac{1}{\sqrt{2\pi}} \frac{y^2 \sqrt{x}}{2} \exp \left[x \left[1 - \ln(2) + 2 \ln(y) + \ln(x) - \frac{2}{3y} - \frac{2}{9y^2} - \frac{44}{405y^3} - \frac{26}{405y^4} + \frac{40}{27y^5} \right. \right. \\ \left. \left. + \frac{179968}{18225y^6} + \frac{4727552}{127575y^7} + \frac{3436796}{32805y^8} + \frac{5492621728}{22143375y^9} + \dots \right] + \ln(2) - 2 \ln(y) - \ln(x) \right] \cdot \\ \cdot \left[1 - \frac{3}{3y} - \frac{1}{18y^2} - \frac{1}{30y^3} + \frac{17207}{3240y^4} + \dots + \frac{1}{x} \left(-\frac{1}{12} + \frac{1}{36y} - \frac{35}{216y^2} + \frac{15029}{3240y^3} + \dots \right) \right. \\ \left. + \frac{1}{x^2} \left(\frac{1}{288} - \frac{1}{864y} + \frac{3527}{5184y^2} + \dots \right) + \mathcal{O} \left(\frac{1}{x^3} \right) \right].$$

Proof: Let us summarize the different steps of the proof. First we compute \tilde{z} and $S(\tilde{z})$ as $S(\tilde{z}) = T_1 T_2$, where T_1 is the dominant term and T_2 is a series of powers of x^{-1} , where each coefficient is a series of powers of y^{-1} . We expand $T_3 := e^{T_2}$. Next the integration procedure leads to $\frac{y^2 \sqrt{x}}{2} T_4$, where T_4 is again a series of powers of x^{-1} , where each coefficient is a series of powers of y^{-1} . We set $T_5 := \frac{1}{\sqrt{2\pi}} \frac{y^2 \sqrt{x}}{2} e^{T_1}$. Finally, we obtain

$$[z^j]\phi_n(z) \sim T_5 T_3 T_4. \quad (5)$$

The first terms in the asymptotics of \tilde{z} are easy to compute: set $\tilde{z} = n\beta$. Equation (4) leads to

$$\psi(n(1 + \beta)) - \psi(n\beta) = \frac{1}{\beta} - \frac{1}{y\beta} + \frac{1}{n\beta}.$$

But $\psi(n) \sim \ln(n)$. So we have

$$\ln\left(1 + \frac{1}{\beta}\right) \sim \frac{1}{\beta} - \frac{1}{y\beta} + \frac{1}{n\beta},$$

or

$$\frac{1}{\beta} - \frac{1}{2\beta^2} \sim \frac{1}{\beta} - \frac{1}{y\beta},$$

or $\beta \sim \frac{y}{2}$.

More generally, we have

$$\beta = \frac{y}{2} \left[1 + \frac{a_1}{y} + \frac{a_2}{y^2} + \frac{a_3}{y^3} + \dots + \frac{1}{x} \left(1 + \frac{b_1}{y} + \frac{b_2}{y^2} + \dots \right) + \frac{1}{x^2} \left(1 + \frac{c_1}{y} + \frac{c_2}{y^2} + \dots \right) + \mathcal{O}\left(\frac{1}{x^3}\right) \right].$$

Note that $\frac{1}{y^3}$ can be of the same order than $\frac{1}{x}$, see below.

By bootstrapping, we obtain (we give the first terms)

$$\begin{aligned} \tilde{z} &= \frac{ny}{2} \left[1 - \frac{4}{3y} + \frac{2}{9y^2} + \frac{8}{135y^3} + \frac{8}{405y^4} + \frac{16}{1701y^5} + \frac{232}{45525y^6} + \frac{64}{18225y^7} + \dots \right. \\ &\quad + \frac{1}{x} \left[1 - \frac{1}{y} + \frac{4}{9y^2} - \frac{16}{135y^3} + \dots \right] \\ &\quad + \frac{1}{x^2} \left[1 - \frac{1}{y} + \frac{0}{y^2} + \dots \right] \\ &\quad + \frac{1}{x^3} [1 + \dots] \\ &\quad \left. + \mathcal{O}\left(\frac{1}{x^4}\right) \right]. \end{aligned} \tag{6}$$

Note that the choice of dominant terms in the bracket of (6) depends on α . For instance, for $\alpha = 3/4$, the dominant terms (in decreasing order) are

$$1, \frac{1}{y}, \frac{1}{y^2}, \left\{ \frac{1}{x}, \frac{1}{y^3} \right\}, \left\{ \frac{1}{xy}, \frac{1}{y^4} \right\}, \left\{ \frac{1}{xy^2}, \frac{1}{y^5} \right\}, \left\{ \frac{1}{x^2}, \frac{1}{xy^3}, \frac{1}{y^6} \right\}, \dots$$

Now we must compute $S(\tilde{z})$ and its asymptotics. First we compute $\ln(\tilde{z} + i)$, take the asymptotics wrt x , sum on i , and again take the asymptotics wrt x (recall that $n = xy$). this leads to

$$\begin{aligned} S_1(\tilde{z}) &= x \left[(-\ln(2) + 2\ln(y) + \ln(x))y - \frac{1}{3} + \frac{4}{405y^2} + \frac{2}{405y^3} + \dots \right] + y - \frac{2}{3} - \frac{2}{3y} - \frac{49}{135y^2} + \dots \\ &\quad + \frac{1}{x} \left(\frac{y}{2} + \frac{1}{6y} + \dots \right) + \frac{1}{x^2} \left(\frac{y}{3} + \dots \right) + \mathcal{O}\left(\frac{y}{x^3}\right). \end{aligned}$$




Fig. 3: $zn/\bar{z}, n = 500$, as function of j , full range




Fig. 4: $zn/\bar{z}, n = 500$, as function of j , restricted range

Here we provide only a few terms but Maple knows more. Next

$$\begin{aligned}
S_2(\tilde{z}) &= x \left[(\ln(2) - 2\ln(y) - \ln(x))y + \frac{4}{3} - \ln(2) + 2\ln(y) + \ln(x) - \frac{2}{3y} - \frac{94}{405y^2} + \dots \right] \\
&\quad - y + \frac{2}{3} + \ln(2) - 2\ln(y) - \ln(x) + \frac{1}{y} + \frac{94}{135y^2} + \dots \\
&\quad + \frac{1}{x} \left(\frac{y}{2} + \frac{1}{6y} + \dots \right) \\
&\quad + \frac{1}{x^2} \left(\frac{y}{3} + \dots \right) \\
&\quad + \mathcal{O}\left(\frac{y}{x^3}\right).
\end{aligned}$$

So, finally

$$\begin{aligned}
S(\tilde{z}) &= S_1(\tilde{z}) + S_2(\tilde{z}) \sim x \left[1 - \ln(2) + 2\ln(y) + \ln(x) - \frac{2}{3y} + \dots \right] \\
&\quad + \ln(2) - 2\ln(y) - \ln(x) + \frac{1}{3y} + \frac{1}{3y^2} + \dots \\
&\quad + \frac{1}{x} \left(-\frac{1}{2} + \frac{1}{3y} - \frac{1}{2y^2} + \dots \right) \\
&\quad + \frac{1}{x^2} \left(-\frac{1}{6} + \frac{19}{18y^2} \dots \right) \\
&\quad + \mathcal{O}\left(\frac{1}{x^3}\right).
\end{aligned}$$

Now we split $S(\tilde{z})$ into two parts:

$$\begin{aligned}
T_1 &= x \left[1 - \ln(2) + 2\ln(y) + \ln(x) - \frac{2}{3y} + \dots \right] + \ln(2) - 2\ln(y) - \ln(x), \\
T_2 &= \frac{1}{3y} + \frac{1}{3y^2} + \dots \\
&\quad + \frac{1}{x} \left(-\frac{1}{2} + \frac{1}{3y} - \frac{1}{2y^2} + \dots \right) \\
&\quad + \frac{1}{x^2} \left(-\frac{1}{6} - \frac{17}{18y^2} \dots \right) \\
&\quad + \mathcal{O}\left(\frac{1}{x^3}\right).
\end{aligned}$$

Note that the dominant term of T_1 is given by

$$T_1 \sim (2 - \alpha)n^\alpha \ln(n). \quad (7)$$

We obtain

$$\exp(S(\tilde{z})) = e^{T_1} e^{T_2} = e^{T_1} T_3,$$

with

$$\begin{aligned} T_3 = e^{T_2} &= 1 + \frac{1}{3y} + \frac{7}{18y^2} + \frac{89}{270y^3} + \frac{18263}{3240y^4} + \frac{98009}{3240y^5} + \frac{9517337}{97200y^6} + \frac{491504273}{2041200y^7} + \dots \\ &+ \frac{1}{x} \left(-\frac{1}{2} + \frac{1}{6y} - \frac{7}{12y^2} + \frac{2311}{540y^3} + \frac{112469}{6480y^4} + \frac{5137}{144y^5} + \dots \right) \\ &+ \frac{1}{x^2} \left(-\frac{1}{24} - \frac{13}{72y} - \frac{557}{932y^2} + \dots \right) \\ &+ \mathcal{O}\left(\frac{1}{x^3}\right). \end{aligned}$$

Here we have given all terms compatible with the expansion (6). Also, with more precision,

$$\begin{aligned} T_1 = x &\left[1 - \ln(2) + 2 \ln(y) + \ln(x) - \frac{2}{3y} - \frac{2}{9y^2} - \frac{44}{405y^3} - \frac{26}{405y^4} + \frac{40}{27y^5} \right. \\ &\left. + \frac{179968}{18225y^6} + \frac{4727552}{127575y^7} + \frac{3436796}{32805y^8} + \frac{5492621728}{22143375y^9} + \dots \right] \\ &+ \ln(2) - 2 \ln(y) - \ln(x). \end{aligned}$$

Now we must consider $S^{(k)}(\tilde{z})$. By direct expansion, this gives the following expressions (again we provide only the first few terms). We must use up to six derivatives to get a sufficient precision (of order x^{-2}) in the Saddle integrals.

$$\begin{aligned} S^{(2)}(\tilde{z}) &= \frac{1}{x} \left[\frac{4}{y^4} + \frac{16}{3y^5} + \dots \right] \\ &+ \frac{1}{x^2} \left[-\frac{12}{y^4} - \frac{40}{3y^5} + \dots \right] \\ &+ \frac{1}{x^3} \left[\frac{12}{y^4} + \frac{8}{y^5} + \dots \right] \\ &+ \frac{1}{x^4} \left[\frac{-4}{y^4} + \dots \right] \\ &+ \mathcal{O}\left(\frac{1}{x^5 y^4}\right), \end{aligned} \tag{8}$$

Note that, with $z = \tilde{z}e^{i\theta}$, this leads to

$$S^{(2)}(\tilde{z}) \frac{(z - \tilde{z})^2}{2} \sim -\frac{1}{2} n^\alpha \theta^2. \tag{9}$$




Fig. 5: $G_n(zn)/G_n(\tilde{z})$, $n = 500$, as function of j




Fig. 6: The quotient of the expression (8) and $S^{(2)}(\tilde{z})$ as function of j , $n = 500$

$$\begin{aligned}
S^{(3)}(\tilde{z}) &= \frac{1}{x^2} \left[-\frac{32}{y^6} + \dots \right] \\
&+ \frac{1}{x^3} \left[\frac{128}{y^6} + \dots \right] \\
&+ \frac{1}{x^4} \left[-\frac{192}{y^6} + \dots \right] \\
&+ \frac{1}{x^5} \left[\frac{128}{y^6} + \dots \right] \\
&+ \mathcal{O}\left(\frac{1}{x^6 y^6}\right),
\end{aligned}$$

$$\begin{aligned}
S^{(4)}(\tilde{z}) &= \frac{1}{x^3} \left[\frac{288}{y^8} + \dots \right] \\
&+ \frac{1}{x^4} \left[-\frac{1440}{y^8} + \dots \right] \\
&+ \frac{1}{x^5} \left[\frac{2880}{y^8} + \dots \right] \\
&+ \frac{1}{x^6} \left[-\frac{2880}{y^8} + \dots \right] \\
&+ \mathcal{O}\left(\frac{1}{x^7 y^8}\right),
\end{aligned}$$

$$\begin{aligned}
S^{(5)}(\tilde{z}) &= \frac{1}{x^4} \left[-\frac{3072}{y^{10}} + \dots \right] \\
&+ \frac{1}{x^5} \left[\frac{18432}{y^{10}} + \dots \right] \\
&+ \mathcal{O}\left(\frac{1}{x^6 y^{10}}\right),
\end{aligned}$$

$$\begin{aligned}
S^{(6)}(\tilde{z}) &= \frac{1}{x^5} \left[\frac{38400}{y^{12}} + \dots \right] \\
&+ \frac{1}{x^6} \left[\frac{268800}{y^{12}} + \dots \right] \\
&+ \mathcal{O}\left(\frac{1}{x^7 y^{12}}\right).
\end{aligned}$$

We proceed now as in Section 2. Again, the justification of the integration procedure is given in the Appendix. This leads to

$$\tau = \frac{y^2 \sqrt{x}}{2} \left[u a_1 + \frac{u^2 a_2}{x^{1/2}} + \frac{u^3 a_3}{x} + \frac{u^4 a_4}{x^{3/2}} + \frac{u^5 a_5}{x^2} + \mathcal{O}\left(\frac{u^6}{x^{5/2}}\right) \right].$$




Fig. 7: The quotient of the expression (8) and $S^{(2)}(\tilde{z})$, as function of j , $n = 500$. Restricted range, $\alpha \leq .84$

We give only a_1 :

$$a_1 = 1 - \frac{2}{3y} - \frac{2}{9y^2} + \dots + \frac{1}{x} \left(\frac{3}{2} - \frac{4}{3y} + \dots \right) + \frac{1}{x^2} \left(\frac{15}{8} - \frac{7}{4y} + \dots \right) + \mathcal{O} \left(\frac{1}{x^3} \right).$$

This leads to

$$\frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-u^2/2} \tau'(u) du = \frac{y^2 \sqrt{x}}{2} T_4,$$

with

$$T_4 = 1 - \frac{2}{3y} - \frac{2}{9y^2} + \dots + \frac{1}{x} \left(\frac{5}{12} - \frac{11}{18y} + \dots \right) + \frac{1}{x^2} \left(\frac{73}{288} - \frac{133}{432y} + \dots \right) + \frac{1}{x^3} \left(\frac{721}{576} + \dots \right) + \mathcal{O} \left(\frac{1}{x^4} \right).$$

Set

$$T_5 := \frac{1}{\sqrt{2\pi}} \frac{y^2 \sqrt{x}}{2} e^{T_1}.$$

This leads to

$$[z^j] \phi_n(z) \sim T_5 T_3 T_4. \quad (10)$$

We can of course combine T_3 and T_4 :

$$T_6 := T_3 T_4 = 1 - \frac{3}{3y} - \frac{1}{18y^2} - \frac{1}{30y^3} + \frac{17207}{3240y^4} + \dots + \frac{1}{x} \left(-\frac{1}{12} + \frac{1}{36y} - \frac{35}{216y^2} + \frac{15029}{3240y^3} + \dots \right) + \frac{1}{x^2} \left(\frac{1}{288} - \frac{1}{864y} + \frac{3527}{5184y^2} + \dots \right) + \mathcal{O} \left(\frac{1}{x^3} \right).$$

□




Fig. 8: The quotient $[z^j]\phi_n(z)/T_8$, two terms in T_4 , as function of j , $n = 500$




Fig. 9: The quotient $[z^j]\phi_n(z)/T_8$, three terms in T_4 , as function of j , $n = 500$

Let us consider the precision of our asymptotics.

The quality of asymptotic (6) is given in Figure 3 and 4, for $n = 500$, and $x \in [\sqrt{n}, n^{0.9}]$ (first range) so that $y \in [n^{0.1}, \sqrt{n}]$. For some values of $j = n - x$, we show \tilde{z}/zn , where, as mentioned, zn is the numerical solution of $S'(zn) = 0$. In the full range $j \in [n - n^{0.9}, n - \sqrt{n}]$, the precision is of order 10^{-5} , in a restricted range, the precision is of order 10^{-6} .

Also a comparison of $G_n(\tilde{z})$ and $G_n(zn)$ is given in Figure 5, showing again a precision of order 10^{-6} .

To check the quality of asymptotic (6), we give in Figure 6 the comparison between the expression (8) and $S^{(2)}(\tilde{z})$. The precision is of order 10^{-2} .

In a restricted range, given in Figure 7, the precision is of order 10^{-4} . $\alpha \leq 0.84$ in this range.

We have made several experiments with (10), with n up to 500. The result is unsatisfactory, only values of x of order \sqrt{n} give reasonable results. Also using e^{T_2} instead of T_3 does not improve the precision. Actually, only very large values of n lead to good precision. So we turn to another formulation: instead of using $e^{T_1 T_3}$ for $e^{S(\tilde{z})}$, we plug directly \tilde{z} into $G_n(z)$, ie we set

$$T_7 = G_n(\tilde{z}),$$

leading to

$$[z^j]\phi_n(z) \sim \frac{1}{\sqrt{2\pi}} \frac{y^2 \sqrt{x}}{2} T_7 T_4 =: T_8 \text{ say.}$$

For $n = 500$, using two and three terms in T_4 , we give in Figures 8 and 9, the quotient $[z^j]\phi_n(z)/T_8$. The precision is of order 10^{-5} .

4 Conclusion

Using an almost mechanized program in Maple, we have obtained some asymptotic expressions for Stirling numbers in central and non-central regions. We intend to use these techniques in other non-central ranges.

Acknowledgement

The pertinent comments of the referee led to improvements in the presentation.

A Appendix. Justification of the integration procedure

A.1 The central region

We proceed as in Flajolet and Sedgewick [3], ch.VIII. We can choose here $\tilde{z} = 1$. This leads, with $z = e^{i\theta}$, to

$$S(z) \sim S_0(z) + \mathcal{O}\left(\sqrt{\ln(n)\theta}\right) + \text{constant term,}$$

with

$$\begin{aligned} S_0(z) &= \sum_{k=0}^{n-1} \ln[e^{i\theta} + k] - H_n i\theta \\ &\sim \sum_{k=0}^{n-1} \frac{1}{1+k} [e^{i\theta} - 1] - \frac{1}{2} \sum_{k=0}^{n-1} \left[\frac{1}{1+k} [e^{i\theta} - 1] \right]^2 - H_n i\theta + \mathcal{O}(\theta^3) \\ &\sim H_n [e^{i\theta} - 1 - i\theta] + \mathcal{O}(\theta^2). \end{aligned}$$

Set

$$h(\theta) := e^{i\theta} - 1 - i\theta.$$

We have

$$h(\theta) \sim -\frac{\theta^2}{2},$$

which conforms to (3).

The function $h(\theta)$ is the same as in [3], Ex.VIII.3, which proves the validity of our integration procedure: we use here $H_n \sim \ln(n)$ instead of n . The complete asymptotic expansion is justified as in [3], Ex.VIII.4.

A.2 The non-central region

We choose here $\tilde{z} = \frac{ny}{2} = \frac{n^{2-\alpha}}{2} := \delta$, say. We have

$$\begin{aligned} \frac{1}{2} &< \alpha < 1, \\ n^\alpha &= \frac{n^2}{2\delta}, \\ n^2 &\gg \delta \gg n \gg n^\alpha. \end{aligned}$$

Set $z = \delta e^{i\theta}$, this leads, with Euler-Maclaurin formula, with the first correction (the other corrections are negligible), to

$$\begin{aligned} S(z) &\sim \sum_{k=0}^{n-1} \ln [\delta e^{i\theta} + k] - (n - n^\alpha) i\theta - (n - n^\alpha) \ln(\delta) \\ &\sim \ln [n + \delta e^{i\theta}] [n + \delta e^{i\theta}] - n - \delta e^{i\theta} \ln [\delta e^{i\theta}] - \left[n - \frac{n^2}{2\delta} \right] (i\theta + \ln(\delta)) - \frac{1}{2} \ln [n + \delta e^{i\theta}] + \frac{1}{2} \ln [\delta e^{i\theta}]. \end{aligned}$$

Set now $n = \rho\delta$, $\rho = 2n^{\alpha-1} \ll 1$ and expand wrt ρ . This gives

$$\begin{aligned} S(z) &\sim \rho \left[-\frac{1}{2} e^{-i\theta} \right] \\ &\quad + \rho^2 \left[\delta \frac{1 + i\theta e^{i\theta}}{2e^{i\theta}} + \frac{1}{4} e^{-2i\theta} + \frac{1}{2} \delta \ln(\delta) \right] \\ &\quad + \rho^3 \left[-\frac{\delta}{6} e^{-2i\theta} - \frac{1}{6} e^{-3i\theta} \right] \\ &\quad + \mathcal{O}(\delta\rho^4). \end{aligned}$$

Note that the dominant constant contribution is given by $\frac{1}{2}\rho^2\delta\ln(\delta) = (2 - \alpha)n^\alpha\ln(n)$, which conforms to (7). The first term gives a variable part $\mathcal{O}(\rho)$. The second term gives a variable part $2n^\alpha h(\theta) + \mathcal{O}(\rho^2)$, with

$$h(\theta) := \frac{1 + i\theta e^{i\theta}}{2e^{i\theta}}.$$

The third term give $\mathcal{O}(n^{2\alpha-1}) \ll n^\alpha$. Note that $2n^\alpha h(\theta) \sim -\frac{1}{2}n^\alpha\theta^2$, which conforms to (9). The function $|e^{h(\theta)}| = e^{\cos(\theta)/2}$ is unimodal with peak at 0 and $h(0) = 1/2$. Let us introduce a splitting value θ_0 such that $n^\alpha\theta_0^2 \rightarrow \infty, n^\alpha\theta_0^3 \rightarrow 0, n \rightarrow \infty$. For instance, we choose $\theta_0 = n^\beta, \beta = -\frac{5\alpha}{12}$. By unimodality property of the cosine, the tail integral

$$K_n^{(1)} := \int_{\theta_0}^{2\pi-\theta_0} e^{2n^\alpha(h(\theta)-1/2)} d\theta$$

is such that

$$|K_n^{(1)}| = \mathcal{O}\left(e^{n^\alpha[\cos(\theta_0)-1]}\right) = \mathcal{O}\left(e^{-Cn^{\alpha/6}}\right)$$

for some $C > 0$. The tail integral is exponentially small.

As $h(\theta) \sim -\frac{\theta^2}{4}$, the central approximation and the tail completion are immediate.

References

- [1] E.A. Bender. Central and local limit theorems applied to asymptotics enumeration. *Journal of Combinatorial Theory*, Series A, 15:91–111, 1973.
- [2] R. Chelluri, L. B. Richmond, and N. M. Temme. Asymptotic estimates for generalized Stirling numbers. Technical report, CWI, MAS-R9923, 1999.
- [3] P. Flajolet and R. Sedgewick. *Analytic combinatorics*. Cambridge University press, 2009.
- [4] R. L. Graham, M. Grötschel, and L. Lovász. *Handbook of Combinatorics*, vol. 2. Elsevier, 1995.
- [5] D. B. Grünberg. On asymptotics, Stirling numbers, Gamma function and polylogs. *Results in Mathematics*, 49(1):89–125, 2006.
- [6] H. K. Hwang. Asymptotic expansions for the Stirling numbers of the first kind. *Journal of Combinatorial Theory*, Series A, 71(2):343–351, 1995.
- [7] H.K. Hwang. Théorèmes limites pour les structures aléatoires et les fonctions arithmétiques. 1994. Thèse, Ecole Polytechnique de Palaiseau.
- [8] H.K. Hwang. On convergence rates in the central limit theorems for combinatorial structures. *European Journal of Combinatorics*, 19:329–343, 1998.
- [9] A. Kyriakoussis and M. G. Vamvakari. On asymptotic for the signless noncentral q-Stirling numbers of the first kind. *Studies in Applied Mathematics*, 117:191–213, 2006.
- [10] L. Moser and M. Wyman. Asymptotic development of the Stirling numbers of the first kind. *Journal of the London Mathematical Society*, 33:133–146, 1958.
- [11] N.M. Temme. Asymptotic estimates of Stirling numbers. *Studies in Applied Mathematics*, 89:233–243, 1993.
- [12] A. N. Timashev. On asymptotic expansions of Stirling numbers of the first and second kind. *Discrete Mathematics and Applications*, 8(5):533–544, 1998.
- [13] E. G. Tsylova. Probabilistic methods for obtaining asymptotic formulas for generalized Stirling numbers. *Discrete Mathematics and Applications*, 75(2):1607–1614, 1995.
- [14] H. S. Wilf. The asymptotic behavior of the Stirling numbers of the first kind. *Journal of Combinatorial Theory*, Series A, 64:344–349, 1993.